



# Oxidation of P700 Ensures Robust Photosynthesis

Ginga Shimakawa<sup>1†</sup> and Chikahiro Miyake<sup>1,2\*</sup>

<sup>1</sup> Department of Biological and Environmental Science, Faculty of Agriculture, Graduate School of Agricultural Science, Kobe University, Kobe, Japan, <sup>2</sup> Core Research for Environmental Science and Technology, Japan Science and Technology Agency, Tokyo, Japan

## OPEN ACCESS

### Edited by:

Marian Brestic,  
Slovak University of Agriculture,  
Slovakia

### Reviewed by:

Yuji Suzuki,  
Iwate University, Japan  
Ko Noguchi,  
Tokyo University of Pharmacy and Life  
Sciences, Japan  
Riichi Oguchi,  
Tohoku University, Japan

### \*Correspondence:

Chikahiro Miyake  
cmiyake@hawk.kobe-u.ac.jp

### †Present address:

Ginga Shimakawa,  
Institute for Integrative Biology of the  
Cell, CEA, CNRS, Université  
Paris-Sud, Université Paris-Saclay,  
Paris, France

### Specialty section:

This article was submitted to  
Plant Abiotic Stress,  
a section of the journal  
Frontiers in Plant Science

**Received:** 21 August 2018

**Accepted:** 18 October 2018

**Published:** 06 November 2018

### Citation:

Shimakawa G and Miyake C  
(2018) Oxidation of P700 Ensures  
Robust Photosynthesis.  
*Front. Plant Sci.* 9:1617.  
doi: 10.3389/fpls.2018.01617

In the light, photosynthetic cells can potentially suffer from oxidative damage derived from reactive oxygen species. Nevertheless, a variety of oxygenic photoautotrophs, including cyanobacteria, algae, and plants, manage their photosynthetic systems successfully. In the present article, we review previous research on how these photoautotrophs safely utilize light energy for photosynthesis without photo-oxidative damage to photosystem I (PSI). The reaction center chlorophyll of PSI, P700, is kept in an oxidized state in response to excess light, under high light and low CO<sub>2</sub> conditions, to tune the light utilization and dissipate the excess photo-excitation energy in PSI. Oxidation of P700 is co-operatively regulated by a number of molecular mechanisms on both the electron donor and acceptor sides of PSI. The strategies to keep P700 oxidized are diverse among a variety of photoautotrophs, which are evolutionarily optimized for their ecological niche.

**Keywords:** P700 oxidation, photosystem I, photoinhibition, reactive oxygen species, photosynthesis

## INTRODUCTION

Nothing ventured, nothing gained. In oxygenic photosynthesis, CO<sub>2</sub> is converted into sugar using H<sub>2</sub>O and light energy. Additionally, O<sub>2</sub> is released from oxygenic photosynthesis as a by-product. Thus, oxygenic photoautotrophs (i.e., cyanobacteria, algae, and plants) support life on Earth with only sun light energy. However, this process is not risk-free, and every photoautotroph is constantly exposed to potential photo-oxidative damage.

Oxygenic photosynthesis is achieved through the assimilation of CO<sub>2</sub> from the atmosphere in the Calvin-Benson cycle (dark reaction) and the utilization of light for the production of nicotinamide adenine dinucleotide phosphate (NADPH) and adenosine triphosphate (ATP) to meet the demand for CO<sub>2</sub> assimilation in the photosynthetic electron transport system (light reaction). Light energy from the sun is absorbed by pigments, such as chlorophyll (Chl), in light-harvesting complexes (LHC) around two photosystems (PSII and PSI) on the thylakoid membrane, and excites the reaction center Chls (P680 and P700 in PSII and PSI, respectively) to drive charge separation and photosynthetic linear electron flow from PSII to PSI via plastoquinone (PQ) pool, cytochrome (Cyt) *b<sub>6</sub>/f* complex, and plastocyanin (PC) or Cyt *c<sub>6</sub>* (Barber and Andersson, 1994). On the electron donor side of PSII, photo-oxidized P680 (P680<sup>+</sup>) oxidizes H<sub>2</sub>O with O<sub>2</sub> evolved with the help of the oxygen-evolving complex (Nathan and Wolfgang, 2015). On the electron acceptor side of PSI, NADP<sup>+</sup> is reduced to NADPH with electrons from P700 via ferredoxin (Fd) and Fd-NADP<sup>+</sup> reductase (FNR). Photo-oxidized P700 (P700<sup>+</sup>) is reduced with electrons from PSII via the Cyt *b<sub>6</sub>/f* complex and PC (or Cyt *c<sub>6</sub>*) in photosynthetic linear electron flow

(Jensen et al., 2007). The proton gradient across the thylakoid membrane ( $\Delta pH$ ), which is the motive force of the chloroplast ATP synthase (ATPase), is established at both the steps of  $H_2O$  oxidation on the luminal side of PSII and electron transport in the Q-cycle of the *Cytb<sub>6</sub>f* complex (Cramer et al., 2006). NADPH and ATP produced in the photosynthetic electron transport system are utilized for driving  $CO_2$  assimilation in the Calvin-Benson cycle (Calvin and Benson, 1948) and are required to be in concordance. However, dynamic natural environmental variations can easily unbalance the production and utilization of NADPH and ATP, resulting in photo-oxidative damage to photosynthetic cells.

Absorption of light energy exceeding the demand for photosynthetic  $CO_2$  assimilation can cause inactivation of these photosystems. This light-dependent inactivation of photosynthesis was first observed in the green alga *Chlorella* sp. more than 50 years ago and was termed photoinhibition by Kok (1956). Based on the characterization of photoinhibition using isolated thylakoid membranes, it has been separately recognized as deriving from the inactivation of PSII or PSI (Satoh, 1970). Photoinhibition of PSII leads to light-dependent degradation of the reaction center subunit (D1 protein), which is observed *in vivo* under various stress conditions, such as high light, high temperature, and drought. Photoinhibited PSII can rapidly recover by replacing the degraded D1 protein with a newly synthesized D1 protein in several hours. The processes that lead to the photo-oxidative damage of PSII are still under debate, although numerous studies have provided extensive and remarkable insights into the mechanisms of PSII photoinhibition (Krause et al., 1985; Aro et al., 1993; Sundby et al., 1993; Andersson and Barber, 1996; Neidhardt et al., 1998; Melis, 1999; Allakhverdiev and Murata, 2004; Hakala et al., 2005; Murata et al., 2007; Tyystjärvi, 2008; Fischer et al., 2013; Pospíšil, 2016; Jimbo et al., 2018).

In this review, we concentrate on the photoinhibition of PSI, which depends on both  $O_2$  and electrons produced by PSII (Satoh, 1970). Compared with studies on PSII photoinhibition, those on PSI photoinhibition are relatively rare, since it hardly occurs *in vivo*, even if oxygenic photoautotrophs are exposed to a stress treatment with excess light (e.g., continuous light illumination with high light) (Critchley, 1981; Powles and Björkman, 1982; Havaux and Eyletters, 1991). Previous studies, using isolated thylakoid membranes and chloroplasts, have suggested that PSI photoinhibition is derived from a dysfunction in the [4Fe-4S] clusters on the acceptor side of PSI (i.e.,  $F_X$ ,  $F_A$ , and  $F_B$ ), caused by reactive oxygen species (ROS) (Satoh, 1970; Inoue et al., 1986). PSI photoinhibition was first observed *in vivo* in the intact leaves of the chilling-sensitive plant *Cucumis sativus* under chilling stress (Terashima et al., 1994). The characterization of PSI photoinhibition under such conditions corroborated previous findings from *in vitro* studies and worked toward establishing the present theory of the mechanisms of PSI photoinhibition (Sonoike, 2011).

On the electron acceptor side of PSI, excess photo-excitation energy can reduce  $O_2$ , generating ROS, including superoxide anion radical ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), and hydroxyl radical ( $\cdot OH$ ) (Mehler, 1951; Asada, 2006;

Rutherford et al., 2012). Owing to their high reactivity, ROS can immediately inactivate PSI (Sonoike, 1996; Sonoike et al., 1997). In comparison with PSII, the damaged PSI takes a long time (days or weeks) to completely recover (Kudoh and Sonoike, 2002; Zivcak et al., 2015b). Therefore, PSI photoinhibition is a lethal event for oxygenic photoautotrophs. That is why PSI photoinhibition hardly occurs *in vivo* except for under specific conditions, such as chilling. The mechanisms of prevention of PSI photoinhibition had remained unknown for long.

## P700 OXIDATION AND ITS PHYSIOLOGICAL SIGNIFICANCE


The generation of ROS in PSI should be strictly suppressed for the purpose of preventing PSI photoinhibition *in vivo*. Based on the simple concept of oxygenic photosynthesis, the electron acceptor side of PSI is expected to be over-reduced when the Calvin-Benson cycle cannot follow the production of NADPH in the photosynthetic electron transport system. Nevertheless, PSI is always kept in an oxidized state in response to situations where the Calvin-Benson cycle is suppressed, which has been observed using *in vivo* spectroscopic measurement techniques for  $P700^+$  (Foyer et al., 1990; Harbinson and Hedley, 1993; Klughammer and Schreiber, 1994; Golding and Johnson, 2003; Miyake et al., 2005). The universal physiological response of oxygenic photoautotrophs is termed “P700 oxidation” and refers to the increase in the ratio of  $P700^+$  to the total amount of photo-oxidizable P700. In comparison with P700,  $P700^+$  cannot drive its photo-oxidation/reduction cycle but directly dissipate the photo-excitation energy as heat (Nuijs et al., 1986; Trissl, 1997; Bukhov and Carpentier, 2003). Therefore, P700 oxidation is expected to be directly linked to the quenching of excess light energy in PSI.

Recently, the impact of P700 oxidation on the alleviation of PSI photoinhibition has been demonstrated by a method to easily and selectively induce PSI photoinhibition in intact plant leaves at room temperature (Sejima et al., 2014). In the method named “repetitive short-pulse (SP) illumination (rSP illumination),” SP light (e.g., 300-1000 ms, 2000-20,000  $\mu mol$  photons  $m^{-2} s^{-1}$ ) is repetitively applied to plant leaves (e.g., every 10 s) under darkness. This experimental procedure is similar to continuously shooting the plants with a camera with a strobe light at night and can be defined as a severe form of artificial fluctuating light. This is different to illumination with continuous light; during rSP illumination, PSI is inactivated significantly faster than PSII in intact plant leaves, depending on the intensity and length of the SP light (Sejima et al., 2014; Zivcak et al., 2015b). Eliminating or limiting  $O_2$  suppress the inactivation of PSI, indicating that PSI photoinhibition during rSP illumination is caused by ROS (Sejima et al., 2014). This observation corresponds to results of previous studies on chilling-induced PSI photoinhibition (Sonoike, 1996; Sonoike et al., 1997). Therefore, rSP illumination is a useful tool to induce PSI photoinhibition *in vivo* (Zivcak et al., 2015a,b; Kono and Terashima, 2016; Takagi et al., 2017b; Mikko and Steffen, 2018).


Sejima et al. (2014) have applied rSP illumination to sunflower leaves under constant actinic light at different intensities, producing different P700 oxidation levels (Figures 1A,B), resulting in a linear relationship of P700 oxidation with the alleviation of PSI photoinhibition (Figure 1C) (Sejima et al., 2014). The effects of P700 oxidation on the protection of PSI against photoinhibition is also evidenced by the kinetics of  $P700^+$  in response to SP light during rSP illumination. In the intact leaves of angiosperms, P700 is excited by SP light and is kept in a reduced state during exposure to SP light (Figure 2A), which suggests that electron transport in PSI is limited on the acceptor side, but not on the donor side, during exposure to SP light. Contrarily, in the presence of a continuously high intensity background light, P700 is kept in an oxidized state during exposure to SP light (Figure 2B), which is due to a change in the limitation step of the electron transport system in PSI from acceptor to donor sides by P700 oxidation system (described in the next chapter). Furthermore, the addition of a far-red light in the background during rSP illumination can also stimulate P700 oxidation in the SP light to suppress PSI photoinhibition (Kono et al., 2017), which might suggest that shaded plants in an understory can efficiently keep P700 oxidized during natural “sunflecks.”

The impact of P700 oxidation on the alleviation of PSI photoinhibition has been observed not only during rSP illumination but also continuous light. In cyanobacteria, the progenitor of oxygenic photosynthesis, P700, is kept in an oxidized state in response to a suppression of photosynthesis under  $CO_2$  limitation, similarly to that in intact plant leaves (Badger and Schreiber, 1993). Shimakawa et al. (2016b) validated the common response of the redox state of PSI to  $CO_2$  limitation in three different cyanobacteria species, *Synechocystis* sp. PCC 6803, *Synechococcus elongatus* PCC 7942, and *Synechococcus* sp. PCC 7002 (Figure 3). Among the mutants deficient in flavodiiron protein (FLV) in each species, only the mutant of *Synechococcus* sp. PCC 7002 cannot keep P700 oxidized under  $CO_2$  limitation and suffers from PSI photoinhibition. These facts led us to suggest that the fate of PSI is determined by whether P700 can be kept in an oxidized state under excess light conditions. Furthermore, the inactivation of PSI in the mutant of *Synechococcus* sp. PCC 7002 has been observed even in the range of photon flux density between 200 and 300  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Shimakawa et al., 2016b), which indicates that PSI has the potential to generate ROS and suffer from PSI photoinhibition even under constant light 10 times less intense than sunlight, if the light exceeds the demand of the electron sinks, such as the Calvin-Benson cycle. Overall, without P700 oxidation, oxygenic photoautotrophs would easily suffer from PSI photoinhibition under natural environmental variation. The diverse strategies to keep P700 oxidized in these cyanobacteria species are further discussed at the section “Flavodiiron Protein (FLV)” in the chapter “Regulatory Mechanisms to Keep P700 in an Oxidized State, P700 Oxidation System.”

Besides being a quencher of light energy to suppress the generation of ROS in PSI,  $P700^+$  is assumed to play other important roles. For example, the charge recombination of  $P700^+$  with the electron acceptors in PSI (e.g.,  $F_X$ ) can occur in the range



**FIGURE 1 |** Relationship of P700 oxidation with the alleviation of the photo-oxidative damage in PSI during exposure to repetitive short-pulse (rSP) illumination. The sunflower plant leaves were exposed to rSP illumination (20,000  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ , 300 ms, every 10 s) in different light intensities producing different P700 oxidation levels (A). Residual photochemical activity of PSI was evaluated as the residual total photo-oxidizable P700 after rSP illumination for 4 h (B), and the decrease in the total photo-oxidizable P700 was plotted against the P700 oxidation levels (C). Data were from Sejima et al. (2014).



of the  $\mu\text{s}$ - or  $\text{ms}$ -order half times (Semenov et al., 2000). That is,  $\text{P700}^+$  can act as an electron sink to oxidize [4Fe–4S] clusters on the acceptor side of PSI, which might suppress the generation of  $\cdot\text{OH}$  (Sonoike, 1996). Additionally,  $\text{P700}^+$  possibly thermally dissipate excess light energy not only in PSI but also around PSII via energy transfer mechanisms, including state transition and/or spillover (Ueno et al., 2018; Yokono and Akimoto, 2018), which ultimately, also has the potential to alleviate PSII photoinhibition.

## REGULATORY MECHANISMS TO KEEP P700 IN AN OXIDIZED STATE, “P700 OXIDATION SYSTEM”

P700 oxidation is strictly regulated by diverse molecular mechanisms (collectively termed P700 oxidation system) in oxygenic photoautotrophs (Figure 4). Importantly, the redox state of P700 depends on both the electron donor and acceptor sides of PSI. There are various regulatory mechanisms functioning on both sides of PSI. P700 oxidation is commonly observed in oxygenic photoautotrophs in response to excess


light conditions, and the strategies to keep P700 in an oxidized state are diverse. Many diverse ways to oxidize P700 have been recognized already in the photosynthetic prokaryote cyanobacteria (Shimakawa et al., 2016b; Figure 3), which have supposedly developed and changed during the evolutionary history of oxygenic photoautotrophs.

## Limiting Electron Transport in Cyt $b_6/f$


On the donor side of PSI, the suppression of electron transport into PSI causes P700 oxidation. Photosynthetic linear electron flow has been recognized as being limited to the oxidation of reduced PQ (i.e., plastoquinol,  $\text{PQH}_2$ ) in Cyt  $b_6/f$  without any specific regulatory mechanisms at moderate lumen pH values (6.5–7.5). This is based on the understanding that the oxidation of  $\text{PQH}_2$  is the slowest step in the photosynthetic electron transport system, and that the amount of Cyt  $b_6/f$  is normally smaller than those of PSII and PSI in plant leaves (Stiehl and Witt, 1969; Anderson, 1992; Schöttler and Tóth, 2014), which is supported by a linear relationship between  $Q_A$  reduction and P700 oxidation (Shimakawa and Miyake, 2018a). The limitation of electron transport in Cyt  $b_6/f$  is likely to be a common strategy to keep P700 in an oxidized state in oxygenic photoautotrophs, considering the following regulatory mechanisms.

Electron transport in Cyt  $b_6/f$  is modulated by a regulatory mechanism, which is believed to be strongly associated with lumen acidification (Nishio and Whitmarsh, 1993). Lumen acidification is linked to photosynthetic linear electron flow since  $\Delta\text{pH}$  is provided by both the oxidation of  $\text{H}_2\text{O}$  at the luminal side of PSII and the Q-cycle in Cyt  $b_6/f$  (Schreiber and Neubauer, 1990; Avenson et al., 2005). Additionally, cyclic electron flow around PSI (CEF) can also promote the formation of  $\Delta\text{pH}$  (Nandha et al., 2007). Contrarily, lumen acidification is stimulated by narrowing the proton efflux in ATPase and ion channels on the thylakoid membrane (Takizawa et al., 2008; Armbruster et al., 2014), leading to P700 oxidation for the alleviation of PSI photoinhibition (Takagi et al., 2017a). These processes are often considered as photosynthetic control (Foyer et al., 1990; Schöttler and Tóth, 2014).

Electron transport in Cyt  $b_6/f$  can be suppressed in response to not only  $\Delta\text{pH}$  but also the reduction in the PQ pool, which has been recently proposed as reduction-induced suppression of electron flow (RISE) in cyanobacteria (Shaku et al., 2016; Shimakawa et al., 2018b). In the Q-cycle (Mitchell, 1966),  $\text{PQH}_2$  donates one electron to Cyt  $f$  through a [2Fe–2S] cluster at  $Q_p$  (or  $Q_o$ ) site in Cyt  $b_6/f$  and the other electron to a PQ at  $Q_n$  (or  $Q_i$ ) site in Cyt  $b_6/f$ . The PQ at the  $Q_n$  site is reduced with the second electron originating from PSII. Therefore, a shortage of PQ can inhibit the operation of the Q-cycle and suppress electron transport in Cyt  $b_6/f$ . Unfortunately, the molecular mechanisms of RISE remain poorly understood and have not yet been properly characterized. However, suppression of electron transport in Cyt  $b_6/f$  is relieved by alternative electron flow mediated by FLV (Shaku et al., 2016) and exogenously added  $\text{H}_2\text{O}_2$  (Shimakawa et al., 2018b). Here, we use the term “alternative electron flow” as the electron transport uncoupled with photosynthesis (i.e., the Calvin-Benson cycle). In cyanobacteria, FLV mediates the electron transport to  $\text{O}_2$  in PSI (Helman et al., 2003) and



**FIGURE 3 |** Relationship of P700 oxidation with the alleviation of the photo-oxidative damage in PSI during exposure to constant light. Both wild types and flavodiiron protein (FLV)-deficient mutants of the three cyanobacteria species grown under high-[CO<sub>2</sub>] conditions show the different responses of the photosynthetic electron transport system to the suppression of the Calvin-Benson cycle under CO<sub>2</sub> limitation: *Synechocystis* sp. PCC 6803 expresses FLV2/4 to mediate O<sub>2</sub>-dependent alternative electron transport but can keep P700 oxidized even without the FLV-mediated alternative electron sink; *Synechococcus elongatus* PCC 7942 suppresses photosynthetic linear electron flow and oxidizes P700 regardless of the existence of FLV; and *Synechococcus* sp. PCC 7002 suppresses photosynthetic linear electron flow and oxidizes P700 with the FLV-mediated alternative electron sink. Among these cyanobacterial cells, PSI photoinhibition is observed only in the mutant of *Synechococcus* sp. PCC 7002 that cannot keep P700 oxidized (Shimakawa et al., 2016b). All arrows indicate electron transport direction, and suppressed electron transport is represented by dashed arrows.



exogenously added H<sub>2</sub>O<sub>2</sub> stimulates the electron transport via peroxidase utilizing the electrons in PSI (Miyake et al., 1991), both of which theoretically consume the electrons on the acceptor

side of PSI but produce ΔpH (Schreiber and Neubauer, 1990). These facts, the electron transport in Cyt b<sub>6</sub>/f is modulated by not only lumen acidification (Trubitsin et al., 2003) but also another

mechanism sensitive to the reduction of the photosynthetic electron transport system in cyanobacteria. The effect of RISE on P700 oxidation is possibly observed also in  $C_3$  plants (Takagi et al., 2016a; Shimakawa and Miyake, 2018a), but further research is required.

## Thylakoid Terminal Oxidases

Terminal oxidases on the thylakoid membrane, including plastid terminal oxidase (PTOX) and cyanobacterial respiratory terminal oxidases (Cox, *aa<sub>3</sub>*-type Cyt *c* oxidase; and Cyd, Cyt *bd* quinol oxidase), are also defined as a P700 oxidation system, on the electron donor side of PSI. These oxidases are localized on the thylakoid membrane and donate electrons from the interchain of the photosynthetic electron transport system: i.e., PQH<sub>2</sub>, PC, and Cyt *c*<sub>6</sub> to O<sub>2</sub> (Pils et al., 1997; Joët et al., 2002). Thus, the electrons on the donor side of PSI can be leaked to O<sub>2</sub>, which has an impact on the redox state of P700 during the transition from dark to light (Bolychevtseva et al., 2015; Feilke et al., 2016).

Particularly in cyanobacteria, the respiratory electron transport system shares the same PQ pool with the photosynthetic electron transport system and can have a large effect on photosynthesis, compared with photosynthetic eukaryotes (Shimakawa et al., 2014; Misumi et al., 2016). Recently, in the cyanobacterium *Synechocystis* sp. PCC 6803, both Cox and Cyd have been found to contribute to P700 oxidation and the alleviation of PSI photoinhibition during rSP illumination (Shimakawa and Miyake, 2018b). Unfortunately, the electron transport capacities of these terminal oxidases are yet to be quantitatively determined to be established as a suitable alternative electron sink for P700 oxidation in the cyanobacterial cells *in vivo* (Schuurmans et al., 2015). Oxidation of the PQ pool by both Cox and Cyd in the darkness is likely to help P700 oxidation to be induced in response to a light illumination. Further, these respiratory terminal oxidases can pump H<sup>+</sup> into the luminal side of the thylakoid membrane (Paumann et al., 2005), possibly stimulating the limitation of electron transport in Cyt *b<sub>6</sub>/f* by photosynthetic control.

The chloroplast of photosynthetic eukaryotes harbors PTOX, a non-heme diiron carboxylate protein showing sequence similarity to the mitochondrial alternative oxidase, which binds strongly to the stromal side of the thylakoid membrane and functions in the process of chlororespiration to oxidize the PQ pool (McDonald et al., 2011; Johnson and Stepien, 2016). It has been reported that the alternative electron flux through PTOX is not functional for the protection of PSII and PSI against photoinhibition at the steady-state photosynthesis (Rosso et al., 2006). At least in the mature tissues of plants the amount of PTOX is 100 times smaller than that of PSII (Lennon et al., 2003). On the other hand, the recombinant PTOX protein of rice harbors the enough O<sub>2</sub> reducing activity comparable to photosynthetic linear electron transport ( $k_{cat}$ , >20 s<sup>-1</sup>; Yu et al., 2014). Additionally, the heterologous expression of *Chlamydomonas reinhardtii* PTOX in the tobacco leads to P700 oxidation (Feilke et al., 2016). These facts indicate that PTOX has the potential to contribute to P700 oxidation for the alleviation of PSI photoinhibition in photosynthetic eukaryotes in the

situations where the relative amount of PTOX to photosystems increases (Rumeau et al., 2007).

## Photorespiration

On the electron acceptor side of PSI, an alternative electron sink, uncoupled with photosynthesis, supports P700 oxidation by relieving limitation of PSI on the acceptor side. Photorespiration is initiated by the oxygenation reaction of ribulose 1,5-bisphosphate (RuBP) carboxylase/oxygenase (so-called Rubisco) with 3-phosphoglycerate and 2-phosphoglycolate produced from RuBP and CO<sub>2</sub> (Berry et al., 1978; Ogren, 1984). In the processes for regeneration of 3-phosphoglycerate from 2-phosphoglycolate in photorespiratory carbon oxidation (PCO) cycle, both reduced Fd and ATP are required. Additionally, both RuBP and CO<sub>2</sub> regenerated by photorespiration are utilized again for CO<sub>2</sub> assimilation in the Calvin-Benson cycle. That is, photorespiration can function as an O<sub>2</sub>-dependent alternative electron sink to dissipate excess light energy (Powles et al., 1979; Kozaki and Takeba, 1996; Takahashi et al., 2007). Indeed, photorespiration functions as the largest alternative electron flow to O<sub>2</sub> (Badger et al., 2000; Ruuska et al., 2000; Driever and Baker, 2011; Sejima et al., 2016), and is responsible for P700 oxidation and the protection of PSI against photoinhibition in  $C_3$  plant leaves (Wiese et al., 1998; Takagi et al., 2016a; Wada et al., 2018).

In contrast to  $C_3$  plant leaves, photorespiration does not function as an alternative electron sink for P700 oxidation in cyanobacteria and algae. Though the genes for PCO cycle enzymes are commonly conserved in oxygenic photoautotrophs, a variety of algae, including cyanobacteria, green algae, and diatoms, show little O<sub>2</sub>-dependent electron sink capacity derived from photorespiration, even under CO<sub>2</sub> limitation (Bidwell and McLachlan, 1985; Weger et al., 1989; Hayashi et al., 2014; Shimakawa et al., 2015, 2016a, 2017b). In aquatic environments the air-equilibrated O<sub>2</sub> concentration is approximately 250 μM at 25°C and the diffusion coefficient of O<sub>2</sub> decreases to approximately 0.01% of that in the atmosphere, which probably makes it difficult to utilize photorespiration, considering that the oxygenation reaction of RuBP catalyzed by Rubisco has significantly low affinity for O<sub>2</sub>. Despite of the large varieties, the  $K_m$  values have been recently reported to be in the range between 100 and 1600 μM at 25°C in diverse oxygenic photoautotrophs except for the Rubisco of Archaea (Tcherkez, 2016; Orr et al., 2016). As mutants of cyanobacteria and algae deficient in the genes for the PCO cycle are impaired in their growth, photorespiration is assumed to play other important roles, rather than acting as an alternative electron sink in these photoautotrophs (Eisenhut et al., 2006; Rademacher et al., 2016). O<sub>2</sub>-dependent electron transport activity with low affinity is observed in *Euglena gracilis* (Euglenophyta) (Shimakawa et al., 2017b); the secondary alga harboring chloroplasts is believed to be derived from green algae (Falkowski et al., 2004), which may suggest that *E. gracilis* uniquely utilizes photorespiration as an electron sink (Yokota and Kitaoka, 1987). Interestingly, photorespiration-derived electron sink comparable to the Calvin-Benson cycle is observed in liverworts, ferns, gymnosperms, and angiosperms except for in  $C_4$  plants (Hanawa et al., 2017). These data indicate that photorespiration had started to function as a

large alternative electron sink since oxygenic photoautotrophs were first exposed to high partial pressures of O<sub>2</sub> in the atmosphere.

## Flavodiiron Protein (FLV)

The protein family of FLV (or FDP) is defined based on two domains: a diiron center and a flavin mononucleotide-binding, and reduces O<sub>2</sub> and NO directly into H<sub>2</sub>O and N<sub>2</sub>O using coenzymes such as rubredoxin and F<sub>420</sub> (Romão et al., 2016). In addition, FLV in oxygenic photoautotrophs harbors a unique domain, similar to a flavin:NAD(P)H oxidoreductase, and therefore has been characterized by an ability to catalyze the reduction of O<sub>2</sub> directly to H<sub>2</sub>O, with NAD(P)H as the electron donor (Vicente et al., 2002). The physiological function of FLV has been well characterized in the cyanobacterium *Synechocystis* sp. PCC 6803 in the pioneering work of Helman et al. (2003). The authors indicate that FLV mediates an O<sub>2</sub>-dependent alternative electron flow, probably on the acceptor side of PSI, and supports P700 oxidation. The electron sink capacity of FLV-mediated electron transport is evidenced by both the measurement of <sup>18</sup>O<sub>2</sub> photoreduction (Helman et al., 2003; Allahverdiyeva et al., 2013; Burlacot et al., 2018) and the simultaneous evaluation of O<sub>2</sub> evolution with Chl fluorescence (Shimakawa et al., 2015). On the contrary, the physiological electron donor for FLV is still unknown. Some recombinant FLV proteins of the cyanobacterium *Synechocystis* sp. PCC 6803 show NAD(P)H-dependent O<sub>2</sub> reduction into H<sub>2</sub>O, but the reduction rates are more than 100 times smaller than those of anaerobic bacteria (Vicente et al., 2002; Di Matteo et al., 2008; Shimakawa et al., 2015). In addition, it has been suggested that FLV interacts with Fd in *Synechocystis* sp. PCC 6803 (Hanke et al., 2011). The molecular mechanisms of FLV still await biochemical validation *in vitro*.

The impact of FLV on P700 oxidation is diversified already in the photosynthetic prokaryote cyanobacteria. Three cyanobacterial species show the different responses of the regulation of photosynthetic electron transport to CO<sub>2</sub> limitation: *Synechocystis* sp. PCC 6803 expresses FLV2/4 to induce the large alternative electron flux to O<sub>2</sub> uncoupled with photosynthesis; *Synechococcus elongatus* PCC 7942 suppresses the electron transport in Cyt *b<sub>6</sub>/f*; and *Synechococcus* sp. PCC 7002 keeps the electron transport capacity with the alternative electron transport to O<sub>2</sub> through FLV1/3 dependent on the CO<sub>2</sub> concentration of the growth conditions (Figure 3; Shimakawa et al., 2016a,b). Among these three cyanobacteria species, only in *Synechococcus* sp. PCC 7002 the FLV-knockout mutant suffers from PSI photoinhibition due to the inability to keep P700 oxidized in the situation where photosynthesis is suppressed under CO<sub>2</sub> limitation (Figure 3; Shimakawa et al., 2016b). That is, FLV is the dominant regulator for the redox state of P700 in this species. On the other word, the other two species can keep P700 oxidized even in the absence of FLV by relying on the other regulatory mechanisms (Figure 3). Unfortunately, it is still unclear what regulatory mechanisms complement the capacity for P700 oxidation in the FLV-knockout mutants of these two species. Shaku et al. (2016) suggests that the limitation of electron transport in Cyt *b<sub>6</sub>/f* by RISE has the large impact on P700

oxidation in *Synechococcus elongatus* PCC 7942. Additionally, from the fact that the genes for Cyt *b<sub>6</sub>/f* are missed in the genome of *Synechococcus* sp. PCC 7002, different from the other two species (Shimakawa et al., 2016b), the terminal oxidase is suggested to be one possibility to complement P700 oxidation in the absence of FLV in *Synechocystis* sp. PCC 6803 and *Synechococcus elongatus* PCC 7942 (Shimakawa and Miyake, 2018b).

Among the oxygenic photoautotrophs, the genes for FLV are conserved in Cyanophyta (cyanobacteria), Chlorophyta (green algae), Bryophyta (liverworts and mosses), Pteridophyta (ferns), gymnosperms, and limited secondary algae (e.g., *Symbiodinium* sp.), in which P700 is rapidly oxidized in response to light exposure in the presence of O<sub>2</sub> to alleviate PSI photoinhibition (Allahverdiyeva et al., 2013; Shirao et al., 2013; Roberty et al., 2014; Gerotto et al., 2016; Shimakawa et al., 2016b, 2017a; Chaux et al., 2017; Ilík et al., 2017; Noridomi et al., 2017; Takagi et al., 2017b). Interestingly, angiosperms have lost FLV at the genetic level (Allahverdiyeva et al., 2015; Yamamoto et al., 2016; Alboresi et al., 2018). Most importantly, P700 oxidation, but not FLV, is essential for oxygenic photoautotrophs to protect PSI against photoinhibition (Shimakawa et al., 2016b). In other words, FLV is not required if P700 can be kept oxidized in excess light conditions without it. Indeed, wild-type plant leaves of angiosperms can rapidly induce P700 oxidation except for during artificial severe stress conditions such as rSP illumination (Takagi et al., 2017b; Shimakawa and Miyake, 2018a). Additionally, most of the red algae (Rhodophyta) and secondary algae that have red plastid, including Cryptophyta, Haptophyta, and Heterokontophyta (diatoms, brown algae, etc.), can rapidly induce P700 oxidation in response to excess light and alleviate PSI photoinhibition during rSP illumination in the absence of FLV (Shimakawa et al., 2018a), implying that FLV is not completely required for oxygenic photoautotrophs already at the time that red algae had birthed. It would not be unexpected for angiosperms to have lost FLV during their evolutionary history. In these oxygenic photoautotrophs without FLV, P700 oxidation should be relying mainly on other regulatory mechanisms as mentioned in this chapter (e.g. limiting electron transport in Cyt *b<sub>6</sub>/f*).

The requirements of FLV are diverse in a variety of oxygenic photoautotrophs, likely depending also upon their ecological niche. The liverwort *M. polymorpha* preferably utilizes the alternative electron sink of FLV, but not photorespiration, when it is submerged (Shimakawa et al., 2017a). Taking the high affinity of the reaction with O<sub>2</sub> (*K<sub>m</sub>*, a few or less μM) (Vicente et al., 2002; Shimakawa et al., 2015) into consideration, FLV probably provides better benefits than photorespiration under water. Additionally, the exposure to a far-red light in the terrestrial fields possibly affect the strategies to utilize FLV in a variety of basal land plants (Kono et al., 2017).

## Proton Gradient Regulation 5 (PGR5)

A number of studies have reported that the 10 kDa thylakoid membrane-associated protein, called PGR5, is essential to keep P700 in an oxidized state in green algae and land plants. A lack of PGR5 creates a profound limitation in PSI on the electron acceptor side, resulting in PSI photoinhibition



under excess light conditions (Munekage et al., 2002). Despite the clear experimental evidence from mutant plants, the molecular mechanisms of PGR5 for P700 oxidation remain poorly understood and controversial. Since the protein was first identified, PGR5 has been proposed to drive CEF together with PGR5-like 1 protein (i.e., PGRL1) for the alleviation of the limitation of PSI on the electron acceptor side and for inducing lumen acidification (Munekage et al., 2002; Yamori and Shikanai, 2016; and references therein). Additionally, a lack of PGR5 impairs the association of FNR with the thylakoid membrane (Mosebach et al., 2017), indicating that PGR5 possibly affects photosynthetic linear electron flow (Takagi and Miyake, 2018). Furthermore, Kanazawa et al. (2017) suggests that PGR5 may function in adjusting the activity of ATPase rather than driving CEF, which is supported by the fact that the profiles of the mutants impaired in ATPase are strikingly similar to those of the PGR5 mutants (Kanazawa et al., 2017). Overall, the relationship of PGR5 with the photosynthetic electron transport system remains controversial.

The impact of PGR5 on P700 oxidation has changed from cyanobacteria to angiosperms. Although PGR5 is essential for P700 oxidation in angiosperms (Munekage et al., 2002), a lack of PGR5 has no effect on cyanobacterial photosynthesis (Allahverdiyeva et al., 2013). The contribution of PGR5 to P700 oxidation is observed in the green alga *C. reinhardtii* (Mosebach et al., 2017), indicating that a PGR5-dependent mechanism started to function as a P700 oxidation system after photosynthetic eukaryotes had evolved.

## Chloroplast NADPH Dehydrogenase (NDH)

Recently, it has been reported that NDH can also function as a P700 oxidation system under fluctuating light in  $C_3$  plant leaves. During a study using artificial, angularly incident, fluctuating light, mutants deficient in NDH showed impaired induction of P700 oxidation in *Arabidopsis thaliana* and *Oryza sativa* (Kono and Terashima, 2016; Yamori et al., 2016). The impact of NDH on P700 oxidation has also been tested during studies using sine-like artificial fluctuating light, named Umibozu, at different frequencies in *A. thaliana*, indicating that NDH is required for P700 oxidation only following a rapid change in light intensity under rapidly fluctuating light (Shimakawa and Miyake, 2018a). In chloroplasts of  $C_3$  plant leaves, NDH has effects on CEF (Shikanai et al., 1998) and chlororespiration (Sazanov et al., 1998). One hypothesis of the mechanism of NDH to support P700 oxidation is that NDH-dependent CEF functions under fluctuating light to produce  $\Delta$ pH, limiting the electron transport in Cyt  $b_6/f$  by photosynthetic control and accelerating the induction of photosynthesis to relieve the limitation of the electron acceptor side of PSI (Martin et al., 2015; Ishikawa et al., 2016; Kono and Terashima, 2016; Yamori et al., 2016). Other is that NDH contributes to oxidation of the chloroplast NADP<sup>+</sup> pool in the darkness or low light in the process of chlororespiration, which can support the rapid start of P700 oxidation in

response to the illumination with a fast fluctuating light (Shimakawa and Miyake, 2018a). Further studies are required on the detailed mechanisms of P700 oxidation by NDH.

In cyanobacteria, NDH can have a large effect on the redox state of both PQ and NADPH pools within cells, compared with that in plant leaves, because NDH also functions in the respiratory electron transport system (Mi et al., 2000; Ogawa et al., 2013). Similar to the case of terminal oxidases, it should be noted that the effects of NDH on P700 oxidation cannot be easily compared between cyanobacteria and photosynthetic eukaryotes.

## RETHINKING THE MECHANISM OF PSI PHOTOINHIBITION AND THE DYNAMICS OF ROS IN PSI

Recent studies have not only supported the hypothetical model of the mechanisms of PSI photoinhibition, established on the basis of the experimental findings of a study based on chilling stress (Sonoike, 2011), but have also provided novel insights into mechanisms for the generation of ROS in PSI. In this review, we revisited the hypothetical model of the mechanisms of PSI photoinhibition and the dynamics of ROS in PSI in oxygenic photoautotrophs. Originally, in chloroplasts the production of the ROS  $O_2^-$  on the electron acceptor side of PSI has been defined as the Mehler reaction (Mehler, 1951). There are four electron acceptors for photo-excited P700 in PSI: chlorophyll  $A_0$  (primary acceptor), phylloquinone  $A_1$ , and [4Fe-4S] clusters  $F_X$  and  $F_A/F_B$ . Both phylloquinones exist asymmetrically in the heterodimeric reaction centers of each of PsaA and PsaB (i.e.,  $A_{0A}$ ,  $A_{0B}$ ,  $A_{1A}$ , and  $A_{1B}$ ) (Joliot and Joliot, 1999). Among these acceptors, the Mehler reaction has been proposed to mainly occur at  $A_{1B}$  (Kruk et al., 2003) or  $F_X$  (Takahashi and Asada, 1988), which is supported by the lower midpoint redox potential (versus NHE) for  $A_{1B}$  (−820 mV) and  $F_X$  (−730 mV) (Brettel and Leibl, 2001; Kozuleva and Ivanov, 2010) than estimated ( $O_2/O_2^-$ ) in the lipid bilayer (from −500 to −600 mV versus NHE) (Wardman, 1990).

The increase in the photo-oxidative damage in PSI during rSP illumination in the range of light intensity for SP light, from 2000 to 20,000  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Sejima et al., 2014), corresponds to the non-light saturation manner of the production of  $O_2^-$  at phylloquinones in PSI (Kozuleva et al., 2014), which implies that  $O_2^-$  produced by  $A_{1B}$  is likely to cause PSI photoinhibition. The production of  $O_2^-$  can occur on both the stromal and luminal sides of PSI (Takahashi and Asada, 1988; Mubarakshina et al., 2006).  $O_2^-$  produced during this process should immediately be disproportionated into  $H_2O_2$  by the oxidation of ascorbate and superoxide dismutase in chloroplasts (Scarpa et al., 1983; Miyake and Asada, 1992). Further,  $H_2O_2$  can react with the reduced [4Fe-4S] centers to produce  $\cdot\text{OH}$  where the acceptor side of PSI is reduced (Youngman and Elstner, 1981; Sonoike et al., 1997). The addition of methyl viologen, which strongly oxidizes the [4Fe-4S] centers and reduces  $O_2$  to  $O_2^-$  on the stromal side of PSI, clearly alleviates PSI photoinhibition, which suggests that the photo-oxidative damage in PSI is caused

by  $\cdot\text{OH}$  produced, depending on the reduced [4Fe–4S] centers (Sonoike, 1996) and/or ROS generated inside PSI (Takagi et al., 2016b).

Although all oxygenic photoautotrophs develop a variety of scavenging enzymes for ROS, including superoxide dismutase, ascorbate peroxidase, and catalase (Asada, 2006), at least in a physiological sense, these scavenging enzymes possibly have no impact on the alleviation of PSI photoinhibition. Originally, Terashima et al. (1998) has found the slight accumulation of  $\text{H}_2\text{O}_2$  with the lower activity of thylakoid-bound ascorbate peroxidase in the  $\text{C}_3$  plant *Cucumis sativus* in the transition to chilling stress where PSI photoinhibition occurs in this plant. Whereas this finding clearly suggests that the ascorbate peroxidase modulates the  $\text{H}_2\text{O}_2$  concentration in the plant leaves, it is still unclear if the accumulated  $\text{H}_2\text{O}_2$  causes PSI photoinhibition. Overall, the effects of P700 oxidation can be lost in the isolated thylakoid membrane. Indeed, it has been proposed that the addition of superoxide dismutase and catalase do not alleviate PSI photoinhibition during a high light-stress treatment in an isolated PSI submembrane (Subramanyam et al., 2005). In the cyanobacterial FLV mutant that cannot keep P700 oxidized under  $\text{CO}_2$  limitation, PSI photoinhibition is rapidly induced even under constant light with approximately  $200 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Shimakawa et al., 2016b). In chloroplasts of plant leaves, PSI photoinhibition is observed during rSP illumination even in the presence of sufficient activities of superoxide dismutase and ascorbate peroxidase (Takagi et al., 2016b). These data clearly suggest that the scavenging enzymes of ROS cannot prevent PSI photoinhibition. That is, once produced,  $\text{O}_2^-$  is supposed to immediately attack PSI and/or trigger the production of  $\cdot\text{OH}$  before it is scavenged, which may be supported by the significantly shorter lives of  $\text{O}_2^-$  (2–4  $\mu\text{s}$ ) and  $\cdot\text{OH}$  (<1  $\mu\text{s}$ ) than that of  $\text{H}_2\text{O}_2$  (1 ms; Van Breusegem et al., 2001). Contrarily, the production and diffusion of  $\text{H}_2\text{O}_2$  can be easily detected in photosynthetic cells (Michelet et al., 2013; Roach et al., 2015), which is reasonable considering that  $\text{H}_2\text{O}_2$  functions as a signaling molecule in oxygenic photoautotrophs (Van Breusegem et al., 2001; Gläßer et al., 2014; Dietz et al., 2016). Overall, we propose that the production of ROS leading to PSI photoinhibition can be completely distinguished from those related to dynamic metabolic and signaling mechanisms. It is possible that the different production site of  $\text{O}_2^-$  causes the different effects of ROS on photosynthetic cells (Takagi et al., 2016b). Unfortunately, identification of the primary site in PSI attacked by ROS is still controversial (Tjus et al., 1999; Subramanyam et al., 2005; Takagi et al., 2016b), and more research is needed to identify the proper dynamics of ROS around PSI *in vivo*. The qualitative and quantitative relationships between the production of ROS and PSI photoinhibition should be addressed in future.

Besides  $\text{O}_2^-$  and  $\cdot\text{OH}$ , singlet  $\text{O}_2$  ( $^1\text{O}_2$ ) has been recently suggested to be generated in PSI and cause PSI photoinhibition. In the core and LHC complexes associated with PSI, triplet Chl can produce  $^1\text{O}_2$  to cause PSI photoinhibition, unless carotenoids such as  $\beta$ -carotene quench the triplet Chl

(Subramanyam et al., 2005; Cazzaniga et al., 2012, 2016), which is considered a potential mechanism for alleviating PSI photoinhibition, in addition to P700 oxidation system. Long-lived triplet P700 suggests that  $^1\text{O}_2$  is unlikely to be generated from triplet P700 (Setif et al., 1981; Rutherford et al., 2012). Nevertheless, the generation of  $^1\text{O}_2$  originating from triplet P700 has recently been suggested during rSP illumination (Takagi et al., 2016b, 2017b). Ultimately, it is difficult to exclude the possibility that  $^1\text{O}_2$  has an impact on the photo-oxidative damage in PSI under severe excess light and in specific mutants.

## CONCLUDING REMARKS

Oxygenic photoautotrophs can safely undergo photosynthesis owing to P700 oxidation system. Despite the current poor understanding of the mechanisms of PSI photoinhibition, the effects of P700 oxidation on the alleviation of PSI photoinhibition discussed herein are likely to be true based on a number of experimental results (e.g., **Figures 1–3**). On the contrary, a recent study has reported an inconsistency between PSI photoinhibition and P700 oxidation in two different shade-established tropical tree species (Huang et al., 2015). Indeed, the degrees of PSI photoinhibition are diverse among a variety of oxygenic photoautotrophs, regardless of P700 oxidation levels (Takagi et al., 2017b), which probably reflects the different levels of robustness of PSI against ROS in each species. Most importantly, P700 oxidation is not directly linked to photosynthesis. Nevertheless, it is impossible for oxygenic photoautotrophs to live without P700 oxidation system because PSI photoinhibition is the lethal event for them (Shimakawa et al., 2016b). These facts reflect that all oxygenic photoautotrophs are confronted with the potential risk of photo-oxidative damage inevitably accompanied with exposure to light and  $\text{O}_2$ . Diverse molecular mechanisms, i.e., P700 oxidation system, support P700 oxidation (**Figure 4**). On the contrary, almost all the agents of P700 oxidation system are still not characterized at the molecular level. There would be various ways to keep P700 oxidized. Intriguingly, the cyanobacterium *Leptolyngbya* sp., the species thriving in the harsh conditions of the desert, has been suggested to induce P700 oxidation by constricting the thylakoid lumen to limit diffusion of PC (Bar-Eyal et al., 2015). Unimaginable diversity of the strategies for P700 oxidation are possibly still unexplored in a variety of oxygenic photoautotrophs.

## AUTHOR CONTRIBUTIONS

CM conceived the project and GS wrote the manuscript.

## FUNDING

This work was supported by the Core Research for Evolutional Science and Technology (CREST) of the Japan Science and Technology Agency (Grant No. AL65D21010 to CM). GS is supported as a JSPS research fellow (Grant No. 16J03443).

## REFERENCES

- Alboresi, A., Storti, M., and Morosinotto, T. (2018). Balancing protection and efficiency in the regulation of photosynthetic electron transport across plant evolution. *New Phytol.* doi: 10.1111/nph.15372 [Epub ahead of print].
- Allahverdiyeva, Y., Isojarvi, J., Zhang, P., and Aro, E. M. (2015). Cyanobacterial oxygenic photosynthesis is protected by flavodiiron proteins. *Life* 5, 716–743. doi: 10.3390/life5010716
- Allahverdiyeva, Y., Mustila, H., Ermakova, M., Bersanini, L., Richaud, P., Ajlani, G., et al. (2013). Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and photosynthesis under fluctuating light. *Proc. Natl. Acad. Sci. U.S.A.* 110, 4111–4116. doi: 10.1073/pnas.1221194110
- Allakhverdiev, S. I., and Murata, N. (2004). Environmental stress inhibits the synthesis de novo of proteins involved in the photodamage–repair cycle of Photosystem II in *Synechocystis* sp. PCC 6803. *Biochim. Biophys. Acta Bioenerg.* 1657, 23–32. doi: 10.1016/j.bbabi.2004.03.003
- Anderson, J. M. (1992). Cytochrome *b<sub>6</sub>f* complex: dynamic molecular organization, function and acclimation. *Photosynth. Res.* 34, 341–357. doi: 10.1007/bf00029810
- Andersson, B., and Barber, J. (1996). “Mechanisms of photodamage and protein degradation during photoinhibition of photosystem II,” in *Photosynthesis and the Environment*, ed. N. R. Baker (Dordrecht: Springer), 101–121.
- Armbruster, U., Carrillo, L. R., Venema, K., Pavlovic, L., Schmidtman, E., Kornfeld, A., et al. (2014). Ion antiport accelerates photosynthetic acclimation in fluctuating light environments. *Nat. Commun.* 5:5439. doi: 10.1038/ncomms6439
- Aro, E. M., McCaffery, S., and Anderson, J. M. (1993). Photoinhibition and D1 protein degradation in peas acclimated to different growth irradiances. *Plant Physiol.* 103, 835–843. doi: 10.1104/pp.103.3.835
- Asada, K. (2006). Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* 141, 391–396. doi: 10.1104/pp.106.082040
- Avenson, T. J., Cruz, J. A., Kanazawa, A., and Kramer, D. M. (2005). Regulating the proton budget of higher plant photosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* 102, 9709–9713. doi: 10.1073/pnas.0503952102
- Badger, M. R., and Schreiber, U. (1993). Effects of inorganic carbon accumulation on photosynthetic oxygen reduction and cyclic electron flow in the cyanobacterium *Synechococcus* PCC7942. *Photosynth. Res.* 37, 177–191. doi: 10.1007/bf00032822
- Badger, M. R., von Caemmerer, S., Ruuska, S., and Nakano, H. (2000). Electron flow to oxygen in higher plants and algae: rates and control of direct photoreduction (Mehler reaction) and rubisco oxygenase. *Philos. Trans. R. Soc. Lond.* 355, 1433–1446. doi: 10.1098/rstb.2000.0704
- Barber, J., and Andersson, B. (1994). Revealing the blueprint of photosynthesis. *Nature* 370, 31–34. doi: 10.1038/370031a0
- Bar-Eyal, L., Eisenberg, I., Faust, A., Raanan, H., Nevo, R., Rappaport, F., et al. (2015). An easily reversible structural change underlies mechanisms enabling desert crust cyanobacteria to survive desiccation. *Biochim. Biophys. Acta Bioenerg.* 1847, 1267–1273. doi: 10.1016/j.bbabi.2015.07.008
- Berry, J. A., Osmond, C. B., and Lorimer, G. H. (1978). Fixation of <sup>18</sup>O<sub>2</sub> during photorespiration. *Plant Physiol.* 62, 954–967. doi: 10.1104/pp.62.6.954
- Bidwell, R. G. S., and McLachlan, J. (1985). Carbon nutrition of seaweeds: photosynthesis, photorespiration and respiration. *J. Exp. Mar. Biol. Ecol.* 86, 15–46. doi: 10.1016/0022-0981(85)90040-1
- Bolychevtseva, Y. V., Kuzminov, F. I., Elanskaya, I. V., Gorbunov, M. Y., and Karapetyan, N. V. (2015). Photosystem activity and state transitions of the photosynthetic apparatus in cyanobacterium *Synechocystis* PCC 6803 mutants with different redox state of the plastoquinone pool. *Biochemistry* 80, 50–60. doi: 10.1134/s000629791501006x
- Brettel, K., and Leibl, W. (2001). Electron transfer in photosystem I. *Biochim. Biophys. Acta Bioenerg.* 1507, 100–114. doi: 10.1016/S0005-2728(01)00202-X
- Bukhov, N. G., and Carpentier, R. (2003). Measurement of photochemical quenching of absorbed quanta in photosystem I of intact leaves using simultaneous measurements of absorbance changes at 830 nm and thermal dissipation. *Planta* 216, 630–638. doi: 10.1007/s00425-002-0886-2
- Burlacot, A., Sawyer, A., Cuiñé, S., Auroy, P., Blangy, S., Happe, T., et al. (2018). Flavodiiron-mediated O<sub>2</sub> photoreduction links H<sub>2</sub> production with CO<sub>2</sub> fixation during the anaerobic induction of photosynthesis. *Plant Physiol.* 177, 1639–1649. doi: 10.1104/pp.18.00721
- Calvin, M., and Benson, A. A. (1948). *The Path of Carbon in Photosynthesis*. Washington, DC: US Atomic Energy Commission. doi: 10.2172/927200
- Cazzaniga, S., Bressan, M., Carbonera, D., Agostini, A., and Dall’Osto, L. (2016). Differential roles of carotenes and xanthophylls in photosystem I photoprotection. *Biochemistry* 55, 3636–3649. doi: 10.1021/acs.biochem.6b00425
- Cazzaniga, S., Li, Z., Niyogi, K. K., Bassi, R., and Dall’Osto, L. (2012). The Arabidopsis *szl1* mutant reveals a critical role of β-carotene in photosystem I photoprotection. *Plant Physiol.* 159, 1745–1758. doi: 10.1104/pp.112.201137
- Chaux, F., Burlacot, A., Mekhalif, M., Auroy, P., Blangy, S., Richaud, P., et al. (2017). Flavodiiron proteins promote fast and transient O<sub>2</sub> photoreduction in *Chlamydomonas*. *Plant Physiol.* 174, 1825–1836. doi: 10.1104/pp.17.00421
- Cramer, W. A., Zhang, H., Yan, J., Kurisu, G., and Smith, J. L. (2006). Transmembrane traffic in the cytochrome *b<sub>6</sub>f* complex. *Annu. Rev. Biochem.* 75, 769–790. doi: 10.1146/annurev.biochem.75.103004.142756
- Critchley, C. (1981). Studies on the mechanism of photoinhibition in higher plants. *Plant Physiol.* 67, 1161–1165. doi: 10.1104/pp.67.6.1161
- Di Matteo, A., Scandurra, F. M., Testa, F., Forte, E., Sarti, P., Brunori, M., et al. (2008). The O<sub>2</sub>-scavenging flavodiiron protein in the human parasite *Giardia intestinalis*. *J. Biol. Chem.* 283, 4061–4068. doi: 10.1074/jbc.M705605200
- Dietz, K.-J., Turkan, I., and Krieger-Liszka, A. (2016). Redox- and reactive oxygen species-dependent signaling into and out of the photosynthesizing chloroplast. *Plant Physiol.* 171, 1541–1550. doi: 10.1104/pp.16.00375
- Driever, S. M., and Baker, N. R. (2011). The water–water cycle in leaves is not a major alternative electron sink for dissipation of excess excitation energy when CO<sub>2</sub> assimilation is restricted. *Plant Cell Environ.* 34, 837–846. doi: 10.1111/j.1365-3040.2011.02288.x
- Eisenhut, M., Kahlon, S., Hasse, D., Ewald, R., Lieman-Hurwitz, J., Ogawa, T., et al. (2006). The plant-like C<sub>2</sub> glycolate cycle and the bacterial-like glycerate pathway cooperate in phosphoglycolate metabolism in cyanobacteria. *Plant Physiol.* 142, 333–342. doi: 10.1104/pp.106.082982
- Falkowski, P. G., Katz, M. E., Knoll, A. H., Quigg, A., Raven, J. A., Schofield, O., et al. (2004). The evolution of modern eukaryotic phytoplankton. *Science* 305, 354–360. doi: 10.1126/science.1095964
- Feilke, K., Streb, P., Cornic, G., Perreau, F., Kruk, J., and Krieger-Liszka, A. (2016). Effect of *Chlamydomonas* plastid terminal oxidase 1 expressed in tobacco on photosynthetic electron transfer. *Plant J.* 85, 219–228. doi: 10.1111/tjp.13101
- Fischer, B. B., Hideg, E., and Krieger-Liszka, A. (2013). Production, detection, and signaling of singlet oxygen in photosynthetic organisms. *Antioxid. Redox Signal.* 18, 2145–2162. doi: 10.1089/ars.2012.5124
- Foyer, C., Furbank, R., Harbinson, J., and Horton, P. (1990). The mechanisms contributing to photosynthetic control of electron transport by carbon assimilation in leaves. *Photosynth. Res.* 25, 83–100. doi: 10.1007/bf00035457
- Gerotto, C., Alboresi, A., Meneghesso, A., Jokel, M., Suorsa, M., Aro, E.-M., et al. (2016). Flavodiiron proteins act as safety valve for electrons in *Physcomitrella patens*. *Proc. Natl. Acad. Sci. U.S.A.* 113, 12322–12327. doi: 10.1073/pnas.1606685113
- Gläßer, C., Haberer, G., Finkemeier, I., Pfanschmidt, T., Kleine, T., Leister, D., et al. (2014). Meta-analysis of retrograde signaling in *Arabidopsis thaliana* reveals a core module of genes embedded in complex cellular signaling networks. *Mol. Plant* 7, 1167–1190. doi: 10.1093/mp/ssu042
- Golding, A. J., and Johnson, G. N. (2003). Down-regulation of linear and activation of cyclic electron transport during drought. *Planta* 218, 107–114. doi: 10.1007/s00425-003-1077-5
- Hakala, M., Tuominen, I., Keränen, M., Tyystjärvi, T., and Tyystjärvi, E. (2005). Evidence for the role of the oxygen-evolving manganese complex in photoinhibition of Photosystem II. *Biochim. Biophys. Acta Bioenerg.* 1706, 68–80. doi: 10.1016/j.bbabi.2004.09.001
- Hanawa, H., Ishizaki, K., Nohira, K., Takagi, D., Shimakawa, G., Sejima, T., et al. (2017). Land plants drive photorespiration as higher electron-sink: comparative study of post-illumination transient O<sub>2</sub>-uptake rates from liverworts to angiosperms through ferns and gymnosperms. *Physiol. Plant* 161, 138–149. doi: 10.1111/ppl.12580
- Hanke, G. T., Satomi, Y., Shinmura, K., Takao, T., and Hase, T. (2011). A screen for potential ferredoxin electron transfer partners uncovers new, redox dependent

- interactions. *Biochim. Biophys. Acta Proteins Proteom.* 1814, 366–374. doi: 10.1016/j.bbapap.2010.09.011
- Harbinson, J., and Hedley, C. L. (1993). Changes in P-700 oxidation during the early stages of the induction of photosynthesis. *Plant Physiol.* 103, 649–660. doi: 10.1104/pp.103.2.649
- Havaux, M., and Eyletters, M. (1991). Is the in vivo photosystem I function resistant to photoinhibition? An answer from photoacoustic and far-red absorbance measurements in intact leaves. *Z. Naturforsch. C* 46, 1038–1044. doi: 10.1515/znc-1991-11-1218
- Hayashi, R., Shimakawa, G., Shaku, K., Shimizu, S., Akimoto, S., Yamamoto, H., et al. (2014). O<sub>2</sub>-dependent large electron flow functioned as an electron sink, replacing the steady-state electron flux in photosynthesis in the cyanobacterium *Synechocystis* sp. PCC 6803, but not in the cyanobacterium *Synechococcus* sp. PCC 7942. *Biosci. Biotechnol. Biochem.* 78, 384–393. doi: 10.1080/09168451.2014.882745
- Helman, Y., Tchernov, D., Reinhold, L., Shibata, M., Ogawa, T., Schwarz, R., et al. (2003). Genes encoding A-type flavoproteins are essential for photoreduction of O<sub>2</sub> in cyanobacteria. *Curr. Biol.* 13, 230–235. doi: 10.1016/S0960-9822(03)00046-0
- Huang, W., Zhang, S.-B., Zhang, J.-L., and Hu, H. (2015). Photoinhibition of photosystem I under high light in the shade-established tropical tree species *Psychotria rubra*. *Front. Plant Sci.* 6:801. doi: 10.3389/fpls.2015.00801
- Ilić, P., Pavlović, A., Kouřil, R., Alboresi, A., Morosinotto, T., Allahverdiyeva, Y., et al. (2017). Alternative electron transport mediated by flavodiiron proteins is operational in organisms from cyanobacteria up to gymnosperms. *New Phytol.* 214, 967–972. doi: 10.1111/nph.14536
- Inoue, K., Sakurai, H., and Hiyama, T. (1986). Photoinactivation sites of photosystem I in isolated chloroplasts. *Plant Cell Physiol.* 27, 961–968. doi: 10.1093/oxfordjournals.pcp.a077213
- Ishikawa, N., Takabayashi, A., Noguchi, K., Tazoe, Y., Yamamoto, H., von Caemmerer, S., et al. (2016). NDH-mediated cyclic electron flow around photosystem I is crucial for C<sub>4</sub> photosynthesis. *Plant Cell Physiol.* 57, 2020–2028. doi: 10.1093/pcp/pcw127
- Jensen, P. E., Bassi, R., Boekema, E. J., Dekker, J. P., Jansson, S., Leister, D., et al. (2007). Structure, function and regulation of plant photosystem I. *Biochim. Biophys. Acta Bioenerg.* 1767, 335–352. doi: 10.1016/j.bbapap.2007.03.004
- Jimbo, H., Yutthanasirikul, R., Nagano, T., Hisabori, T., Hihara, Y., and Nishiyama, Y. (2018). Oxidation of translation factor EF-Tu inhibits the repair of photosystem II. *Plant Physiol.* 176, 2691–2699. doi: 10.1104/pp.18.00037
- Joët, T., Genty, B., Josse, E.-M., Kuntz, M., Cournac, L., and Peltier, G. (2002). Involvement of a plastid terminal oxidase in plastoquinone oxidation as evidenced by expression of the *Arabidopsis thaliana* enzyme in tobacco. *J. Biol. Chem.* 277, 31623–31630. doi: 10.1074/jbc.M203538200
- Johnson, G. N., and Stepien, P. (2016). Plastid terminal oxidase as a route to improving plant stress tolerance: known knowns and known unknowns. *Plant Cell Physiol.* 57, 1387–1396. doi: 10.1093/pcp/pcw042
- Joliot, P., and Joliot, A. (1999). In vivo analysis of the electron transfer within photosystem I: are the two phylloquinones involved? *Biochemistry* 38, 11130–11136. doi: 10.1021/bi990857c
- Kanazawa, A., Ostendorf, E., Kohzuma, K., Hoh, D., Strand, D. D., Sato-Cruz, M., et al. (2017). Chloroplast ATP synthase modulation of the thylakoid proton motive force: implications for photosystem I and photosystem II photoprotection. *Front. Plant Sci.* 8:719. doi: 10.3389/fpls.2017.00719
- Klughammer, C., and Schreiber, U. (1994). An improved method, using saturating light pulses, for the determination of photosystem I quantum yield via P700<sup>+</sup>-absorbance changes at 830 nm. *Planta* 192, 261–268. doi: 10.1007/bf01089043
- Kok, B. (1956). On the inhibition of photosynthesis by intense light. *Biochim. Biophys. Acta* 21, 234–244. doi: 10.1016/0006-3002(56)90003-8
- Kono, M., and Terashima, I. (2016). Elucidation of photoprotective mechanisms of PSI against fluctuating light photoinhibition. *Plant Cell Physiol.* 57, 1405–1414. doi: 10.1093/pcp/pcw103
- Kono, M., Yamori, W., Suzuki, Y., and Terashima, I. (2017). Photoprotection of PSI by far-red light against the fluctuating light-induced photoinhibition in *Arabidopsis thaliana* and field-grown plants. *Plant Cell Physiol.* 58, 35–45. doi: 10.1093/pcp/pcw215
- Kozaki, A., and Takeba, G. (1996). Photorespiration protects C<sub>3</sub> plants from photooxidation. *Nature* 384, 557–560. doi: 10.1038/384557a0
- Kozuleva, M. A., and Ivanov, B. N. (2010). Evaluation of the participation of ferredoxin in oxygen reduction in the photosynthetic electron transport chain of isolated pea thylakoids. *Photosynth. Res.* 105, 51–61. doi: 10.1007/s1120-010-9565-5
- Kozuleva, M. A., Petrova, A. A., Mamedov, M. D., Semenov, A. Y., and Ivanov, B. N. (2014). O<sub>2</sub> reduction by photosystem I involves phylloquinone under steady-state illumination. *FEBS Lett.* 588, 4364–4368. doi: 10.1016/j.febslet.2014.10.003
- Krause, G. H., Köster, S., and Wong, S. C. (1985). Photoinhibition of photosynthesis under anaerobic conditions studied with leaves and chloroplasts of *Spinacia oleracea* L. *Planta* 165, 430–438. doi: 10.1007/bf00392242
- Kruk, J., Jemiola-Rzemińska, M., Burda, K., Schmid, G. H., and Strzałka, K. (2003). Scavenging of superoxide generated in photosystem I by plastoquinone and other prenyllipids in thylakoid membranes. *Biochemistry* 42, 8501–8505. doi: 10.1021/bi034036q
- Kudoh, H., and Sonoike, K. (2002). Irreversible damage to photosystem I by chilling in the light: cause of the degradation of chlorophyll after returning to normal growth temperature. *Planta* 215, 541–548. doi: 10.1007/s00425-002-0790-9
- Lennon, A. M., Prommeenate, P., and Nixon, P. J. (2003). Location, expression and orientation of the putative chlororespiratory enzymes, Ndh and IMMUTANS, in higher-plant plastids. *Planta* 218, 254–260. doi: 10.1007/s00425-003-1111-7
- Martin, M., Noarbe, D. M., Serrot, P. H., and Sabater, B. (2015). The rise of the photosynthetic rate when light intensity increases is delayed in *ndh* gene-defective tobacco at high but not at low CO<sub>2</sub> concentrations. *Front. Plant Sci.* 6:34. doi: 10.3389/fpls.2015.00034
- McDonald, A. E., Ivanov, A. G., Bode, R., Maxwell, D. P., Rodermerl, S. R., and Hüner, N. P. A. (2011). Flexibility in photosynthetic electron transport: the physiological role of plastoquinol terminal oxidase (PTOX). *Biochim. Biophys. Acta Bioenerg.* 1807, 954–967. doi: 10.1016/j.bbapap.2010.10.024
- Mehler, A. H. (1951). Studies on reactions of illuminated chloroplasts. *Arch. Biochem. Biophys.* 33, 65–77. doi: 10.1016/0003-9861(51)90082-3
- Melis, A. (1999). Photosystem-II damage and repair cycle in chloroplasts: what modulates the rate of photodamage in vivo? *Trends Plant Sci.* 4, 130–135. doi: 10.1016/S1360-1385(99)01387-4
- Mi, H., Klughammer, C., and Schreiber, U. (2000). Light-induced dynamic changes of NADPH fluorescence in *Synechocystis* PCC 6803 and its *ndhB*-defective mutant M55. *Plant Cell Physiol.* 41, 1129–1135. doi: 10.1093/pcp/pcd038
- Michelet, L., Roach, T., Fischer, B. B., Bedhomme, M., Lemaire, S. D., and Krieger-Liszky, A. (2013). Down-regulation of catalase activity allows transient accumulation of a hydrogen peroxide signal in *Chlamydomonas reinhardtii*. *Plant Cell Environ.* 36, 1204–1213. doi: 10.1111/pce.12053
- Mikko, T., and Steffen, G. (2018). Switching off photoprotection of photosystem I – a novel tool for gradual PSI photoinhibition. *Physiol. Plant.* 162, 156–161. doi: 10.1111/ppl.12618
- Misumi, M., Katoh, H., Tomo, T., and Sonoike, K. (2016). Relationship between photochemical quenching and non-photochemical quenching in six species of cyanobacteria reveals species difference in redox state and species commonality in energy dissipation. *Plant Cell Physiol.* 57, 1510–1517. doi: 10.1093/pcp/pcv185
- Misumi, M., and Sonoike, K. (2017). Characterization of the influence of chlororespiration on the regulation of photosynthesis in the glaucophyte *Cyanophora paradoxa*. *Sci. Rep.* 7:46100. doi: 10.1038/srep46100
- Mitchell, P. (1966). Chemiosmotic coupling in oxidative and photosynthetic phosphorylation. *Biol. Rev. Cambridge Philos. Soc.* 41, 445–501. doi: 10.1111/j.1469-185X.1966.tb01501.x
- Miyake, C., and Asada, K. (1992). Thylakoid-bound ascorbate peroxidase in spinach chloroplasts and photoreduction of its primary oxidation product monodehydroascorbate radicals in thylakoids. *Plant Cell Physiol.* 33, 541–553. doi: 10.1093/oxfordjournals.pcp.a078288
- Miyake, C., Michihata, F., and Asada, K. (1991). Scavenging of hydrogen peroxide in prokaryotic and eukaryotic algae: acquisition of ascorbate peroxidase during the evolution of cyanobacteria. *Plant Cell Physiol.* 32, 33–43. doi: 10.1093/oxfordjournals.pcp.a078050
- Miyake, C., Miyata, M., Shinzaki, Y., and Tomizawa, K. (2005). CO<sub>2</sub> response of cyclic electron flow around PSI (CEF-PSI) in tobacco leaves—relative electron fluxes through PSI and PSII determine the magnitude of non-photochemical

- quenching (NPQ) of Chl fluorescence. *Plant Cell Physiol.* 46, 629–637. doi: 10.1093/pcp/pci067
- Mosebach, L., Heilmann, C., Mutoh, R., Gäbelein, P., Steinbeck, J., Happe, T., et al. (2017). Association of Ferredoxin:NADP<sup>+</sup> oxidoreductase with the photosynthetic apparatus modulates electron transfer in *Chlamydomonas reinhardtii*. *Photosynth. Res.* 134, 291–306. doi: 10.1007/s11120-017-0408-5
- Mubarakshina, M., Khorobrykh, S., and Ivanov, B. (2006). Oxygen reduction in chloroplast thylakoids results in production of hydrogen peroxide inside the membrane. *Biochim. Biophys. Acta Bioenerg.* 1757, 1496–1503. doi: 10.1016/j.bbabi.2006.09.004
- Munekage, Y., Hojo, M., Meurer, J., Endo, T., Tasaka, M., and Shikanai, T. (2002). PGR5 is involved in cyclic electron flow around photosystem I and is essential for photoprotection in *Arabidopsis*. *Cell* 110, 361–371. doi: 10.1016/S0092-8674(02)00867-X
- Murata, N., Takahashi, S., Nishiyama, Y., and Allakhverdiev, S. I. (2007). Photoinhibition of photosystem II under environmental stress. *Biochim. Biophys. Acta Bioenerg.* 1767, 414–421. doi: 10.1016/j.bbabi.2006.11.019
- Nandha, B., Finazzi, G., Joliot, P., Hald, S., and Johnson, G. N. (2007). The role of PGR5 in the redox poisoning of photosynthetic electron transport. *Biochim. Biophys. Acta Bioenerg.* 1767, 1252–1259. doi: 10.1016/j.bbabi.2007.07.007
- Nathan, N., and Wolfgang, J. (2015). Structure and energy transfer in photosystems of oxygenic photosynthesis. *Annu. Rev. Biochem.* 84, 659–683. doi: 10.1146/annurev-biochem-092914-041942
- Neidhardt, J., Benemann, J. R., Zhang, L., and Melis, A. (1998). Photosystem-II repair and chloroplast recovery from irradiance stress: relationship between chronic photoinhibition, light-harvesting chlorophyll antenna size and photosynthetic productivity in *Dunaliella salina* (green algae). *Photosynth. Res.* 56, 175–184. doi: 10.1023/a:1006024827225
- Nishio, J. N., and Whitmarsh, J. (1993). Dissipation of the proton electrochemical potential in intact chloroplasts (II. The pH gradient monitored by cytochrome *f* reduction kinetics). *Plant Physiol.* 101, 89–96. doi: 10.1104/pp.101.1.89
- Noridomi, M., Nakamura, S., Tsuyama, M., Futamura, N., and Vladkova, R. (2017). Opposite domination of cyclic and pseudocyclic electron flows in short-illuminated dark-adapted leaves of angiosperms and gymnosperms. *Photosynth. Res.* 134, 149–164. doi: 10.1007/s11120-017-0419-2
- Nuijs, A. M., Shuvalov, V. A., van Gorkom, H. J., Plijter, J. J., and Duysens, L. N. M. (1986). Picosecond absorbance difference spectroscopy on the primary reactions and the antenna-excited states in Photosystem I particles. *Biochim. Biophys. Acta Bioenerg.* 850, 310–318. doi: 10.1016/0005-2728(86)90186-6
- Ogawa, T., Harada, T., Ozaki, H., and Sonoike, K. (2013). Disruption of the *ndhF1* gene affects Chl fluorescence through state transition in the cyanobacterium *Synechocystis* sp. PCC 6803, resulting in apparent high efficiency of photosynthesis. *Plant Cell Physiol.* 54, 1164–1171. doi: 10.1093/pcp/ptc068
- Ogren, W. L. (1984). Photorespiration: pathways, regulation, and modification. *Annu. Rev. Plant Physiol.* 35, 415–442. doi: 10.1146/annurev.pp.35.060184.002215
- Orr, D. J., Alcántara, A., Kapralov, M. V., Andralojc, P. J., Carmo-Silva, E., and Parry, M. A. J. (2016). Surveying Rubisco diversity and temperature response to improve crop photosynthetic efficiency. *Plant Physiol.* 172, 707–717. doi: 10.1104/pp.16.00750
- Paumann, M., Regelsberger, G., Obinger, C., and Peschek, G. A. (2005). The bioenergetic role of dioxygen and the terminal oxidase(s) in cyanobacteria. *Biochim. Biophys. Acta Bioenerg.* 1707, 231–253. doi: 10.1016/j.bbabi.2004.12.007
- Pils, D., Gregor, W., and Schmetterer, G. (1997). Evidence for *in vivo* activity of three distinct respiratory terminal oxidases in the cyanobacterium *Synechocystis* sp. strain PCC6803. *FEMS Microbiol. Lett.* 152, 83–88. doi: 10.1111/j.1574-6968.1997.tb10412.x
- Pospišil, P. (2016). Production of reactive oxygen species by photosystem II as a response to light and temperature stress. *Front. Plant Sci.* 7:1950. doi: 10.3389/fpls.2016.01950
- Powles, S. B., and Björkman, O. (1982). Photoinhibition of photosynthesis: effect on chlorophyll fluorescence at 77K in intact leaves and in chloroplast membranes of *Nerium oleander*. *Planta* 156, 97–107. doi: 10.1007/bf00395424
- Powles, S. B., Osmond, C. B., and Thorne, S. W. (1979). Photoinhibition of intact attached leaves of C3 plants illuminated in the absence of both carbon dioxide and of photorespiration. *Plant Physiol.* 64, 982–988. doi: 10.1104/pp.64.6.982
- Rademacher, N., Kern, R., Fujiwara, T., Mettler-Altmann, T., Miyagishima, S., Hagemann, M., et al. (2016). Photorespiratory glycolate oxidase is essential for the survival of the red alga *Cyanidioschyzon merolae* under ambient CO<sub>2</sub> conditions. *J. Exp. Bot.* 67, 3165–3175. doi: 10.1093/jxb/erw118
- Roach, T., Na, C. S., and Krieger-Liszka, A. (2015). High light-induced hydrogen peroxide production in *Chlamydomonas reinhardtii* is increased by high CO<sub>2</sub> availability. *Plant J.* 81, 759–766. doi: 10.1111/tpj.12768
- Roberty, S., Bailleul, B., Berne, N., Franck, F., and Cardol, P. (2014). PSI Mehler reaction is the main alternative photosynthetic electron pathway in *Symbiodinium* sp., symbiotic dinoflagellates of cnidarians. *New Phytol.* 204, 81–91. doi: 10.1111/nph.12903
- Romão, C. V., Vicente, J. B., Borges, P. T., Frazão, C., and Teixeira, M. (2016). The dual function of flavodiiron proteins: oxygen and/or nitric oxide reductases. *J. Biol. Inorg. Chem.* 21, 39–52. doi: 10.1007/s00775-015-1329-4
- Rosso, D., Ivanov, A. G., Fu, A., Geisler-Lee, J., Hendrickson, L., Geisler, M., et al. (2006). IMMUTANS does not act as a stress-induced safety valve in the protection of the photosynthetic apparatus of *Arabidopsis* during steady-state photosynthesis. *Plant Physiol.* 142, 574–585. doi: 10.1104/pp.106.08.5886
- Rumeau, D., Peltier, G., and Cournac, L. (2007). Chlororespiration and cyclic electron flow around PSI during photosynthesis and plant stress response. *Plant Cell Environ.* 30, 1041–1051. doi: 10.1111/j.1365-3040.2007.01675.x
- Rutherford, A. W., Osyczka, A., and Rappaport, F. (2012). Back-reactions, short-circuits, leaks and other energy wasteful reactions in biological electron transfer: redox tuning to survive life in O<sub>2</sub>. *FEBS Lett.* 586, 603–616. doi: 10.1016/j.febslet.2011.12.039
- Ruuska, S. A., Badger, M. R., Andrews, T. J., and von Caemmerer, S. (2000). Photosynthetic electron sinks in transgenic tobacco with reduced amounts of Rubisco: little evidence for significant Mehler reaction. *J. Exp. Bot.* 51, 357–368. doi: 10.1093/jexbot/51.suppl\_1.357
- Satoh, K. (1970). Mechanism of photoinactivation in photosynthetic systems II. The occurrence and properties of two different types of photoinactivation. *Plant Cell Physiol.* 11, 29–38. doi: 10.1093/oxfordjournals.pcp.a074493
- Sazanov, L. A., Burrows, P. A., and Nixon, P. J. (1998). The chloroplast Ndh complex mediates the dark reduction of the plastoquinone pool in response to heat stress in tobacco leaves. *FEBS Lett.* 429, 115–118. doi: 10.1016/S0014-5793(98)00573-0
- Scarpa, M., Stevanato, R., Viglino, P., and Rigo, A. (1983). Superoxide ion as active intermediate in the autoxidation of ascorbate by molecular oxygen. Effect of superoxide dismutase. *J. Biol. Chem.* 258, 6695–6697.
- Schöttler, M. A., and Tóth, S. Z. (2014). Photosynthetic complex stoichiometry dynamics in higher plants: environmental acclimation and photosynthetic flux control. *Front. Plant Sci.* 5:188. doi: 10.3389/fpls.2014.00188
- Schreiber, U., and Neubauer, C. (1990). O<sub>2</sub>-dependent electron flow, membrane energization and the mechanism of non-photochemical quenching of chlorophyll fluorescence. *Photosynth. Res.* 25, 279–293. doi: 10.1007/bf0033169
- Schuermans, R. M., van Alphen, P., Schuurmans, J. M., Matthijs, H. C. P., and Hellingwerf, K. J. (2015). Comparison of the photosynthetic yield of cyanobacteria and green algae: different methods give different answers. *PLoS One* 10:e0139061. doi: 10.1371/journal.pone.0139061
- Sejima, T., Hanawa, H., Shimakawa, G., Takagi, D., Suzuki, Y., Fukayama, H., et al. (2016). Post-illumination transient O<sub>2</sub>-uptake is driven by photorespiration in tobacco leaves. *Physiol. Plant.* 156, 227–238. doi: 10.1111/ppl.12388
- Sejima, T., Takagi, D., Fukayama, H., Makino, A., and Miyake, C. (2014). Repetitive short-pulse light mainly inactivates photosystem I in sunflower leaves. *Plant Cell Physiol.* 55, 1184–1193. doi: 10.1093/pcp/pcu061
- Semenov, A. Y., Vassiliev, I. R., van der Est, A., Mamedov, M. D., Zybailov, B., Shen, G., et al. (2000). Recruitment of a foreign quinone into the A1 site of photosystem I: altered kinetics of electron transfer in phyloquinone biosynthetic pathway mutants studied by time-resolved optical, EPR, and electrometric techniques. *J. Biol. Chem.* 275, 23429–23438. doi: 10.1074/jbc.M000508200
- Setif, P., Hervo, G., and Mathis, P. (1981). Flash-induced absorption changes in Photosystem I, Radical pair or triplet state formation? *Biochim. Biophys. Acta Bioenerg.* 638, 257–267. doi: 10.1016/0005-2728(81)90235-8
- Shaku, K., Shimakawa, G., Hashiguchi, M., and Miyake, C. (2016). Reduction-induced suppression of electron flow (RISE) in the photosynthetic electron

- transport system of *Synechococcus elongatus* PCC 7942. *Plant Cell Physiol.* 57, 1443–1453. doi: 10.1093/pcp/pcv198
- Shikanai, T., Endo, T., Hashimoto, T., Yamada, Y., Asada, K., and Yokota, A. (1998). Directed disruption of the tobacco *ndhB* gene impairs cyclic electron flow around photosystem I. *Proc. Natl. Acad. Sci. U.S.A.* 95, 9705–9709. doi: 10.1073/pnas.95.16.9705
- Shimakawa, G., Akimoto, S., Ueno, Y., Wada, A., Shaku, K., Takahashi, Y., et al. (2016a). Diversity in photosynthetic electron transport under [CO<sub>2</sub>]-limitation: the cyanobacterium *Synechococcus* sp. PCC 7002 and green alga *Chlamydomonas reinhardtii* drive an O<sub>2</sub>-dependent alternative electron flow and non-photochemical quenching of chlorophyll fluorescence during CO<sub>2</sub>-limited photosynthesis. *Photosynth. Res.* 130, 293–305. doi: 10.1007/s1120-016-0253-y
- Shimakawa, G., Shaku, K., and Miyake, C. (2016b). Oxidation of P700 in photosystem I is essential for the growth of cyanobacteria. *Plant Physiol.* 172, 1443–1450. doi: 10.1104/pp.16.01227
- Shimakawa, G., Hasunuma, T., Kondo, A., Matsuda, M., Makino, A., and Miyake, C. (2014). Respiration accumulates Calvin cycle intermediates for the rapid start of photosynthesis in *Synechocystis* sp. PCC 6803. *Biosci. Biotechnol. Biochem.* 78, 1997–2007. doi: 10.1080/09168451.2014.943648
- Shimakawa, G., Ishizaki, K., Tsukamoto, S., Tanaka, M., Sejima, T., and Miyake, C. (2017a). The liverwort, *Marchantia*, drives alternative electron flow using a flavodiiron protein to protect PSI. *Plant Physiol.* 173, 1636–1647. doi: 10.1104/pp.16.01038
- Shimakawa, G., Matsuda, Y., Nakajima, K., Tamoi, M., Shigeoka, S., and Miyake, C. (2017b). Diverse strategies of O<sub>2</sub> usage for preventing photo-oxidative damage under CO<sub>2</sub> limitation during algal photosynthesis. *Sci. Rep.* 7:41022. doi: 10.1038/srep41022
- Shimakawa, G., and Miyake, C. (2018a). Changing frequency of fluctuating light reveals the molecular mechanism for P700 oxidation in plant leaves. *Plant Direct* 2:e00073. doi: 10.1002/pld3.73
- Shimakawa, G., and Miyake, C. (2018b). Respiratory terminal oxidases alleviate photo-oxidative damage in photosystem I during repetitive short-pulse illumination in *Synechocystis* sp. PCC 6803. *Photosynth. Res.* 137, 241–250. doi: 10.1007/s1120-018-0495-y
- Shimakawa, G., Murakami, A., Niwa, K., Matsuda, Y., Wada, A., and Miyake, C. (2018a). Comparative analysis of strategies to prepare electron sinks in aquatic photoautotrophs. *Photosynth. Res.* doi: 10.1007/s1120-018-0522-z [Epub ahead of print].
- Shimakawa, G., Shaku, K., and Miyake, C. (2018b). Reduction-induced suppression of electron flow (RISE) is relieved by non-ATP-consuming electron flow in *Synechococcus elongatus* PCC 7942. *Front. Microbiol.* 9:886. doi: 10.3389/fmicb.2018.00886
- Shimakawa, G., Shaku, K., Nishi, A., Hayashi, R., Yamamoto, H., Sakamoto, K., et al. (2015). FLAVODIIRON2 and FLAVODIIRON4 proteins mediate an oxygen-dependent alternative electron flow in *Synechocystis* sp. PCC 6803 under CO<sub>2</sub>-limited conditions. *Plant Physiol.* 167, 472–480. doi: 10.1104/pp.114.249987
- Shirao, M., Kuroki, S., Kaneko, K., Kinjo, Y., Tsuyama, M., Förster, B., et al. (2013). Gymnosperms have increased capacity for electron leakage to oxygen (Mehler and PTOX reactions) in photosynthesis compared with angiosperms. *Plant Cell Physiol.* 54, 1152–1163. doi: 10.1093/pcp/pt066
- Sonoike, K. (1996). Degradation of *psaB* gene product, the reaction center subunit of photosystem I, is caused during photoinhibition of photosystem I: possible involvement of active oxygen species. *Plant Sci.* 115, 157–164. doi: 10.1016/0168-9452(96)04341-5
- Sonoike, K. (2011). Photoinhibition of photosystem I. *Physiol. Plant.* 142, 56–64. doi: 10.1111/j.1399-3054.2010.01437.x
- Sonoike, K., Kamo, M., Hihara, Y., Hiyama, T., and Enami, I. (1997). The mechanism of the degradation of *PSAB* gene product, one of the photosynthetic reaction center subunits of Photosystem I, upon photoinhibition. *Photosynth. Res.* 53, 55–63. doi: 10.1023/a:1005852330671
- Stiehl, H. H., and Witt, H. T. (1969). Quantitative treatment of the function of plastoquinone in photosynthesis. *Z. Naturforsch. B* 24, 1588–1598. doi: 10.1515/znb-1969-1219
- Subramanyam, R., David, J., Alain, G., Marc, B., and Robert, C. (2005). Protective effect of active oxygen scavengers on protein degradation and photochemical function in photosystem I submembrane fractions during light stress. *FEBS J.* 272, 892–902. doi: 10.1111/j.1742-4658.2004.04512.x
- Sundby, C., Chow, W. S., and Anderson, J. M. (1993). Effects on photosystem II function, photoinhibition, and plant performance of the spontaneous mutation of serine-264 in the photosystem II reaction center D1 protein in triazine-resistant *Brassica napus* L. *Plant Physiol.* 103, 105–113. doi: 10.1104/pp.103.1.105
- Takagi, D., Amako, K., Hashiguchi, M., Fukaki, H., Ishizaki, K., Goh, T., et al. (2017a). Chloroplastic ATP synthase builds up a proton motive force preventing production of reactive oxygen species in photosystem I. *Plant J.* 91, 306–324. doi: 10.1111/tpj.13566
- Takagi, D., Ishizaki, K., Hanawa, H., Mabuchi, T., Shimakawa, G., Yamamoto, H., et al. (2017b). Diversity of strategies for escaping reactive oxygen species production within photosystem I among land plants: P700 oxidation system is prerequisite for alleviating photoinhibition in photosystem I. *Physiol. Plant.* 161, 56–74. doi: 10.1111/ppl.12562
- Takagi, D., Hashiguchi, M., Sejima, T., Makino, A., and Miyake, C. (2016a). Photorespiration provides the chance of cyclic electron flow to operate for the redox-regulation of P700 in photosynthetic electron transport system of sunflower leaves. *Photosynth. Res.* 129, 279–290. doi: 10.1007/s1120-016-0267-5
- Takagi, D., Takumi, S., Hashiguchi, M., Sejima, T., and Miyake, C. (2016b). Superoxide and singlet oxygen produced within the thylakoid membranes both cause photosystem I photoinhibition. *Plant Physiol.* 171, 1626–1634. doi: 10.1104/pp.16.00246
- Takagi, D., and Miyake, C. (2018). PROTON GRADIENT REGULATION 5 supports linear electron flow to oxidize photosystem I. *Physiol. Plant.* 164, 337–348. doi: 10.1111/ppl.12723
- Takahashi, M., and Asada, K. (1988). Superoxide production in aprotic interior of chloroplast thylakoids. *Arch. Biochem. Biophys.* 267, 714–722. doi: 10.1016/0003-9861(88)90080-X
- Takahashi, S., Bauwe, H., and Badger, M. (2007). Impairment of the photorespiratory pathway accelerates photoinhibition of photosystem II by suppression of repair but not acceleration of damage processes in *Arabidopsis*. *Plant Physiol.* 144, 487–494. doi: 10.1104/pp.107.097253
- Takizawa, K., Kanazawa, A., and Kramer, D. M. (2008). Depletion of stromal Pi induces high ‘energy-dependent’ antenna exciton quenching (qE) by decreasing proton conductivity at CFo-CF1 ATP synthase. *Plant Cell Environ.* 31, 235–243. doi: 10.1111/j.1365-3040.2007.01753.x
- Tcherkez, G. (2016). The mechanism of Rubisco-catalysed oxygenation. *Plant Cell Environ.* 39, 983–997. doi: 10.1111/pce.12629
- Terashima, I., Funayama, S., and Sonoike, K. (1994). The site of photoinhibition in leaves of *Cucumis sativus* L. at low temperatures is photosystem I, not photosystem II. *Planta* 193, 300–306. doi: 10.1007/bf00192544
- Terashima, I., Noguchi, K., Itoh-Nemoto, T., Park, Y.-M., Kuhn, A., and Tanaka, K. (1998). The cause of PSI photoinhibition at low temperatures in leaves of *Cucumis sativus*, a chilling-sensitive plant. *Physiol. Plant.* 103, 295–303. doi: 10.1034/j.1399-3054.1998.1030301.x
- Tjus, S. E., Møller, B. L., and Scheller, H. V. (1999). Photoinhibition of Photosystem I damages both reaction centre proteins PSI-A and PSI-B and acceptor-side located small Photosystem I polypeptides. *Photosynth. Res.* 60, 75–86. doi: 10.1023/a:1006283618695
- Trissl, H.-W. (1997). Determination of the quenching efficiency of the oxidized primary donor of Photosystem I, P700<sup>+</sup>: implications for the trapping mechanism. *Photosynth. Res.* 54, 237–240. doi: 10.1023/a:1005981016835
- Trubitsin, B. V., Mamedov, M. D., Vitukhnovskaya, L. A., Semenov, A. Y., and Tikhonov, A. N. (2003). EPR study of light-induced regulation of photosynthetic electron transport in *Synechocystis* sp. strain PCC 6803. *FEBS Lett.* 544, 15–20. doi: 10.1016/S0014-5793(03)00429-0
- Tyystjärvi, E. (2008). Photoinhibition of Photosystem II and photodamage of the oxygen evolving manganese cluster. *Coord. Chem. Rev.* 252, 361–376. doi: 10.1016/j.ccr.2007.08.021
- Ueno, Y., Shimakawa, G., Miyake, C., and Akimoto, S. (2018). Light-harvesting strategy during CO<sub>2</sub>-dependent photosynthesis in the green alga *Chlamydomonas reinhardtii*. *J. Phys. Chem. Lett.* 9, 1028–1033. doi: 10.1021/acs.jpclett.7b03404

- Van Breusegem, F., Vranová, E., Dat, J. F., and Inzé, D. (2001). The role of active oxygen species in plant signal transduction. *Plant Sci.* 161, 405–414. doi: 10.1016/S0168-9452(01)00452-6
- Vicente, J. B., Gomes, C. M., Wasserfallen, A., and Teixeira, M. (2002). Module fusion in an A-type flavoprotein from the cyanobacterium *Synechocystis* condenses a multiple-component pathway in a single polypeptide chain. *Biochem. Biophys. Res. Commun.* 294, 82–87. doi: 10.1016/S0006-291X(02)00434-5
- Wada, S., Suzuki, Y., Takagi, D., Miyake, C., and Makino, A. (2018). Effects of genetic manipulation of the activity of photorespiration on the redox state of photosystem I and its robustness against excess light stress under CO<sub>2</sub>-limited conditions in rice. *Photosynth. Res.* doi: 10.1007/s11120-018-0515-y [Epub ahead of print].
- Wardman, P. (1990). Bioreductive activation of quinones: redox properties and thiol reactivity. *Free Radical Res. Commun.* 8, 219–229. doi: 10.3109/10715769009053355
- Weger, H. G., Herzig, R., Falkowski, P. G., and Turpin, D. H. (1989). Respiratory losses in the light in a marine diatom: measurements by short-term mass spectrometry. *Limnol. Oceanogr.* 34, 1153–1161. doi: 10.4319/lo.1989.34.7.1153
- Wiese, C., Shi, L., and Heber, U. (1998). Oxygen reduction in the Mehler reaction is insufficient to protect photosystems I and II of leaves against photoinactivation. *Physiol. Plant.* 102, 437–446. doi: 10.1034/j.1399-3054.1998.1020312.x
- Yamamoto, H., Takahashi, S., Badger, M. R., and Shikanai, T. (2016). Artificial remodelling of alternative electron flow by flavodiiron proteins in *Arabidopsis*. *Nat. Plants* 2:16012. doi: 10.1038/nplants.2016.12
- Yamori, W., Makino, A., and Shikanai, T. (2016). A physiological role of cyclic electron transport around photosystem I in sustaining photosynthesis under fluctuating light in rice. *Sci. Rep.* 6:20147. doi: 10.1038/srep20147
- Yamori, W., and Shikanai, T. (2016). Physiological functions of cyclic electron transport around photosystem I in sustaining photosynthesis and plant growth. *Annu. Rev. Plant Biol.* 67, 81–106. doi: 10.1146/annurev-arplant-043015-112002
- Yokono, M., and Akimoto, S. (2018). Energy transfer and distribution in photosystem super/megacomplexes of plants. *Curr. Opin. Biotechnol.* 54, 50–56. doi: 10.1016/j.copbio.2018.01.001
- Yokota, A., and Kitaoka, S. (1987). Rates of glycolate synthesis and metabolism during photosynthesis of *Euglena* and microalgae grown on low CO<sub>2</sub>. *Planta* 170, 181–189. doi: 10.1007/bf00397886
- Youngman, R. J., and Elstner, E. F. (1981). Oxygen species in paraquat toxicity: the crypto-OH radical. *FEBS Lett.* 129, 265–268. doi: 10.1016/0014-5793(81)80180-9
- Yu, Q., Feilke, K., Krieger-Liszka, A., and Beyer, P. (2014). Functional and molecular characterization of plastid terminal oxidase from rice (*Oryza sativa*). *Biochim. Biophys. Acta Bioenerg.* 1837, 1284–1292. doi: 10.1016/j.bbabi.2014.04.007
- Zivcak, M., Brestic, M., Kunderlikova, K., Olsovska, K., and Allakhverdiev, S. I. (2015a). Effect of photosystem I inactivation on chlorophyll *a* fluorescence induction in wheat leaves: does activity of photosystem I play any role in OJIP rise? *J. Photochem. Photobiol. B* 152, 318–324. doi: 10.1016/j.jphotobiol.2015.08.024
- Zivcak, M., Brestic, M., Kunderlikova, K., Sytar, O., and Allakhverdiev, S. I. (2015b). Repetitive light pulse-induced photoinhibition of photosystem I severely affects CO<sub>2</sub> assimilation and photoprotection in wheat leaves. *Photosynth. Res.* 126, 449–463. doi: 10.1007/s11120-015-0121-1

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer YS declared a past co-authorship with the authors GS and CM to the handling Editor.

Copyright © 2018 Shimakawa and Miyake. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.