



Oxidative Stress and Neonatal Respiratory Extracorporeal Membrane Oxygenation

Genny Raffaeli^{1,2}, Stefano Ghirardello¹, Sofia Passera^{1,2}, Fabio Mosca^{1,2} and Giacomo Cavallaro^{1*}

¹ NICU, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy, ² Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy

OPEN ACCESS

Edited by:

Murugesan Velayutham,
University of Pittsburgh, United States

Reviewed by:

Hitesh Singh Sandhu,
The University of Tennessee Health
Science Center (UTHSC),
United States
Ana Denicola,
Universidad de la República, Uruguay

*Correspondence:

Giacomo Cavallaro
giacomo.cavallaro@mangiagalli.it

Specialty section:

This article was submitted to
Oxidant Physiology,
a section of the journal
Frontiers in Physiology

Received: 17 September 2018

Accepted: 19 November 2018

Published: 04 December 2018

Citation:

Raffaeli G, Ghirardello S,
Passera S, Mosca F and Cavallaro G
(2018) Oxidative Stress and Neonatal
Respiratory Extracorporeal Membrane
Oxygenation. *Front. Physiol.* 9:1739.
doi: 10.3389/fphys.2018.01739

Oxidative stress is a frequent condition in critically ill patients, especially if exposed to extracorporeal circulation, and it is associated with worse outcomes and increased mortality. The inflammation triggered by the contact of blood with a non-endogenous surface, the use of high volumes of packed red blood cells and platelets transfusion, the risk of hyperoxia and the impairment of antioxidation systems contribute to the increase of reactive oxygen species and the imbalance of the redox system. This is responsible for the increased production of superoxide anion, hydrogen peroxide, hydroxyl radicals, and peroxynitrite resulting in increased lipid peroxidation, protein oxidation, and DNA damage. The understanding of the pathophysiologic mechanisms leading to redox imbalance would pave the way for the future development of preventive approaches. This review provides an overview of the clinical impact of the oxidative stress during neonatal extracorporeal support and concludes with a brief perspective on the current antioxidant strategies, with the aim to focus on the potential oxidative stress-mediated cell damage that has been implicated in both short and long-term outcomes.

Keywords: antioxidants, cardiopulmonary bypass, CRRT, ECMO, extracorporeal circulation, oxidative stress

Abbreviations: AKI, acute kidney injury; ARDS, acute respiratory distress syndrome; ATP, adenosine triphosphate; CAT, catalase; CPB, cardiopulmonary bypass; CRRT, Continuous Renal Replacement Therapies; Cys-S⁻, cysteine thiolate anion; Cys-SO₂H, cysteine sulfinic acid; Cys-SO₃H, cysteine sulfonic acid; Cys-SOH, cysteine sulfenic acid; d-ROMs, reactive oxygen metabolism and derivatives; ECC, extracorporeal circulation; ECMO, extracorporeal membrane oxygenation; ELSO, extracorporeal life support organization; Fe²⁺, ferrous ions; Fe³⁺, ferric ions; FO, fluid overload; GPx, glutathione peroxidases; Grx, glutaredoxin; GSH, glutathione; H₂O₂, hydrogen peroxide; hMSCs, hypoxia-preconditioned mesenchymal stem cells; IFN β , interferon β ; IL, interleukin; MAS, meconium aspiration syndrome; MDA, malondialdehyde; MOF, multi-organ failure; NAC, N-acetylcysteine; NADP⁺, nicotinamide adenine dinucleotide phosphate oxidized form; NADPH, nicotinamide adenine dinucleotide phosphate reduced form; NEC, necrotizing enterocolitis; NO, Nitric oxide; NOS, nitric oxide synthase; NOX, NADPH oxidase; O₂, oxygen; O₂⁻, superoxide anion; OH, hydroxyl radicals; ONO₂⁻, peroxynitrite; PCs, platelet concentrates; PFHb, plasma free hemoglobin; PPH, persistent pulmonary hypertension; PRBC, packed red blood cells; PRx, peroxiredoxins; PUFA, polyunsaturated fatty acid; pVWF, plasmatic Von Willebrand factor; RBCs, red blood cells; RDS, respiratory distress syndrome; RNS, reactive nitrogen species; ROS, reactive oxygen species; Se, selenium; SIRS, systemic inflammatory response syndrome; SOD, superoxide dismutases; TAC, total antioxidant capacity; TF, tissue factor; TNF α , tumor necrosis factor α ; TR, thioredoxin reductase; Trx, thioredoxin; VA-ECMO, venous-arterial ECMO; VV-ECMO, venous-venous ECMO.

INTRODUCTION

Extracorporeal membrane oxygenation is life support for reversible heart and/or lung failure refractory to standard treatment, and it is used as bridge-to-recovery, bridge-to-bridge, bridge-to-transplant, or bridge-to-decision (Brogan et al., 2017). This technique was developed by the CPB technology, and it is based on a partial/full bypass of the native cardiac and/or pulmonary function. Neonatal respiratory ECMO was introduced by Robert Bartlett during the 1970s for severe hypoxemic respiratory failure secondary to MAS, PPH and RDS (Bartlett et al., 1976). Despite the progress in neonatal care following the introduction of new therapies and ventilatory strategies, ECMO remains life-saving support for about 600–800 neonates each year worldwide (Thiagarajan et al., 2017). Across the world, from 1989 to 2017, about 27,238 newborns needed an ECMO procedure for respiratory failure, 7,592 for cardiac problems, 1,694 for extracorporeal cardiopulmonary resuscitation; the overall survival rate is 83, 64, and 66%, respectively (ELSO). Although the overall survival is high, there are consistent differences in the incidence rates between the diseases. Since 1979 the survival of ECMO patients with congenital diaphragmatic hernia and sepsis is stable at around 51 and 48%, respectively. In contrast, PPH, RDS, and MAS have a significantly higher survival of 73, 76, and 92%, each. Furthermore, cardiomyopathy and myocarditis have better survival (65 and 50%, respectively), compared to congenital heart disease and cardiogenic shock (44 and 47%, respectively) (ELSO). Nevertheless, the mortality of infants and children with congenital heart disease has reduced in recent decades as a result of the improvement of CPB techniques and myocardial protective strategies (Gilboa et al., 2010; Cheng et al., 2014; Erikssen et al., 2015). Nowadays, although the clinical management of ECMO patients has remarkably progressed over the years, the rate of complications (hemostatic disorders, brain damage, AKI, sepsis) still have a high impact on patients' outcome (ELSO).

Oxidative stress plays a critical role in the genesis of tissue damage during an ECMO procedure, and newborns are particularly prone to the toxic effects of free radicals, due to the immaturity of their antioxidant systems resulting in lowered detoxification capacity (Comporti et al., 2004; Castellheim et al., 2005; Saugstad, 2005; Mishra et al., 2008; Signorini et al., 2008; Afolayan et al., 2012, 2015; Annagür et al., 2015; Asci et al., 2015; Mokra et al., 2015; Aras-López et al., 2016; Mikolka et al., 2016; Mátyás and Zaharie, 2018; Moore et al., 2018; Perrone et al., 2018). The inflammatory response following the introduction of ECC generates a high amount of ROS contemporarily reducing endogenous antioxidants (Descamps-Latscha, 2001; Goodyear-Bruch and Pierce, 2002; Karu et al., 2010). Various studies associated the exposure to extracorporeal circuits (CPB, ECMO, and CRRT) with the increased oxidative stress that may negatively impact on morbidity and mortality (McDonald et al., 2014a). Redox imbalance is mainly induced in the early stage of ECMO, especially after the CPB, while with prolonged exposure (beyond 24 h) the TAC gradually restores (Wan et al., 1997; Babior, 2000; Desborough, 2000; Clermont et al., 2002; Levy and Tanaka, 2003; Kouchoukos et al., 2012; Chen Q. et al., 2014;

Zakkar et al., 2015a,b; Fudulu and Angelini, 2016; Petersen et al., 2018). The modulation of oxidative stress would seem to be of great clinical importance. Indeed, its manipulation through the use of exogenous antioxidants, cardioplegia during CPB, blood priming strategies, or miniaturization of the extracorporeal circuit could mitigate the ROS production (Fudulu and Angelini, 2016). However, data are scant, especially for neonatal ECMO, and are mainly related to pre-clinical evidence. This review focuses on the mechanisms of oxidative stress during neonatal ECMO, involved in the pathophysiology of both short and long-term disorders as MOF, hemostatic derangements up to neurodegenerative, cardiovascular and cancer disease in adults. Lastly, a brief overview of currently available antioxidant strategies is provided (Halliwell and Gutteridge, 2015).


REDOX BIOLOGY AND OXIDATIVE STRESS

Energy is produced by mitochondrial metabolism through glucose, amino acids and lipids combustion in the presence of O_2 (Semenza, 2009; Vento, 2014). Approximately 98% of O_2 is reduced to H_2O by the conventional combustion process, while only 2% of remaining O_2 loses its electrons with the rear O_2^- and ROS production (Maltepe and Saugstad, 2009; Vento, 2014).

The “old definition” of oxidative stress was referred to conditions in which the balance between ROS production and endogenous antioxidants break down resulting in tissue and cellular injury through lipid peroxidation, protein oxidation, and DNA damage (Schieber and Chandel, 2014). Currently, the “new definition” is much more complex, referring to an imbalance between oxidants and antioxidants in favor of the oxidants, that cause an arrest of redox signaling and/or molecular damage (Sies, 2018). More recently the term “reductive stress” has been introduced, but its exact definition is still unclear (Gutteridge and Halliwell, 2018). It would seem to be due to high levels of reduced GSH and NADPH, unbalancing the redox system to the opposite oxidative extreme, which has shown to play an essential role in the pathogenesis of dilated cardiomyopathy in cases of lamin-associated muscular dystrophy (Margaritelis et al., 2014; Dialynas et al., 2015; Stevens et al., 2017; Bellezza et al., 2018).

The O_2^- , H_2O_2 , OH^\bullet , and ONO_2^- are the most common ROS, produced by aerobic metabolism (Schieber and Chandel, 2014). The intracellular O_2^- arises from the one-electron reduction of molecular O_2 that is primarily produced by the oxidation of NADPH by NOX enzymes or by electron loss from aerobic respiration in mitochondria (Lambeth, 2004; Brand, 2010; Sahoo et al., 2016). Superoxide anion is rapidly converted by SOD into H_2O_2 , and its storage is prevented by SODs, which are mainly located in the cytosol and mitochondria. Conversely, the increase in H_2O_2 triggers autophagy and cell death (Figure 1; Fridovich, 1997; Chen Y. et al., 2009).

Cysteine residues within proteins are oxidized by H_2O_2 to sulfenic form (Cys-SOH) and reduced to its basic thiolate anion (Cys-S⁻) form by the disulfide reductases Trx and Grx (Winterbourn and Hampton, 2008). The excess of H_2O_2 further oxidizes thiolate anions to sulfinic (Cys-SO₂H) or sulfonic



(Cys-SO₃H) classes. The sulfinic and sulfonic oxidation is irreversible and could lead to permanent protein damage that is prevented by PRx, GPx and CAT (Figures 1, 2; Winterbourn and Hampton, 2008; Finkel, 2012). In contrast, the OH produced by Fenton reaction in the presence of Fe²⁺ has a detrimental effect on lipids, proteins, and DNA (Figure 1; Dizdaroglu and Jaruga, 2012).

Nitric oxide is one of the most important and ubiquitous intracellular messengers that mediates several basic physiological processes including neurotransmission, vasodilatation, angiogenesis and host cell defense (MacMicking et al., 1997; Egginton, 2009; Bogdan, 2015; Philippu, 2016; Kraehling and Sessa, 2017). NO is synthesized from L-arginine by several forms of NOS, and its reaction with O_2^- produces ONO_2^- which alters mitochondrial function and causes apoptosis (Figure 1; Stuehr, 1999; Pacher et al., 2007; Radi, 2013).

Lipid Peroxidation

Lipid peroxidation, caused by increased ROS and RNS, is linked to many acute and chronic diseases of adulthood such as MOF, hemorrhagic and septic shock, ARDS, acute lung injury, cardiovascular pathology requiring surgery, aging, atherosclerosis, Parkinson, Alzheimer, diabetes mellitus, cataracts and rheumatoid arthritis, and several diseases of the neonatal

period (Takeda et al., 1984; Hammond and Hess, 1985; Bertrand et al., 1989; Richard et al., 1990; Quinlan et al., 1996; Carpenter et al., 1998; Lamb et al., 1999; Motoyama et al., 2003; Comporti et al., 2004; Quinlan et al., 2004; Fläring et al., 2005; Laplace et al., 2005; Niki et al., 2005; Saugstad, 2005; Huet et al., 2007; Signorini et al., 2008; Giustarini et al., 2009; Briones and Touyz, 2010; Giacco and Brownlee, 2010; Khaper et al., 2010; Romano et al., 2010; Barrera et al., 2018; Maiorino et al., 2018; Mátyás and Zaharie, 2018; Moore et al., 2018; Orban and Singer, 2018; Perrone et al., 2018; Romano et al., 2018).

Lipid peroxidation is the oxidative decay of Puffs, which involves the abstraction of hydrogen from a carbon molecule with O_2 insertion and consequent production of lipid peroxy radicals and hydroperoxides (Tappel, 1970; Yin et al., 2011; Al-Dalaen and Al-Qtaitat, 2014). Hydroperoxides are very unstable and easily decompose into MDAs and 4-hydroxy-2-nonenal, while peroxidation of arachidonic acid produces isoprostanes (Montuschi et al., 2004; Lakshminrusimha et al., 2006; Al-Dalaen and Al-Qtaitat, 2014). The response of the cell membrane to lipid peroxidation depends on the degree of peroxidation achieved. If lipid peroxidation rate is low, the damage is repaired by anti-oxidant defense systems. On the other hand, if the lipid peroxidation rate is high, the tissue self-repairing capacity is lost,




FIGURE 2 | Thioredoxin and Peroxiredoxin system. Reduced Trx-SH catalyzes the reduction of disulfide (s-s) bridges within oxidized cell proteins. This process causes the oxidation of Trx-SH to Trx-disulfide which is subsequently reduced by TR at the expense of NADPH. The TR in its catalytic Trx reduction process uses Se as a cofactor. Reduced PRx-SH promotes the separation of two H₂O molecules from H₂O₂ through a “two-step mechanism.” The peroxidic cysteine sulfhydryl group of a Prx subunit is oxidized to sulfenic acid (-SOH). Subsequently, sulfenic acid condenses with the reduced cysteine group present in the other subunit to form a disulfide bond between the units. This two-step process releases two H₂O molecules from H₂O₂ with the oxidation of PRx-SH to PRx-SOH to PRx-disulfide. The oxidized PRx is in turn reduced by the Trx-SH. H₂O₂, hydrogen peroxide; NADP⁺, nicotinamide adenine dinucleotide phosphate oxidated form; NADPH, nicotinamide adenine dinucleotide phosphate reduced form; PRx, peroxiredoxin; TR, thioredoxin reductase; Trx, thioredoxin.

and apoptosis or programmed cell death is induced (Chaudhary et al., 2010; Ayala et al., 2014).

Oxidative Protein Damage

Protein oxidation damage does affect not only the cell membranes but also the enzyme system because it is exclusively constituted by proteins (Goodyear-Bruch and Pierce, 2002). The amino acid fragmentation, the following generation of cross-links between proteins and the oxidation of the body of the protein molecule

with loss of specific (contractile, enzymatic, and transport) function is caused by ROS and RNS (Dalle-Donne et al., 2003a,b; Giustarini et al., 2009). Protein Carbonyl groups (aldehydes and ketones) are produced generally by the direct oxidation of proline, arginine, lysine, and threonine residues, or by oxidative cleavage of the protein backbone, and indirectly by lipid peroxidation or reaction of reducing sugars or their oxidation products (Dalle-Donne et al., 2003a, 2006). The damaged proteins, in turn, contribute to many disorders and diseases

(Dalle-Donne et al., 2003a). Oxidized or aldehyde-modified proteins would be perceived as “foreign proteins” by the immune system, with the production of self-antibodies in rheumatoid arthritis, lupus, and scleroderma in adults (Mantle et al., 1999; Dimon-Gadal et al., 2000; Renke et al., 2000; Dalle-Donne et al., 2003a; Halliwell and Gutteridge, 2015). Moreover, always in adults, the increase of oxidized proteins has been associated with multiple diseases, like cancer, heart failure, pre-eclampsia, ARDS, Parkinson, Alzheimer, and sepsis (Floor and Wetzel, 1998; Lenz et al., 1999; Winterbourn et al., 2000; Butterfield et al., 2001; Zusterzeel et al., 2001; Abu-Zidan et al., 2002; Castegna et al., 2002). In the neonatal period, the oxidation of proteins plays a key role in some diseases such as intraventricular hemorrhage, retinopathy of prematurity, bronchopulmonary dysplasia, NEC (Gladstone and Levine, 1994; Ramsay et al., 2001; Perrone et al., 2009, 2010, 2014, 2018; Couroucli, 2017).

Oxidative DNA Damage

Oxidative DNA damage is generated by high OH[•] concentrations (Cooke et al., 2003). ROS can damage both purines and pyrimidines in DNA and in the free cell nucleotide pool and generates 8-hydroxy-2'-deoxyguanosine that can be used as a biomarker of the injury (Suliman et al., 2003; Bayr, 2005; Li et al., 2014). Despite the damage, the cell continues the replication process, replicating the damage and inducing the new cells to apoptosis (Roos and Kaina, 2006). Furthermore, oxidative DNA damage would be associated with other diseases such as sepsis, carcinogenesis, neurodegenerative diseases, cardiovascular diseases, and aging (Cooke et al., 2003; Niki et al., 2005; Giustarini et al., 2009; Halliwell and Gutteridge, 2015; Bahar et al., 2018; Gutteridge and Halliwell, 2018; Klaunig and Je, 2018).

OXIDATIVE STRESS AND ECMO

To date, evidence of oxidative stress during ECMO are limited and are mainly related to pre-clinical studies (Möller et al., 1993; Trittenwein et al., 1999). Möller et al. (1993) observed a reduction of the antioxidant capacity of SOD and GPx on a lamb ECMO model, while increased lipid peroxidation was demonstrated on a rabbit model of ECMO (Trittenwein et al., 1999). One study on pediatric patients described two different oxidative stress patterns during ECC; in non-survivors, lipid peroxidation augmented during the first 96 h of ECMO whereas it reached a plateau after 24 h in the survivor's group (Hirthler et al., 1992).

No data are currently available on critically ill neonates undergoing respiratory ECMO. Despite this gap of knowledge, previous evidence, mainly derived from studies on CPB and critically ill patients, suggested that oxidative stress is a relevant contributor to morbidity and mortality (Möller et al., 1993; Trittenwein et al., 1999; Brix-Christensen et al., 2001, 2002; Goodyear-Bruch and Pierce, 2002; Levy and Tanaka, 2003). ECMO shares with both CPB and critical illness common potential triggers for increased ROS generation and redox unbalance: exposure to the extracorporeal circuit and related SIRS, transfusion burden, hyperoxia, CRRT, hemolysis, and

sequestration of antioxidants into the ECMO circuit (McDonald et al., 2014a). We describe hereafter the possible role of these triggers in the pathogenesis of oxidative stress-related morbidity during ECC.

ECMO and Inflammation

A SIRS-like syndrome is typically observed from the first hours after starting an ECMO procedure, as a consequence of the blood exposure to the exogenous surface of the extracorporeal device (Davies and Hagen, 1997). Despite the use of biocompatible equipment, the interaction between biological fluids and the circuit activates both the coagulation cascade and the host inflammatory response (Tayama et al., 2000; Palatianos et al., 2003; Wiesenack et al., 2004; Thiara et al., 2010). The humoral and cell-mediated immune response triggers the release of pro-inflammatory cytokines (IL-1 β , TNF α , and IFN β), leading to endothelial damage, disrupted microcirculation and multi-organ dysfunction, in case of suboptimal compensatory anti-inflammatory response (Hocker et al., 1991; Hirthler et al., 1992; Depuydt et al., 1993; Plötz et al., 1993; Fortenberry et al., 1996; Adrian et al., 1998; Graulich et al., 2002; Golej et al., 2003; Mildner et al., 2005; Adib-Conquy and Cavaillon, 2009; Mcilwain et al., 2010; He et al., 2014; Shi et al., 2014; Rungatscher et al., 2015; Wang et al., 2015). The redox biology regulates the homeostasis of innate and adaptive immune response through ROS production. Many concurrent factors (hyperoxia, blood products, sepsis, mechanical ventilation) contribute to the increased ROS generation, thus promoting an overwhelming release of pro-inflammatory cytokines (IL-1 β , TNF α , and IFN β) and eventually the SIRS-like (Wiesenack et al., 2004; Cruz et al., 2007; Chen and Nuñez, 2010; Iwasaki and Medzhitov, 2010; Bulua et al., 2011; West et al., 2011; Mittal et al., 2014).

Furthermore, the intrinsic and extrinsic coagulation cascade could be directly activated by the contact of blood with the circuit surface or lesion of endothelium, or indirectly by ROS mediation (Davie et al., 1991; Görlach, 2004, 2005; Herkert et al., 2004; Mackman et al., 2007; Oliver, 2009; Fall et al., 2015). The intrinsic coagulative pathway would be activated exclusively during the respiratory ECMO with low thrombin generation; while the release of high amount of TF and air-blood interface would stimulate both the intrinsic and extrinsic way with high thrombin generation during the CPB (Levy and Tanaka, 2003; Oliver, 2009; Kraft et al., 2015). Indeed, TF binds to Factor VII, converting it into Factor VIIa and activating the common coagulation cascade (Davie et al., 1991; Levy and Tanaka, 2003; Mackman et al., 2007; Oliver, 2009). The intrinsic pathway is activated through the conversion of Factor XII into Factor XIIa within 10 min from the beginning of the extracorporeal procedure and is associated with an increase in Bradykinin, Kallikrein, and activation of Factor IX (Wendel et al., 1999; Millar et al., 2016). Bradykinin stimulates NO, TNF α , and IL-10 production, while Kallikrein directly activates neutrophils (Rodell et al., 1995; Wachtfogel et al., 1995). Factor IXa induces the activation of Factor X which in turn causes the conversion of prothrombin into thrombin and fibrinogen in fibrin resulting in thrombus formation (Millar et al., 2016). Thrombin stimulates the endothelial production of P-selectin, E-selectin, and platelet

wounds. According to the ELSO registry, these latter events represents about 19% of all hemorrhagic complications during neonatal respiratory ECMO, impacting on survival (ELSO). Overall, the mortality rate is increased among neonates exposed to large volumes of PRBC transfusions (Smith et al., 2013; Fiser et al., 2014; Ghirardello et al., 2017). Dos Santos et al. observed a 27.8% mortality in critically ill infants exposed to transfusions during the hospital stay, with an increased relative risk of death of 1.49 in those infants who received at least one transfusion compared to controls (Dos Santos et al., 2011). In a cohort of extremely low birth weight infants, the incidence of mortality was 24% in patients transfused compared to 8.5% in the control group (Ghirardello et al., 2017). Similarly, Smith and co-authors observed a 24% increase in the odds of hospital death associated with every 10 ml/kg/day of PRBC transfusion in patients during respiratory ECMO (Smith et al., 2013).

In addition to PRBC, newborns on ECMO require approximately 25 ml/kg/day of PCs (Stiller et al., 2004). Platelet count is reduced by 40–50% during ECMO, partly due to the activation of the coagulation cascade (Nawab and Williams, 2017). Moreover, PCs are a source of ROS, especially O_2^- , which modulate platelet activation and aggregation (Krötz et al., 2002; Begonja et al., 2005). The increase in ROS, whose impact on the patient's oxidation state is still unknown, is mainly related to the storage process (Sonego et al., 2017).

ECMO and Hemolysis

ECMO-related hemolysis is a common phenomenon among pediatric patients and represents an independent predictor of mortality (Omar et al., 2015; Dalton et al., 2018). It consists of the RBCs lysis with following release of hemoglobin into the plasma (Rother et al., 2005). The shear stress is the main causative mechanism for hemolysis during mechanical support, and some risk factors have been identified, such as the negative inlet pressure, pump speed, pump and oxygenator type, cavitation and priming solution (Steinhorn et al., 1989; Lou et al., 2014; Rother et al., 2005). In addition, oxidative stress is a determinant of hemolysis (Boyle et al., 1997; Walski et al., 2018). Lipid peroxidation and protein damage of cell membranes from ROS cause enzymatic inactivation, depolarization of the cell membrane and a change in its permeability, resulting in cell lysis (Matés et al., 1999; Kirkham and Rahman, 2006). The mechanical and oxidative stress during ECMO leads to the release of the PFHb, which is cleared by hemoglobin scavengers (Dalton et al., 2018). However, in case of severe hemolysis, the hemoglobin-scavenging mechanisms become saturated, resulting in hemoglobinemia and hemoglobinuria (Rother et al., 2005). The PFHb may exert either a direct toxic effect (i.e., renal damage) or may cause endothelial dysfunction, by binding NO. The NO consumption leads to increased systemic and pulmonary resistance, platelet activation and aggregation, endothelial adhesion, fibrin deposition and thrombin formation (Rother et al., 2005).

Moreover, the shear stress induces procoagulant conformational changes of the pVWF, which interacts with circulating platelets enhancing their adhesion and aggregation properties (Cruz et al., 2018).

Furthermore, high levels of PFHb inhibit the cleavage of pVWF by ADAMTS-13, contributing to increased clotting within the mechanical devices (Da et al., 2015). Eventually, hyperbilirubinemia and iron release constitute additional drivers of the hemolysis-induced morbidity and mortality (Davis et al., 1999; Kaplan et al., 2014). Besides hemolysis, the mechanical stress may hesitate into sublethal RBCs damage, leading to increased fragility and decreased deformability of RBCs (Kameneva et al., 1999). Their consequent failure to enter small capillaries causes end-organ dysfunction, through the impairment of tissue oxygenation and microcirculation (Watanabe et al., 2007). Based on the concepts mentioned above, ECMO providers should monitor the hemolysis, as it is strictly associated with increased mortality and morbidity (kidney injury, vascular impairment, need for PRBC and PCs transfusion, thrombosis).

Moreover, efforts should focus on the technological progress, because of specific components of the circuit act as main drivers of the ECMO-induced hemolysis, such as the oxygenator (Williams et al., 2015). The use of smaller pediatric oxygenators was associated with greater shear stress and, hence, increased hemolysis, when compared to larger adult oxygenators (Williams et al., 2015). Although the relative contribution of each circuit component on the hemolysis and related oxidative stress is unknown, the use of a centrifugal pump and small size connectors should be considered as potential sites of increased hemolysis (Lou et al., 2014; Williams et al., 2015). Furthermore, the use of red-to-infrared radiation (R/NIR) to stabilize erythrocyte membranes with increased resistance to oxidative stress seems promising (Itoh et al., 2000; Komorowska et al., 2001; Chludzińska et al., 2005; Walski et al., 2015, 2018).

Technology advances should target the development of new, more bio-compatible, circuit-blood interfaces in order to reduce hemolysis and oxidative stress.

ECMO and Hyperoxia

The antioxidant capacity of tissues is overtaken with O_2^- production during hyperoxia. Oxygen toxicity in the perinatal period is well established, and many efforts have been made to define safety targets during neonatal resuscitation and care (Tin and Gupta, 2007; Auten and Davis, 2009; Finer and Leone, 2009; Askie et al., 2011; Saugstad et al., 2012; Vento et al., 2012; Schmidt et al., 2013; Saugstad and Aune, 2014; Manja et al., 2015).

Nevertheless, hyperoxia remains a significant driver for radical O_2 species generation, especially during ECC (Cashen et al., 2018). Hyperoxia is frequent during cardiac surgery in CPB and seems to play a key role in the development of oxidative stress as a consequence of post-ischemic reperfusion (Cavarocchi et al., 1986; Börgermann et al., 2007). The clamp of the aorta induces heart and lung ischemia while declamping generates reperfusion, and either action produces an inflammatory response. On-pump cardiac surgery is associated with increased oxidative stress if compared to off-pump surgery (Matata et al., 2000; Cavalca et al., 2006). The post-ischemic reperfusion, rather than hyperoxia *per se*, seems to play a role; there is evidence from pediatric cardiac surgery that isolated hyperoxia does not increase oxidative stress and inflammation if compared with normoxia (Hövels-Gürich

et al., 2002; Caputo et al., 2005, 2009). Although ECMO does not directly cause ischemia/reperfusion, patients requiring ECMO experience prolonged periods of pre-ECMO hypoxia causing an increase in oxidative stress (Crimi et al., 2006; Hayes et al., 2013; Chen Q. et al., 2014). ROS generation following hyperoxia is significantly exaggerated when hyperoxia is subsequent a period of hypoxia, due to the dysfunction of the post-ischemic mitochondrial electron transport chain (Neumar, 2011). Sznycer-Taub et al. (2016) identified hyperoxia within 48 h of ECMO following cardiac surgery as an independent risk factor for mortality at 30 days post-operatively, while Trittenwein et al. (1999) observed how VV-ECMO more than VA-ECMO would increase pulmonary lipid peroxidation following endotoxemia-induced hypoxia. Lipid peroxidation of the lung is a known consequence of sepsis and inflammation, but it has also been described as a consequence of tissue reperfusion (Demling et al., 1991; Horton and Walker, 1993; Christie et al., 1994; Chow et al., 2003). In the study by Trittenwein et al. (1999) hyperoxic reperfusion of the hypoxic lung following VV-ECMO, caused a significant increase of plasma MDA as an index of lipid peroxidation, higher than that observed in VA-ECMO where the hyperoxic blood supply to the lung is less.

More recently, similar results have been observed by Cashen et al. (2018) that showed higher in-hospital mortality among hyperoxic patients submitted to ECMO compared to normoxics, ascribing the high mortality to the effects of ROS on poor cardiac output rather than the SIRS-like syndrome, hemostatic activation and MOF.

ECMO and Continuous Renal Replacement Therapies

Oxidative stress is a common condition in kidney diseases (Witko-Sarsat et al., 1996; Hasselwander and Young, 1998); it is induced by renal pathology itself, reduced antioxidant intake associated with malnutrition, CRRT with the loss of antioxidant molecules during filtration and the contact of blood with CRRT circuit (Wolf et al., 1951; Hasselwander and Young, 1998; Cano, 2001; Sosa et al., 2006; Varan et al., 2010; Libetta et al., 2011; Coombes and Fassett, 2012; Ozbek, 2012). The SIRS-like syndrome, which typically occurs in the early hours of ECMO, is associated with hypotension, oligo-anuria, decreased lung compliance, pulmonary edema, increased plasma concentrations of pro-inflammatory cytokines and neutrophils' activation (Hocker et al., 1991; Davies and Hagen, 1997). AKI and FO complicate neonatal ECMO runs in more than half of cases, with a negative impact on survival rates (Heiss et al., 1987; Weber et al., 1990; Swaniker et al., 2000; Gadepalli et al., 2011; Zwieters et al., 2013). Therefore, hemofiltration and hemodialysis have become a standard of care in the management of AKI during ECMO (Heiss et al., 1987; Meyer et al., 2001; Paden et al., 2011; Chen H. et al., 2014). During CRRT, inflammation is activated by an increase of ROS and inflammatory cytokines (TNF α , IL-1, and IL-6) (Tarakçoğlu et al., 2003; Borazan et al., 2004) and a reduction of antioxidant proteins as hydrosoluble vitamin C and uric acids (Hegbrant and Hultkvist, 1999; Yang et al., 2006; Libetta et al., 2011). The use of biocompatible filtration

systems by using vitamin E-coated filters reduces the oxidative stress linked to the contact of blood with exogenous materials (Yang et al., 2014). Selewski et al. (2012) suggested an early filtration treatment to anticipate FO, and reduce mortality rate. As reported by ELSO registry, about 33% of neonatal respiratory ECMO and 38% of cardiac ECMO require renal replacement therapy (ELSO). As a result of the combination of two ECC systems, the hemolysis-related oxidative stress would increase (Chen H. et al., 2014). The in-line use of CRRT would limit the ECMO-related inflammation by reducing the concentrations of TNF α , IL-1 β , IL-6, and IL-8 (Jialiang et al., 2014).

ECMO Post-cardiopulmonary Bypass

Adult patients with cardiovascular diseases and neonatal patients with congenital cyanotic heart diseases have an increased level of oxidative stress, even before heart surgery, if compared to the healthy population (Ercan et al., 2009; Giustarini et al., 2009; Karu et al., 2010; Small et al., 2012).

Moreover, cardiac surgery is always associated with both systemic inflammation and a variable degree of oxidative stress, which might contribute to post-operative complications, failure of single and multiple organs and mortality (Kim et al., 2008; Billings et al., 2011; Antoniadis et al., 2012; Plicner et al., 2014; Wu et al., 2015). Plicner et al. (2014) observed that the increase of asymmetric dimethylarginine and 8-iso-prostaglandin F $_{2\alpha}$ was associated with an unfavorable outcome of patients undergoing coronary artery bypass grafting. Similarly, Antoniadis et al. (2012) has shown how the increase of O $_2^-$ and ONO $_2^-$ concentration was closely related to atrial fibrillation, need for inotropic drugs and lengthening of hospital stay during post-cardiac surgery.

The main cause of oxidative stress during cardiac surgery is the exposure to the foreign surface of the extracorporeal circuit, hypothermia, and circulatory arrest associated with systemic inflammation, ischemia-reperfusion injury, altered hemostatic status (Prasad et al., 1992; Edmunds, 1998; Clermont et al., 2002; Levy and Tanaka, 2003; Ulus et al., 2003; Gessler et al., 2004; Karu et al., 2010). Approximately 3–5% of patients undergoing congenital cardiac surgery require post-CPB extracorporeal support, due to the weaning failure following myocardial dysfunction, low cardiac output state, cardiac or respiratory arrest, pulmonary hypertension, or shunt occlusion (Aharon et al., 2001; Salvin et al., 2008; Mascio et al., 2014). The long duration of ECMO post-CPB might impair the organ function, thus predisposing to an increased risk of complications (i.e., sepsis) and mortality (Duncan et al., 1999; Chaturvedi et al., 2004; Salvin et al., 2008; Kumar et al., 2010; Rood et al., 2011; ELSO, 2018).

CLINICAL RELEVANCE OF OXIDATIVE STRESS DURING NEONATAL ECMO

The role of continuing oxidative stress has been established in the long term, in a variety of chronic diseases, such as diabetes mellitus, atherosclerosis and neurodegenerative illness (Haidara et al., 2006). Currently, the interest is moving toward the effects

of the redox unbalance in the acute illness (Bar-Or et al., 2015). Here, we summarize the available evidence in relation to the role of the oxidative stress in critical conditions, which often occur during ECMO (Figure 4).

Sepsis and Multi-Organ Failure

A wealth of studies has shown that the normal redox balance is altered in septic shock and multiple organ failure, due to both an increase in O_2^- derived free radicals and a decrease in the plasma antioxidant potential (Ogilvie et al., 1991; Goode et al., 1995). Adult septic patients have higher plasma CAT and SOD if compared to healthy controls (Warner et al., 1995). Moreover, non-survivors of sepsis have shown higher levels of SOD and a reduced plasmatic antioxidant status, which was associated with unfavorable outcomes (Cowley et al., 1996; Guerreiro et al., 2010). The sepsis-related MOF, which remains the major cause of death in intensive care units, is associated with oxidative stress and mitochondrial dysfunction (Brealey et al., 2002; Galley, 2011). Two antioxidants targeting the protection of mitochondria (MitoQ and MitoVitE) through the reduction of oxidative stress have been beneficial in a rat model of acute sepsis (Lowe et al., 2013). Other potential oxidative-induced mechanisms for MOF development have been proposed, as a result of the anti-oxidant impairment secondary to GSH and Se depletion, suggesting a positive effect of Se supplementation (Brealey et al., 2002; Manzanares et al., 2012; Broman et al., 2018).

Hemostatic Derangement and Anticoagulation Management


Maintaining the hemostatic homeostasis during ECMO is extremely challenging and, despite technical advances, bleeding and thrombosis remain serious complications (Murphy et al., 2015). The hemostatic perturbation during ECMO is a multifactorial process, and the redox imbalance is one of the main contributing factors. Besides the dilutional coagulopathy, which depends on priming compositions and affects both

clotting factors and platelets, the shear stress has a key role as it induces platelet dysfunction, fibrinolysis activation, acquired von Willebrand syndrome and intravascular hemolysis (Görlach et al., 2002; Görlach, 2004, 2005; Fall et al., 2015). As previously mentioned, the oxidative stress is strictly associated with the shear stress, resulting in endothelial activation (see section “ECMO and Inflammation”) and enhanced hemolysis (see section “ECMO and Hemolysis”) (Görlach et al., 2002; Görlach, 2004, 2005; Herkert et al., 2004; Djordjevic et al., 2005; Millar et al., 2016).

Moreover, the exposure to the exogenous circuits triggers a SIRS-like syndrome, which is associated with the ROS generation, the release of pro-inflammatory cytokines, the activation of the intrinsic coagulation pathway with a potential procoagulant effect (Millar et al., 2016). The clot formation triggers the consumption of coagulation factors and platelets, thus increasing the risk of disseminated intravascular coagulopathy and bleeding (Millar et al., 2016). The anticoagulation management should be modulated, by taking into account the drivers of hemostatic unbalance mentioned above.

Biomarkers of the Redox State

Based on the relevant clinical implication of the redox unbalance during ECMO, the identification of diagnostic markers would help in the detection, monitoring, and follow-up of the redox potential in ECMO neonates. However, the profile of biomarkers of both oxidative stress and antioxidant defenses during neonatal ECMO has not been characterized yet. Evidence can be derived from previous pre-clinical and clinical studies, which evaluate the pro-oxidative potential during and after the exposure to an extracorporeal circuit (i.e., CRRT, CPB) (Takouli et al., 2010; McDonald et al., 2012). As most of the research has been performed on the adult population, the evaluation of the redox status in neonatal systemic diseases (i.e., septic shock) may provide further insights into the age-related antioxidant deficiency (Batra et al., 2000). Among potential pro-oxidant



biomarkers, the d-ROMs have been evaluated in a dialyzer, while the MDA has been applied in a porcine ECMO model (Takouli et al., 2010; Chen Q. et al., 2014).

Additionally, as nitro-oxidative stress is involved in a large variety of pathological mechanisms, the biomarkers validated for the detection of nitrosation and nitration would be of interest (Cipak Gasparovic et al., 2017). However, their application in the extra-corporeal setting is still lacking. The anti-oxidant response may be evaluated through the detection of the TAC, SOD, and GSx activity (Takouli et al., 2010; McDonald et al., 2012). Moreover, the reduction of key trace elements, such as Se, copper, and zinc might suggest a reduced anti-oxidant activity (McDonald et al., 2012). In clinical practice, the oxidative stress may be indirectly presumed through the monitoring of inflammatory markers, such as the increased levels of IL-6 and IL-8 or by evaluating the hemolysis with the PFHb levels (Rother et al., 2005; Takouli et al., 2010; Cruz et al., 2018).

STRATEGIES TO REDUCE OXIDATIVE STRESS: UPDATE ON ANTIOXIDANTS

The ROS production is regulated by intra and extracellular antioxidant systems, which are impaired during the perinatal period and critical illness (Manzanares et al., 2012; McDonald et al., 2012). ECMO further complicates the issue, by adding specific sources for redox state unbalance (Manzanares et al., 2012; McDonald et al., 2012, 2014a; Jialiang et al., 2014; Millar et al., 2016; Bocca et al., 2017). The antioxidant system of the newborns is immature with both a reduction of antioxidant enzymes (SOD, CAT, and GPx) and non-enzymatic antioxidants (vitamin E, β -carotene, melatonin, ceruloplasmin, transferrin, coenzyme Q) while ascorbic acid and bilirubin are present only for a short period after birth (Gopinathan et al., 1994; Gitto et al., 2009; Bocca et al., 2017). Free iron levels are significantly higher than in adults, causing an increased Fenton reaction, which in turn stimulates the production of OH^\bullet (Saugstad, 2005). Furthermore both the SIRS-like and the sequestration into the ECMO/hemofilter circuit reduce the level of trace elements (copper, manganese, zinc, iron, Se as cofactors for SOD, GPx, and CAT function) and non-enzymatic antioxidants (albumin, uric acid, vitamins C and E) (Cano, 2001; Yamamoto et al., 2003; Yamawaki and Berk, 2006; Berndt et al., 2007; Tonelli et al., 2009; Hoffmann et al., 2011; Visser et al., 2011; Manzanares et al., 2012; Wołonciej et al., 2016; Ciapetti et al., 2018).

Based on the paucity of data regarding patients in ECMO both neonatal and adult, we will illustrate previous data regarding the use of anti-oxidants in critically ill adult patients undergoing surgery, CRRT or CPB. Moreover, taking into account the peculiarity of the perinatal period, we will mention some data on anti-oxidant strategies studied in neonatal conditions where oxidative stress is one of the pathogenic mechanisms, such as sepsis or NEC (Ozdemir et al., 2012; Poggi and Dani, 2018) (**Supplementary Table S1**). Among others, vitamin A, vitamin E, vitamin C, SOD, and NAC have been studied to reduce tissue damage mediated by oxidative stress with contrasting results

(Tyson et al., 1999; Suresh et al., 2001; Wardle et al., 2001; Brion et al., 2003; Darlow et al., 2016).

Antioxidants have two main mechanisms of action: the prevention, inactivating the free radical present in the systems and the scavenging of the active radical, suppressing the chain initiation or breaking the chain propagation. They may exert their effect by many actions including electron donation, hydrogen donor, metal ion chelation, co-antioxidants, radical scavenger, peroxide decomposer, singlet oxygen quencher, enzyme inhibitor or by gene expression regulation (Lobo et al., 2010).

Selenoproteins contribute to the function of many enzymes among which GSx, TRs, and methionine sulfoxide reductase and seem to protect the cardiovascular system against oxidative stress (Rotruck et al., 1973; Kim and Gladyshev, 2004; Hoffmann and Berry, 2005; Berndt et al., 2007; Hoffmann et al., 2011). Based on the reduction of Se levels in critically ill patients or after cardiac surgery, Se supplementation has been investigated in this contexts with contradictory results (Manzanares et al., 2012; Stoppe et al., 2013; Broman et al., 2018). To date, there are no data to support its extended use (Allingstrup and Afshari, 2015; Manzanares et al., 2016). Although not conclusive, neonatal supplementation with oral or parenteral Se in the first weeks of life to preterm infants led to a reduction in the proportion of infants having one or more episodes of sepsis (Darlow and Austin, 2003). The administration of pre and post-operative *vitamin C* in cardiac surgical adults has led to a reduction of post-operative atrial fibrillation, which seems to be associated with increased atrial oxidative stress and ONO_2^- formation (Carnes et al., 2001).

Moreover, supplementation with ascorbic acid caused a reduction of plasma levels of oxidative stress during hemodialysis and restoration of endothelial function in critically ill patients (Eiselt et al., 2001; Wilson, 2009).

Similarly, *vitamin E*-coated dialysis membranes used in adult patients on hemodialysis led to a decrease in reactive oxygen metabolites and derivatives, an increase of TAC and superoxide dismutase (Takouli et al., 2010). In addition, a decrease in post-operative atrial fibrillation has been documented after the supplementation of pre and post-operative NAC in surgical adult patients (Ozaydin et al., 2008). Ozdemir et al. (2012) have shown a beneficial effect of NAC in a murine NEC model in terms of reduced levels of oxidative stress (MDA) and inflammation (TNF α), with increased antioxidant activity of the SOD. Also concerning the supplementation of *glutamine*, data are not conclusive. Indeed, previous studies showed that infusion of glutamine during and for 3 days after cardio-surgery in adult patients led to an increase of glutamine blood levels and GSH activity without clinical improvement (Engel et al., 2009). Enteral glutamine either alone or in combination with arginine has been tested in healthy newborn rats showing a reduction of oxidative stress. These results may suggest a potential benefit of glutamine and arginine supplementation in the prevention of NEC in premature neonates with insufficient oxidative resistance (Kul et al., 2009). The evidence is building on the neonatal use of the ROS scavenger *melatonin*. Its supplementation in newborns with sepsis within the first 12 h after diagnosis has led to a reduction of lipid peroxidation products (MDA) and improvement of clinical outcomes (Gitto et al., 2001). The post-operative administration

of melatonin reduced the proinflammatory cytokines and NO levels in newborns undergoing surgery (Gitto et al., 2004).

Moreover, a reduction of late-onset sepsis risk ratio has been shown in newborn supplemented with *lactoferrin* and a reduction of the risk of antibiotic treatment failure in septic newborns exposed to *zinc* (Bhatnagar et al., 2012; Pammi and Suresh, 2017). Additional promising agents are the *apocynin*, an NOX inhibitor applied in the animal model of stroke and brain injury induced by ischemia-reperfusion, and the α -*lipoic acid*, a GSH-mediated antioxidant used in adults undergoing surgery during ECC (Packer et al., 1997; Kahles et al., 2007; Lambeth et al., 2008; Aly et al., 2009; Chen H. et al., 2009; Choi et al., 2010; Xia et al., 2010; Simonyi et al., 2012; Uyar et al., 2013). Novel antioxidant molecules, such as the ROS scavenger *edaravone*, showed promising results. In a pre-clinical model of neonatal sepsis the administration of edaravone 30 min after the injury has led to both a biochemical and clinical improvement (Kato et al., 2009). Other agents like *hyperbaric oxygen* and *medical ozone* have been suggested to reduce oxidative stress by enhancing the antioxidant system. The technique based on the use of O₂/ozone mix was previously applied in CRRT with promising results, even if its mechanism of action remains still unveiled (Bocci et al., 1999, 2001; Bocci and Paolo, 2004; Di Paolo et al., 2005). Data on the use of ozone in ECMO are lacking and might be a future area of interest. Lastly, it is worth mentioning the potential role of hMSCs in ischemia/reperfusion-induced lung injury, by acting through anti-oxidant, anti-inflammatory, and anti-apoptotic defense pathways (Ortiz et al., 2003; Gupta et al., 2007; Xu et al., 2007; Lee et al., 2009; Gotts and Matthay, 2011; Liu et al., 2017).

CONCLUSIONS AND FUTURE PERSPECTIVES

The imbalance of the redox system induced by the ECC is a field of active research, but many aspects of neonatal ECMO and its potential harm remain unsolved. Key points for both caregivers and researchers of the perinatal area addressing the burden of ECMO should include the following:

- Improve our knowledge of the mechanisms of oxidative stress during neonatal ECMO, both in pre-clinical and clinical settings, with a focus on the biochemical and cellular mechanism that could contribute to multi-organ damage.
- Define biomarkers for oxidative stress that could be a potential target for pharmacological approaches.
- Support technological research to improve ECMO circuit miniaturization and biocompatibility.
- Prevent the redox state unbalance during ECMO, through the promotion of rational use of blood products, the maintenance of well-defined O₂ targets, the prevention of hemolysis and AKI, the timely resort to in-line renal replacement therapy, the simplification of ECMO circuit lines.

AUTHOR CONTRIBUTIONS

GC, GR, SG, SP, and FM conceived and designed the review. GC, GR, and SG wrote the first draft of the manuscript. All authors contributed to manuscript critical revision, read, and approved the submitted version.

ACKNOWLEDGMENTS

The authors would like to thank all the Neonatal ECMO Team Mangiagalli of the Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico: nurses and neonatologists of the NICU, surgeons of the Department of Pediatric Surgery, anesthesiologists of the Pediatric Anesthesiology and Intensive Care Unit, nurses of the operating room. A special thank to Dr. Gabriele Zuanetti for the graphical support.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2018.01739/full#supplementary-material>

REFERENCES

- Abu-Zidan, F. M., Plank, L. D., and Windsor, J. A. (2002). Proteolysis in severe sepsis is related to oxidation of plasma protein. *Eur. J. Surg.* 168, 119–123. doi: 10.1080/11024150252884359
- Adib-Conquy, M., and Cavaiillon, J.-M. (2009). Compensatory anti-inflammatory response syndrome. *Thromb. Haemost.* 101, 36–47. doi: 10.1160/TH08-07-0421
- Adrian, K., Mellgren, K., Skogby, M., Friberg, L. G., Mellgren, G., and Wadenvik, H. (1998). Cytokine release during long-term extracorporeal circulation in an experimental model. *Artif. Organs* 22, 859–863. doi: 10.1046/j.1525-1594.1998.06121.x
- Afolayan, A. J., Eis, A., Alexander, M., Michalkiewicz, T., Teng, R.-J., Lakshminrusimha, S., et al. (2015). Decreased endothelial nitric oxide synthase expression and function contribute to impaired mitochondrial biogenesis and oxidative stress in fetal lambs with persistent pulmonary hypertension. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 310, L40–L49. doi: 10.1152/ajplung.00392.2014
- Afolayan, A. J., Eis, A., Teng, R.-J., Bakhtashvili, I., Kaul, S., Davis, J. M., et al. (2012). Decreases in manganese superoxide dismutase expression and activity contribute to oxidative stress in persistent pulmonary hypertension of the newborn. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 303, L870–L879. doi: 10.1152/ajplung.00098.2012
- Aharon, A. S., Drinkwater, D. C. Jr., Churchwell, K. B., Quisling, S. V., Reddy, V. S., Taylor, M., et al. (2001). Extracorporeal membrane oxygenation in children after repair of congenital cardiac lesions. *Ann. Thorac. Surg.* 72, 2095–2102. doi: 10.1016/S0003-4975(01)03209-X
- Al-Dalaen, S., and Al-Qatait, A. (2014). Review article: oxidative stress versus antioxidants. *Am. J. Biosci. Bioeng.* 2, 60–71. doi: 10.11648/j.bio.20140205.11
- Allingstrup, M., and Afshari, A. (2015). Selenium supplementation for critically ill adults. *Cochrane Database Syst. Rev.* 7:CD003703. doi: 10.1002/14651858.CD003703.pub3
- Aly, H. A., Lightfoot, D. A., and El-Shemy, H. A. (2009). Modulatory role of lipoic acid on lipopolysaccharide-induced oxidative stress in adult rat Sertoli cells in vitro. *Chem. Biol. Interact.* 182, 112–118. doi: 10.1016/j.cbi.2009.08.013

- Ang, A., Teo, D., Lim, C., Leou, K., Tien, S., and Koh, M. (2009). Blood transfusion requirements and independent predictors of increased transfusion requirements among adult patients on extracorporeal membrane oxygenation—a single centre experience. *Vox Sang.* 96, 34–43. doi: 10.1111/j.1423-0410.2008.01110.x
- Annagür, A., Örs, R., Altunhan, H., Kurban, S., Ertuğrul, S., Konak, M., et al. (2015). Total antioxidant and total oxidant states, and serum paraoxonase-1 in neonatal sepsis. *Pediatr. Int.* 57, 608–613. doi: 10.1111/ped.12557
- Antoniades, C., Demosthenous, M., Reilly, S., Margaritis, M., Zhang, M.-H., Antonopoulos, A., et al. (2012). Myocardial redox state predicts in-hospital clinical outcome after cardiac surgery: effects of short-term pre-operative statin treatment. *J. Am. Coll. Cardiol.* 59, 60–70. doi: 10.1016/j.jacc.2011.08.062
- Aras-López, R., Tovar, J., and Martínez, L. (2016). Possible role of increased oxidative stress in pulmonary hypertension in experimental diaphragmatic hernia. *Pediatr. Surg. Int.* 32, 141–145. doi: 10.1007/s00383-015-3826-5
- Asci, A., Surmeli-Onay, O., Erkekoglu, P., Yigit, S., Yurdakok, M., and Kocergumusel, B. (2015). Oxidant and antioxidant status in neonatal proven and clinical sepsis according to selenium status. *Pediatr. Int.* 57, 1131–1137. doi: 10.1111/ped.12698
- Askie, L. M., Brocklehurst, P., Darlow, B. A., Finer, N., Schmidt, B., and Tarnow-Mordi, W. (2011). NeOProm: neonatal oxygenation prospective M etanalysis Collaboration study protocol. *BMC Pediatr.* 11:6. doi: 10.1186/1471-2431-11-6
- Auten, R. L., and Davis, J. M. (2009). Oxygen toxicity and reactive oxygen species: the devil is in the details. *Pediatr. Res.* 66, 121–127. doi: 10.1203/PDR.0b013e3181a9eafb
- Ayala, A., Muñoz, M. F., and Argüelles, S. (2014). Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid. Med. Cell. Longev.* 2014:360438. doi: 10.1155/2014/360438
- Babior, B. M. (2000). Phagocytes and oxidative stress. *Am. J. Med.* 109, 33–44. doi: 10.1016/S0002-9343(00)00481-2
- Bahar, I., Elay, G., Başkol, G., Sungur, M., and Donmez-Altuntas, H. (2018). Increased DNA damage and augmented apoptosis and necrosis in patients with severe sepsis and septic shock. *J. Crit. Care* 43, 271–275. doi: 10.1016/j.jccr.2017.09.035
- Bar-Or, D., Bar-Or, R., Rael, L. T., and Brody, E. N. (2015). Oxidative stress in severe acute illness. *Redox Biol.* 4, 340–345. doi: 10.1016/j.redox.2015.01.006
- Barrera, G., Pizzimenti, S., Daga, M., Dianzani, C., Arcaro, A., Cetrangolo, G., et al. (2018). Lipid peroxidation-derived aldehydes, 4-hydroxynonenal and malondialdehyde in aging-related disorders. *Antioxidants* 7:E102. doi: 10.3390/antiox7080102
- Bartlett, R. H., Gazzaniga, A. B., Jefferies, M. R., Huxtable, R. F., Haiduc, N., and Fong, S. (1976). Extracorporeal membrane oxygenation (ECMO) cardiopulmonary support in infancy. *ASAIO J.* 22, 80–92.
- Batra, S., Kumar, R., Seema, Kapoor, A. K., and Ray, G. (2000). Alterations in antioxidant status during neonatal sepsis. *Ann. Trop. Paediatr.* 20, 27–33. doi: 10.1080/02724930092039
- Bayr, H. (2005). Reactive oxygen species. *Crit. Care Med.* 33, S498–S501. doi: 10.1097/01.CCM.0000186787.64500.12
- Begonja, A. J., Gambaryan, S., Geiger, J., Aktas, B., Pozgajova, M., Nieswandt, B., et al. (2005). Platelet NAD (P) H-oxidase-generated ROS production regulates α IIb β 3-integrin activation independent of the NO/cGMP pathway. *Blood* 106, 2757–2760. doi: 10.1182/blood-2005-03-1047
- Bellezza, I., Giambanco, I., Minelli, A., and Donato, R. (2018). Nrf2-Keap1 signaling in oxidative and reductive stress. *Biochim. Biophys. Acta Mol. Cell Res.* 1865, 721–733. doi: 10.1016/j.bbmc.2018.02.010
- Berndt, C., Lillig, C. H., and Holmgren, A. (2007). Thiol-based mechanisms of the thioredoxin and glutaredoxin systems: implications for diseases in the cardiovascular system. *Am. J. Physiol. Heart Circ. Physiol.* 292, H1227–H1236. doi: 10.1152/ajpheart.01162.2006
- Bertrand, Y., Pincemail, J., Hanique, G., Denis, B., Leenaerts, L., Vankeerberghen, L., et al. (1989). Differences in tocopherol-lipid ratios in ARDS and non-ARDS patients. *Intensive Care Med.* 15, 87–93. doi: 10.1007/BF00295983
- Bhatnagar, S., Wadhwa, N., Aneja, S., Lodha, R., Kabra, S. K., Natchu, U. C. M., et al. (2012). Zinc as adjunct treatment in infants aged between 7 and 120 days with probable serious bacterial infection: a randomised, double-blind, placebo-controlled trial. *Lancet* 379, 2072–2078. doi: 10.1016/S0140-6736(12)60477-2
- Billings, F. T., Ball, S. K., Roberts, I., L. J., and Pretorius, M. (2011). Postoperative acute kidney injury is associated with hemoglobinemia and an enhanced oxidative stress response. *Free Radic. Biol. Med.* 50, 1480–1487. doi: 10.1016/j.freeradbiomed.2011.02.011
- Bocca, B., Ciccarelli, S., Agostino, R., and Alimonti, A. (2017). Trace elements, oxidative status and antioxidant capacity as biomarkers in very low birth weight infants. *Environ. Res.* 156, 705–713. doi: 10.1016/j.envres.2017.04.027
- Bocci, V., Di, N. P., Borrelli, E., Larini, A., and Cappelletti, F. (2001). Ozonation of blood during extracorporeal circulation. II. Comparative analysis of several oxygenator-ozonators and selection of one type. *Int. J. Artif. Organs* 24, 890–897. doi: 10.1177/039139880102401202
- Bocci, V., Di, N. P., Garosi, G., Aldinucci, C., Borrelli, E., Valacchi, G., et al. (1999). Ozonation of blood during extracorporeal circulation. I. Rationale, methodology and preliminary studies. *Int. J. Artif. Organs* 22, 645–651. doi: 10.1177/039139889902200910
- Bocci, V., and Paolo, N. D. (2004). Oxygenation-ozonization of blood during extracorporeal circulation (EBOO). Part III: a new medical approach. *Ozone Sci. Eng.* 26, 195–205. doi: 10.1080/01919510490439564
- Bogdan, C. (2015). Nitric oxide synthase in innate and adaptive immunity: an update. *Trends Immunol.* 36, 161–178. doi: 10.1016/j.it.2015.01.003
- Borazan, A., Aydemir, S., Sert, M., and Yilmaz, A. (2004). The effects of hemodialysis and peritoneal dialysis on serum homocysteine and C-reactive protein levels. *Mediators Inflamm.* 13, 361–364. doi: 10.1080/09629350400008786
- Börgermann, J., Scheubel, R., Simm, A., Silber, R., and Friedrich, I. (2007). Inflammatory response in on-versus off-pump myocardial revascularization: is ECC really the culprit? *Thorac. Cardiovasc. Surg.* 55, 473–480. doi: 10.1055/s-2007-965631
- Boyle, E. M., Pohlman, T. H., Johnson, M. C., and Verrier, E. D. (1997). Endothelial cell injury in cardiovascular surgery: the systemic inflammatory response. *Ann. Thorac. Surg.* 63, 277–284.
- Brand, M. D. (2010). The sites and topology of mitochondrial superoxide production. *Exp. Gerontol.* 45, 466–472. doi: 10.1016/j.exger.2010.01.003
- Brealey, D., Brand, M., Hargreaves, I., Heales, S., Land, J., Smolenski, R., et al. (2002). Association between mitochondrial dysfunction and severity and outcome of septic shock. *Lancet* 360, 219–223. doi: 10.1016/S0140-6736(02)09459-X
- Brion, L. P., Bell, E. F., and Raghuvver, T. S. (2003). Vitamin E supplementation for prevention of morbidity and mortality in preterm infants. *Cochrane Database Syst. Rev.* 4:CD003665. doi: 10.1002/14651858.CD003665
- Briones, A. M., and Touyz, R. M. (2010). Oxidative stress and hypertension: current concepts. *Curr. Hypertens. Rep.* 12, 135–142. doi: 10.1007/s11906-010-0100-z
- Brix-Christensen, V., Petersen, T., Ravn, H. B., Hjortdal, V. E., Andersen, N. T., and Tønnesen, E. (2001). Cardiopulmonary bypass elicits a pro- and anti-inflammatory cytokine response and impaired neutrophil chemotaxis in neonatal pigs. *Acta Anaesthesiol. Scand.* 45, 407–413. doi: 10.1034/j.1399-6576.2001.045004407.x
- Brix-Christensen, V., Tønnesen, E., Hjortdal, V. E., Chew, M., Flø, C., Marqvorsen, J., et al. (2002). Neutrophils and platelets accumulate in the heart, lungs, and kidneys after cardiopulmonary bypass in neonatal pigs. *Crit. Care Med.* 30, 670–676. doi: 10.1097/00003246-200203000-00029
- Brogan, T., Laurance Lequier, M. D., Lorusso, R., Maclaren, G., and Peek, G. (eds). (2017). *Extracorporeal Life Support: the ELSO Red Book*, 5th Edn. Ann Arbor, MI: ELSO.
- Broman, M., Lindfors, M., Norberg, Å., Hebert, C., Rooyackers, O., Wernerman, J., et al. (2018). Low serum selenium is associated with the severity of organ failure in critically ill children. *Clin. Nutr.* 37, 1399–1405. doi: 10.1016/j.clnu.2017.06.014
- Bulua, A. C., Simon, A., Maddipati, R., Pelletier, M., Park, H., Kim, K.-Y., et al. (2011). Mitochondrial reactive oxygen species promote production of proinflammatory cytokines and are elevated in TNFR1-associated periodic syndrome (TRAPS). *J. Exp. Med.* 208, 519–533. doi: 10.1084/jem.20102049
- Butch, S., Knafl, P., Oberman, H., and Bartlett, R. (1996). Blood utilization in adult patients undergoing extracorporeal membrane oxygenated therapy. *Transfusion* 36, 61–63. doi: 10.1046/j.1537-2995.1996.36196190517.x

- Butterfield, D. A., Drake, J., Pocernich, C., and Castegna, A. (2001). Evidence of oxidative damage in Alzheimer's disease brain: central role for amyloid β -peptide. *Trends Mol. Med.* 7, 548–554. doi: 10.1016/S1471-4914(01)02173-6
- Cano, N. (2001). Hemodialysis, inflammation and malnutrition. *Nefrologia* 21, 437–442.
- Caputo, M., Bays, S., Rogers, C., Pawade, A., Parry, A., and Suleiman, M. (2005). The effects of normothermic and hypothermic cardiopulmonary bypass on myocardial injury, oxidative stress, and inflammatory response in paediatric open-heart surgery: a prospective randomized study. *Ann. Thorac. Surg.* 80, 982–988. doi: 10.1016/j.athoracsur.2005.03.062
- Caputo, M., Mokhtari, A., Rogers, C. A., Panayiotou, N., Chen, Q., Ghorbel, M. T., et al. (2009). The effects of normoxic versus hyperoxic cardiopulmonary bypass on oxidative stress and inflammatory response in cyanotic pediatric patients undergoing open cardiac surgery: a randomized controlled trial. *J. Thorac. Cardiovasc. Surg.* 138, 206–214. doi: 10.1016/j.jtcvs.2008.12.028
- Carnes, C. A., Chung, M. K., Nakayama, T., Nakayama, H., Baliga, R. S., Piao, S., et al. (2001). Ascorbate attenuates atrial pacing-induced peroxynitrite formation and electrical remodeling and decreases the incidence of postoperative atrial fibrillation. *Circ. Res.* 89, e32–e38. doi: 10.1161/hh1801.097644
- Carpenter, C. T., Price, P. V., and Christman, B. W. (1998). Exhaled breath condensate isoprostanes are elevated in patients with acute lung injury or ARDS. *Chest* 114, 1653–1659. doi: 10.1378/chest.114.6.1653
- Cashen, K., Reeder, R., Dalton, H. J., Berg, R. A., Shanley, T. P., Newth, C. J., et al. (2018). Hyperoxia and hypocapnia during pediatric extracorporeal membrane oxygenation: associations with complications, mortality, and functional status among survivors. *Pediatr. Crit. Care Med.* 19, 245–253. doi: 10.1097/PCC.0000000000001439
- Castegna, A., Aksenov, M., Thongboonkerd, V., Klein, J. B., Pierce, W. M., Booze, R., et al. (2002). Proteomic identification of oxidatively modified proteins in Alzheimer's disease brain. Part II: dihydropyrimidinase-related protein 2, α -enolase and heat shock cognate 71. *J. Neurochem.* 82, 1524–1532. doi: 10.1046/j.1471-4159.2002.01103.x
- Castellheim, A., Pharo, A., Fung, M., Saugstad, O. D., and Mollnes, T. E. (2005). Complement C5a is a key mediator of meconium-induced neutrophil activation. *Pediatr. Res.* 57, 242–247. doi: 10.1203/01.PDR.0000150725.78971.30
- Cavallo, V., Sisillo, E., Veglia, F., Tremoli, E., Cighetti, G., Salvi, L., et al. (2006). Isoprostanes and oxidative stress in off-pump and on-pump coronary bypass surgery. *Ann. Thorac. Surg.* 81, 562–567. doi: 10.1016/j.athoracsur.2005.08.019
- Cavarocchi, N. C., England, M. D., Schaff, H. V., Russo, P., Orszulak, T. A., Schnell, W. Jr., et al. (1986). Oxygen free radical generation during cardiopulmonary bypass: correlation with complement activation. *Circulation* 74, III130–III133.
- Chaturvedi, R., Macrae, D., Brown, K., Schindler, M., Smith, E., Davis, K., et al. (2004). Cardiac ECMO for biventricular hearts after paediatric open heart surgery. *Heart* 90, 545–551. doi: 10.1136/hrt.2002.003509
- Chaudhary, P., Sharma, R., Sharma, A., Vatsyayan, R., Yadav, S., Singhal, S. S., et al. (2010). Mechanisms of 4-hydroxy-2-nonenal induced pro- and anti-apoptotic signaling. *Biochemistry* 49, 6263–6275. doi: 10.1021/bi100517x
- Chen, G. Y., and Nuñez, G. (2010). Sterile inflammation: sensing and reacting to damage. *Nat. Rev. Immunol.* 10, 826–837. doi: 10.1038/nri2873
- Chen, H., Song, Y. S., and Chan, P. H. (2009). Inhibition of NADPH oxidase is neuroprotective after ischemia—reperfusion. *J. Cereb. Blood Flow Metab.* 29, 1262–1272. doi: 10.1038/jcbfm.2009.47
- Chen, Y., Azad, M., and Gibson, S. (2009). Superoxide is the major reactive oxygen species regulating autophagy. *Cell Death Differ.* 16, 1040–1052. doi: 10.1038/cdd.2009.49
- Chen, H., Yu, R.-G., Yin, N.-N., and Zhou, J.-X. (2014). Combination of extracorporeal membrane oxygenation and continuous renal replacement therapy in critically ill patients: a systematic review. *Crit. Care* 18:675. doi: 10.1186/s13054-014-0675-x
- Chen, Q., Yu, W., Shi, J., Shen, J., Hu, Y., Gong, J., et al. (2014). The effect of extracorporeal membrane oxygenation therapy on systemic oxidative stress injury in a porcine model. *Artif. Organs* 38, 426–431. doi: 10.1111/aor.12198
- Cheng, H. H., Wypij, D., Laussen, P. C., Bellinger, D. C., Stopp, C. D., Soul, J. S., et al. (2014). Cerebral blood flow velocity and neurodevelopmental outcome in infants undergoing surgery for congenital heart disease. *Ann. Thorac. Surg.* 98, 125–132. doi: 10.1016/j.athoracsur.2014.03.035
- Chludzińska, L., Ananicz, E., Jarosławska, A., and Komorowska, M. (2005). Near-infrared radiation protects the red cell membrane against oxidation. *Blood Cells Mol. Dis.* 35, 74–79. doi: 10.1016/j.bcmd.2005.04.003
- Choi, D. K., Koppula, S., Choi, M., and Suk, K. (2010). Recent developments in the inhibitors of neuroinflammation and neurodegeneration: inflammatory oxidative enzymes as a drug target. *Expert Opin. Ther. Pat.* 20, 1531–1546. doi: 10.1517/13543776.2010.525220
- Chow, C.-W., Herrera Abreu, M. T., Suzuki, T., and Downey, G. P. (2003). Oxidative stress and acute lung injury. *Am. J. Respir. Cell Mol. Biol.* 29, 427–431. doi: 10.1165/rcmb.F278
- Christie, N. A., Smith, D. E., Decampos, K. N., Slutsky, A. S., Patterson, G. A., and Tanswell, A. K. (1994). Lung oxidant injury in a model of lung storage and extended reperfusion. *Am. J. Respir. Crit. Care Med.* 150, 1032–1037. doi: 10.1164/ajrccm.150.4.7921433
- Ciapetti, M., Mancinelli, P., Cecchi, A., Borrelli, E., Bocci, V., and Peris, A. (2018). Reduction of non-enzymatic antioxidants in plasma during ECMO-treatment in ARDS by influence A H1N1. *J. Crit. Care* 43, 220–224. doi: 10.1016/j.jcrc.2017.08.005
- Cipak Gasparovic, A., Zarkovic, N., Zarkovic, K., Semen, K., Kaminsky, D., Yelisyeyeva, O., et al. (2017). Biomarkers of oxidative and nitro-oxidative stress: conventional and novel approaches. *Br. J. Pharmacol.* 174, 1771–1783. doi: 10.1111/bph.13673
- Clermont, G., Vergely, C., Jazayeri, S., Lahet, J.-J., Goudeau, J.-J., Lecour, S., et al. (2002). Systemic free radical activation is a major event involved in myocardial oxidative stress related to cardiopulmonary bypass. *Anesthesiology* 96, 80–87. doi: 10.1097/00000542-200201000-00019
- Collard, K., Godeck, S., Holley, J., and Quinn, M. (2004). Pulmonary antioxidant concentrations and oxidative damage in ventilated premature babies. *Arch. Dis. Child. Fetal Neonatal Ed.* 89, F412–F416. doi: 10.1136/adc.2002.016717
- Collard, K. J., Godeck, S., and Holley, J. E. (2005). Blood transfusion and pulmonary lipid peroxidation in ventilated premature babies. *Pediatr. Pulmonol.* 39, 257–261. doi: 10.1002/ppul.20190
- Comporti, M., Signorini, C., Leoncini, S., Buonocore, G., Rossi, V., and Ciccoli, L. (2004). Plasma F2-isoprostanes are elevated in newborns and inversely correlated to gestational age. *Free Radic. Biol. Med.* 37, 724–732. doi: 10.1016/j.freeradbiomed.2004.06.007
- Cooke, M. S., Evans, M. D., Dizdaroglu, M., and Lunec, J. (2003). Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J.* 17, 1195–1214. doi: 10.1096/fj.02-0752rev
- Coombes, J. S., and Fasset, R. G. (2012). Antioxidant therapy in hemodialysis patients: a systematic review. *Kidney Int.* 81, 233–246. doi: 10.1038/ki.2011.341
- Courouclis, X. I. (2017). Oxidative stress in the retina: implications for Retinopathy of Prematurity. *Curr. Opin. Toxicol.* 7, 102–109. doi: 10.1016/j.cotox.2017.11.008
- Cowley, H. C., Bacon, P. J., Goode, H. F., Webster, N. R., Jones, J. G., and Menon, D. K. (1996). Plasma antioxidant potential in severe sepsis: a comparison of survivors and nonsurvivors. *Crit. Care Med.* 24, 1179–1183. doi: 10.1097/00003246-199607000-00019
- Crimi, E., Sica, V., Williams-Ignarro, S., Zhang, H., Slutsky, A. S., Ignarro, L. J., et al. (2006). The role of oxidative stress in adult critical care. *Free Radic. Biol. Med.* 40, 398–406. doi: 10.1016/j.freeradbiomed.2005.10.054
- Cruz, C. M., Rinna, A., Forman, H. J., Ventura, A. L., Persechini, P. M., and Ojcius, D. M. (2007). ATP activates a reactive oxygen species-dependent oxidative stress response and secretion of proinflammatory cytokines in macrophages. *J. Biol. Chem.* 282, 2871–2879. doi: 10.1074/jbc.M608083200
- Cruz, M. A., Yee, A., and Valladolid, C. (2018). von Willebrand factor, free hemoglobin and thrombosis in ECMO. *Front. Med.* 5:228. doi: 10.3389/fmed.2018.00228
- Da, Q., Teruya, M., Guchhait, P., Teruya, J., Olson, J. S., and Cruz, M. A. (2015). Free hemoglobin increases von Willebrand factor-mediated platelet adhesion in vitro: implications on circulatory devices. *Blood* 126, 2338–2341. doi: 10.1182/blood-2015-05-648030
- Dalle-Donne, I., Aldini, G., Carini, M., Colombo, R., Rossi, R., and Milzani, A. (2006). Protein carbonylation, cellular dysfunction, and disease progression. *J. Cell. Mol. Med.* 10, 389–406. doi: 10.1111/j.1582-4934.2006.tb00407.x
- Dalle-Donne, I., Giustarini, D., Colombo, R., Rossi, R., and Milzani, A. (2003a). Protein carbonylation in human diseases. *Trends Mol. Med.* 9, 169–176. doi: 10.1016/S1471-4914(03)00031-5

- Dalle-Donne, I., Rossi, R., Giustarini, D., Milzani, A., and Colombo, R. (2003b). Protein carbonyl groups as biomarkers of oxidative stress. *Clin. Chim. Acta* 329, 23–38.
- Dalton, H. J., Cashen, K., Reeder, R. W., Berg, R. A., Shanley, T. P., Newth, C. J., et al. (2018). Hemolysis during pediatric extracorporeal membrane oxygenation: associations with circuitry, complications, and mortality. *Pediatr. Crit. Care Med.* 19, 1067–1076. doi: 10.1097/PCC.0000000000001709
- Dani, C., Reali, M., Bertini, G., Martelli, E., Pezzati, M., and Rubaltelli, F. (2001). The role of blood transfusions and iron intake on retinopathy of prematurity. *Early Hum. Dev.* 62, 57–63. doi: 10.1016/S0378-3782(01)00115-3
- Darlow, B. A., and Austin, N. (2003). Selenium supplementation to prevent short-term morbidity in preterm neonates. *Cochrane Database Syst. Rev.* 4:CD003312. doi: 10.1002/14651858.CD003312
- Darlow, B. A., Graham, P., and Rojas-Reyes, M. X. (2016). Vitamin A supplementation to prevent mortality and short-and long-term morbidity in very low birth weight infants. *Cochrane Database Syst. Rev.* 8:CD000501. doi: 10.1002/14651858.CD000501.pub4
- Davie, E. W., Fujikawa, K., and Kisiel, W. (1991). The coagulation cascade: initiation, maintenance, and regulation. *Biochemistry* 30, 10363–10370. doi: 10.1021/bi00107a001
- Davies, M., and Hagen, P. O. (1997). Systemic inflammatory response syndrome. *Br. J. Surg.* 84, 920–935. doi: 10.1002/bjs.1800840707
- Davis, C. L., Kausz, A. T., Zager, R. A., Kharasch, E. D., and Cochran, R. P. (1999). Acute renal failure after cardiopulmonary bypass is related to decreased serum ferritin levels. *J. Am. Soc. Nephrol.* 10, 2396–2402.
- Demling, R. H., La Londe, C., Daryani, R., Zhu, D., Knox, J., and Youn, Y.-K. (1991). Relationship between the lung and systemic response to endotoxin: comparison of physiologic change and the degree of lipid peroxidation. *Circ. Shock* 34, 364–370.
- Depuydt, L. E., Schuit, K. E., and Smith, S. D. (1993). Effect of extracorporeal membrane oxygenation on neutrophil function in neonates. *Crit. Care Med.* 21, 1324–1327. doi: 10.1097/00003246-199309000-00015
- Desborough, J. (2000). The stress response to trauma and surgery. *Br. J. Anaesth.* 85, 109–117. doi: 10.1093/bja/85.1.109
- Di Paolo, N., Bocci, V., Salvo, D., Palasciano, F., Biagioli, M., Meini, S., et al. (2005). *Extracorporeal Blood Oxygenation and Ozonation (EBOO): a Controlled Trial in Patients with Peripheral Artery Disease*. London: SAGE Publications.
- Dialynas, G., Shrestha, O. K., Ponce, J. M., Zwerger, M., Thiemann, D. A., Young, G. H., et al. (2015). Myopathic lamin mutations cause reductive stress and activate the nrf2/keap-1 pathway. *PLoS Genet.* 11:e1005231. doi: 10.1371/journal.pgen.1005231
- Dimon-Gadal, S., Gerbaud, P., Guibourdenche, J., Evain-Brion, D., Raynaud, F., Thérond, P., et al. (2000). Increased oxidative damage to fibroblasts in skin with and without lesions in psoriasis. *J. Investig. Dermatol.* 114, 984–989. doi: 10.1046/j.1523-1747.2000.00962.x
- Dizdaroğlu, M., and Jaruga, P. (2012). Mechanisms of free radical-induced damage to DNA. *Free Radic. Res.* 46, 382–419. doi: 10.3109/10715762.2011.653969
- Djordjevic, T., Pogrebniak, A., Belaiba, R. S., Bonello, S., Wotzlaw, C., Acker, H., et al. (2005). The expression of the NADPH oxidase subunit p22phox is regulated by a redox-sensitive pathway in endothelial cells. *Free Radic. Biol. Med.* 38, 616–630. doi: 10.1016/j.freeradbiomed.2004.09.036
- Dos Santos, A. M. N., Guinsburg, R., De Almeida, M. F. B., Procianny, R. S., Leone, C. R., Marba, S. T. M., et al. (2011). Red blood cell transfusions are independently associated with intra-hospital mortality in very low birth weight preterm infants. *J. Pediatr.* 159, 371–376.
- Descamps-Latscha, B., Drüeke, T., and Witko-sarsat, V. (2001). Dialysis-induced oxidative stress: biological aspects, clinical consequences, and therapy. *Semin. Dial.* 14, 193–199. doi: 10.1046/j.1525-139X.2001.00052.x
- Duncan, B. W., Hraska, V., Jonas, R. A., Wessel, D. L., Del Nido, P. J., Laussen, P. C., et al. (1999). Mechanical circulatory support in children with cardiac disease. *J. Thorac. Cardiovasc. Surg.* 117, 529–542. doi: 10.1016/S0022-5223(99)70333-8
- Edmunds, L. H. Jr. (1998). Inflammatory response to cardiopulmonary bypass. *Ann. Thorac. Surg.* 66, S12–S16. doi: 10.1016/S0003-4975(98)00967-9
- Egginton, S. (2009). Invited review: activity-induced angiogenesis. *Pflügers Arch.* 457, 963–977. doi: 10.1007/s00424-008-0563-9
- Eiselt, J., Racek, J., Trefil, L., and Opatrný, K. (2001). Effects of a vitamin E-modified dialysis membrane and vitamin C infusion on oxidative stress in hemodialysis patients. *Artif. Organs* 25, 430–436. doi: 10.1046/j.1525-1594.2001.025006430.x
- ELSO (2018). *ELSO Registry Report. International Summary January 2018*. Ann Arbor, MI: ELSO.
- Engel, J., Pitz, S., Mühling, J., Menges, T., Martens, F., Kwapisz, M., et al. (2009). Role of glutamine administration on T-cell derived inflammatory response after cardiopulmonary bypass. *Clin. Nutr.* 28, 15–20. doi: 10.1016/j.clnu.2008.08.007
- Ercan, S., Çakmak, A., Kösecik, M., and Erel, Ö. (2009). The oxidative state of children with cyanotic and acyanotic congenital heart disease. *Anadolu Kardiyol. Derg.* 9, 486–490.
- Erikssen, G., Liestøl, K., Seem, E., Birkeland, S., Saatvedt, K. J., Hoel, T. N., et al. (2015). Achievements in congenital heart defect surgery: a prospective, 40-year study of 7038 patients. *Circulation* 131, 337–346. doi: 10.1161/CIRCULATIONAHA.114.012033
- Fall, L., New, K. J., Evans, K. A., and Bailey, D. M. (2015). Arterial hypoxaemia and its impact on coagulation: significance of altered redox homeostasis. *J. Clin. Pathol.* 68, 752–754. doi: 10.1136/jclinpath-2015-202952
- Finer, N., and Leone, T. (2009). Oxygen saturation monitoring for the preterm infant: the evidence basis for current practice. *Pediatr. Res.* 65, 375–380. doi: 10.1203/PDR.0b013e318199386a
- Finkel, T. (2012). From sulfonylation to sulfhydration: what a thiolate needs to tolerate. *Sci. Signal.* 5:e10. doi: 10.1126/scisignal.2002943
- Fiser, R. T., Irby, K., Ward, R. M., Tang, X., Mckemie, W., Prophan, P., et al. (2014). RBC transfusion in pediatric patients supported with extracorporeal membrane oxygenation: is there an impact on tissue oxygenation? *Pediatr. Crit. Care Med.* 15, 806–813. doi: 10.1097/PCC.0000000000000222
- Fläring, U., Rooyackers, O., Hebert, C., Bratel, T., Hammarqvist, F., and Wernerman, J. (2005). Temporal changes in whole-blood and plasma glutathione in ICU patients with multiple organ failure. *Intensive Care Med.* 31, 1072–1078. doi: 10.1007/s00134-005-2687-0
- Floor, E., and Wetzel, M. G. (1998). Increased protein oxidation in human substantia nigra pars compacta in comparison with basal ganglia and prefrontal cortex measured with an improved dinitrophenylhydrazine assay. *J. Neurochem.* 70, 268–275. doi: 10.1046/j.1471-4159.1998.70010268.x
- Fortenberry, J. D., Bhardwaj, V., Niemer, P., Cornish, J. D., Wright, J. A., and Bland, L. (1996). Neutrophil and cytokine activation with neonatal extracorporeal membrane oxygenation. *J. Pediatr.* 128, 670–678. doi: 10.1016/S0022-3476(96)80133-8
- Fridovich, I. (1997). Superoxide anion radical (O₂^{•-}), superoxide dismutases, and related matters. *J. Biol. Chem.* 272, 18515–18517. doi: 10.1074/jbc.272.30.18515
- Fudulu, D., and Angelini, G. (2016). Oxidative stress after surgery on the immature heart. *Oxid. Med. Cell. Longev.* 2016:1971452. doi: 10.1155/2016/1971452
- Gadepalli, S. K., Selewski, D. T., Drongowski, R. A., and Mychaliska, G. B. (2011). Acute kidney injury in congenital diaphragmatic hernia requiring extracorporeal life support: an insidious problem. *J. Pediatr. Surg.* 46, 630–635. doi: 10.1016/j.jpedsurg.2010.11.031
- Galley, H. F. (2011). Oxidative stress and mitochondrial dysfunction in sepsis. *Br. J. Anaesth.* 107, 57–64. doi: 10.1093/bja/aer093
- Gessler, P., Pretre, R., Hohl, V., Rousson, V., Fischer, J., and Dahinden, C. (2004). CX-C-chemokine stimulation of neutrophils correlates with plasma levels of myeloperoxidase and lactoferrin and contributes to clinical outcome after pediatric cardiac surgery. *Shock* 22, 513–520. doi: 10.1097/01.shk.0000145939.54838.51
- Ghirardello, S., Dusi, E., Cortinovis, I., Villa, S., Fumagalli, M., Agosti, M., et al. (2017). Effects of red blood cell transfusions on the risk of developing complications or death: an observational study of a cohort of very low birth weight infants. *Am. J. Perinatol.* 34, 88–95. doi: 10.1055/s-0036-1584300
- Giacco, F., and Brownlee, M. (2010). Oxidative stress and diabetic complications. *Circ. Res.* 107, 1058–1070. doi: 10.1161/CIRCRESAHA.110.223545
- Gilboa, S. M., Salemi, J. L., Nembhard, W. N., Fixler, D. E., and Correa, A. (2010). Mortality resulting from congenital heart disease among children and adults in the United States, 1999 to 2006. *Circulation* 122, 2254–2263. doi: 10.1161/CIRCULATIONAHA.110.947002
- Gitto, E., Karbownik, M., Reiter, R. J., Tan, D. X., Cuzzocrea, S., Chiuazzi, P., et al. (2001). Effects of melatonin treatment in septic newborns. *Pediatr. Res.* 50, 756–760. doi: 10.1203/00006450-200112000-00021
- Gitto, E., Pellegrino, S., D'arrigo, S., Barberi, I., and Reiter, R. (2009). Oxidative stress in resuscitation and in ventilation of newborns. *Eur. Respir. J.* 34, 1461–1469. doi: 10.1183/09031936.00032809

- Gitto, E., Reiter, R. J., Cordaro, S. P., La Rosa, M., Chiurazzi, P., Trimarchi, G., et al. (2004). Oxidative and inflammatory parameters in respiratory distress syndrome of preterm newborns: beneficial effects of melatonin. *Am. J. Perinatol.* 21, 209–216. doi: 10.1055/s-2004-828610
- Giustarini, D., Dalle-Donne, I., Tskias, D., and Rossi, R. (2009). Oxidative stress and human diseases: origin, link, measurement, mechanisms, and biomarkers. *Crit. Rev. Clin. Lab. Sci.* 46, 241–281. doi: 10.3109/10408360903142326
- Gladstone, I. M., and Levine, R. L. (1994). Oxidation of proteins in neonatal lungs. *Pediatriatrics* 93, 764–768.
- Golej, J., Winter, P., Schöffmann, G., Kahlbacher, H., Stoll, E., Boigner, H., et al. (2003). Impact of extracorporeal membrane oxygenation modality on cytokine release during rescue from infant hypoxia. *Shock* 20, 110–115. doi: 10.1097/01.shk.0000075571.93053.2c
- Goode, H. F., Cowley, H. C., Walker, B. E., Howdle, P. D., and Webster, N. R. (1995). Decreased antioxidant status and increased lipid peroxidation in patients with septic shock and secondary organ dysfunction. *Crit. Care Med.* 23, 646–651. doi: 10.1097/00003246-199504000-00011
- Goodyear-Bruch, C., and Pierce, J. D. (2002). Oxidative stress in critically ill patients. *Am. J. Crit. Care* 11, 543–551.
- Gopinathan, V., Miller, N. J., Milner, A. D., and Rice-Evans, C. A. (1994). Bilirubin and ascorbate antioxidant activity in neonatal plasma. *FEBS Lett.* 349, 197–200. doi: 10.1016/0014-5793(94)00666-0
- Görlach, A. (2004). *Redox Control of Blood Coagulation*. New Rochelle, NY: Mary Ann Liebert, Inc.
- Görlach, A. (2005). Redox regulation of the coagulation cascade. *Antioxid. Redox Signal.* 7, 1398–1404. doi: 10.1089/ars.2005.7.1398
- Görlach, A., Kietzmann, T., and Hess, J. (2002). Redox signaling through NADPH oxidases: involvement in vascular proliferation and coagulation. *Ann. N. Y. Acad. Sci.* 973, 505–507. doi: 10.1111/j.1749-6632.2002.tb04691.x
- Gotts, J. E., and Matthay, M. A. (2011). Mesenchymal stem cells and acute lung injury. *Crit. Care Clin.* 27, 719–733. doi: 10.1016/j.ccc.2011.04.004
- Graulich, J., Sonntag, J., Marcinkowski, M., Bauer, K., Kössel, H., Bühner, C., et al. (2002). Complement activation by in vivo neonatal and in vitro extracorporeal membrane oxygenation. *Mediators Inflamm.* 11, 69–73. doi: 10.1080/09629350220131908
- Guerreiro, M. O., Petronilho, F., Andrades, M., Constantino, L., Mina, F. G., Moreira, J. C. F., et al. (2010). Plasma superoxide dismutase activity and mortality in patients with septic. *J. Trauma Acute Care Surg.* 69, E102–E106. doi: 10.1097/TA.0b013e3181dbb289
- Gupta, N., Su, X., Popov, B., Lee, J. W., Serikov, V., and Matthay, M. A. (2007). Intrapulmonary delivery of bone marrow-derived mesenchymal stem cells improves survival and attenuates endotoxin-induced acute lung injury in mice. *J. Immunol.* 179, 1855–1863. doi: 10.4049/jimmunol.179.3.1855
- Gutteridge, J. M., and Halliwell, B. (2018). Mini-Review: oxidative stress, redox stress or redox success? *Biochem. Biophys. Res. Commun.* 502, 183–186. doi: 10.1016/j.bbrc.2018.05.045
- Haidara, M. A., Yassin, H. Z., Rateb, M., Ammar, H., and Zorkani, M. A. (2006). Role of oxidative stress in development of cardiovascular complications in diabetes mellitus. *Curr. Vasc. Pharmacol.* 4, 215–227. doi: 10.2174/15701610677698469
- Halliwell, B., and Gutteridge, J. M. (2015). *Free Radicals in Biology and Medicine*. New York, NY: Oxford University Press. doi: 10.1093/acprof:oso/9780198717478.001.0001
- Hammond, B., and Hess, M. L. (1985). The oxygen free radical system: potential mediator of myocardial injury. *J. Am. Coll. Cardiol.* 6, 215–220. doi: 10.1016/S0735-1097(85)80278-3
- Hasselwander, O., and Young, I. S. (1998). Oxidative stress in chronic renal failure. *Free Radic. Res.* 29, 1–11. doi: 10.1080/10715769800300011
- Hayes, R., Shekar, K., and Fraser, J. (2013). Is hyperoxaemia helping or hurting patients during extracorporeal membrane oxygenation? Review of a complex problem. *Perfusion* 28, 184–193. doi: 10.1177/0267659112473172
- He, C., Yang, S., Yu, W., Chen, Q., Shen, J., Hu, Y., et al. (2014). Effects of continuous renal replacement therapy on intestinal mucosal barrier function during extracorporeal membrane oxygenation in a porcine model. *J. Cardiothorac. Surg.* 9:72. doi: 10.1186/1749-8090-9-72
- Hegbrant, J., and Hultkvist, B. U. (1999). Vitamin C and E as antioxidants in hemodialysis patients. *Int. J. Artif. Organs* 22, 69–73. doi: 10.1177/039139889902200203
- Heiss, K. F., Pettit, B., Hirschl, R. B., Cilley, R. E., Chapman, R., and Bartlett, R. H. (1987). Renal insufficiency and volume overload in neonatal ECMO managed by continuous ultrafiltration. *ASAIO J.* 33, 557–560.
- Herkert, O., Djordjevic, T., Belaiba, R. S., and Görlach, A. (2004). Insights into the redox control of blood coagulation: role of vascular NADPH oxidase-derived reactive oxygen species in the thrombogenic cycle. *Antioxid. Redox Signal.* 6, 765–776. doi: 10.1089/1523086041361695
- Hirthler, M., Simoni, J., and Dickson, M. (1992). Elevated levels of endotoxin, oxygen-derived free radicals, and cytokines during extracorporeal membrane oxygenation. *J. Pediatr. Surg.* 27, 1199–1202. doi: 10.1016/0022-3468(92)90787-8
- Hocker, J. R., Wellhausen, S. R., Ward, R. A., Simpson, P. M., and Cook, L. N. (1991). Effect of extracorporeal membrane oxygenation on leukocyte function in neonates. *Artif. Organs* 15, 23–28. doi: 10.1111/j.1525-1594.1991.tb00755.x
- Hoffmann, F. W., Hashimoto, A. S., Lee, B. C., Rose, A. H., Shohet, R. V., and Hoffmann, P. R. (2011). Specific antioxidant selenoproteins are induced in the heart during hypertrophy. *Arch. Biochem. Biophys.* 512, 38–44. doi: 10.1016/j.abb.2011.05.007
- Hoffmann, P. R., and Berry, M. J. (2005). Selenoprotein synthesis: a unique translational mechanism used by a diverse family of proteins. *Thyroid* 15, 769–775. doi: 10.1089/thy.2005.15.769
- Horton, J. W., and Walker, P. B. (1993). Oxygen radicals, lipid peroxidation, and permeability changes after intestinal ischemia and reperfusion. *J. Appl. Physiol.* 74, 1515–1520. doi: 10.1152/jappl.1993.74.4.1515
- Hövels-Gürich, H. H., Vazquez-Jimenez, J. F., Silvestri, A., Schumacher, K., Minkenber, R., Duchateau, J., et al. (2002). Production of proinflammatory cytokines and myocardial dysfunction after arterial switch operation in neonates with transposition of the great arteries. *J. Thorac. Cardiovasc. Surg.* 124, 811–820. doi: 10.1067/jmtc.2002.122308
- Huet, O., Obata, R., Aubron, C., Spraul-Davit, A., Charpentier, J., Laplace, C., et al. (2007). Plasma-induced endothelial oxidative stress is related to the severity of septic shock. *Crit. Care Med.* 35, 821–826. doi: 10.1097/01.CCM.0000257464.79067.AF
- Itoh, T., Murakami, H., Orihashi, K., Sueda, T., Kusumoto, Y., Kakehashi, M., et al. (2000). Low power laser protects human erythrocytes in an in vitro model of artificial heart-lung machines. *Artif. Organs* 24, 870–873. doi: 10.1046/j.1525-1594.2000.06624.x
- Iwasaki, A., and Medzhitov, R. (2010). Regulation of adaptive immunity by the innate immune system. *Science* 327, 291–295. doi: 10.1126/science.1183021
- Jialiang, S., Juanhong, S., Qiyi, C., Wenkui, Y., Changsheng, H., Yimin, H., et al. (2014). In-line hemofiltration minimized extracorporeal membrane oxygenation-related inflammation in a porcine model. *Perfusion* 29, 526–533. doi: 10.1177/0267659114529320
- Kahles, T., Luedike, P., Endres, M., Galla, H.-J., Steinmetz, H., Busse, R., et al. (2007). NADPH oxidase plays a central role in blood-brain barrier damage in experimental stroke. *Stroke* 38, 3000–3006. doi: 10.1161/STROKEAHA.107.489765
- Kameneva, M. V., Undar, A., Antaki, J. F., Watach, M. J., Calhoon, J. H., and Borovetz, H. S. (1999). Decrease in red blood cell deformability caused by hypothermia, hemodilution, and mechanical stress: factors related to cardiopulmonary bypass. *ASAIO J.* 45, 307–310. doi: 10.1097/00002480-199907000-00010
- Kaplan, M., Bromiker, R., and Hammerman, C. (2014). Hyperbilirubinemia, hemolysis, and increased bilirubin neurotoxicity. *Semin. Perinatol.* 38, 429–437. doi: 10.1053/j.semperi.2014.08.006
- Kaplanski, G., Fabrigoule, M., Boulay, V., Dinarello, C. A., Bongrand, P., Kaplanski, S., et al. (1997). Thrombin induces endothelial type II activation in vitro: IL-1 and TNF-alpha-independent IL-8 secretion and E-selectin expression. *J. Immunol.* 158, 5435–5441.
- Karu, I., Taal, G., Zilmer, K., Pruunsild, C., Starkopf, J., and Zilmer, M. (2010). Inflammatory/oxidative stress during the first week after different types of cardiac surgery. *Scand. Cardiovasc. J.* 44, 119–124. doi: 10.3109/14017430903490981
- Kato, S., Hussein, M. H., Kakita, H., Goto, T., Daoud, G. A., Kato, T., et al. (2009). Edaravone, a novel free radical scavenger, reduces high-mobility group box 1 and prolongs survival in a neonatal sepsis model. *Shock* 32, 586–592. doi: 10.1097/SHK.0b013e3181a2b886

- Khaper, N., Bryan, S., Dhingra, S., Singal, R., Bajaj, A., Pathak, C. M., et al. (2010). Targeting the vicious inflammation-oxidative stress cycle for the management of heart failure. *Antioxid. Redox Signal.* 13, 1033–1049. doi: 10.1089/ars.2009.2930
- Kim, H.-Y., and Gladyshev, V. N. (2004). Methionine sulfoxide reduction in mammals: characterization of methionine-R-sulfoxide reductases. *Mol. Biol. Cell* 15, 1055–1064. doi: 10.1091/mbc.e03-08-0629
- Kim, Y. M., Kattach, H., Ratnatunga, C., Pillai, R., Channon, K. M., and Casadei, B. (2008). Association of atrial nicotinamide adenine dinucleotide phosphate oxidase activity with the development of atrial fibrillation after cardiac surgery. *J. Am. Coll. Cardiol.* 51, 68–74. doi: 10.1016/j.jacc.2007.07.085
- Kirkham, P., and Rahman, I. (2006). Oxidative stress in asthma and COPD: antioxidants as a therapeutic strategy. *Pharmacol. Ther.* 111, 476–494. doi: 10.1016/j.pharmthera.2005.10.015
- Klaunig, and Je, W. Z. (2018). Oxidative stress in carcinogenesis. *Curr. Opin. Toxicol.* 7, 116–121. doi: 10.1016/j.cotox.2017.11.014
- Koch, C. G., Li, L., Sessler, D. I., Figueroa, P., Hoeltge, G. A., Mihaljevic, T., et al. (2008). Duration of red-cell storage and complications after cardiac surgery. *N. Engl. J. Med.* 358, 1229–1239. doi: 10.1056/NEJMoa070403
- Komorowska, M., Cui, A., Czarnoleski, A., and Bialas, W. (2001). Erythrocyte response to near infrared radiation. *Cell. Mol. Biol. Lett.* 6, 212–212.
- Kotani, N., Hashimoto, H., Sessler, D. I., Muraoka, M., Wang, J.-S., O'Connor, M. F., et al. (2000). Neutrophil number and interleukin-8 and elastase concentrations in bronchoalveolar lavage fluid correlate with decreased arterial oxygenation after cardiopulmonary bypass. *Anesth. Analg.* 90, 1046–1051. doi: 10.1097/00000539-200005000-00009
- Kouchoukos, N. T., Blackstone, E. H., Hanley, F. L., and Kirklin, J. K. (2012). *Kirklin/Barratt-Boyes Cardiac Surgery E-Book*. Amsterdam: Elsevier Health Sciences.
- Kraehling, J. R., and Sessa, W. C. (2017). Contemporary approaches to modulating the nitric oxide-cgmp pathway in cardiovascular disease. *Circ. Res.* 120, 1174–1182. doi: 10.1161/CIRCRESAHA.117.303776
- Kraft, F., Schmidt, C., Van Aken, H., and Zarbock, A. (2015). Inflammatory response and extracorporeal circulation. *Best Pract. Res. Clin. Anaesthesiol.* 29, 113–123. doi: 10.1016/j.bpa.2015.03.001
- Krötz, F., Sohn, H. Y., Gloe, T., Zahler, S., RieXinger, T., Schiele, T. M., et al. (2002). NAD (P) H oxidase-dependent platelet superoxide anion release increases platelet recruitment. *Blood* 100, 917–924. doi: 10.1182/blood.V100.3.917
- Kul, M., Vurucu, S., Demirkaya, E., Tunc, T., Aydin, S., Meral, C., et al. (2009). Enteral glutamine and/or arginine supplementation have favorable effects on oxidative stress parameters in neonatal rat intestine. *J. Pediatr. Gastroenterol. Nutr.* 49, 85–89. doi: 10.1097/MPG.0b013e318198cd36
- Kumar, T. S., Zurakowski, D., Dalton, H., Talwar, S., Allard-Picou, A., Duebener, L. F., et al. (2010). Extracorporeal membrane oxygenation in postcardiotomy patients: factors influencing outcome. *J. Thorac. Cardiovasc. Surg.* 140, 330–336.e2. doi: 10.1016/j.jtcvs.2010.02.034
- Lakshminrusimha, S., Russell, J. A., Wedgwood, S., Gugino, S. F., Kazzaz, J. A., Davis, J. M., et al. (2006). Superoxide dismutase improves oxygenation and reduces oxidation in neonatal pulmonary hypertension. *Am. J. Respir. Crit. Care Med.* 174, 1370–1377. doi: 10.1164/rccm.200605-676OC
- Lamb, N. J., Quinlan, G. J., Westerman, S. T., Gutteridge, J. M., and Evans, T. W. (1999). Nitration of proteins in bronchoalveolar lavage fluid from patients with acute respiratory distress syndrome receiving inhaled nitric oxide. *Am. J. Respir. Crit. Care Med.* 160, 1031–1034. doi: 10.1164/ajrccm.160.3.9810048
- Lambeth, J. D. (2004). NOX enzymes and the biology of reactive oxygen. *Nat. Rev. Immunol.* 4, 181–189. doi: 10.1038/nri1312
- Lambeth, J. D., Krause, K.-H., and Clark, R. A. (2008). NOX enzymes as novel targets for drug development. *Semin. Immunopathol.* 30, 339–363. doi: 10.1007/s00281-008-0123-6
- Laplace, C., Huet, O., Vicaut, E., Ract, C., Martin, L., Benhamou, D., et al. (2005). Endothelial oxidative stress induced by serum from patients with severe trauma hemorrhage. *Intensive Care Med.* 31, 1174–1180. doi: 10.1007/s00134-005-2737-7
- Lee, J. W., Fang, X., Gupta, N., Serikov, V., and Matthay, M. A. (2009). Allogeneic human mesenchymal stem cells for treatment of *E. coli* endotoxin-induced acute lung injury in the ex vivo perfused human lung. *Proc. Natl. Acad. Sci. U.S.A.* 106, 16357–16362. doi: 10.1073/pnas.0907996106
- Lenz, A., Jorens, P., Meyer, B., De Backer, W., Van Overveld, F., Bossaert, L., et al. (1999). Oxidatively modified proteins in bronchoalveolar lavage fluid of patients with ARDS and patients at-risk for ARDS. *Eur. Respir. J.* 13, 169–174. doi: 10.1034/j.1399-3003.1999.13a31.x
- Levy, J. H., and Tanaka, K. A. (2003). Inflammatory response to cardiopulmonary bypass. *Ann. Thorac. Surg.* 75, S715–S720. doi: 10.1016/S0003-4975(02)04701-X
- Li, P., Gu, Y., Yu, S., Li, Y., Yang, J., and Jia, G. (2014). Assessing the suitability of 8-OHdG and micronuclei as genotoxic biomarkers in chromate-exposed workers: a cross-sectional study. *BMJ Open* 4:e005979. doi: 10.1136/bmjopen-2014-005979
- Libetta, C., Sepe, V., Esposito, P., Galli, F., and Dal Canton, A. (2011). Oxidative stress and inflammation: implications in uremia and hemodialysis. *Clin. Biochem.* 44, 1189–1198. doi: 10.1016/j.clinbiochem.2011.06.988
- Liu, Y.-Y., Chiang, C.-H., Hung, S.-C., Chian, C.-F., Tsai, C.-L., Chen, W.-C., et al. (2017). Hypoxia-preconditioned mesenchymal stem cells ameliorate ischemia/reperfusion-induced lung injury. *PLoS One* 12:e0187637. doi: 10.1371/journal.pone.0187637
- Lobo, V., Patil, A., Phatak, A., and Chandra, N. (2010). Free radicals, antioxidants and functional foods: impact on human health. *Pharmacogn. Rev.* 4, 118–126. doi: 10.4103/0973-7847.70902
- Lou, S., Maclaren, G., Best, D., Delzoppo, C., and Butt, W. (2014). Hemolysis in pediatric patients receiving centrifugal-pump extracorporeal membrane oxygenation: prevalence, risk factors, and outcomes. *Crit. Care Med.* 42, 1213–1220. doi: 10.1097/CCM.0000000000000128
- Lowes, D., Webster, N., Murphy, M., and Galley, H. (2013). Antioxidants that protect mitochondria reduce interleukin-6 and oxidative stress, improve mitochondrial function, and reduce biochemical markers of organ dysfunction in a rat model of acute sepsis. *Br. J. Anaesth.* 110, 472–480. doi: 10.1093/bja/ae577
- Luban, N. L. (1995). Massive transfusion in the neonate. *Transfus. Med. Rev.* 9, 200–214. doi: 10.1016/S0887-7963(05)80110-5
- Mackman, N., Tilley, R. E., and Key, N. S. (2007). Role of the extrinsic pathway of blood coagulation in hemostasis and thrombosis. *Arterioscler. Thromb. Vasc. Biol.* 27, 1687–1693. doi: 10.1161/ATVBAHA.107.141911
- MacMicking, J., Xie, Q.-W., and Nathan, C. (1997). Nitric oxide and macrophage function. *Annu. Rev. Immunol.* 15, 323–350. doi: 10.1146/annurev.immunol.15.1.323
- Maiorino, M., Conrad, M., and Ursini, F. (2018). GPx4, lipid peroxidation, and cell death: discoveries, rediscoveries, and open issues. *Antioxid. Redox Signal.* 29, 61–74. doi: 10.1089/ars.2017.7115
- Maltepe, E., and Saugstad, O. D. (2009). Oxygen in health and disease: regulation of oxygen homeostasis-clinical implications. *Pediatr. Res.* 65, 261–268. doi: 10.1203/PDR.0b013e3181818f83f
- Manja, V., Lakshminrusimha, S., and Cook, D. J. (2015). Oxygen saturation target range for extremely preterm infants: a systematic review and meta-analysis. *JAMA Pediatr.* 169, 332–340. doi: 10.1001/jamapediatrics.2014.3307
- Mantle, D., Falkous, G., and Walker, D. (1999). Quantification of protease activities in synovial fluid from rheumatoid and osteoarthritis cases: comparison with antioxidant and free radical damage markers. *Clin. Chim. Acta* 284, 45–58. doi: 10.1016/S0009-8981(99)00055-8
- Manzanares, W., Dhaliwal, R., Jiang, X., Murch, L., and Heyland, D. K. (2012). Antioxidant micronutrients in the critically ill: a systematic review and meta-analysis. *Crit. Care* 16:R66. doi: 10.1186/cc11316
- Manzanares, W., Lemieux, M., Elke, G., Langlois, P. L., Bloos, F., and Heyland, D. K. (2016). High-dose intravenous selenium does not improve clinical outcomes in the critically ill: a systematic review and meta-analysis. *Crit. Care* 20:356. doi: 10.1186/s13054-016-1529-5
- Margaritelis, N., Kyparos, A., Paschalis, V., Theodorou, A., Panayiotou, G., Zafeiridis, A., et al. (2014). Reductive stress after exercise: the issue of redox individuality. *Redox Biol.* 2, 520–528. doi: 10.1016/j.redox.2014.02.003
- Mascio, C. E., Austin, E. H., Jacobs, J. P., Jacobs, M. L., Wallace, A. S., He, X., et al. (2014). Perioperative mechanical circulatory support in children: an analysis of the Society of Thoracic Surgeons Congenital Heart Surgery Database. *J. Thorac. Cardiovasc. Surg.* 147, 658–665. doi: 10.1016/j.jtcvs.2013.09.075
- Matata, B. M., Sosnowski, A. W., and Galiñanes, M. (2000). Off-pump bypass graft operation significantly reduces oxidative stress and inflammation. *Ann. Thorac. Surg.* 69, 785–791. doi: 10.1016/S0003-4975(99)01420-4

- Matés, J. M., Pérez-Gómez, C., and De Castro, I. N. (1999). Antioxidant enzymes and human diseases. *Clin. Biochem.* 32, 595–603. doi: 10.1016/S0009-9120(99)00075-2
- Mátyás, M., and Zaharie, G. (2018). “Particularities of oxidative stress in newborns,” in *Novel Prospects in Oxidative and Nitrosative Stress*, ed. P. Atukeren (London: IntechOpen).
- McCoy-Pardington, D., Judd, W., Knafl, P., Abruzzo, L. V., Coombes, K., Butch, S., et al. (1990). Blood use during extracorporeal membrane oxygenation. *Transfusion* 30, 307–309. doi: 10.1046/j.1537-2995.1990.30490273436.x
- McDonald, C. I., Fraser, J. F., Coombes, J. S., and Fung, Y. L. (2014a). Oxidative stress during extracorporeal circulation. *Eur. J. Cardiothorac. Surg.* 46, 937–943. doi: 10.1093/ejcts/ezt637
- McDonald, C. I., Fraser, J. F., Shekar, K., Dunster, K. R., Thom, O., and Fung, Y. L. (2014b). Transfusion of packed red blood cells reduces selenium levels and increases lipid peroxidation in an in vivo ovine model. *Transfus. Med.* 24, 50–54.
- McDonald, C. I., Fung, Y. L., and Fraser, J. F. (2012). Antioxidant trace element reduction in an in vitro cardiopulmonary bypass circuit. *ASAIO J.* 58, 217–222. doi: 10.1097/MAT.0b013e31824cc856
- McIlwain, R. B., Timpa, J. G., Kurundkar, A. R., Holt, D. W., Kelly, D. R., Hartman, Y. E., et al. (2010). Plasma concentrations of inflammatory cytokines rise rapidly during ECMO-related SIRS due to the release of preformed stores in the intestine. *Lab. Invest.* 90, 128–139. doi: 10.1038/labinvest.2009.119
- Meyer, R. J., Brophy, P. D., Bunchman, T. E., Annich, G. M., Maxvold, N. J., Mottes, T. A., et al. (2001). Survival and renal function in pediatric patients following extracorporeal life support with hemofiltration. *Pediatr. Crit. Care Med.* 2, 238–242. doi: 10.1097/00130478-200107000-00009
- Mikolka, P., Kopincova, J., Mikusiakova, L. T., Kosutova, P., Calkovska, A., and Mokra, D. (2016). Antiinflammatory effect of N-acetylcysteine combined with exogenous surfactant in meconium-induced lung injury. *Adv. Exp. Med. Biol.* 934, 63–75. doi: 10.1007/5584_2016_15
- Mildner, R. J., Taub, N., Vyas, J. R., Killer, H. M., Firmin, R. K., Field, D. J., et al. (2005). Cytokine imbalance in infants receiving extracorporeal membrane oxygenation for respiratory failure. *Neonatology* 88, 321–327. doi: 10.1159/000087630
- Millar, J. E., Fanning, J. P., McDonald, C. I., McAuley, D. F., and Fraser, J. F. (2016). The inflammatory response to extracorporeal membrane oxygenation (ECMO): a review of the pathophysiology. *Crit. Care* 20:387. doi: 10.1186/s13054-016-1570-4
- Mishra, O. P., Pooniya, V., Ali, Z., Upadhyay, R. S., and Prasad, R. (2008). Antioxidant status of children with acute renal failure. *Pediatr. Nephrol.* 23:2047. doi: 10.1007/s00467-008-0875-1
- Mittal, M., Siddiqui, M. R., Tran, K., Reddy, S. P., and Malik, A. B. (2014). Reactive oxygen species in inflammation and tissue injury. *Antioxid. Redox Signal.* 20, 1126–1167. doi: 10.1089/ars.2012.5149
- Mokra, D., Drgova, A., Mokry, J., Antosova, M., Durdik, P., and Calkovska, A. (2015). N-acetylcysteine effectively diminished meconium-induced oxidative stress in adult rabbits. *J. Physiol. Pharmacol.* 66, 101–110.
- Möller, J., Gilman, J., Sussman, J., Raszynski, A., and Wolfsdorf, J. (1993). Changes in plasma levels of oxygen radical scavenging enzymes during extracorporeal membrane oxygenation in a lamb model. *Neonatology* 64, 134–139. doi: 10.1159/000243983
- Montuschi, P., Barnes, P. J., and Roberts, L. J. (2004). Isoprostanes: markers and mediators of oxidative stress. *FASEB J.* 18, 1791–1800. doi: 10.1096/fj.04-2330rev
- Moore, T. A., Ahmad, I. M., and Zimmerman, M. C. (2018). Oxidative stress and preterm birth: an integrative review. *Biol. Res. Nurs.* 20, 497–512. doi: 10.1177/1099800418791028
- Motoyama, T., Okamoto, K., Kukita, I., Hamaguchi, M., Kinoshita, Y., and Ogawa, H. (2003). Possible role of increased oxidant stress in multiple organ failure after systemic inflammatory response syndrome. *Crit. Care Med.* 31, 1048–1052. doi: 10.1097/01.CCM.0000055371.27268.36
- Murphy, D. A., Hockings, L. E., Andrews, R. K., Aubron, C., Gardiner, E. E., Pellegrino, V. A., et al. (2015). Extracorporeal membrane oxygenation—hemostatic complications. *Transfus. Med. Rev.* 29, 90–101. doi: 10.1016/j.tmr.2014.12.001
- Nawab, U., and Williams, S. B. (2017). “Transfusion considerations for neonatal extracorporeal membrane oxygenation (ECMO),” in *Neonatal Transfusion Practices*, ed. D. A. Sesok-Pizzini (Berlin: Springer), 103–121. doi: 10.1007/978-3-319-42764-5_7
- Neumar, R. W. (2011). Optimal oxygenation during and after cardiopulmonary resuscitation. *Curr. Opin. Crit. Care* 17, 236–240. doi: 10.1097/MCC.0b013e3283454c8c
- Niki, E., Yoshida, Y., Saito, Y., and Noguchi, N. (2005). Lipid peroxidation: mechanisms, inhibition, and biological effects. *Biochem. Biophys. Res. Commun.* 338, 668–676. doi: 10.1016/j.bbrc.2005.08.072
- Ogilvie, A., Groeneveld, A., Straub, J., and Thijs, L. (1991). Plasma lipid peroxides and antioxidants in human septic shock. *Intensive Care Med.* 17, 40–44. doi: 10.1007/BF01708408
- Oliver, W. C. (2009). Anticoagulation and coagulation management for ECMO. *Semin. Cardiothorac. Vasc. Anesth.* 13, 154–175. doi: 10.1177/1089253209347384
- Omar, H. R., Mirsaedi, M., Socias, S., Sprenger, C., Caldeira, C., Camporesi, E. M., et al. (2015). Plasma free hemoglobin is an independent predictor of mortality among patients on extracorporeal membrane oxygenation support. *PLoS One* 10:e0124034. doi: 10.1371/journal.pone.0124034
- Orban, J.-C., and Singer, M. (2018). “Oxygen and oxidative stress,” in *Metabolic Disorders and Critically Ill Patients*, eds C. Ichai, H. Quintard, and J.-C. Orban (Berlin: Springer), 431–439. doi: 10.1007/978-3-319-64010-5_20
- Ortiz, L. A., Gambelli, F., McBride, C., Gaupp, D., Baddoo, M., Kaminski, N., et al. (2003). Mesenchymal stem cell engraftment in lung is enhanced in response to bleomycin exposure and ameliorates its fibrotic effects. *Proc. Natl. Acad. Sci. U.S.A.* 100, 8407–8411. doi: 10.1073/pnas.1432929100
- Ozaydin, M., Peker, O., Erdogan, D., Kapan, S., Turker, Y., Varol, E., et al. (2008). N-acetylcysteine for the prevention of postoperative atrial fibrillation: a prospective, randomized, placebo-controlled pilot study. *Eur. Heart J.* 29, 625–631. doi: 10.1093/eurheartj/ehn011
- Ozbek, E. (2012). Induction of oxidative stress in kidney. *Int. J. Nephrol.* 2012:465897. doi: 10.1155/2012/465897
- Ozdemir, R., Yurttutan, S., Sarı, F. N., Uysal, B., Unverdi, H. G., Canpolat, F. E., et al. (2012). Antioxidant effects of N-acetylcysteine in a neonatal rat model of necrotizing enterocolitis. *J. Pediatr. Surg.* 47, 1652–1657. doi: 10.1016/j.jpedsurg.2012.02.016
- Pacher, P., Beckman, J. S., and Liaudet, L. (2007). Nitric oxide and peroxynitrite in health and disease. *Physiol. Rev.* 87, 315–424. doi: 10.1152/physrev.00029.2006
- Packer, L., Tritschler, H. J., and Wessel, K. (1997). Neuroprotection by the metabolic antioxidant α -lipoic acid. *Free Radic. Biol. Med.* 22, 359–378. doi: 10.1016/S0891-5849(96)00269-9
- Paden, M. L., Warshaw, B. L., Heard, M. L., and Fortenberry, J. D. (2011). Recovery of renal function and survival after continuous renal replacement therapy during extracorporeal membrane oxygenation. *Pediatr. Crit. Care Med.* 12, 153–158. doi: 10.1097/PCC.0b013e3181e2a596
- Palatianos, G. M., Foroulis, C. N., Vassili, M. I., Astras, G., Triantafyllou, K., Papadakis, E., et al. (2003). A prospective, double-blind study on the efficacy of the bioline surface-heparinized extracorporeal perfusion circuit. *Ann. Thorac. Surg.* 76, 129–135. doi: 10.1016/S0003-4975(03)00338-2
- Pammi, M., and Suresh, G. (2017). Enteral lactoferrin supplementation for prevention of sepsis and necrotizing enterocolitis in preterm infants. *Cochrane Database Syst. Rev.* 6:CD007137. doi: 10.1002/14651858.CD007137.pub5
- Perrone, S., Luisa Tataranno, M., Santacroce, A., Negro, S., and Buonocore, G. (2014). The role of oxidative stress on necrotizing enterocolitis in very low birth weight infants. *Curr. Pediatr. Rev.* 10, 202–207.
- Perrone, S., Santacroce, A., Longini, M., Proietti, F., Bazzini, F., and Buonocore, G. (2018). The free radical diseases of prematurity: from cellular mechanisms to bedside. *Oxid. Med. Cell. Longev.* 2018:7483062. doi: 10.1155/2018/7483062
- Perrone, S., Tataranno, M. L., Negro, S., Longini, M., Marzocchi, B., Proietti, F., et al. (2010). Early identification of the risk for free radical-related diseases in preterm newborns. *Early Hum. Dev.* 86, 241–244. doi: 10.1016/j.earlhumdev.2010.03.008
- Perrone, S., Vezzosi, P., Longini, M., Marzocchi, B., Paffetti, P., Bellieni, C. V., et al. (2009). Biomarkers of oxidative stress in babies at high risk for retinopathy of prematurity. *Front. Biosci.* 1, 547–552.
- Petersen, J., Kazakov, A., Böhm, M., Schäfers, H.-J., Laufs, U., and Abdul-Khalik, H. (2018). Cardiopulmonary bypass reduces myocardial oxidative stress, inflammation and increases c-kit+ CD45- cell population in newborns. *J. Transl. Med.* 16:111. doi: 10.1186/s12967-018-1478-7

- Philippu, A. (2016). Nitric Oxide: a universal modulator of brain function. *Curr. Med. Chem.* 23, 2643–2652. doi: 10.2174/0929867323666160627120408
- Plicner, D., Mazur, P., Sadowski, J., and Undas, A. (2014). Asymmetric dimethylarginine and oxidative stress following coronary artery bypass grafting: associations with postoperative outcome. *Eur. J. Cardiothorac. Surg.* 45, e136–e141. doi: 10.1093/ejcts/etz646
- Plötz, F., Van Oeveren, W., Bartlett, R., and Wildevuur, C. (1993). Blood activation during neonatal extracorporeal life support. *J. Thorac. Cardiovasc. Surg.* 105, 823–832.
- Poggi, C., and Dani, C. (2018). Sepsis and oxidative stress in the newborn: from pathogenesis to novel therapeutic targets. *Oxid. Med. Cell. Longev.* 2018:9390140. doi: 10.1155/2018/9390140
- Prasad, K., Kalra, J., Bharadwaj, B., and Chaudhary, A. K. (1992). Increased oxygen free radical activity in patients on cardiopulmonary bypass undergoing aortocoronary bypass surgery. *Am. Heart J.* 123, 37–45. doi: 10.1016/0002-8703(92)90744-G
- Prescott, S. M., Zimmerman, G. A., and McIntyre, T. M. (1984). Human endothelial cells in culture produce platelet-activating factor (1-alkyl-2-acetyl-sn-glycero-3-phosphocholine) when stimulated with thrombin. *Proc. Natl. Acad. Sci. U.S.A.* 81, 3534–3538. doi: 10.1073/pnas.81.11.3534
- Quinlan, G. J., Lamb, N. J., Evans, T. W., and Gutteridge, J. M. (1996). Plasma fatty acid changes and increased lipid peroxidation in patients with adult respiratory distress syndrome. *Crit. Care Med.* 24, 241–246. doi: 10.1097/00003246-199602000-00010
- Quinlan, G. J., Mumby, S., Martin, G. S., Bernard, G. R., Gutteridge, J. M., and Evans, T. W. (2004). Albumin influences total plasma antioxidant capacity favorably in patients with acute lung injury. *Crit. Care Med.* 32, 755–759. doi: 10.1097/01.CCM.0000114574.18641.5D
- Radi, R. (2013). Peroxynitrite, a stealthy biological oxidant. *J. Biol. Chem.* 288, 26464–26472. doi: 10.1074/jbc.R113.472936
- Ramsay, P. L., Demayo, F. J., Hegemier, S. E., Wearden, M. E., Smith, C. V., and Welty, S. E. (2001). Clara cell secretory protein oxidation and expression in premature infants who develop bronchopulmonary dysplasia. *Am. J. Respir. Crit. Care Med.* 164, 155–161. doi: 10.1164/ajrccm.164.1.2008022
- Renke, J., Popadiuk, S., Korzon, M., Bugajczyk, B., and Woźniak, M. (2000). Protein carbonyl groups' content as a useful clinical marker of antioxidant barrier impairment in plasma of children with juvenile chronic arthritis. *Free Radic. Biol. Med.* 29, 101–104. doi: 10.1016/S0891-5849(00)00288-4
- Richard, C., Lemonnier, F., Thibault, M., Couturier, M., and Auzepy, P. (1990). Vitamin E deficiency and lipoperoxidation during adult respiratory distress syndrome. *Crit. Care Med.* 18, 4–9. doi: 10.1097/00003246-199001000-00002
- Rodell, T. C., Naidoo, Y., and Bhoola, K. D. (1995). Role of kinins in inflammatory responses. *Clin. Immunother.* 3, 352–361. doi: 10.1007/BF03259501
- Romano, A., Serviddio, G., De Mattheis, A., and Bellanti, F., Vendemiale, G. (2010). *Oxidative stress and aging. J. Nephrol.* 23, S29–S36.
- Romano, R., Cristescu, S. M., Risby, T. H., and Marczin, N. (2018). Lipid peroxidation in cardiac surgery: towards consensus on biomonitoring, diagnostic tools and therapeutic implementation. *J. Breath Res.* 12:027109. doi: 10.1088/1752-7163/aa9856
- Rood, K. L., Teele, S. A., Barrett, C. S., Salvin, J. W., Rycus, P. T., Fynn-Thompson, F., et al. (2011). Extracorporeal membrane oxygenation support after the Fontan operation. *J. Thorac. Cardiovasc. Surg.* 142, 504–510. doi: 10.1016/j.jtcvs.2010.11.050
- Roos, W. P., and Kaina, B. (2006). DNA damage-induced cell death by apoptosis. *Trends Mol. Med.* 12, 440–450. doi: 10.1016/j.molmed.2006.07.007
- Rosa, S. D., Bristot, M. L., Tapanotti, M. F., Tomasi, C. D., Felisberto, F., Vuolo, F. S., et al. (2011). Effect of red blood cell transfusion on parameters of inflammation and oxidative stress in critically ill patients. *Rev. Bras. Ter. Intensiva* 23, 30–35. doi: 10.1590/S0103-507X2011000100006
- Rother, R. P., Bell, L., Hillmen, P., and Gladwin, M. T. (2005). The clinical sequelae of intravascular hemolysis and extracellular plasma hemoglobin: a novel mechanism of human disease. *JAMA* 293, 1653–1662. doi: 10.1001/jama.293.13.1653
- Rotruck, J. T., Pope, A. L., Ganther, H., Swanson, A., Hafeman, D. G., and Hoekstra, W. (1973). Selenium: biochemical role as a component of glutathione peroxidase. *Science* 179, 588–590. doi: 10.1126/science.179.4073.588
- Rungtatscher, A., Tessari, M., Stranieri, C., Solani, E., Linardi, D., Milani, E., et al. (2015). Oxygenator is the main responsible for leukocyte activation in experimental model of extracorporeal circulation: a cautionary tale. *Mediators Inflamm.* 2015:484979. doi: 10.1155/2015/484979
- Sahoo, S., Meijles, D. N., and Pagano, P. J. (2016). NADPH oxidases: key modulators in aging and age-related cardiovascular diseases? *Clin. Sci.* 130, 317–335. doi: 10.1042/CS20150087
- Salvin, J. W., Laussen, P. C., and Thiagarajan, R. R. (2008). Extracorporeal membrane oxygenation for postcardiotomy mechanical cardiovascular support in children with congenital heart disease. *Pediatr. Anesth.* 18, 1157–1162. doi: 10.1111/j.1460-9592.2008.02795.x
- Saugstad, O. D. (2005). Oxidative stress in the newborn—a 30-year perspective. *Neonatology* 88, 228–236. doi: 10.1159/000087586
- Saugstad, O. D., and Aune, D. (2014). Optimal oxygenation of extremely low birth weight infants: a meta-analysis and systematic review of the oxygen saturation target studies. *Neonatology* 105, 55–63. doi: 10.1159/000356561
- Saugstad, O. D., Sejersted, Y., Solberg, R., Wollen, E. J., and Bjørås, M. (2012). Oxygenation of the newborn: a molecular approach. *Neonatology* 101, 315–325. doi: 10.1159/000337345
- Schieber, M., and Chandel, N. S. (2014). ROS function in redox signaling and oxidative stress. *Curr. Biol.* 24, R453–R462. doi: 10.1016/j.cub.2014.03.034
- Schmidt, B., Whyte, R. K., Asztalos, E. V., Moddemann, D., Poets, C., Rabi, Y., et al. (2013). Effects of targeting higher vs lower arterial oxygen saturations on death or disability in extremely preterm infants: a randomized clinical trial. *JAMA* 309, 2111–2120. doi: 10.1001/jama.2013.5555
- Selewski, D. T., Cornell, T. T., Blatt, N. B., Han, Y. Y., Mottes, T., Kommareddi, M., et al. (2012). Fluid overload and fluid removal in pediatric patients on extracorporeal membrane oxygenation requiring continuous renal replacement therapy. *Crit. Care Med.* 40, 2694. doi: 10.1097/CCM.0b013e318258ff01
- Semenza, G. L. (2009). Regulation of oxygen homeostasis by hypoxia-inducible factor 1. *Physiology* 24, 97–106. doi: 10.1152/physiol.00045.2008
- Shi, J., Chen, Q., Yu, W., Shen, J., Gong, J., He, C., et al. (2014). Continuous renal replacement therapy reduces the systemic and pulmonary inflammation induced by venovenous extracorporeal membrane oxygenation in a porcine model. *Artif. Organs* 38, 215–223. doi: 10.1111/aor.12154
- Shohat, M., Reischer, S. H., Krikler, R., Nissenkorn, I., Yassur, Y., and Ben-Sira, I. (1983). Retinopathy of prematurity: incidence and risk factors. *Pediatrics* 72, 159–163.
- Sies, H. (2018). On the history of oxidative stress: concept and some aspects of current development. *Curr. Opin. Toxicol.* 7, 122–126. doi: 10.1016/j.cotox.2018.01.002
- Signorini, C., Perrone, S., Sgherri, C., Ciccoli, L., Buonocore, G., Leoncini, S., et al. (2008). Plasma esterified F 2-isoprostanes and oxidative stress in newborns: role of nonprotein-bound iron. *Pediatr. Res.* 63, 287–291. doi: 10.1203/PDR.0b013e318163a1fd
- Silvers, K., Gibson, A., Russell, J., and Powers, H. (1998). Antioxidant activity, packed cell transfusions, and outcome in premature infants. *Arch. Dis. Child Fetal Neonatal Ed.* 78, F214–F219. doi: 10.1136/fn.78.3.F214
- Simonyi, A., Serfozo, P., Lehmidi, T. M., Cui, J., Gu, Z., Lubahn, D. B., et al. (2012). The neuroprotective effects of apocynin. *Front. Biosci.* 4, 2183–2193. doi: 10.2741/e535
- Small, D. M., Coombes, J. S., Bennett, N., Johnson, D. W., and Gobe, G. C. (2012). Oxidative stress, anti-oxidant therapies and chronic kidney disease. *Nephrology* 17, 311–321. doi: 10.1111/j.1440-1797.2012.01572.x
- Smith, A., Hardison, D., Bridges, B., and Pietsch, J. (2013). Red blood cell transfusion volume and mortality among patients receiving extracorporeal membrane oxygenation. *Perfusion* 28, 54–60. doi: 10.1177/0267659112457969
- Sonogo, G., Abonnenc, M., Tissot, J.-D., Prudent, M., and Lion, N. (2017). Redox proteomics and platelet activation: understanding the redox proteome to improve platelet quality for transfusion. *Int. J. Mol. Sci.* 18:E387. doi: 10.3390/ijms18020387
- Sosa, M. A., Balk, E. M., Lau, J., Liangos, O., Balakrishnan, V. S., Madias, N. E., et al. (2006). A systematic review of the effect of the Excebrane dialyser on biomarkers of lipid peroxidation. *Nephrol. Dial. Transplant.* 21, 2825–2833. doi: 10.1093/ndt/gfl376
- Steinhorn, R. H., Isham-Schopf, B., Smith, C., and Green, T. P. (1989). Hemolysis during long-term extracorporeal membrane oxygenation. *J. Pediatr.* 115, 625–630. doi: 10.1016/S0022-3476(89)80299-9
- Stevens, K. L., Black, A. L., Wells, K. M., Yeo, K. B., Steuart, R. F., Stirling, C. J., et al. (2017). Diminished Ost3-dependent N-glycosylation of the BiP nucleotide

- exchange factor S11 is an adaptive response to reductive ER stress. *Proc. Natl. Acad. Sci. U.S.A.* 114, 12489–12494. doi: 10.1073/pnas.1705641114
- Stiller, B., Lemmer, J., Merkle, F., Alexi-Meskishvili, V., Weng, Y., Hübler, M., et al. (2004). Consumption of blood products during mechanical circulatory support in children: comparison between ECMO and a pulsatile ventricular assist device. *Intensive Care Med.* 30, 1814–1820. doi: 10.1007/s00134-004-2352-z
- Stoppe, C., Spillner, J., Rossaint, R., Coburn, M., Schälte, G., Wildenhues, A., et al. (2013). Selenium blood concentrations in patients undergoing elective cardiac surgery and receiving perioperative sodium selenite. *Nutrition* 29, 158–165. doi: 10.1016/j.nut.2012.05.013
- Stuehr, D. J. (1999). Mammalian nitric oxide synthases. *Biochim. Biophys. Acta* 1411, 217–230. doi: 10.1016/S0005-2728(99)00016-X
- Suliman, H. B., Carraway, M. S., and Piantadosi, C. A. (2003). Postlipopolysaccharide oxidative damage of mitochondrial DNA. *Am. J. Respir. Crit. Care Med.* 167, 570–579. doi: 10.1164/rccm.200206-518OC
- Suresh, G., Davis, J. M., and Soll, R. (2001). Superoxide dismutase for preventing chronic lung disease in mechanically ventilated preterm infants. *Cochrane Database Syst. Rev.* 1:CD001968. doi: 10.1002/14651858.CD001968
- Swaniker, F., Kolla, S., Moler, F., Custer, J., Grams, R., Bartlett, R., et al. (2000). Extracorporeal life support outcome for 128 pediatric patients with respiratory failure. *J. Pediatr. Surg.* 35, 197–202. doi: 10.1016/S0022-3468(00)90009-5
- Szyncer-Taub, N. R., Lowery, R., Yu, S., Owens, S. T., Hirsch-Romano, J. C., and Owens, G. E. (2016). Hyperoxia is associated with poor outcomes in pediatric cardiac patients supported on venoarterial extracorporeal membrane oxygenation. *Pediatr. Crit. Care Med.* 17, 350–358. doi: 10.1097/PCC.0000000000000655
- Takeda, K., Shimada, Y., Amano, M., Sakai, T., Okada, T., and Yoshiya, I. (1984). Plasma lipid peroxides and alpha-tocopherol in critically ill patients. *Crit. Care Med.* 12, 957–959. doi: 10.1097/00003246-198411000-00007
- Takouli, L., Hadjiyannakos, D., Metaxaki, P., Sideris, V., Filiopoulos, V., Anogiati, A., et al. (2010). Vitamin E-coated cellulose acetate dialysis membrane: long-term effect on inflammation and oxidative stress. *Ren. Fail.* 32, 287–293. doi: 10.3109/08860221003615795
- Tappel, A. (1970). Biological antioxidant protection against lipid peroxidation damage. *Am. J. Clin. Nutr.* 23, 1137–1139. doi: 10.1093/ajcn/23.8.1137
- Tarakçoğlu, M., Erbağcı, A. B., Usalan, C., Deveci, R., and Kocabaş, R. (2003). Acute effect of hemodialysis on serum levels of the proinflammatory cytokines. *Mediators Inflamm.* 12, 15–19. doi: 10.1080/0962935031000096935
- Tayama, E., Hayashida, N., Akasu, K., Kosuga, T., Fukunaga, S., Akashi, H., et al. (2000). Biocompatibility of heparin-coated extracorporeal bypass circuits: new heparin bonded bioline system. *Artif. Organs* 24, 618–623. doi: 10.1046/j.1525-1594.2000.06615.x
- Thiagarajan, R. R., Barbaro, R. P., Rycus, P. T., McMullan, D. M., Conrad, S. A., Fortenberry, J. D., et al. (2017). Extracorporeal life support organization registry international report 2016. *ASAIO J.* 63, 60–67. doi: 10.1097/MAT.0000000000000475
- Thiara, A., Andersen, V., Videm, V., Mollnes, T., Svennevig, K., Hoel, T., et al. (2010). Comparable biocompatibility of Physio-and Bioline-coated cardiopulmonary bypass circuits indicated by the inflammatory response. *Perfusion* 25, 9–16. doi: 10.1177/0267659110362822
- Tin, W., and Gupta, S. (2007). Optimum oxygen therapy in preterm babies. *Arch. Dis. Child. Fetal Neonatal Ed.* 92, F143–F147. doi: 10.1136/adc.2005.092726
- Tonelli, M., Wiebe, N., Hemmelgarn, B., Klarenbach, S., Field, C., Manns, B., et al. (2009). Trace elements in hemodialysis patients: a systematic review and meta-analysis. *BMC Med.* 7:25. doi: 10.1186/1741-7015-7-25
- Trittenwein, G., Rotta, A. T., Gunnarsson, B., and Steinhorn, D. M. (1999). Lipid peroxidation during initiation of extracorporeal membrane oxygenation after hypoxia in endotoxemic rabbits. *Perfusion* 14, 49–57. doi: 10.1177/026765919901400108
- Tyson, J. E., Wright, L. L., Oh, W., Kennedy, K. A., Mele, L., Ehrenkranz, R. A., et al. (1999). Vitamin A supplementation for extremely-low-birth-weight infants. *N. Engl. J. Med.* 340, 1962–1968. doi: 10.1056/NEJM199906243402505
- Ulus, A., Aksoyok, A., Ozkan, M., Katircioğlu, S., and Basu, S. (2003). Cardiopulmonary bypass as a cause of free radical-induced oxidative stress and enhanced blood-borne isoprostanes in humans. *Free Radic. Biol. Med.* 34, 911–917. doi: 10.1016/S0891-5849(03)00030-3
- Uyar, I. S., Akpınar, M. B., Sahin, V., Uguz, A. C., Onal, S., Gonen, I., et al. (2013). Alpha lipoic acid attenuates inflammatory response during extracorporeal circulation. *Cardiovasc. J. Afr.* 24, 322–326. doi: 10.5830/CVJA-2013-067
- Varan, H. I., Dursun, B., Dursun, E., Ozben, T., and Suleymanlar, G. (2010). Acute effects of hemodialysis on oxidative stress parameters in chronic uremic patients: comparison of two dialysis membranes. *Int. J. Nephrol. Renovasc. Dis.* 3, 39–45.
- Vento, M. (2014). Oxygen supplementation in the neonatal period: changing the paradigm. *Neonatology* 105, 323–331. doi: 10.1159/000360646
- Vento, M., Escobar, J., Cernada, M., Escrig, R., and Aguar, M. (2012). The use and misuse of oxygen during the neonatal period. *Clin. Perinatol.* 39, 165–176. doi: 10.1016/j.clp.2011.12.014
- Visser, J., Labadarios, D., and Blaauw, R. (2011). Micronutrient supplementation for critically ill adults: a systematic review and meta-analysis. *Nutrition* 27, 745–758. doi: 10.1016/j.nut.2010.12.009
- Wachtfogel, Y. T., Hack, C. E., Nuijens, J., Kettner, C., Reilly, T. M., Knabb, R. M., et al. (1995). Selective kallikrein inhibitors alter human neutrophil elastase release during extracorporeal circulation. *Am. J. Physiol. Heart Circ. Physiol.* 268, H1352–H1357. doi: 10.1152/ajpheart.1995.268.3.H1352
- Walski, T., Drohomirecka, A., Bujok, J., Czernski, A., Wąż, G., Trochanowska, N., et al. (2018). Low-level light therapy protects red blood cells against oxidative stress and haemolysis during extracorporeal circulation. *Front. Physiol.* 9:647. doi: 10.3389/fphys.2018.00647
- Walski, T., Dyrda, A., Dzik, M., Chludzińska, L., Tomków, T., Mehl, J., et al. (2015). Near infrared light induces post-translational modifications of human red blood cell proteins. *Photochem. Photobiol. Sci.* 14, 2035–2045. doi: 10.1039/c5pp00203f
- Wan, S., Leclerc, J.-L., and Vincent, J.-L. (1997). Inflammatory response to cardiopulmonary bypass: mechanisms involved and possible therapeutic strategies. *Chest* 112, 676–692. doi: 10.1378/chest.112.3.676
- Wang, D., Sun, J., Solomon, S. B., Klein, H. G., and Natanson, C. (2012). Transfusion of older stored blood and risk of death: a meta-analysis. *Transfusion* 52, 1184–1195. doi: 10.1111/j.1537-2995.2011.03466.x
- Wang, S., Krawiec, C., Patel, S., Kunselman, A. R., Song, J., Lei, F., et al. (2015). Laboratory evaluation of hemolysis and systemic inflammatory response in neonatal nonpulsatile and pulsatile extracorporeal life support systems. *Artif. Organs* 39, 774–781. doi: 10.1111/aor.12466
- Wardle, S., Drury, J., Garr, R., and Weindling, A. (2002). Effect of blood transfusion on lipid peroxidation in preterm infants. *Arch. Dis. Child. Fetal Neonatal Ed.* 86, F46–F48. doi: 10.1136/fn.86.1.F46
- Wardle, S., Hughes, A., Chen, S., and Shaw, N. (2001). Randomised controlled trial of oral vitamin A supplementation in preterm infants to prevent chronic lung disease. *Arch. Dis. Child. Fetal Neonatal Ed.* 84, F9–F13. doi: 10.1136/fn.84.1.F9
- Warner, A., Bencosme, A., Healy, D., and Verme, C. (1995). Prognostic role of antioxidant enzymes in sepsis: preliminary assessment. *Clin. Chem.* 41, 867–871.
- Watanabe, N., Sakota, D., Ohuchi, K., and Takatani, S. (2007). Deformability of red blood cells and its relation to blood trauma in rotary blood pumps. *Artif. Organs* 31, 352–358. doi: 10.1111/j.1525-1594.2007.00392.x
- Weber, T. R., Connors, R. H., Tracy, T. F., Bailey, P. V., Stephens, C., and Keenan, W. (1990). Prognostic determinants in extracorporeal membrane oxygenation for respiratory failure in newborns. *Ann. Thorac. Surg.* 50, 720–723. doi: 10.1016/0003-4975(90)90669-W
- Wendel, H. P., Scheule, A. M., Eckstein, F. S., and Ziemer, G. (1999). Haemocompatibility of paediatric membrane oxygenators with heparin-coated surfaces. *Perfusion* 14, 21–28. doi: 10.1177/026765919901400104
- West, A. P., Brodsky, I. E., Rahner, C., Woo, D. K., Erdjument-Bromage, H., Tempst, P., et al. (2011). TLR signalling augments macrophage bactericidal activity through mitochondrial ROS. *Nature* 472, 476–480. doi: 10.1038/nature09973
- Wiesnack, C., Liebold, A., Philipp, A., Ritzka, M., Koppenberg, J., Birnbaum, D. E., et al. (2004). Four years' experience with a miniaturized extracorporeal circulation system and its influence on clinical outcome. *Artif. Organs* 28, 1082–1088. doi: 10.1111/j.1525-1594.2004.00030.x
- Williams, D. C., Turi, J. L., Hornik, C. P., Bonnadonna, D. K., Williford, W. L., Walczak, R. J., et al. (2015). Circuit oxygenator contributes to extracorporeal membrane oxygenation-induced hemolysis. *ASAIO J.* 61, 190–195. doi: 10.1097/MAT.0000000000000173

- Wilson, J. X. (2009). Mechanism of action of vitamin C in sepsis: ascorbate modulates redox signaling in endothelium. *Biofactors* 35, 5–13. doi: 10.1002/biof.7
- Winterbourn, C. C., Buss, I. H., Chan, T. P., Plank, L. D., Clark, M. A., and Windsor, J. A. (2000). Protein carbonyl measurements show evidence of early oxidative stress in critically ill patients. *Crit. Care Med.* 28, 143–149. doi: 10.1097/00003246-200001000-00024
- Winterbourn, C. C., and Hampton, M. B. (2008). Thiol chemistry and specificity in redox signaling. *Free Radic. Biol. Med.* 45, 549–561. doi: 10.1016/j.freeradbiomed.2008.05.004
- Witko-Sarsat, V., Friedlander, M., Capeillère-Blandin, C., Nguyen-Khoa, T., Nguyen, A. T., Zingraff, J., et al. (1996). Advanced oxidation protein products as a novel marker of oxidative stress in uremia. *Kidney Int.* 49, 1304–1313. doi: 10.1038/ki.1996.186
- Wolf, A., Remp, D. G., Kiley, J. E., and Currie, G. D. (1951). Artificial kidney function: kinetics of hemodialysis. *J. Clin. Invest.* 30, 1062–1070. doi: 10.1172/JCI102526
- Wołonciej, M., Milewska, E., and Roszkowska-Jakimiec, W. (2016). Trace elements as an activator of antioxidant enzymes. *Postepy Hig. Med. Dosw.* 70, 1483–1498. doi: 10.5604/17322693.1229074
- Wu, J. H., Marchioli, R., Silletta, M. G., Masson, S., Sellke, F. W., Libby, P., et al. (2015). Oxidative stress biomarkers and incidence of postoperative atrial fibrillation in the omega-3 fatty acids for prevention of postoperative atrial fibrillation (OPERA) Trial. *J. Am. Heart Assoc.* 4:e001886. doi: 10.1161/JAHA.115.001886
- Xia, W., Han, J., Huang, G., and Ying, W. (2010). Inflammation in ischaemic brain injury: current advances and future perspectives. *Clin. Exp. Pharmacol. Physiol.* 37, 253–258. doi: 10.1111/j.1440-1681.2009.05279.x
- Xu, J., Woods, C. R., Mora, A. L., Joodi, R., Brigham, K. L., Iyer, S., et al. (2007). Prevention of endotoxin-induced systemic response by bone marrow-derived mesenchymal stem cells in mice. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 293, L131–L141. doi: 10.1152/ajplung.00431.2006
- Yamamoto, M., Yang, G., Hong, C., Liu, J., Holle, E., Yu, X., et al. (2003). Inhibition of endogenous thioredoxin in the heart increases oxidative stress and cardiac hypertrophy. *J. Clin. Invest.* 112, 1395–1406. doi: 10.1172/JCI200317700
- Yamawaki, H., and Berk, B. C. (2006). Thioredoxin in the cardiovascular system. *J. Mol. Med.* 84, 997–1003. doi: 10.1007/s00109-006-0109-6
- Yang, C.-C., Hsu, S.-P., Wu, M.-S., Hsu, S.-M., and Chien, C.-T. (2006). Effects of vitamin C infusion and vitamin E-coated membrane on hemodialysis-induced oxidative stress. *Kidney Int.* 69, 706–714. doi: 10.1038/sj.ki.5000109
- Yang, S.-K., Xiao, L., Xu, B., Xu, X.-X., Liu, F.-Y., and Sun, L. (2014). Effects of vitamin E-coated dialyzer on oxidative stress and inflammation status in hemodialysis patients: a systematic review and meta-analysis. *Ren. Fail.* 36, 722–731. doi: 10.3109/0886022X.2014.890858
- Yin, H., Xu, L., and Porter, N. A. (2011). Free radical lipid peroxidation: mechanisms and analysis. *Chem. Rev.* 111, 5944–5972. doi: 10.1021/cr200084z
- Zakkar, M., Ascione, R., James, A., Angelini, G., and Suleiman, M. (2015a). Inflammation, oxidative stress and postoperative atrial fibrillation in cardiac surgery. *Pharmacol. Ther.* 154, 13–20. doi: 10.1016/j.pharmthera.2015.06.009
- Zakkar, M., Guida, G., Suleiman, M., and Angelini, G. D. (2015b). Cardiopulmonary bypass and oxidative stress. *Oxid. Med. Cell. Longev.* 2015:189863. doi: 10.1155/2015/189863
- Zimmerman, G. A., Mcintyre, T. M., and Prescott, S. M. (1985). Thrombin stimulates the adherence of neutrophils to human endothelial cells in vitro. *J. Clin. Invest.* 76, 2235–2246. doi: 10.1172/JCI112232
- Zusterzeel, P., Rütten, H., Roelofs, H., Peters, W., and Steegers, E. (2001). Protein carbonyls in decidua and placenta of pre-eclamptic women as markers for oxidative stress. *Placenta* 22, 213–219. doi: 10.1053/plac.2000.0606
- Zwiers, A. J., De Wildt, S. N., Hop, W. C., Dorresteijn, E. M., Giscler, S. J., Tibboel, D., et al. (2013). Acute kidney injury is a frequent complication in critically ill neonates receiving extracorporeal membrane oxygenation: a 14-year cohort study. *Crit. Care* 17:R151. doi: 10.1186/cc12830

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Raffaelli, Ghirardello, Passera, Mosca and Cavallaro. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.