

Packing subgroups in relatively hyperbolic groups

G CHRISTOPHER HRUSKA
DANIEL T WISE

We introduce the bounded packing property for a subgroup of a countable discrete group G . This property gives a finite upper bound on the number of left cosets of the subgroup that are pairwise close in G . We establish basic properties of bounded packing and give many examples; for instance, every subgroup of a countable, virtually nilpotent group has bounded packing. We explain several natural connections between bounded packing and group actions on CAT(0) cube complexes.

Our main result establishes the bounded packing of relatively quasiconvex subgroups of a relatively hyperbolic group, under mild hypotheses. As an application, we prove that relatively quasiconvex subgroups have finite height and width, properties that strongly restrict the way families of distinct conjugates of the subgroup can intersect. We prove that an infinite, nonparabolic relatively quasiconvex subgroup of a relatively hyperbolic group has finite index in its commensurator. We also prove a virtual malnormality theorem for separable, relatively quasiconvex subgroups, which is new even in the word hyperbolic case.

[20F65](#); [20F67](#), [20F69](#)

1 Introduction

In treating groups as geometric objects, it is natural to emphasize certain subgroups for special treatment, and these “quasiconvex” subgroups play a role akin to convex subspaces of a geodesic metric space. While for arbitrary groups the quasiconvexity of a subgroup is not even well-defined, for word-hyperbolic groups quasiconvexity is independent of the choice of generating system, and there is a useful theory of such subgroups which endows them with the properties of the ambient group, and then examines the favorable way in which they are embedded.

There has been substantial progress in the past few years generalizing properties of word-hyperbolic groups to properties of relatively hyperbolic groups with hypotheses on the peripheral subgroups. This paper is part of this trend where we focus on generalizations of ideas related to a certain important property of a quasiconvex subgroup. Let us first discuss the background of this property before indicating our generalizations.

An infinite quasiconvex subgroup H of a word-hyperbolic group G cannot be normal unless it is of finite index (see Short [1]). More generally, there are substantial limits on the way it intersects its conjugates. The definitive result in this direction, obtained by Gitik–Mitra–Rips–Sageev [8], states that a quasiconvex subgroup has “finite width”. This roughly means that there is an upper bound on the number of conjugates whose pairwise intersection is infinite. More precisely the *width* of H in G equals $(n - 1)$ if n is the smallest number with the property that for any n distinct right cosets g_1H, \dots, g_nH the intersection $g_iHg_i^{-1} \cap g_jHg_j^{-1}$ is finite for some i, j . It is unknown whether having finite width characterizes quasiconvexity, but it seems unlikely that one could jump from an algebraic hypothesis to a geometric conclusion in this context.

1.1 Bounded packing

The geometric mechanism lying behind the finite width conclusion is what we call “bounded packing.” In a metric space (X, d) the distance between two subsets Y and Z is the infimum of distances between points $y \in Y$ and $z \in Z$. If H is a subgroup of G , then H has *bounded packing* in G if for each D , there is a bound $M = M(D)$ on the number of distinct cosets g_iH such that $d(g_iH, g_jH) < D$ for each i, j . (A more precise statement can be found in Section 2 below.) The connection between width and the bounded packing of a κ -quasiconvex subgroup H of the δ -hyperbolic group G , is that if $g_iHg_i^{-1} \cap g_jHg_j^{-1}$ is infinite, then $d(g_iH, g_jH) < K = K(\kappa, \delta)$.

The goal of this paper is to prove a bounded packing statement in the relatively hyperbolic context, and to deduce from this an appropriate finite width consequence.

Bounded packing is the fundamental notion operating behind proofs about the widths of subgroups and appears implicitly in Gitik–Mitra–Rips–Sageev [8]. It is a natural algebraic generalization of the Finite Plane Intersection Property (FPIP) for universal covers of surfaces in 3-manifolds (see, for instance, Rubinstein–Sageev [18]). It is also the key point used by Sageev [21] to prove the finite dimensionality of the cube complex arising from Sageev’s construction applied to a codimension–1 quasiconvex subgroup of a word-hyperbolic group, as discussed in Section 3 below. In [11], the authors apply the bounded packing property to generalize Sageev’s finite dimensional cubulation result to codimension–1 relatively quasiconvex subgroups of relatively hyperbolic groups. Our main theorem about bounded packing is the following result about relatively hyperbolic groups.

Theorem 1.1 *Let H be a relatively quasiconvex subgroup of a relatively hyperbolic group G . Suppose $H \cap gPg^{-1}$ has bounded packing in gPg^{-1} for each conjugate of each peripheral subgroup P . Then H has bounded packing in G .*

This theorem generalizes a result of Gitik–Mitra–Rips–Sageev [8] on quasiconvex subgroups of a word hyperbolic group and also generalizes a result of Rubinstein–Sageev [18] on geometrically finite surfaces in 3–manifolds whose toral decomposition has only hyperbolic components.

We note that the condition about intersections with peripheral subgroups is necessary. Indeed, every group G is hyperbolic relative to itself, and in that case every subgroup of G is relatively quasiconvex.

We have collected together a number of basic properties of bounded packing in Section 2, and we state a variety of problems there for further thought. In general, we would like to know:

Question 1.2 Which countable groups have the property that all their (finitely generated) subgroups have bounded packing?

We also show in Section 3 that under mild hypotheses an action of a group G on a CAT(0) cube complex C gives rise to natural subgroups of G with bounded packing. These subgroups arise as stabilizers of hyperplanes in C .

1.2 Pairwise close spaces

Niblo–Reeves [13] proved that given n , κ , D and δ there exists $M = M(n, \kappa, D, \delta)$ such that the following thinness condition holds: If A_1, \dots, A_n are κ –quasiconvex subspaces of a δ –hyperbolic space X such that $d(A_i, A_j) < D$ for all i, j , then there is a point $x \in X$ with $d(A_i, x) < M$ for all i . This statement is also implicit in Sageev’s work in [21], and plays a fundamental role in both these papers in proving the cocompactness of the cube complex associated with a finite set of quasiconvex codimension–1 subgroups.

We generalize this statement in Proposition 7.7, which contains a slightly stronger conclusion than the following assertion that there is either a point or a peripheral coset nearby.

Proposition 1.3 *Let G be a relatively hyperbolic group. Choose positive constants κ and D and an integer $n \geq 1$. Let \mathcal{A} be an arbitrary set of κ –relatively quasiconvex subspaces of G such that for each $A, A' \in \mathcal{A}$ the distance $d(A, A')$ is less than D . Then there is a constant $M = M(\kappa, D, n)$ so that at least one of the following holds.*

- (1) *For every set $\{A_1, \dots, A_n\}$ of n distinct elements of \mathcal{A} , there is a point $x \in X$ such that $d(x, A_i) < M$ for each i .*
- (2) *There is a peripheral coset gP such that $d(gP, A) < M$ for all $A \in \mathcal{A}$.*

Proposition 7.7 plays a significant role in our proof of Theorem 1.1.

1.3 Bounded height and width

For relatively hyperbolic groups, the “small” subgroups are either “elliptic,” meaning finite, or “parabolic,” which means conjugate into a peripheral subgroup. In this context, we therefore redefine the *height* of H in G to be $(n - 1)$ if n is the smallest number with the property that for any n distinct left cosets g_1H, \dots, g_nH , the intersection $\bigcap_{1 \leq i \leq n} g_iHg_i^{-1}$ is elliptic or parabolic. Similarly, the *width* of H in G equals $(n - 1)$ if n is the smallest number with the property that for any n distinct left cosets g_1H, \dots, g_nH the intersection $g_iHg_i^{-1} \cap g_jHg_j^{-1}$ is elliptic or parabolic for some i, j .

The following is our main theorem about finite height and width:

Theorem 1.4 *Let G be a relatively hyperbolic group, and let H be a relatively quasiconvex subgroup. Then H has finite height.*

Suppose, furthermore that $H \cap gPg^{-1}$ has bounded packing in gPg^{-1} for each conjugate of each peripheral subgroup P . Then H has finite width.

We hope that further work in this area will resolve the following:

Problem 1.5 *Let G be relatively hyperbolic. Does every relatively quasiconvex subgroup of G have finite width?*

If H is a subgroup of G , the *commensurator* of H in G is the subgroup of all $g \in G$ such that $H \cap gHg^{-1}$ has finite index in both H and gHg^{-1} . We prove the following result on commensurators of relatively quasiconvex subgroups using ideas related to [Theorem 1.4](#).

Theorem 1.6 *Let H be a relatively quasiconvex subgroup of a relatively hyperbolic group G . Suppose H is infinite and nonparabolic. Then H has finite index in its commensurator.*

We use similar techniques to show that a separable (relatively) quasiconvex subgroup H of a (relatively) hyperbolic group G is (relatively) malnormal in a finite index subgroup K of G (see [Theorem 9.3](#)). This result is new even in the hyperbolic case.

1.4 Overview

In [Section 2](#) we define the notion of bounded packing and prove a number of basic results about this concept. We also give a number of examples and collect several problems.

In [Section 3](#) we discuss the relation between bounded packing and actions on CAT(0) cube complexes. By a result of Sageev, a codimension–1 subgroup $H < G$ with bounded packing gives rise to an action of G on a finite dimensional CAT(0) cube complex. In this section, we give a partial converse to Sageev’s result.

In [Section 4](#) we prove the bounded width theorem of Gitik–Mitra–Rips–Sageev [8] in the word hyperbolic setting. The reader is advised to understand this proof first, as our relatively hyperbolic generalization will follow it closely, albeit with certain aspects substantially more complicated.

In [Section 5](#) we recount essential facts about the geometry of the Cayley graph of a relatively hyperbolic group with a finite generating set that were established by Druţu–Sapir in [5]. We use these results to prove two auxiliary results about triangles and quadrilaterals with the property that each side lies near a peripheral subspace.

In [Section 6](#) we discuss the geometry of relative Cayley graphs, concentrating on the interplay between the usual (proper) metric and the relative metric as applied to thinness conditions.

In [Section 7](#) we analyze collections of relatively quasiconvex subspaces in a relatively hyperbolic group. In particular, we prove [Proposition 7.7](#), an important technical result underlying this work that was described above.

In [Section 8](#) we prove [Theorem 1.4](#), our main result on bounded packing and bounded width for a relatively quasiconvex subgroup of a relatively hyperbolic group. We also prove [Theorem 1.6](#) on the commensurator of a relatively quasiconvex subgroup.

[Section 9](#) contains a brief proof of the relative malnormality result [Theorem 9.3](#) mentioned above, using results from the previous section.

Acknowledgements The first author’s contribution to this material is based upon work supported by the National Science Foundation under Grant Nos. DMS-0505659, DMS-0754254 and DMS-0808809. The second author is supported by grants from NSERC.

2 Bounded packing: First properties and examples

Definition 2.1 (Bounded packing) Let G be a discrete group with a left invariant metric d . Suppose also that d is *proper* in the sense that every metric ball is finite. A subgroup H has *bounded packing* in G (with respect to d) if, for each constant D , there is a number $N = N(G, H, D)$ so that for any collection of N distinct cosets gH in G , at least two are separated by a distance of at least D .

A discrete group admitting a proper metric is necessarily countable. On the other hand, if G is finitely generated then the word metric for any finite generating set is a proper, equivariant metric. Restricting the word metric on G to a subgroup H gives a proper, left invariant metric on H . Since every countable group is a subgroup of a finitely generated group, it follows that a discrete group G admits a proper, left invariant metric if and only if G is countable.

Lemma 2.2 *If G is countable, then bounded packing of a subgroup H in G is independent of the choice of proper, left invariant metric on G . In other words, if d_1, d_2 are proper, left invariant metrics on G , then H has bounded packing in G with respect to d_1 if and only if it has bounded packing with respect to d_2 .*

Proof Define $\rho: \mathbb{N} \rightarrow \mathbb{N}$ so that $\rho(n)$ is the supremum of $d_1(1, g)$ over all g with $d_2(1, g) < n$. Note that there are only finitely many such g since d_2 is proper, so $\rho(n)$ is finite for each n . By the left invariance of d_1 and d_2 , it follows that

$$d_1(g_1, g_2) \leq \rho(d_2(g_1, g_2))$$

for all $g_1, g_2 \in G$.

Now if \mathcal{H} is a collection of left cosets of H with pairwise d_2 -distances at most D , their pairwise d_1 -distances are at most $\rho(D)$. Thus if H has bounded packing with respect to d_1 , it also has bounded packing with respect to d_2 . \square

The following result follows easily from the definition of bounded packing.

Lemma 2.3 *Any finite index subgroup K of a countable group G has bounded packing in G .*

Proof The definition of bounded packing is vacuously satisfied once N is larger than $[G : H]$, since there are no collections of N distinct cosets of H in G . \square

Lemma 2.4 *Suppose that $H \leq K \leq G$ and G is countable.*

- (1) *If H has bounded packing in G then H has bounded packing in K .*
- (2) *If H has bounded packing in K , and K has bounded packing in G then H has bounded packing in G .*

Proof We may assume the metric on K is a restriction of the metric on G , so that (1) is immediate.

Let \mathcal{H} is a collection of left cosets of H in G with pairwise distances at most D . The bounded packing of H in K bounds the number of elements of \mathcal{H} that lie in each left coset gK . Furthermore, the bounded packing of K in G bounds the number of cosets gK that contain elements of \mathcal{H} . Thus the size of \mathcal{H} is bounded in terms of D , establishing (2). \square

Proposition 2.5 (Commensurability) *Let G be a countable group.*

- (1) *Suppose $H \leq K \leq G$ and $[K : H] < \infty$. Then H has bounded packing in G if and only if K has bounded packing in G .*
- (2) *Suppose $H \leq K \leq G$ and $[G : K] < \infty$. Then H has bounded packing in K if and only if H has bounded packing in G .*
- (3) *Suppose $H, K \leq G$ and $[G : K] < \infty$. Then $H \cap K$ has bounded packing in K if and only if H has bounded packing in G .*

Proof Let d be a proper, left invariant metric on G . To see (1) suppose $[K : H] < \infty$. Then there is a constant C such that in the metric d each left coset gK is at most a Hausdorff distance C from a left coset gH (the choice of g is irrelevant). Suppose H has bounded packing in G . Let \mathcal{K} be a collection of left cosets of K with pairwise distances at most D . Replacing each coset gK with a corresponding coset gH gives a collection \mathcal{H} of left cosets of H with the same cardinality as \mathcal{K} and with pairwise distances at most $D + 2C$. The bounded packing of H in G bounds the size of \mathcal{H} , and hence also the size of \mathcal{K} , as a function of D . Thus K has bounded packing in G .

Conversely suppose K has bounded packing in G . By Lemma 2.3, we know that H has bounded packing in K . Therefore H has bounded packing in G by Lemma 2.4(2), completing the proof of (1).

Assertion (2) follows immediately from Lemmas 2.3 and 2.4.

Suppose now that $H, K \leq G$ and $[G : K] \leq \infty$. Then $H \cap K$ has finite index in H . Observe that (3) is an immediate consequence of (1) and (2), since $H \cap K$ has bounded packing in K if and only if $H \cap K$ has bounded packing in G if and only if H has bounded packing in G . \square

Corollary 2.6 *If G is countable, any finite subgroup $K \leq G$ has bounded packing in G .*

Proof Fix a proper, left invariant metric d on G . By Proposition 2.5(1) it suffices to show that the trivial subgroup has bounded packing in G . A left coset of the trivial subgroup is just a single element of G . If \mathcal{H} is any collection of pairwise D -close

left cosets of the trivial subgroup, (in other words, elements of G) then \mathcal{H} is a subset of the ball of radius D centered at any of its elements. The number of elements in such a ball is finite since d is proper, and this number depends only on D since d is left-invariant. □

Proposition 2.7 *Suppose H and K have bounded packing in a countable group G . Then $H \cap K$ has bounded packing in G .*

Proof Fix a proper, left invariant metric d on G . By Lemmas 2.4(2) and 2.2 it suffices to show that $L := H \cap K$ has bounded packing in H with respect to d . Let \mathcal{L} be a set of left cosets of L in H whose pairwise d -distances are at most D . If hL and $h'L$ are distinct cosets of L in H then $hL = H \cap hK$ and $h'L = H \cap h'K$ for distinct cosets hK and $h'K$ in G . Thus \mathcal{L} is in one-to-one correspondence with a set $\hat{\mathcal{L}}$ of left cosets of K in G whose pairwise distances are at most D . But the bounded packing of K in G bounds the size of $\hat{\mathcal{L}}$ as a function of D . □

Lemma 2.8 *Let $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ be a short exact sequence of countable groups. Let H be a subgroup of G which projects to the subgroup \bar{H} of Q . Then \bar{H} has bounded packing in Q if and only if HN has bounded packing in G .*

Proof For each $g \in G$, let \bar{g} denote the image of g in Q . Fix a proper, left invariant metric d on G . Define $\bar{d}: Q \rightarrow \mathbb{R}$ so that $\bar{d}(q_1, q_2)$ is the infimum of $d(g_1, g_2)$ over the set of all g_1, g_2 with $q_i = \bar{g}_i$. A straightforward argument shows that \bar{d} is a proper, left invariant metric on Q .

The projection $G \rightarrow Q$ induces a one-to-one correspondence between left cosets of HN in G and left cosets of \bar{H} in Q . It is clear that $d(g_1, g_2) \geq \bar{d}(\bar{g}_1, \bar{g}_2)$ for all $g_1, g_2 \in G$. Therefore

$$d(xHN, yHN) \geq \bar{d}(\bar{x}\bar{H}, \bar{y}\bar{H})$$

for all $x, y \in G$.

Conversely, any element $x \in Q$ with $\bar{d}(x, \bar{1}) = a$ lifts to an element $y \in G$ with $d(y, 1) = a$. If $\bar{d}(\bar{g}_1\bar{H}, \bar{g}_2\bar{H}) = n$ then there is an element $x \in Q$ with $\bar{d}(x, \bar{1}) = a$ such that $\bar{g}_1\bar{H}\bar{w}$ intersects $\bar{g}_2\bar{H}$. If y is a lift of x with $d(y, 1) = a$, then $g_1HyN = g_1HNy$ intersects g_2HN , so $d(g_1HN, g_2HN) \leq a$.

Therefore the distance between left cosets of HN in G is equal to the distance between corresponding left cosets of \bar{H} in Q . Evidently HN has bounded packing in G if and only if \bar{H} has bounded packing in Q . □

Corollary 2.9 *Each normal subgroup N of a countable group G has bounded packing.*

Proof Apply [Lemma 2.8](#) with H equal to the trivial subgroup. □

Remark 2.10 In fact, a normal subgroup N has the stronger property that for each D there exists $M = M(D)$ such that at most M left cosets gN satisfy $d(N, gN) < D$.

Corollary 2.11 *Every subgroup of a countable abelian group has bounded packing.* □

Theorem 2.12 *Let N be a countable, virtually nilpotent group. Then each subgroup of N has bounded packing in N .*

We note that this can be also be proven using either [Lemma 2.17](#) or [Corollary 2.15](#).

Proof By [Proposition 2.5\(3\)](#) it suffices to prove the theorem when N is a countable nilpotent group. The theorem is obvious if N is the trivial group. We proceed by induction on the length of the lower central series. Let H be a subgroup of N . Let Z denote the center of N . Let $\bar{N} := N/Z$ and let \bar{H} be the image of H in \bar{N} . By induction, \bar{H} has bounded packing in \bar{N} . Therefore by [Lemma 2.8](#), HZ has bounded packing in N . Because Z is central, H is normal in HZ and so H has bounded packing in HZ by [Corollary 2.9](#). Finally, by [Lemma 2.4\(2\)](#), H has bounded packing in G since, H has bounded packing in HZ and HZ has bounded packing in G . □

Question 2.13 Let S be a solvable group. Does every subgroup of S have bounded packing in S ?

It seems too much to expect an affirmative answer to [Question 2.13](#), however we do expect the following to hold:

Conjecture 2.14 *Let P be virtually polycyclic. Then each subgroup of P has bounded packing in P .*

The following is an immediate consequence of [Corollary 2.9](#) together with [Lemma 2.4\(2\)](#).

Corollary 2.15 *Let H be a subgroup of a countable group G . If H is subnormal then H has bounded packing in G .* □

Example 2.16 For $n \neq 0$, let $\text{BS}(1, n)$ be the group presented by $\langle a, t \mid tat^{-1} = a^n \rangle$. It is well-known that for $n \neq \pm 1$, the subgroup $\langle a \rangle$ is not quasi-isometrically embedded; in fact, it is exponentially distorted. Nevertheless $\langle a \rangle$ has bounded packing in $\text{BS}(1, n)$. Indeed, there is a short exact sequence $1 \rightarrow A \rightarrow \text{BS}(1, n) \rightarrow \mathbb{Z} \rightarrow 1$ with $\langle a \rangle \leq A$ such that A is isomorphic to $\mathbb{Z}[1/n]$. Since A is abelian, $\langle a \rangle$ is subnormal in $\text{BS}(1, n)$ and hence has bounded packing by [Corollary 2.15](#).

It is interesting to note that A is not finitely generated. (Indeed, A is locally cyclic but not itself cyclic.) Thus consideration of bounded packing in non-finitely generated groups can give us useful information about finitely generated subgroups of a finitely generated group.

As the following Lemma shows, the group $F_2 \times \mathbb{Z}$ has bounded packing with respect to each of its finitely generated subgroups.

Lemma 2.17 *Let $1 \rightarrow C \rightarrow G \rightarrow Q \rightarrow 1$ be a short exact sequence with C central in the countable group G . If every (finitely generated) subgroup of Q has bounded packing then every (finitely generated) subgroup of G has bounded packing.*

Proof Let H be a [finitely generated] subgroup of G . By hypothesis, its image \bar{H} has bounded packing in Q . Now H has bounded packing in CH by [Corollary 2.9](#), and CH has bounded packing in G by [Lemma 2.8](#). Therefore H has bounded packing in G by [Lemma 2.4](#). \square

Example 2.18 Let M be a Seifert-fibered 3-manifold. Then there is a short exact sequence $1 \rightarrow \mathbb{Z} \rightarrow \pi_1 M \rightarrow \pi_1 S \rightarrow 1$, where S is a 2-dimensional orbihedron. Since every finitely generated subgroup of $\pi_1 S$ has bounded packing (by local quasiconvexity and [Theorem 4.8](#) below) it follows from [Lemma 2.17](#) that every finitely generated subgroup of $\pi_1 M$ has bounded packing.

Let us now examine subgroups that do not have bounded packing. At present surprisingly few such examples are known. In fact, the authors know of only one basic example, due to Rubinstein–Wang. The group is the fundamental group of a 3-dimensional graph manifold; that is, a manifold formed by gluing Seifert fibered manifolds together along tori. In light of the preceding example, we see that the bounded packing property is not preserved by very simple graphs of groups.

Example 2.19 Rubinstein–Wang [19] constructed a finitely generated surface subgroup of the fundamental group of a graph manifold that doesn't have bounded packing. Their example is an immersed π_1 -injective surface S in a 3-dimensional graph

manifold M . Let $\tilde{S} \rightarrow \tilde{M}$ denote a lift of $S \rightarrow M$ to a map between the universal covers. The various lifts correspond to translates $g\tilde{S}$ where g varies over the various cosets $g\pi_1 S$. Remarkably, any two such lifts $g_i\tilde{S}$ and $g_j\tilde{S}$ intersect each other. Consequently any two cosets $g_i\pi_1 S$ and $g_j\pi_1 S$ are a uniformly bounded distance from each other in $\pi_1 M$.

It seems reasonable to expect subgroups without bounded packing to appear in the fundamental groups of other graph manifolds. The following problem is a reasonable test case for this phenomenon.

Problem 2.20 Determine exactly which subgroups of the following group G have bounded packing:

$$G := \langle a, b, c, d \mid [a, b], [b, c], [c, d] \rangle$$

(The group G is the fundamental group of a graph manifold.)

Since there are subgroups of countable groups without bounded packing, it follows from Lemma 2.8 that (non-finitely generated) subgroups of the free group F_2 do not always have bounded packing. Notice that, since finitely generated subgroups of F_2 are quasiconvex, Theorem 4.8 below implies that finitely generated subgroups of F_2 always have bounded packing.

Rips gave the following construction in [16]:

Proposition 2.21 Let Q be a finitely presented group. Then there exists a short exact sequence $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ such that N is finitely generated, and G is a finitely presented $C'(\frac{1}{6})$ group.

Example 2.22 Using Rips’s construction, we will construct a word hyperbolic group G and a finitely generated subgroup K such that K does not have bounded packing in G . Let Q be a finitely presented group with a subgroup H that fails to have bounded packing in Q . Let $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ be the short exact sequence from Proposition 2.21. Let K be the preimage of H in G , and note that K is finitely generated since both N and H are. By Lemma 2.8, K does not have bounded packing in G . Observe that G is word hyperbolic, since it is finitely presented and $C'(\frac{1}{6})$.

Remark 2.23 We can choose H to be a codimension-1 subgroup without bounded packing to make the group G in Example 2.22 have an exotic action on an infinite dimensional cube complex.

Problem 2.24 Give an example of a cyclic subgroup Z of a finitely generated group G such that Z does not have bounded packing in G .

Question 2.25 Are there finitely generated subgroups of $F_2 \times F_2$ without bounded packing?

Question 2.26 Let $1 \rightarrow N \rightarrow G \rightarrow A \rightarrow 1$ be an extension of finitely generated groups. Suppose every subgroup of N has bounded packing in N , and A is abelian. Does every subgroup of G have bounded packing in G ?

Problem 2.27 Let H be a retract of the finitely generated group G . Does H have bounded packing in G ?

3 Bounded packing and actions on CAT(0) cube complexes

Suppose G is a group with a finite generating set S . A subgroup $H < G$ is *codimension-1* if there is a constant $L > 0$ so that the L -neighborhood of H in $\text{Cayley}(G, S)$ has at least two “deep” complementary components. A complementary component is *deep* if it contains elements whose distance from H is arbitrarily large.

A theorem of Sageev [20], together with a result proved independently by Gerasimov [7] and Niblo–Roller [14], shows that a group G with a codimension-1 subgroup H has an action on a CAT(0) cube complex C without a global fixed point. All hyperplanes in C lie in a single orbit under the action of G , and H is the stabilizer of some hyperplane Λ . The dimension of C is the maximal size of a set of pairwise transverse hyperplanes. The relation to bounded packing is the following basic fact, observed by Sageev [21]: If H is finitely generated and hyperplanes $g_0\Lambda$ and $g_1\Lambda$ in C are transverse, then $d_S(g_0H, g_1H) < M$ for some universal constant M .

Corollary 3.1 (Sageev) *Suppose H is a finitely generated codimension-1 subgroup of a finitely generated group G . If H has bounded packing in G , then the corresponding CAT(0) cube complex C is finite dimensional.*

The goal of this section is to prove the following converse, of sorts, to the preceding result.

Theorem 3.2 *Suppose the countable discrete group G acts on a CAT(0) cube complex C , and $H \leq G$ is the stabilizer of a hyperplane Λ .*

- (1) *If C is locally finite, then H has bounded packing in G .*
- (2) *Let $V(\Lambda)$ be the set of vertices incident to the edges that cross Λ . If G acts metrically properly on C , and H acts on $V(\Lambda)$ with only finitely many orbits of vertices, then H has bounded packing in G .*

In a sense this theorem is optimal, as shown by the following example.

Example 3.3 Recall that [Example 2.19](#) gives a group G and a subgroup H such that H does not have bounded packing in G . The group H is the fundamental group of an immersed surface S in a 3-dimensional graph manifold M with $G = \pi_1(M)$. Since H is a codimension-1 subgroup of G , it follows that G has an action on a CAT(0) cube complex C such that H is the stabilizer of a hyperplane Λ . Condition (1) fails resolutely since C is an infinite dimensional cube, as any two translates of the surface S cross. The authors believe failure of (2) can be traced to a failure of the cocompactness of the action of H on $V(\Lambda)$. Indeed even if the action of G on C is not proper, the authors believe this can be remedied with the addition of further codimension-1 subgroups leading to a metrically proper action on a new cube complex C' . The action of H on the new hyperplane Λ' would necessarily fail to be cocompact by [Theorem 3.2](#), as H does not have bounded packing in G .

[Theorem 3.2](#) follows from several results about convex subsets of a CAT(0) cube complex C . Let C^1 denote the 1-skeleton of a CAT(0) cube complex with its induced path metric d . It is well-known that the metric d on the vertices coincides with the “wall metric.” That is to say, $d(v, w)$ is equal to the number of hyperplanes separating v from w . The interval $[v, w]$ is the set of all vertices u that lie on some d -geodesic from v to w . A subset $S \subseteq C^0$ is d -convex if every interval connecting two elements of S is contained in S .

We also find it useful to consider the graph C^Δ obtained from C^1 by including an edge between two vertices if they lie in a common cube. Let d^Δ be the path metric on C^Δ .

Lemma 3.4 *Let C be a CAT(0) cube complex, and let $[r, s]$ and $[t, u]$ be intervals such that t and u lie in the 1-neighborhood of $[r, s]$ in C^Δ . Then each vertex of $[t, u]$ lies in the 1-neighborhood of $[r, s]$ in C^Δ as well.*

Proof It is enough to show that any geodesic c from t to u lies in the 1-neighborhood of $[r, s]$ in C^Δ if its endpoints lie in the 1-neighborhood of $[r, s]$ in C^Δ . If $d(t, u) \leq 1$, the result is trivial, so we induct on $\ell := d(t, u)$ for $\ell \geq 2$.

By hypothesis, there is a cube A containing t and intersecting $[r, s]$. Let us choose A minimal with this property. Then the hyperplanes $\Lambda_1, \dots, \Lambda_n$ separating t from $[r, s]$ are the same as the hyperplanes transverse to A . (In particular, A is an n -cube.)

Let v be the vertex of c adjacent to t , and let Π be the hyperplane separating v from t . If $\Pi = \Lambda_i$ for some i , then v is a vertex of the cube A , and we are done by induction.

If Π is transverse with Λ_i for all i , then the collection $\{\Pi, \Lambda_1, \dots, \Lambda_n\}$ is pairwise transverse. Since Π and the Λ_i are each transverse to edges incident to t , there is an $(n+1)$ -cube A' containing A as a subcube, such that Π and each Λ_i are transverse to A' . Since A intersects $[r, s]$ and A' contains v , the result follows from the inductive hypothesis.

The only remaining possibility is that Π is nested with some $\Lambda_i \neq \Pi$. Then Λ_i separates $[r, s]$ from $\{t, v\}$ and Π separates $[r, s] \cup \{t\}$ from $\{v\}$. In this case, any path in C^1 from v to $[r, s]$ must cross Π first and then Λ_i . Since c is a geodesic in the wall metric, it crosses Π only once, so Π lies in the pairwise transverse family $\{\Pi_1, \dots, \Pi_m\}$ of hyperplanes separating u from $[r, s]$, corresponding to a minimal cube B that contains u and intersects $[r, s]$. But then Λ_i is also among the Π_j , contradicting our assumption that Λ_i and Π are nested. \square

Corollary 3.5 *Let C be a CAT(0) cube complex, and let K be a d -convex subset of C^0 . For each $n \geq 0$, the n -neighborhood of K in C^Δ is also d -convex.*

Proof It is clearly sufficient to prove the result when $n = 1$, since the general result follows by induction on n . But the case $n = 1$ is an immediate consequence of [Lemma 3.4](#). \square

A collection of subsets \mathcal{F} of a space has the *Helly property* if, whenever $\mathcal{F}_0 \subseteq \mathcal{F}$ is a finite collection of pairwise intersecting subsets, the total intersection $\bigcap_{F \in \mathcal{F}_0} F$ is nonempty.

Corollary 3.6 *If C is any CAT(0) cube complex, then the family of d -convex subsets of C^0 has the Helly property.*

Proof It has been shown by Gerasimov [\[7\]](#), Roller [\[17\]](#) and Chepoi [\[4\]](#), independently, that the 1-skeleton of a CAT(0) cube complex is a median graph, meaning that for any three vertices x , y and z , the intervals $[x, y]$, $[y, z]$ and $[x, z]$ have triple intersection consisting of a single vertex. In a median graph, it is clear from the definition that any collection of three pairwise intersecting convex sets has a nonempty triple intersection. The Helly property now follows by an elementary induction argument. \square

The previous result generalizes a result of Bandelt and van de Vel [\[2\]](#) in the setting of median graphs stating that the collection of all balls in C^Δ has the Helly property. Note that d^Δ -balls are d -convex by [Corollary 3.5](#). Graphs in which the family of balls has the Helly property are known as *Helly graphs*.

We also record the following well-known fact about hyperplanes, which we derive from results above.

Proposition 3.7 *Let Λ be a hyperplane in a CAT(0) cube complex, and let $V(\Lambda)$ be the set of vertices of the edges crossing Λ . Then $V(\Lambda)$ is d -convex.*

Proof Each halfspace of Λ is clearly d -convex. But $V(\Lambda)$ is the set of points within a d^Δ -distance 1 of both halfspaces of Λ . Since d^Δ -neighborhoods of d -convex sets are convex and intersections of d -convex sets are d -convex, the result is obvious. \square

We are now ready to prove [Theorem 3.2](#).

Proof of Theorem 3.2 Suppose H stabilizes the hyperplane Λ in C . Let $V(\Lambda)$ be the convex set of vertices of the edges that cross Λ . Fix a proper, left invariant metric d_G on G and a constant $D > 0$. Suppose \mathcal{H} is a finite collection of cosets of H whose pairwise distances in G are at most D .

Choose a basepoint $x \in V(\Lambda)$, and let

$$D' := \sup d^\Delta(x, g(x))$$

where the supremum ranges over the finite set of group elements whose d_G -distance from 1 is at most D . If cosets gH and $g'H$ are at a distance at most D in G , then the sets $gH(x) \subset V(g\Lambda)$ and $g'H(x) \subset V(g'\Lambda)$ are at a d^Δ -distance at most D' . Therefore the translates $V(g\Lambda)$ and $V(g'\Lambda)$ are at a d^Δ -distance at most D' as well. It follows immediately that the D' -neighborhoods of the translates $V(g\Lambda)$ are pairwise intersecting d -convex sets in C . Hence there is a point p within a d^Δ -distance D' of every $V(g\Lambda)$.

If C is uniformly locally finite, then there is a uniform bound on the number of hyperplanes intersecting any d^Δ -ball of radius D' in C . Hence the cardinality of \mathcal{H} is bounded as well.

Now suppose G acts metrically properly on C , and H acts on $V(\Lambda)$ with a finite quotient. Then $V(\Lambda)$ lies in the R -neighborhood of the orbit Hx in C^Δ for some $R > 0$. Thus the d^Δ -ball centered at p with radius $D' + R$ intersects $gH(x)$ for each coset $gH \in \mathcal{H}$. Since G acts metrically properly on C , there is a finite upper bound on the size of \mathcal{H} , as desired. \square

4 Bounded packing in hyperbolic groups

In this section, we give a self-contained proof in the word hyperbolic setting that quasiconvex subgroups have bounded packing.

Definition 4.1 Let $H \leq G$ be a subgroup. The *height* of H in G , denoted $\text{height}_G(H)$, is the maximal number of distinct cosets $g_i H$ so that the intersection $\bigcap_i g_i H g_i^{-1}$ is infinite. The *width* of H in G , denoted $\text{width}_G(H)$, is the maximal number of distinct cosets $g_i H$ so that for all i, j the intersection $g_i H g_i^{-1} \cap g_j H g_j^{-1}$ is infinite. In case there is no maximum then we say that the height/width is infinite.

Finally, note that the height and width of a finite subgroup are both equal to zero.

Lemma 4.2 Suppose $H \leq G$ has height $0 < n < \infty$ in G . Choose $g \in G$ so that $gH \neq H$, and let $K := H \cap gHg^{-1}$. Then $\text{height}_H(K) < n$.

Proof Choose $h_1, \dots, h_n \in H$ so that the cosets $h_i K$ are distinct. We will show that the intersection $\bigcap h_i K h_i^{-1}$ is finite. Note that

$$\bigcap h_i K h_i^{-1} = \bigcap h_i (H \cap gHg^{-1}) h_i^{-1} = H \cap \left(\bigcap h_i gHg^{-1} h_i^{-1} \right)$$

is an intersection of $n + 1$ conjugates of H in G . Since $\text{height}_G(H) = n$, it suffices to show that the elements $1, h_1 g, \dots, h_n g$ represent distinct left cosets of H in G .

First note that $H \neq h_i g H$ since $H \neq g H$. Now suppose two cosets $h_i g H$ and $h_j g H$ are equal. Then we have $g H = h_i^{-1} h_j g H$, which implies that $h_i^{-1} h_j \in g H g^{-1}$. But $h_i^{-1} h_j \in H$ as well, so $h_i^{-1} h_j \in H \cap g H g^{-1} = K$. Thus $h_i K = h_j K$, and we must have $i = j$. □

Lemma 4.3 Let Y be a κ -quasiconvex subspace of a δ -hyperbolic space X . Then there exists $\kappa' = \kappa'(\kappa, \delta)$ so that any geodesic line A in a finite neighborhood of Y lies in a κ' -neighborhood of Y .

Proof Let c be an arbitrary line in a finite neighborhood of Y . Then the endpoints c^+ and c^- of c lie in the limit set of Y . Let c' be a line connecting c^+ and c^- that is a limit of geodesic segments connecting points of Y . In a δ -hyperbolic space, two geodesic lines with the same endpoints at infinity are at a Hausdorff distance at most 2δ . But by quasiconvexity, $c' \subseteq \mathcal{N}_\kappa(Y)$, so $c \subseteq \mathcal{N}_{\kappa+2\delta}(Y)$. □

Lemma 4.4 Let H be a κ -quasiconvex subgroup of a δ -hyperbolic group G . Then $\text{height}_G(H)$ is finite.

Proof Let $g_1 H, \dots, g_n H$ be distinct cosets in G , and suppose $\bigcap g_i H g_i^{-1}$ contains an infinite order (hyperbolic) element x . Let A be a geodesic axis for x . Then A lies in a finite neighborhood of each coset $g_i H$, so by Lemma 4.3 the size of these neighborhoods is uniformly bounded by κ' . In particular, each coset $g_i H$ intersects the ball of radius κ' about some point $a \in A$. But there is a uniform bound N on the number of cosets of H intersecting any metric ball of radius κ' . □

Lemma 4.5 *Let G have a finite generating set \mathcal{A} , and suppose xH and yK are arbitrary left cosets of subgroups of G . For each constant L there is a constant $L' = L'(G, \mathcal{A}, xH, yK)$ so that in G with the word metric $d_{\mathcal{A}}$ we have:*

$$\mathcal{N}_L(xH) \cap \mathcal{N}_L(yK) \subseteq \mathcal{N}_{L'}(xHx^{-1} \cap yKy^{-1})$$

Proof If there is no such L' , then there is a sequence (z_i) in G so that z_i is in the L -neighborhood of both xH and yK , but $i < d(z_i, xHx^{-1} \cap yKy^{-1})$ for each i . It follows that $z_i = xh_i p_i = yk_i q_i$ for some $h_i \in H, k_i \in K$ and $p_i, q_i \in G$ with $|p_i|_{\mathcal{A}}$ and $|q_i|_{\mathcal{A}}$ both less than L . Passing to a subsequence if necessary, we may assume that p_i and q_i are constants p and q , so that for each i we have $z_i = xh_i p = yk_i q$. Therefore

$$z_i z_1^{-1} = xh_i h_1^{-1} x^{-1} = yk_i k_1^{-1} y^{-1} \in xHx^{-1} \cap yKy^{-1}.$$

It follows that the distance between z_i and $xHx^{-1} \cap yKy^{-1}$ is at most $|z_1|_{\mathcal{A}}$ for all i , contradicting our choice of (x_i) . □

The following result follows immediately by induction from [Lemma 4.5](#).

Corollary 4.6 *Let G be a finitely generated group with a fixed generating set \mathcal{A} . For each finite set of cosets $g_1 H_1, \dots, g_n H_n$ and each $M > 0$ there is a constant $M' > 0$ such that:*

$$\mathcal{N}_M(g_1 H_1) \cap \dots \cap \mathcal{N}_M(g_n H_n) \subseteq \mathcal{N}_{M'}(g_1 H_1 g_1^{-1} \cap \dots \cap g_n H_n g_n^{-1}) \quad \square$$

Lemma 4.7 *Let X be a δ -hyperbolic space, and let Y be a κ -quasiconvex subspace. For any $\xi > 0$ there is an $\eta = \eta(\delta, \kappa, \xi)$ so that the ξ -neighborhood of Y is η -quasiconvex.*

Proof Choose points $x, y \in \mathcal{N}_{\xi}(Y)$, and $z, w \in Y$ so that $d(x, z)$ and $d(y, w)$ are less than ξ . Since quadrilaterals in X are 2δ -thin, the geodesic $[x, y]$ lies in the 2δ -neighborhood of $[x, z] \cup [z, w] \cup [w, y]$. But this piecewise geodesic lies in the $(\kappa + \eta)$ -neighborhood of Y . □

Theorem 4.8 (cf Gitik–Mitra–Rips–Sageev [\[8\]](#)) *Let H be a quasiconvex subgroup of a δ -hyperbolic group G . Then H has bounded packing in G .*

Before giving the proof of [Theorem 4.8](#), we record the following:

Corollary 4.9 (Gitik–Mitra–Rips–Sageev [\[8\]](#)) *If H is a quasiconvex subgroup of a δ -hyperbolic group G , then $\text{width}_G(H)$ is finite.*

Proof As in [Lemma 4.4](#), apply [Lemma 4.3](#) to see that any two conjugates with infinite intersection have cosets uniformly close together. □

Proof of Theorem 4.8 Fix finite generating sets \mathcal{A} and \mathcal{B} for G and H respectively. By Lemma 4.4, we know that $\text{height}_G(H)$ is finite. We will prove the theorem by induction on height. The case where H has height zero is easy, since in this case H is finite and the result follows from Corollary 2.6.

Now assume by induction that the theorem holds for every hyperbolic group G' and quasiconvex subgroup H' with $\text{height}_{G'}(H') < \text{height}_G(H)$. Let \mathcal{H} be a set of left cosets gH whose pairwise distances are at most D . Our goal is to show that \mathcal{H} is finite and to bound the cardinality of \mathcal{H} as a function of D . Translating \mathcal{H} if necessary, we may assume that $H \in \mathcal{H}$. Observe that if $d(gH, H) < D$ then the coset hgH is within a distance D of the identity for some $h \in H$. Since the metric $d_{\mathcal{A}}$ is proper on G , the ball of radius D centered at the identity is finite. It follows that the left cosets gH intersecting $\mathcal{N}_D(H)$ lie in at most f distinct H -orbits for some $f = f(D) < \infty$. Thus it suffices to bound the number of elements of \mathcal{H} in the orbit $H(gH)$ for each fixed $g \notin H$.

If we let $K := H \cap gHg^{-1}$, then Lemma 4.2 shows that

$$\text{height}_H(K) < \text{height}_G(H).$$

Since K is a quasiconvex subgroup of the hyperbolic group H , the inductive hypothesis applied to $K \leq H$ gives for each D' a number $M' = M'(D') < \infty$ so that any collection of M' distinct cosets hK in H contains a pair separated by a \mathcal{B} -distance at least D' . Furthermore, the proof of Lemma 4.2 shows that there is a well-defined map $hgH \rightarrow hK$ taking left cosets of H in the orbit of gH to left cosets of K . A similar argument shows that this map is bijective.

In order to complete the proof, we will show that D -closeness of distinct cosets h_0gH and h_1gH in $(G, d_{\mathcal{A}})$ implies D' -closeness of the corresponding cosets h_0K and h_1K in $(H, d_{\mathcal{B}})$ for some D' depending on D . Roughly speaking, this claim is proved by considering a δ -thin triangle in G whose sides are close to the three cosets H , h_0gH and h_1gH , and applying Lemma 4.5. More precisely, suppose we have points x, y and z such that

$$x \in \mathcal{N}_D(H) \cap \mathcal{N}_D(h_0gH),$$

$$y \in \mathcal{N}_D(H) \cap \mathcal{N}_D(h_1gH)$$

and
$$z \in \mathcal{N}_D(h_0gH) \cap \mathcal{N}_D(h_1gH).$$

By Lemma 4.7 we have

$$[x, y] \subset \mathcal{N}_{D_1}(H), \quad [x, z] \subset \mathcal{N}_{D_1}(h_0gH) \quad \text{and} \quad [y, z] \subset \mathcal{N}_{D_1}(h_1gH)$$

for some D_1 depending on D . By δ -hyperbolicity of $(G, d_{\mathcal{A}})$, there is a point $w \in [x, y]$ within a distance δ of both $[x, z]$ and $[y, z]$. Applying the isometry h_i to the conclusion of Lemma 4.5, we see that

$$\mathcal{N}_{D_1}(H) \cap \mathcal{N}_{D_1+\delta}(h_i g H) \subseteq \mathcal{N}_{D_2}(h_i K),$$

where D_2 depends on D_1 but is independent of the choice of $h_i \in H$. In other words, w lies within an \mathcal{A} -distance D_2 of both $h_1 K$ and $h_2 K$ in G . Since H is an undistorted subgroup of G , it follows that the \mathcal{B} -distance between $h_1 K$ and $h_2 K$ in H is similarly bounded in terms of D , as desired. \square

5 Relatively hyperbolic groups and the word metric

Various equivalent formulations of relatively hyperbolic groups have been introduced and studied by Gromov [9], Farb [6], Bowditch [3], Druţu–Sapir [5] and Osin [15].

In this section we discuss geometric features of the Cayley graph of a relatively hyperbolic group with respect to a finite generating set. In particular, we prove two results about triangles and quadrilaterals with the property that each side lies near a peripheral subspace.

We begin by recalling the definition of a relatively hyperbolic group.

Definition 5.1 (Relatively hyperbolic) Let G be a finitely generated group and \mathbb{P} a finite collection of subgroups of G . Suppose G acts on a δ -hyperbolic graph K with finite edge stabilizers and finitely many orbits of edges. Suppose K is *fine* in the sense that for each n each edge of K is contained in only finitely many circuits of length n . Suppose also that \mathbb{P} is a set of representatives of the conjugacy classes of infinite vertex stabilizers. Then the pair (G, \mathbb{P}) is *relatively hyperbolic*. The subgroups $P \in \mathbb{P}$ are the *peripheral subgroups* of (G, \mathbb{P}) , and their left cosets gP are *peripheral cosets*.

Throughout this section all paths and distances are taken in the Cayley graph of a relatively hyperbolic group with respect to a fixed finite generating set.

We now collect several results due to Druţu–Sapir on the geometry of the word metric for a relatively hyperbolic group. The first states that peripheral cosets are “isolated.”

Theorem 5.2 (Druţu–Sapir [5, Theorem 4.1]) *Suppose (G, \mathbb{P}) is relatively hyperbolic. For each $\rho < \infty$ there is a constant $\kappa = \kappa(\rho) < \infty$ so that for any two peripheral cosets $gP \neq g'P'$ we have*

$$\text{diam}(\mathcal{N}_\rho(gP) \cap \mathcal{N}_\rho(g'P')) < \kappa.$$

The next result quantifies the quasiconvexity of peripheral cosets with respect to quasigeodesics.

Theorem 5.3 (Druţu–Sapir [5, Lemma 4.15]) *Suppose (G, \mathbb{P}) is relatively hyperbolic. Given positive constants ϵ and ν , there is a constant $\tau = \tau(\epsilon, \nu)$ so that the following holds. Let c be an ϵ -quasigeodesic with endpoints in the ν -neighborhood of a peripheral coset gP . Then $c \subseteq \mathcal{N}_\tau(gP)$.*

The following proposition roughly states that a pair of quasigeodesics beginning in the same peripheral coset and ending in the same peripheral coset must leave the former coset and enter the latter coset close together.

Proposition 5.4 (Druţu–Sapir [5, Lemma 8.11]) *Let (G, \mathbb{P}) be relatively hyperbolic. There is a constant ν_0 such that for each $\nu > \nu_0$ there is a constant $D_1 = D_1(\nu)$ such that the following holds. Let gP and $g'P'$ be two distinct peripheral cosets and let c_1 and c_2 be two geodesics such that c_i has endpoints x_i and y_i . If $c_i \cap \overline{\mathcal{N}_\nu(gP)} = \{x_i\}$ and $c_i \cap \overline{\mathcal{N}_\nu(g'P')} = \{y_i\}$ then $d(x_1, x_2)$ and $d(y_1, y_2)$ are both less than D_1 .*

Druţu–Sapir also establish the following geometric description of quasigeodesic triangles.

Theorem 5.5 (Druţu–Sapir [5, Section 8.1.3]) *Let (G, \mathbb{P}) be relatively hyperbolic. For each ϵ there are constants $\delta = \delta(\epsilon)$ and $\mu = \mu(\epsilon)$ such that the following holds. Let Δ be an ϵ -quasigeodesic triangle with sides c_0, c_1 and c_2 . Then either*

- (1) *there is a point w such that the ball $B(w, \delta)$ intersects all three sides of Δ , or*
- (2) *there is a peripheral coset gP such that the neighborhood $\mathcal{N}_\delta(gP)$ intersects all three sides of Δ .*

In the second case, illustrated in [Figure 1](#), let c'_i be the smallest subpath of c_i containing $c_i \cap \mathcal{N}_\delta(gP)$. Then the terminal endpoint of c'_i and the initial endpoint of c'_{i+1} are mutually within a distance μ (indices modulo 3).

Roughly speaking, the following lemma deals with a triangle each of whose sides lies close to a peripheral coset. The conclusion is that either all three peripheral cosets are equal or one of the sides of the triangle is short.

Lemma 5.6 *Let (G, \mathbb{P}) be relatively hyperbolic. For each $\eta > 0$, there is a constant $\zeta > 0$ such that the following holds. Let g_0P_0, g_1P_1 and g_2P_2 be peripheral cosets such that $g_0P_0 \notin \{g_1P_1, g_2P_2\}$. Suppose for each $\{i, j, k\} = \{0, 1, 2\}$ there is a point*

$$y_i \in \mathcal{N}_\eta(g_jP_j) \cap \mathcal{N}_\eta(g_kP_k).$$

Then $d(y_1, y_2) < \zeta$.




Figure 1: A quasigeodesic triangle in a relatively hyperbolic group with a peripheral coset at the center

Proof For each choice of $\{i, j, k\} = \{0, 1, 2\}$, pick a point $y_i \in \mathcal{N}_\eta(g_j P_j) \cap \mathcal{N}_\eta(g_k P_k)$. Consider a geodesic triangle $\Delta := c_0 \cup c_1 \cup c_2 \cup \Delta$ such that each c_i has endpoints y_j and y_k . By [Theorem 5.3](#), there is a constant τ depending on η such that c_i lies in the τ -neighborhood of $g_i P_i$.

Let $\delta = \delta(1)$ and $\mu = \mu(1)$ be the constants given by [Theorem 5.5](#). Then there is either a point z or a peripheral coset gP whose δ -neighborhood intersects all three sides of Δ .

We will see that we must be in the former case, for a possibly larger constant δ' . Suppose $\mathcal{N}_\delta(gP)$ intersects all three sides of Δ . For each i , let c'_i denote the smallest subpath of c_i containing $c_i \cap \overline{\mathcal{N}_\delta(gP)}$.

Choose $i \in \{0, 1, 2\}$ so that $gP \neq g_i P_i$. Since $c'_i \subseteq \mathcal{N}_\delta(g_i P_i)$, it follows from [Theorem 5.2](#) that c'_i has length less than $\kappa = \kappa(2\delta)$. By [Theorem 5.5](#), the endpoints z_j and z_k of c'_i are within a distance μ of the segments c_j and c_k respectively. Thus we have

$$d(z_j, c_i) = 0, \quad d(z_j, c_j) < \mu \quad \text{and} \quad d(z_j, c_k) < \kappa + \mu.$$

In other words, the ball of radius $\kappa + \mu$ centered at z_j intersects all three sides of Δ .

It follows that, in all cases, there exists a point z such that the δ' -ball centered at z intersects all three sides of Δ , where $\delta' = \delta + \kappa + \mu$. Now for each choice of $\{j, k\} = \{1, 2\}$ we have

$$z \in \mathcal{N}_{\delta'+\tau}(g_0 P_0) \cap \mathcal{N}_{\delta'+\tau}(g_j P_j)$$

and
$$y_k \in \mathcal{N}_\tau(g_0 P_0) \cap \mathcal{N}_\tau(g_j P_j).$$

Consequently, [Theorem 5.2](#) gives an upper bound $\kappa' := \kappa(\delta' + \tau)$ on the distance $d(y_k, z)$. Varying $k \in \{1, 2\}$, we see that $d(y_1, y_2)$ is less than $2\kappa'$. □

The following result is analogous to the previous lemma, but for quadrilaterals instead of triangles. If each side of a quadrilateral lies close to a peripheral coset, then either all four cosets are equal or some pair of vertices of the quadrilateral is close together.

Lemma 5.7 *Let (G, \mathbb{P}) be relatively hyperbolic. For each $\eta > 0$, there is a constant $\xi > 0$ such that the following holds. Let $g_0 P_0, g_1 P_1, g_2 P_2$ and $g_3 P_3$ be peripheral cosets. Suppose for each $i \in \{0, 1, 2, 3\}$ there is a point*

$$y_i \in \mathcal{N}_\eta(g_{i-1} P_{i-1}) \cap \mathcal{N}_\eta(g_i P_i) \quad (\text{indices modulo } 4).$$

If the pairwise distances between the points $\{y_0, y_1, y_2, y_3\}$ are all at least ξ , then all four peripheral cosets are equal.

Proof Suppose the four peripheral cosets are not all equal. We will show that one of the pairwise distances between the y_i is bounded above by an appropriate constant ξ .

For each i , let c_i be a geodesic in X from y_i to y_{i+1} . Then by [Theorem 5.3](#), we have

$$c_i \subseteq \mathcal{N}_\tau(g_i P_i)$$

for some τ depending on η .

If $g_i P_i = g_{i+1} P_{i+1}$ for some i , the result follows immediately from [Lemma 5.6](#). Now suppose $g_i P_i = g_{i+2} P_{i+2}$ for some i , but $g_{i+1} P_{i+1} \neq g_i P_i$. Then c_{i+1} lies in the τ -neighborhood of both $g_i P_i$ and $g_{i+1} P_{i+1}$. But then [Theorem 5.2](#) gives an upper bound $\kappa_0 := \kappa(\tau)$ on $d(y_{i+1}, y_{i+2})$, completing the proof. Thus it suffices to assume that the four peripheral cosets are all different.

By hypothesis, the points y_0 and y_1 lie in $\mathcal{N}_\eta(g_0 P_0)$. Let u be the point where c_3 first enters the closed η -neighborhood of $g_0 P_0$ when traversed from y_3 to y_0 . Similarly, let v be the point where c_1 first enters the closed η -neighborhood of $g_0 P_0$ when traversed from y_2 to y_1 . Since $g_0 P_0 \notin \{g_1 P_1, g_3 P_3\}$, [Theorem 5.2](#) gives an upper bound $\kappa_1 := \kappa(\eta)$ on the distances $d(y_0, u)$ and $d(y_1, v)$. We may assume without loss of generality that η is greater than the constant ν_0 given by [Proposition 5.4](#). Since $g_0 P_0 \neq g_2 P_2$, [Proposition 5.4](#) gives an upper bound $D_1 = D_1(\eta)$ on the distance $d(u, v)$. Thus we have

$$d(y_0, y_1) < D_1 + 2\kappa_1$$

completing the proof of the lemma. □

6 The geometry of relative Cayley graphs

The results in this section and the next involve the interplay between two different metrics on a relatively hyperbolic G with a fixed finite generating set \mathcal{S} . When a metric is not specified, all distances are assumed to refer to the word metric $d_{\mathcal{S}}$ or (equivalently) in the Cayley graph $\text{Cayley}(G, d_{\mathcal{S}})$. Let \mathcal{P} be the union of all the peripheral subgroups of G . We will also use the word metric $d_{\mathcal{S} \cup \mathcal{P}}$ and the Cayley graph $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ for the (typically infinite) generating set $\mathcal{S} \cup \mathcal{P}$. Whenever we use this “relative metric” or “relative Cayley graph,” we will explicitly call attention to its use. In particular, if $A \subset G$ the notation $\mathcal{N}_{\epsilon}(A)$ always refers to the ϵ -neighborhood of A using the metric $d_{\mathcal{S}}$, and the notation $B(x, \epsilon)$ refers to an \mathcal{S} -metric ball.

Geodesics in the relative Cayley graph are sometimes referred to as *relative geodesics in G* , etc. Observe that $\text{Cayley}(G, \mathcal{S})$ is a subgraph of $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ containing all the vertices but omitting the edges labelled by elements of \mathcal{P} .

The next result is a relatively hyperbolic analogue of the Morse Lemma, dealing with a pair of relative geodesics whose respective endpoints are close together in $d_{\mathcal{S}}$.

Proposition 6.1 (Osin [15, Proposition 3.15]) *Let (G, \mathbb{P}) be relatively hyperbolic. For each $k \geq 0$, there is a constant $\rho = \rho(k)$ such that the following holds. Let c and c' be relative geodesics in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ such that their initial endpoints are within an \mathcal{S} -distance k and their terminal endpoints are also within an \mathcal{S} -distance k . Then the set of vertices of c and the set of vertices of c' are within a Hausdorff \mathcal{S} -distance ρ .*

The following result is an analogue of [Theorem 5.5](#) for relative geodesic triangles.

Theorem 6.2 (Osin [15, Theorem 3.26]) *Let (G, \mathbb{P}) be relatively hyperbolic. Then there is a constant $\nu > 0$ such that if $c_0 \cup c_1 \cup c_2$ is a relative geodesic triangle in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ then for each vertex v of c_0 there is a vertex u in the union $c_1 \cup c_2$ such that*

$$d_{\mathcal{S}}(u, v) < \nu.$$

Subdividing an n -gon into $n - 2$ triangles and applying the previous theorem, we get the following corollary. As an aside we note that the linear function $n - 2$ in the conclusion of the corollary can be improved to a logarithmic function of n by subdividing more carefully.

Corollary 6.3 *Let (G, \mathbb{P}) and ν be as in the previous theorem. If $c_1 \cup \dots \cup c_n$ is a relative geodesic n -gon in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$, then for each vertex v on c_1 there is a vertex u on the union $c_2 \cup \dots \cup c_n$ such that*

$$d_{\mathcal{S}}(u, v) < (n - 2)\nu.$$

The notion of saturation of a quasigeodesic was introduced by Druţu–Sapir [5]. The saturation of an arbitrary subset is defined similarly, as follows.

Definition 6.4 Suppose (G, \mathbb{P}) is relatively hyperbolic, and let Y be a subset of G . For each $\nu > 0$ the ν -saturation of Y , denoted $\text{Sat}_\nu(Y)$, is the union of Y and every peripheral coset intersecting $\mathcal{N}_\nu(Y)$.

We conclude the section with a mild generalization of Proposition 6.1.

Proposition 6.5 Let (G, \mathbb{P}) be relatively hyperbolic. For each $\tau > 0$ there is a constant $\lambda = \lambda(\tau) > 0$ such that the following holds. Let c and c' be relative geodesic segments in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$, such that the endpoints x_0 and x_1 of c' lie within an \mathcal{S} -distance τ of $\text{Sat}_\tau(\text{Vert}(c))$. Let v be a vertex of c' . Then v lies within an \mathcal{S} -distance λ of either x_0 , x_1 , or a vertex of c .

Proof For each $i \in \{0, 1\}$, we define a path c_i in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ from x_i to c as follows. If x_i is within an \mathcal{S} -distance τ of a vertex of c , then let c_i be an \mathcal{S} -geodesic of shortest length from x_i to c . Otherwise, x_i is within an \mathcal{S} -distance τ of a left coset $g_i P_i$ that is within an \mathcal{S} -distance τ of a vertex of c . In the latter case, let c_i be the concatenation of an \mathcal{S} -geodesic q_i of shortest length from x_i to $g_i P_i$ followed by a peripheral edge e_i in $g_i P_i$ followed by an \mathcal{S} -geodesic r_i of shortest length from $g_i P_i$ to a vertex of c . Let $a_i \in C$ denote the terminal vertex of c_i , and let \bar{c} denote the portion of c from a_0 to a_1 .

Note that c_i is a concatenation of at most $2\tau + 1$ edges, each of which is a relative geodesic of length one. Thus $c' \cup c_0 \cup \bar{c} \cup c_1$ is a relative geodesic polygon with at most $4\tau + 4$ sides. Consequently, by Corollary 6.3 each vertex of c' is within an \mathcal{S} -distance $(4\tau + 2)\nu$ of some vertex of $c_0 \cup \bar{c} \cup c_1$.

Choose a vertex v of c' such that v is at an \mathcal{S} -distance more than $(4\tau + 2)\nu + \tau$ from x_0 . If c_0 is an \mathcal{S} -geodesic of length at most τ , then, by the triangle inequality, no vertex of c_0 can be within an \mathcal{S} -distance $(4\tau + 2)\nu$ of v . Thus v is within an \mathcal{S} -distance $(4\tau + 2)\nu$ of some vertex of $\bar{c} \cup c_1$. On the other hand, if c_0 is a concatenation $q_0 \cup e_0 \cup r_0$ as described above, then q_0 has \mathcal{S} -length at most τ . By a similar argument, it follows that v is within an \mathcal{S} -distance $(4\tau + 2)\nu$ of some vertex of $r_0 \cup \bar{c} \cup c_1$. Since r_0 has \mathcal{S} -length at most τ , each of its vertices lies within an \mathcal{S} -distance τ of the vertex a_0 . Thus in either case, v is within an \mathcal{S} -distance $(4\tau + 2)\nu + \tau$ of some vertex of $\bar{c} \cup c_1$.

Interchanging the roles of c_0 and c_1 , we see that if v is also at least an \mathcal{S} -distance $(4\tau + 2)\nu + \tau$ from x_1 then it is within an \mathcal{S} -distance $(4\tau + 2)\nu + \tau$ from some vertex of \bar{c} . Setting $\lambda = (4\tau + 2)\nu + \tau$ completes the proof. \square

Definition 6.6 Let (G, \mathbb{P}) be relatively hyperbolic with a finite generating set \mathcal{S} . Let c be a geodesic of $\text{Cayley}(G, \mathcal{S})$ and let ϵ and R be positive constants. A point $x \in c$ is (ϵ, R) -deep in a peripheral coset gP (with respect to c) if x is not within an \mathcal{S} -distance R of an endpoint of c and if $B(x, R) \cap c$ lies in $\mathcal{N}_\epsilon(gP)$. If x is not (ϵ, R) -deep in any gP then x is an (ϵ, R) -transition point of c .

Proposition 6.7 (Hruska [10, Proposition 8.13]) *Let (G, \mathbb{P}) be a relatively hyperbolic group with a finite generating set \mathcal{S} . There exist constants ϵ, R and L such that the following holds. Let c be any geodesic of $\text{Cayley}(G, \mathcal{S})$ with endpoints in G , and let c' be a relative geodesic in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ with the same endpoints as c . Then the set of vertices of c' is at a Hausdorff \mathcal{S} -distance at most L from the set of (ϵ, R) -transition points of c .*

Furthermore, let \bar{c} be a connected component of the set of all (ϵ, R) -deep points of c . Then there is a peripheral coset gP such that each $x \in \bar{c}$ is (ϵ, R) -deep in gP and is not (ϵ, R) -deep in any other peripheral coset.

7 Relatively quasiconvex subspaces

In the theory of word hyperbolic groups, the most natural subgroups (and subspaces) are the quasiconvex subgroups (and subspaces). In this section, we examine “relatively quasiconvex” subspaces, which play an analogous role in the theory of relatively hyperbolic groups. The definition of relative quasiconvexity given below was introduced by Osin [15].

Definition 7.1 Let (G, \mathbb{P}) be relatively hyperbolic such that G is generated by a finite set \mathcal{S} . For $\sigma \geq 0$, a subset $A \leq G$ is σ -relatively quasiconvex in G if the following condition holds: Let c be a relative geodesic in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ connecting two points of Y . Then each vertex v of c lies in the σ -neighborhood of A .

A subset $A \subseteq G$ is relatively quasiconvex if it is σ -relatively quasiconvex for some $\sigma \geq 0$.

Proposition 7.2 (Osin [15, Proposition 4.10]) *Relative quasiconvexity of a subset $A \subseteq G$ is independent of the choice of finite generating set \mathcal{S} for G .*

The following result analogous to [Theorem 5.5](#) follows immediately from the definition of relative quasiconvexity.

Proposition 7.3 *Each peripheral coset gP in a relatively hyperbolic group is 0-relatively quasiconvex.* □

The following result was proved independently by Hruska [10, Theorem 9.1] and Martinez-Pedroza [12, Proposition 1.5].

Proposition 7.4 *Let (G, \mathbb{P}) be relatively hyperbolic and let $H \leq G$ be σ -relatively quasiconvex. The infinite maximal parabolic subgroups $Q = H \cap gPg^{-1}$ of H lie in finitely many H -conjugacy classes, and the corresponding peripheral cosets gP in G lie in finitely many H -orbits.*

Proposition 7.5 *Let (G, \mathbb{P}) be relatively hyperbolic. Choose positive constants σ and τ . Then there is a constant $R = R(\sigma, \tau)$ such that the following holds. Choose $n \geq 3$, and let $\mathcal{A} = \{A_1, \dots, A_n\}$ be a collection of σ -relatively quasiconvex subsets of G . Suppose for each $i = 1, \dots, n$ there is a point*

$$x_i \in \bigcap \{ \mathcal{N}_\tau(A_j) \mid 1 \leq j \leq n, j \neq i \}.$$

Then we have the following consequences:

- (1) *There exists a vertex*

$$x_0 \in \mathcal{N}_R(\text{Sat}_R(A_1)) \cap \mathcal{N}_R(A_2) \cap \dots \cap \mathcal{N}_R(A_n).$$

- (2) *If A_1 is a left coset of a peripheral subgroup, then there exists a vertex*

$$x_0 \in \mathcal{N}_R(A_1) \cap \dots \cap \mathcal{N}_R(A_n).$$

- (3) *If A_0 is another σ -relatively quasiconvex subset of G such that x_i is contained in $\mathcal{N}_\tau(\text{Sat}_\tau(A_0))$ for all $i = 1, \dots, n$, then there exists a vertex x_0 satisfying (2) such that $x_0 \in \mathcal{N}_R(\text{Sat}_R(A_0))$.*

Proof Choose relative geodesics

$$c_1 = [x_2, x_3], \quad c_2 = [x_3, x_1] \quad \text{and} \quad c_3 = [x_1, x_2]$$

in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$, and consider the triangle $\Delta := c_1 \cup c_2 \cup c_3$. By Proposition 6.1 and the σ -relative quasiconvexity of A_ℓ , there is a constant $\rho = \rho(\tau)$ such that if $i \in \{1, 2, 3\}$ and $\ell \in \{i\} \cup \{4, \dots, n\}$ then each vertex of c_i lies within an \mathcal{S} -distance $\sigma + \rho$ of A_ℓ .

By Theorem 6.2, there is a constant ν such that for each choice of $\{i, j, k\} = \{1, 2, 3\}$ each vertex of c_i lies within an \mathcal{S} -distance ν of a vertex of $c_j \cup c_k$. It follows that c_i contains a pair of adjacent vertices v_{ij} and v_{ik} connected by an edge e_i such that v_{ij} is within an \mathcal{S} -distance ν of a vertex of c_j and v_{ik} is within an \mathcal{S} -distance ν of a vertex of c_k .

If for some choice of $\{i, j, k\} = \{1, 2, 3\}$, the edge e_i is labelled by a generator in \mathcal{S} , then for all $\ell \in \{i\} \cup \{4, \dots, n\}$ we have:

$$(7-1) \quad v_{ij} \in \mathcal{N}_{1+\nu+\sigma+\rho}(A_k) \cap \mathcal{N}_{\nu+\sigma+\rho}(A_j) \cap \mathcal{N}_{\sigma+\rho}(A_\ell)$$

If not, then e_3 is labelled by a generator in \mathcal{P} and v_{32} and v_{31} are vertices of a peripheral coset gP . Since $d_{\mathcal{S}}(v_{31}, A_1) < \nu + \sigma + \rho$, the coset gP is contained in $\text{Sat}_{\nu+\sigma+\rho}(A_1)$. Whenever $\ell = 3, \dots, n$, we now have:

$$v_{32} \in \text{Sat}_{\nu+\sigma+\rho}(A_1) \cap \mathcal{N}_{\nu+\sigma+\rho}(A_2) \cap \mathcal{N}_{\sigma+\rho}(A_\ell)$$

In either case, choosing $x_0 := v_{32}$ and $R \geq 1 + \nu + \sigma + \rho$ completes the proof of (1).

Now let us consider assertion (2). Suppose the subspace A_1 is a peripheral coset gP . If for some $i \in \{1, 2, 3\}$ the edge e_i corresponds to a generator in \mathcal{S} then we are done, since (7-1) shows that v_{ij} is within an \mathcal{S} -distance $1 + \nu + \sigma + \rho$ of A_ℓ for all $\ell = 1, \dots, n$.

It therefore suffices to assume that each e_i corresponds to a generator from \mathcal{P} such that $v_{ij}, v_{ik} \in g_i P_i$. for some peripheral coset $g_i P_i$. If $gP \neq g_1 P_1$, then by Theorem 5.2, the \mathcal{S} -distance between v_{12} and v_{13} is at most $\kappa = \kappa(\sigma + \rho)$. In this case, a computation similar to (7-1) shows that v_{12} is within an \mathcal{S} -distance $\kappa + \nu + \sigma + \rho$ of A_ℓ for all $\ell = 1, \dots, n$. Similarly if the cosets $g_1 P_1, g_2 P_2$ and $g_3 P_3$ are not all the same, then by Lemma 5.6 the \mathcal{S} -distance between some pair v_{ij} and v_{ik} is bounded above by $\zeta = \zeta(\nu)$. Therefore v_{ij} is within an \mathcal{S} -distance $\zeta + \nu + \sigma + \rho$ of A_ℓ for all $\ell = 1, \dots, n$.

The only remaining possibility is that the four cosets $gP, g_1 P_1, g_2 P_2$ and $g_3 P_3$ are all equal. In particular, we have $A_1 = gP = g_3 P_3$. It follows that

$$d_{\mathcal{S}}(v_{32}, A_1) = 0, \quad d_{\mathcal{S}}(v_{32}, A_2) < \nu + \sigma + \rho \quad \text{and} \quad d_{\mathcal{S}}(v_{32}, A_\ell) < \sigma + \rho$$

for all $\ell = 3, \dots, n$. In all possible cases, to complete the proof of (2) it suffices to choose

$$R > 1 + \kappa + \zeta + \nu + \sigma + \rho.$$

Finally, we turn our attention to assertion (3). We have shown that assertions (1) and (2) hold for sufficiently large R and for some vertex x_0 lying on one of the sides c_i of the triangle Δ . Suppose the endpoints x_j and x_k of c_i lie within an \mathcal{S} -distance τ of $\text{Sat}_\tau(A_0)$. Choose points u and v in A_0 such that $x_j \in \mathcal{N}_\tau(\text{Sat}_\tau(u))$ and $x_k \in \mathcal{N}_\tau(\text{Sat}_\tau(v))$, and let \bar{c} be a geodesic in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ from u to v .

It follows from Proposition 6.5 that the vertex x_0 of c_i lies within a distance $\lambda = \lambda(\tau)$ of either x_j, x_k , or a vertex of \bar{c} . Since A_0 is σ -quasiconvex, it follows that x_0 lies

within a distance $\lambda + \sigma$ of $\{x_j, x_k\} \cup A_0$. In particular, x_0 is within an \mathcal{S} -distance $\lambda + \sigma + \tau$ of $\text{Sat}_\tau(A_0)$, establishing (3) for all $R > \lambda + \sigma + \tau$. \square

Corollary 7.6 *Let (G, \mathbb{P}) be relatively hyperbolic. Choose positive constants σ and D and an integer $n \geq 1$. There is a constant $R_1 = R_1(\sigma, D, n)$ such that the following holds. Let $\mathcal{A} = \{A_1, \dots, A_n\}$ be a set of σ -relatively quasiconvex subsets of G . Suppose the following intersection is nonempty:*

$$\bigcap_{i=1}^n \mathcal{N}_D(A_i)$$

If there is a peripheral coset gP such that for all i the intersection $\mathcal{N}_D(A_i) \cap \mathcal{N}_D(gP)$ is nonempty, then the following intersection is nonempty:

$$\mathcal{N}_{R_1}(gP) \cap \mathcal{N}_{R_1}(A_1) \cap \dots \cap \mathcal{N}_{R_1}(A_n)$$

Proof Recall that gP is 0-relatively hyperbolic. When $n = 1$, the result is trivial. We proceed by induction on n for $n \geq 2$. Let $\tau := R_1(\sigma, D, n - 1)$ denote the constant given by the inductive hypothesis. By assumption, $\bigcap_{i=1}^n \mathcal{N}_D(A_i)$ is nonempty. By induction, for each $i = 1, \dots, n$ the following intersection is nonempty:

$$\mathcal{N}_\tau(gP) \cap \bigcap \{ \mathcal{N}_\tau(A_j) \mid 1 \leq j \leq n, j \neq i \}$$

Applying Proposition 7.5(2) to the collection $\{gP, A_1, \dots, A_n\}$ of σ -quasiconvex subspaces, the result now follows. \square

Proposition 7.7 *Let (G, \mathbb{P}) be relatively hyperbolic. Choose positive constants σ and D and an integer $n \geq 1$. Let \mathcal{A} be an arbitrary set of σ -relatively quasiconvex subsets of G such that for each $A, A' \in \mathcal{A}$ we have $d_{\mathcal{S}}(A, A') < D$. Then there is a constant $M_0 = M_0(\sigma, D, n)$ so that whenever $M \geq M_0$ at least one of the following holds.*

- (1) *For every subset $\{A_1, \dots, A_n\} \subseteq \mathcal{A}$, the following intersection is nonempty:*

$$\bigcap_{\ell=1}^n \mathcal{N}_M(A_\ell)$$

- (2) *There is a peripheral coset gP such that $d_{\mathcal{S}}(gP, A) < M$ for all $A \in \mathcal{A}$. Furthermore, for all $A, A' \in \mathcal{A}$, the following triple intersection is nonempty:*

$$\mathcal{N}_M(gP) \cap \mathcal{N}_M(A) \cap \mathcal{N}_M(A')$$

Proof Recall that every peripheral coset gP is 0–quasiconvex, as observed in [Proposition 7.3](#). In order to establish (2), it is sufficient to find a constant M and a peripheral coset gP such that $d_S(gP, A) < M$ for all $A \in \mathcal{A}$. For then applying [Proposition 7.5\(2\)](#) to each collection $\{gP, A, A'\}$ with $A, A' \in \mathcal{A}$ gives a larger constant M' so that the following triple intersection is nonempty:

$$\mathcal{N}_{M'}(gP) \cap \mathcal{N}_{M'}(A) \cap \mathcal{N}_{M'}(A')$$

The proposition is trivial when $n \leq 2$. We proceed by induction on n for $n \geq 3$. Let τ denote the constant $M_0(\sigma, D, n - 1)$ given by the inductive hypothesis. We will show that the proposition holds whenever M is sufficiently large (in terms of σ and τ).

Suppose $\{A_0, \dots, A_{n-1}\}$ is a subset of \mathcal{A} such that

$$(7-2) \quad \bigcap_{\ell=0}^{n-1} \mathcal{N}_M(A_\ell) = \emptyset.$$

If no such set exists, there is nothing to prove. Furthermore, if there is a peripheral coset gP such that $d_S(P, A) < \nu$ for all $A \in \mathcal{A}$, then we are done. Thus we may assume that for every collection of $n - 1$ distinct elements of \mathcal{A} , their ν –neighborhoods have a nonempty intersection.

Applying [Proposition 7.5\(1\)](#) to the collection $\{A_0, \dots, A_{n-1}\}$ gives $R_0 = R_0(\sigma, \tau)$ and a point

$$x_n \in \mathcal{N}_{R_0}(\text{Sat}_{R_0}(A_0)) \cap \mathcal{N}_{R_0}(A_1) \cap \dots \cap \mathcal{N}_{R_0}(A_{n-1}).$$

If we choose M greater than R_0 , then (7-2) implies that $d_S(x_n, A_0)$ is at least R_0 . Thus $x_n \in \mathcal{N}_{R_0}(g_0P_0)$ for some peripheral coset g_0P_0 such that $g_0P_0 \subseteq \text{Sat}_{R_0}(A_0)$. Note that $d_S(g_0P_0, A_\ell) < 2R_0$ for all $\ell = 0, \dots, n - 1$. If $\mathcal{A} = \{A_0, \dots, A_{n-1}\}$, we are done. Otherwise, choose an arbitrary $A_n \in \mathcal{A} - \{A_0, \dots, A_{n-1}\}$. To complete the proof, we will bound the \mathcal{S} –distance from A_n to g_0P_0 in terms of σ and τ , provided that M is sufficiently large.

For each $i = 1, \dots, n - 1$, applying [Proposition 7.5\(1\)](#) to the collection

$$\{A_0, \dots, A_{i-1}, A_{i+1}, \dots, A_n\}$$

gives a point

$$x_i \in \mathcal{N}_{R_0}(\text{Sat}_{R_0}(A_0)) \cap \mathcal{N}_{R_0}(A_1) \cap \dots \cap \mathcal{N}_{R_0}(A_{i-1}) \cap \mathcal{N}_{R_0}(A_{i+1}) \cap \dots \cap \mathcal{N}_{R_0}(A_n).$$

Notice that the points x_1, \dots, x_n and subspaces A_0, A_1, \dots, A_n satisfy the hypothesis of [Proposition 7.5\(3\)](#). Therefore there exist a constant $R_1 := R(\sigma, R_0)$ and a point

$$x_0 \in \mathcal{N}_{R_1}(\text{Sat}_{R_1}(A_0)) \cap \mathcal{N}_{R_1}(A_1) \cap \dots \cap \mathcal{N}_{R_1}(A_n).$$

Choose a relative geodesic c from x_0 to x_n . Since the endpoints of c lie within a distance $R_0 + R_1$ of A_ℓ for each $\ell = 1, \dots, n-1$, we know from [Proposition 6.1](#) and the σ -relative quasiconvexity of A_ℓ that each vertex of c lies within an \mathcal{S} -distance $\sigma + \rho$ of A_ℓ , where $\rho = \rho(R_0 + R_1)$. However, by [\(7-2\)](#) there is no vertex within a distance M of every A_ℓ for $\ell = 0, \dots, n$.

Roughly speaking, it follows that once M is sufficiently large, each vertex of c will be far from A_0 . We have already used this principle to conclude that x_n is within an \mathcal{S} -distance R_0 of g_0P_0 , which is itself within an \mathcal{S} -distance R_0 of A_0 , once M is sufficiently large. Let y_n be a vertex of A_0 within an \mathcal{S} -distance R_0 of g_0P_0 . A similar argument shows that x_0 is within a distance R_1 of a peripheral coset g_1P_1 , which is itself within a distance R_1 of A_0 , for sufficiently large M . Let y_0 be a vertex of A_0 within an \mathcal{S} -distance R_1 of g_1P_1 . Choose a relative geodesic \bar{c} of Cayley($G, \mathcal{S} \cup \mathcal{P}$) from y_0 to y_n .

Since x_0 and x_n are both contained in $\mathcal{N}_{R_0+R_1}(\text{Sat}_{R_0+R_1}(\bar{c}))$, it follows from [Proposition 6.5](#) that each vertex of c lies within an \mathcal{S} -distance λ of either x_0 , x_n , or a vertex of \bar{c} . By the σ -relative quasiconvexity of A_0 , each vertex of \bar{c} lies within an \mathcal{S} -distance σ of A_0 . Thus each vertex of c either lies within an \mathcal{S} -distance λ of $\{x_0, x_n\}$ or lies within an \mathcal{S} -distance $\lambda + \sigma$ of A_0 for some $\lambda = \lambda(R_0 + R_1)$. Suppose v is a vertex of c whose \mathcal{S} -distance from A_0 is less than $\lambda + \sigma$. Once $M > \lambda + \sigma + \rho$, the existence of such a vertex v contradicts [\(7-2\)](#). Therefore we may assume that each vertex v of c lies within an \mathcal{S} -distance λ of $\{x_0, x_n\}$. A similar argument shows that each vertex v of \bar{c} lies within an \mathcal{S} -distance λ of $\{y_0, y_n\}$, since otherwise v would be close to both A_0 and a vertex of c , contradicting [\(7-2\)](#) once M is sufficiently large.

Let e be an edge of c with endpoints v_0 and v_n such that v_0 is within an \mathcal{S} -distance λ of x_0 and v_n is within an \mathcal{S} -distance λ of x_n . If e corresponds to a generator in \mathcal{S} , then $d_{\mathcal{S}}(x_0, x_n) < 2\lambda + 1$. Since $d_{\mathcal{S}}(x_n, g_0P_0) < R_0$ and $d_{\mathcal{S}}(x_0, A_n) < R_1$, it follows that $d_{\mathcal{S}}(A_n, g_0P_0) < 2\lambda + 1 + R_0 + R_1$ and we are done. Thus we may assume that e corresponds to a generator in \mathcal{P} and that $v_0, v_n \in g_2P_2$ for some peripheral coset g_2P_2 .

Now let \bar{e} be an edge of \bar{c} with endpoints w_0 and w_n within a distance λ of y_0 and y_n respectively. We will see that \bar{e} must also correspond to a generator in \mathcal{P} . Suppose instead that \bar{e} corresponds to a generator in \mathcal{S} . Then $d_{\mathcal{S}}(y_0, y_n)$ is less than $2\lambda + 1$. In this case, consider the three peripheral cosets g_0P_0 , g_1P_1 , and g_2P_2 . Recall that x_0 lies within a distance $R_1 + \lambda$ of both g_1P_1 and g_2P_2 . If $g_0P_0 \in \{g_1P_1, g_2P_2\}$, then $d(x_0, g_0P_0) < R_1 + \lambda$, completing the proof as before since $d(x_0, A_n) < R_0$. On the other hand, if $g_0P_0 \notin \{g_1P_1, g_2P_2\}$, we can apply [Lemma 5.6](#) to g_0P_0 , g_1P_1 and g_2P_2 to produce an upper bound ζ on the distance from x_n to A_0 , contradicting [\(7-2\)](#)

once M is at least $\zeta + \sigma + \rho$ as before. Therefore we may assume that \bar{e} corresponds to a generator in \mathcal{P} and that $w_0, w_n \in g_3 P_3$ for some peripheral coset $g_3 P_3$.

We now have a cycle of four peripheral cosets $g_0 P_0, g_2 P_2, g_1 P_1$, and $g_3 P_3$ such that the following hold for $\eta := R_0 + R_1 + \lambda$:

$$x_n \in \mathcal{N}_\eta(g_0 P_0) \cap \mathcal{N}_\eta(g_2 P_2)$$

$$x_0 \in \mathcal{N}_\eta(g_2 P_2) \cap \mathcal{N}_\eta(g_1 P_1)$$

$$y_0 \in \mathcal{N}_\eta(g_1 P_1) \cap \mathcal{N}_\eta(g_3 P_3)$$

and
$$y_n \in \mathcal{N}_\eta(g_3 P_3) \cap \mathcal{N}_\eta(g_0 P_0)$$

Now Lemma 5.7 implies that either all four cosets are equal, or some pair from the set $\{x_n, x_0, y_0, y_n\}$ has distance less than ξ , for some constant $\xi = \xi(\eta)$.

We can now complete the proof using arguments similar to the ones used above. As we have seen, it suffices to assume that $g_0 P_0 \notin \{g_1 P_1, g_2, P_2\}$. Therefore some pair from $\{x_n, x_0, y_0, y_n\}$ is within a distance ξ . If $d(y_0, y_n) < \xi$, then the result follows from an application of Lemma 5.6. If $d(x_0, x_n) < \xi$, then $d(A_n, g_0 P_0) < R_0 + R_1 + \xi$. If $d(\{x_0, x_n\}, \{y_0, y_n\}) < \xi$ then, since $y_0, y_n \in A_0$, it follows that either x_0 or x_n is within a distance ξ of A_0 and within a distance $R_0 + R_1$ of A_1, \dots, A_{n-1} , contradicting (7-2) once $M > \xi + R_0 + R_1$. □

Proposition 7.8 (Close to two peripherals) *Let (G, \mathbb{P}) be relatively hyperbolic. For each $M > 0$ and $\sigma > 0$ there exists $M_1 = M_1(M, \sigma)$ such that the following holds. Suppose \mathcal{A} is a collection of σ -relatively quasiconvex subsets and $g_0 P_0 \neq g_1 P_1$ are distinct peripheral cosets. If $d_S(g_i P_i, A) < M$ for all $i \in \{0, 1\}$ and for all $A \in \mathcal{A}$, then the following intersection is nonempty:*

$$\bigcap_{A \in \mathcal{A}} \mathcal{N}_{M_1}(A)$$

Proof Let ϵ, R and L be the constants given by Proposition 6.7, let ν_0 be the constant given by Proposition 5.4, let $\nu := \nu_0 + M + \epsilon$, and let $D_1 = D_1(\nu)$ be given by Proposition 5.4. Let $\tau = \tau(\nu)$ be given by Theorem 5.3, and let $\kappa = \kappa(\epsilon + \tau)$ be given by Theorem 5.2.

For each $A \in \mathcal{A}$ there exists a geodesic c in $\text{Cayley}(G, \mathcal{S})$ whose endpoints x_0, x_1 satisfy $d_S(x_i, g_i P_i) < M$. Choose $y \in c$ to be the last point of c that lies in $\overline{\mathcal{N}_\nu(g_0 P_0)}$ (when c is traversed from x_0 to x_1).

We will see that y is within an S -distance κ of an (ϵ, R) -transition point of c . Suppose y is not itself an (ϵ, R) -transition point. Then by Proposition 6.7 there is

a subgeodesic \bar{c} of c containing y each of whose interior points is (ϵ, R) -deep in a fixed peripheral coset gP and whose endpoints are (ϵ, R) -transition points. It is clear that $gP \neq g_0P_0$, since $\epsilon \leq \nu$. Let $[x_0, y]$ denote the portion of c from x_0 to y . Since $M \leq \nu$, the points x_0 and y each lie in $\mathcal{N}_\nu(g_0P_0)$. Therefore, $[x_0, y]$ lies in $\mathcal{N}_\tau(g_0P_0)$. On the other hand, \bar{c} lies in $\mathcal{N}_\epsilon(gP)$. Thus $\bar{c} \cap [x_0, y]$ lies in $\mathcal{N}_\epsilon(gP) \cap \mathcal{N}_\tau(g_0P_0)$ and hence has length at most κ . In other words, y is within an \mathcal{S} -distance κ of an (ϵ, R) -transition point z of c , as desired.

By [Proposition 6.7](#) this point z is within an \mathcal{S} -distance L of a vertex w of \hat{c} , where \hat{c} is any relative geodesic in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ with the same endpoints as c . Since A is σ -relatively quasiconvex, it follows that $d_{\mathcal{S}}(w, A) < \sigma$. Therefore, $d_{\mathcal{S}}(y, A) < \kappa + L + \sigma$.

If we now vary the choice of A , the above argument produces for each $A \in \mathcal{A}$ a point $y = y(A)$ within an \mathcal{S} -distance $\kappa + L + \sigma$ of A . Since y is the last point of c that lies in $\overline{\mathcal{N}_\nu(g_0P_0)}$, it follows from [Proposition 5.4](#) that for each $A, A' \in \mathcal{A}$ we have $d_{\mathcal{S}}(y(A), y(A')) < D_1$. In particular, if we fix $A \in \mathcal{A}$, then every $A' \in \mathcal{A}$ comes within an \mathcal{S} -distance $D_1 + \kappa + L + \sigma$ of the point $y(A)$. Setting $M_1 \geq D_1 + \kappa + L + \sigma$ completes the proof. □

The following corollary essentially states that, if we increase the constants given by [Proposition 7.7](#), then the peripheral coset appearing in conclusion (2) is unique.

Corollary 7.9 (Uniqueness of peripheral coset) *Let (G, \mathbb{P}) be relatively hyperbolic. Choose σ, D, n and \mathcal{A} satisfying the hypothesis of [Proposition 7.7](#), and let $M_0 = M_0(\sigma, D, n)$ be the constant given by [Proposition 7.7](#). Then for each $M \geq M_0$ there exists $M_2 = M_2(\sigma, D, n, M)$ such that at least one of the following holds:*

- (1) For every $\{A_1, \dots, A_n\} \subseteq \mathcal{A}$, the following intersection is nonempty:

$$\bigcap_{\ell=1}^n \mathcal{N}_{M_2}(A_\ell)$$

- (2) There is a unique peripheral coset gP such that $d_{\mathcal{S}}(gP, A) < M$ for all $A \in \mathcal{A}$. Furthermore, for all $A, A' \in \mathcal{A}$ the following intersection is nonempty:

$$\mathcal{N}_M(gP) \cap \mathcal{N}_M(A) \cap \mathcal{N}_M(A')$$

Proof Choose $M \geq M_0$ arbitrary. If there does not exist a peripheral coset gP such that for all $A \in \mathcal{A}$ we have $d_{\mathcal{S}}(gP, A) < M$, then [Proposition 7.7](#) implies that (1) holds for any $M_2 \geq M$.

On the other hand, suppose there are two peripheral cosets $g_0P_0 \neq g_1P_1$ such that $d_S(g_iP_i, A) < M$ for all $i \in \{0, 1\}$ and all $A \in \mathcal{A}$. In this case, Proposition 7.8 implies that

$$\bigcap_{A \in \mathcal{A}} \mathcal{N}_{M_2}(A)$$

is nonempty once M_2 is larger than the constant $M_1 = M_1(M, \sigma)$. Condition (1) follows immediately. \square

8 Bounded packing in relatively hyperbolic groups

Let (G, \mathbb{P}) be relatively hyperbolic. An element $f \in G$ is *elliptic* if it has finite order. If f has infinite order, then f is *parabolic* if it is contained in a conjugate of a peripheral subgroup $P \in \mathbb{P}$ and *loxodromic* otherwise. A subgroup $H \leq G$ is *elliptic* if it is finite, and *parabolic* if it is infinite and contained in a conjugate of a peripheral subgroup $P \in \mathbb{P}$. Otherwise H is *loxodromic*. Every loxodromic subgroup contains a loxodromic element.

Definition 8.1 Let $\text{height}_{(G, \mathbb{P})}(H) = n$ if n is the maximal number of distinct cosets g_iH such that the intersection $\bigcap g_iHg_i^{-1}$ is loxodromic. Let $\text{width}_{(G, \mathbb{P})}(H) = n$ if n is the maximal number of distinct cosets g_iH such that for each i, j the intersection $g_iHg_i^{-1} \cap g_jHg_j^{-1}$ is loxodromic. Note that an elliptic or parabolic subgroup has both height and width zero.

Lemma 8.2 Let (G, \mathbb{P}) be relatively hyperbolic, and let \mathcal{S} be a finite generating set for G . There is a universal constant $L \geq 0$ so that the following holds. Let c and c' be relative geodesic lines in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ with the same endpoints at infinity. Then the set of vertices of c and the set of vertices of c' are within a Hausdorff distance L with respect to the metric d_S .

Proof Since c and c' have the same endpoints at infinity, they are at a finite Hausdorff distance, say D , in the relative metric $d_{\mathcal{S} \cup \mathcal{P}}$. Let ν be the constant given by Theorem 6.2. Choose a vertex v of c . We will show that the \mathcal{S} -distance from v to some vertex v' of c' is bounded above by a constant that does not depend on D .

Let x and y be vertices of c on either side of v , at an $(\mathcal{S} \cup \mathcal{P})$ -distance at least $D + 2\nu$ from v . Choose vertices x' and y' of c' such that $d_{\mathcal{S} \cup \mathcal{P}}(x, x') < D$ and $d_{\mathcal{S} \cup \mathcal{P}}(y, y') < D$. Let \bar{c} be the subpath of c' from x' to y' , and choose relative geodesics c_0 from x to x' and c_1 from y' to y . By Corollary 6.3, the vertex v is within an \mathcal{S} -distance 2ν of a vertex v' of $c_0 \cup \bar{c} \cup c_1$. However, by the triangle inequality, the $(\mathcal{S} \cup \mathcal{P})$ -distance from v to any vertex of $c_0 \cup c_1$ is greater than 2ν . Therefore v' is a vertex of \bar{c} . \square

Lemma 8.3 *Let (G, \mathbb{P}) be relatively hyperbolic with a finite generating set \mathcal{S} . Suppose $H \leq G$ is σ -relatively quasiconvex. If f is a loxodromic element of H , then there exists a bi-infinite relative geodesic c in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ such that the following hold:*

- (1) *The Hausdorff \mathcal{S} -distance between the set of vertices of c and the subgroup $\langle f \rangle$ is finite.*
- (2) *Each vertex v of c lies in the σ -neighborhood of H .*

Proof For each $n = 1, 2, 3, \dots$ choose a relative geodesic c_n connecting f^{-n} with f^n . Since H is σ -relatively quasiconvex, each vertex of c_n lies in the σ -neighborhood of H . To complete the proof, it suffices to show that the sequence $\{c_n\}$ has a subsequence that converges pointwise to a relative geodesic line c . In order to ensure the existence of such a subsequence we will show that the vertices of c_n track \mathcal{S} -close to the set $\{f^i \mid -n \leq i \leq n\}$.

The subgroup $\langle f \rangle$ is L -relatively quasiconvex in G for some $L = L(f)$, and $\langle f \rangle \cap gPg^{-1}$ is finite for all conjugates of peripheral subgroups gP by Osin [15, Theorem 4.19]. Furthermore, there exists $\epsilon = \epsilon(f)$ such that the inclusion $\langle f \rangle \rightarrow \text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$ is an ϵ -quasi-isometric embedding with respect to the word metric on $\langle f \rangle$ for the standard generating set $\{f\}$ (see Osin [15, Corollary 4.20]). Let D be an upper bound on the \mathcal{S} -diameters of the finitely many finite subgroups $\langle f \rangle \cap gPg^{-1}$ with $|g|_{\mathcal{S}} < L$.

By quasiconvexity, each vertex of c_n lies \mathcal{S} -near $\langle f \rangle$. We need to show that if $-n \leq i \leq n$ the vertex f^i lies \mathcal{S} -near some vertex of c_n . Indeed, quasiconvexity guarantees that some of the f^i lie within an \mathcal{S} -distance σ' of vertices of c_n , so we only need to bound the size of the “gaps” between these elements f^i .

Let us begin with a pair of vertices v and w in the L -neighborhood of $\langle f \rangle$ connected by an edge e in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$. Translate by an element of $\langle f \rangle$ so that v and w are within an \mathcal{S} -distance L of 1 and f^k for some $k \geq 0$. If e is labelled by a generator $s \in \mathcal{S}$ then $|f^k|_{\mathcal{S}}$ is at most $2L + 1$. On the other hand, suppose e is labelled by a parabolic element $p \in P$ for some $P \in \mathbb{P}$. Then both endpoints of e lie in one of the finitely many peripheral cosets gP with $|g|_{\mathcal{S}} < L$. By Lemma 4.5 we have

$$\{1, f^k\} \subseteq \mathcal{N}_L(\langle f \rangle) \cap \mathcal{N}_L(gP) \subseteq \mathcal{N}_{L'}(\langle f \rangle \cap gPg^{-1})$$

for some constant L' . Maximizing over the finitely many possibilities for gP , we can assume that L' does not depend on the choice of edge e . Observe that $|f^k|_{\mathcal{S}} < 2L' + D$. Thus in all possible cases, we have

$$|f^k|_{\mathcal{S}} < M := \max\{2L + 1, 2L' + D\}.$$

It follows that:

$$0 \leq k \leq \epsilon |f^k|_{\mathcal{S} \cup \mathcal{P}} + \epsilon \leq \epsilon |f^k|_{\mathcal{S}} + \epsilon \leq \epsilon M + \epsilon$$

On the other hand, whenever $0 \leq i \leq k$ we have:

$$|f^i|_{\mathcal{S}} \leq i |f|_{\mathcal{S}} \leq k |f|_{\mathcal{S}} \leq (\epsilon M + \epsilon) |f|_{\mathcal{S}}$$

Therefore the \mathcal{S} -distance from f^i to $\{v, w\}$ is bounded above by the constant:

$$R := L + (\epsilon M + \epsilon) |f|_{\mathcal{S}}$$

By quasiconvexity, each vertex of c_n lies within an \mathcal{S} -distance L of some element of $\langle f \rangle$. Since c_n is an edge path, we can apply the above argument to the set of vertices of the path c_n to see that whenever $-n \leq i \leq n$ the vertex f^i lies within an \mathcal{S} -distance R of some vertex of c_n . Since $L \leq R$, the Hausdorff distance between the vertices of c_n and the set $\{f^i \mid -n \leq i \leq n\}$ is at most R . An easy diagonal argument shows that the sequence of relative geodesics $\{c_n\}$ has a subsequence converging pointwise to a relative geodesic line c .

Clearly the Hausdorff \mathcal{S} -distance between the vertices of c and the subgroup $\langle f \rangle$ is at most R . Since the vertices of c_n lie in the σ -neighborhood of H , the same holds for the vertices of c , completing the proof of the lemma. □

Lemma 8.4 (Loxodromic virtual intersection implies close) *Let (G, \mathbb{P}) be relatively hyperbolic, let \mathcal{S} be a finite generating set for G , and choose $\sigma \geq 0$. There is a constant C so that the following holds. Let H be a σ -relatively quasiconvex subgroup of (G, \mathbb{P}) containing a loxodromic element f . Suppose $\mathcal{A} = \{g_i H \mid i \in I\}$ is a collection of distinct cosets of H such that each conjugate $g_i H g_i^{-1}$ contains some power f^{k_i} of f . Then there exists a vertex $z \in G$ such that the ball of radius C in $\text{Cayley}(G, \mathcal{S})$ intersects every coset $g_i H \in \mathcal{A}$.*

Proof By Osin [15, Corollary 4.20], the map $n \mapsto f^n$ is a quasigeodesic line in $\text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$. Therefore the set $\{f^n \mid n \in \mathbb{Z}\}$ has exactly two limit points $f^{\pm\infty}$ in $\partial \text{Cayley}(G, \mathcal{S} \cup \mathcal{P})$. If $g_i H \in \mathcal{A}$ then $f^{k_i} \in g_i H g_i^{-1}$ for some $k_i \in \mathbb{Z}$. Thus $f^{nk_i} g_i \in g_i H$ for all $n \in \mathbb{Z}$. The sets $\{f^{nk_i} g_i \mid n \in \mathbb{Z}\}$ and $\{f^n \mid n \in \mathbb{Z}\}$ are at a finite Hausdorff \mathcal{S} -distance and thus have the same limit points $f^{\pm\infty}$.

Translating $g_i H$ by g_i^{-1} , we see that the subgroup $\langle g_i^{-1} f^{k_i} g_i \rangle$ of H has limit points $g_i^{-1} f^{\pm\infty}$. By Lemma 8.3 the points $g_i^{-1} f^{\pm\infty}$ can be connected by a relative geodesic line c_i such that each vertex of c_i lies within an \mathcal{S} -distance σ of H . Translating back by g_i produces a relative geodesic line $g_i c_i$ with endpoints $f^{\pm\infty}$ such that each vertex of $g_i c_i$ lies in the σ -neighborhood of $g_i H$.

For each $i, j \in I$ the relative geodesics $g_i c_i$ and $g_j c_j$ have the same endpoints at infinity. By [Lemma 8.2](#) the Hausdorff \mathcal{S} -distance between the vertex sets of $g_i c_i$ and $g_j c_j$ is at most L . Therefore for each vertex z of $g_i c_i$, the coset $g_j H$ intersects the ball of radius $L + \sigma$ in $\text{Cayley}(G, \mathcal{S})$ centered at z . \square

The following special case of [Lemma 8.4](#) is the main connection between height/width and packing. Roughly speaking, a collection of conjugates with loxodromic intersection comes mutually close together in $\text{Cayley}(G, \mathcal{S})$.

Corollary 8.5 (Loxodromic intersection implies close) *Let (G, \mathbb{P}) be relatively hyperbolic, let \mathcal{S} be a finite generating set for G , and choose $\sigma \geq 0$. There is a constant C so that the following holds. Let H be a σ -relatively quasiconvex subgroup of (G, \mathbb{P}) . Suppose $\{g_i H \mid i \in I\}$ is a collection of distinct cosets of H such that $\bigcap g_i H g_i^{-1}$ is loxodromic. Then there exists a vertex $z \in G$ such that the ball of radius C in $\text{Cayley}(G, \mathcal{S})$ intersects every coset $g_i H$.* \square

The following corollary is an immediate consequence of the preceding result together with the local finiteness of $\text{Cayley}(G, \mathcal{S})$.

Corollary 8.6 *Let (G, \mathbb{P}) be relatively hyperbolic and let $H \leq G$ be relatively quasiconvex. Then $\text{height}_{(G, \mathbb{P})}(H)$ is finite.* \square

[Corollary 8.5](#) also has the following easy consequence.

Corollary 8.7 *Let (G, \mathbb{P}) be relatively hyperbolic and let $H \leq G$ be a relatively quasiconvex subgroup. There are only finitely many double cosets $H g_1 H, \dots, H g_n H$ such that $H \cap g_i H g_i^{-1}$ is loxodromic.*

Proof Suppose $\{g_i H \mid i \in I\}$ is a collection of cosets such that $H \cap g_i H g_i^{-1}$ is loxodromic for each i . By [Corollary 8.5](#), the distances $d_{\mathcal{S}}(H, g_i H)$ are bounded above by a constant C . Thus we can translate $g_i H$ by an element of H to obtain a coset $h g_i H$ intersecting the ball of radius C in $\text{Cayley}(G, \mathcal{S})$ centered at the identity. Since this ball is finite, it follows that the cosets $g_i H$ lie in only finitely many double cosets $H g_i H$. \square

[Theorem 1.6](#) also follows from [Lemma 8.4](#).

Proof of [Theorem 1.6](#) Let $\mathcal{A} = \{g_i H \mid i \in I\}$ be the set of all cosets $g_i H$ such that $g_i H g_i^{-1} \cap H$ has finite index in both $g_i H g_i^{-1}$ and H . Then the union of all cosets in \mathcal{A} is the commensurator of H in G , and the cardinality of \mathcal{A} is equal to the index of H

in the commensurator. Since H is loxodromic, it contains a loxodromic element f . If $g_i H \in \mathcal{A}$ then $f^{k_i} \in g_i H g_i^{-1}$ for some $k_i \in \mathbb{Z}$. Thus by Lemma 8.4 there is a vertex z such that the ball of radius C in $\text{Cayley}(G, \mathcal{S})$ centered at z intersects each coset $g_i H \in \mathcal{A}$. The finite cardinality of this ball is an upper bound for the size of \mathcal{A} . \square

Proposition 8.8 (Packing peripheral subgroups) *If (G, \mathbb{P}) is relatively hyperbolic then each peripheral subgroup $P \in \mathbb{P}$ has bounded packing in G .*

Proof Choose a finite generating set \mathcal{S} for G . Let \mathcal{A} be a collection of left cosets of P whose pairwise distances are less than D . Consider three distinct elements $g_0 P, g_1 P, g_2 P$ of \mathcal{A} . Choose points $x_0 \in \mathcal{N}_D(g_1 P) \cap \mathcal{N}_D(g_2 P)$ and $x_1 \in \mathcal{N}_D(g_0 P) \cap \mathcal{N}_D(g_2 P)$. Then by Lemma 5.6, there is a constant $\zeta = \zeta(D)$ such that $d(x_0, x_1) < \zeta$. Therefore $g_0 P$ intersects the ball of radius $\zeta + D$ centered at x_0 . Keeping $g_1 P$ and $g_2 P$ fixed and varying $g_0 P$ over all elements of $\mathcal{A} - \{g_1 P, g_2 P\}$, we see that every element of \mathcal{A} intersects this ball. Since the metric $d_{\mathcal{S}}$ is proper and left invariant, the size of \mathcal{A} is bounded in terms of D . \square

Theorem 8.9 (Packing relative to peripheral subgroups) *Let (G, \mathbb{P}) be relatively hyperbolic, choose a finite generating set \mathcal{S} for G , and let \mathcal{H} be a finite collection of σ -relatively quasiconvex subgroups of G . For each constant D , there are constants R and M_3 such that the following holds. Let \mathcal{A} be any set of left cosets gH with $g \in G$ and $H \in \mathcal{H}$ such that for all $gH, g'H' \in \mathcal{A}$ we have $d_{\mathcal{S}}(gH, g'H') < D$. Suppose the following intersection is empty:*

$$(8-1) \quad \bigcap_{gH \in \mathcal{A}} \mathcal{N}_{M_3}(gH) = \emptyset$$

Then we have the following:

- (1) *There is a unique peripheral coset gP such that for all $aH \in \mathcal{A}$ the intersection $\mathcal{N}_R(gP) \cap \mathcal{N}_R(aH)$ is nonempty.*
- (2) *$\mathcal{N}_R(gP) \cap \mathcal{N}_R(aH) \cap \mathcal{N}_R(a'H')$ is nonempty for all $aH, a'H' \in \mathcal{A}$.*
- (3) *$\mathcal{N}_R(gP) \cap \mathcal{N}_R(aH)$ is unbounded for all $aH \in \mathcal{A}$.*
- (4) *$gPg^{-1} \cap aHa^{-1}$ is infinite for all $aH \in \mathcal{A}$.*

Proof We will first prove assertions (1) and (2). Let

$$n = 1 + \sum_{H \in \mathcal{H}} \text{height}_{(G, \mathbb{P})}(H),$$

which is known to be finite by Corollary 8.6. Let $M_0 = M_0(\sigma, D, n+1)$ be the constant given by Proposition 7.7, and let $M_2 = M_2(\sigma, D, n+1, M_0)$ be the constant given by

Corollary 7.9. If there exist $n + 1$ elements of \mathcal{A} whose M_2 -neighborhoods have empty intersection, then [Corollary 7.9](#) gives (1) and (2) immediately, using $R = M_0$. Thus it suffices to assume that for all $g_1 H_1, \dots, g_{n+1} H_{n+1} \in \mathcal{A}$ the following intersection is nonempty:

$$\mathcal{N}_{M_2}(g_1 H_1) \cap \dots \cap \mathcal{N}_{M_2}(g_{n+1} H_{n+1})$$

Fix n distinct cosets $g_1 H_1, \dots, g_n H_n \in \mathcal{A}$. (If we choose $M_3 \geq M_2$, then (8-1) implies that \mathcal{A} has at least $n + 2$ elements.) Consider the following intersection of conjugates:

$$K := g_1 H_1 g_1^{-1} \cap \dots \cap g_n H_n g_n^{-1}$$

By our choice of n , some $H \in \mathcal{H}$ appears at least $1 + \text{height}(H)$ times in the list H_1, \dots, H_n . Thus K is contained in the intersection of at least $1 + \text{height}(H)$ distinct conjugates of H , so that K is either elliptic or parabolic.

Case 1 Suppose K is elliptic. Once M_3 is chosen sufficiently large, we will reach a contradiction with (8-1). Translating \mathcal{A} by an element of G , we can assume, without loss of generality, that the cosets $g_1 H_1, \dots, g_n H_n$ all intersect the ball of radius M_2 centered at the identity. Since there are only finitely many possibilities for these cosets, our choice of M_3 is permitted to depend on $g_1 H_1, \dots, g_n H_n$. (Maximizing over all such cosets would give a uniform constant M_3 .)

By [Corollary 4.6](#) we have

$$(8-2) \quad \mathcal{N}_{M_2}(g_1 H_1) \cap \dots \cap \mathcal{N}_{M_2}(g_n H_n) \subseteq \mathcal{N}_{M'}(K)$$

for some M' depending on M_2 and the cosets $g_1 H_1, \dots, g_n H_n$. If aH is any left coset in \mathcal{A} , the following intersection is nonempty by hypothesis:

$$\mathcal{N}_{M_2}(aH) \cap (\mathcal{N}_{M_2}(g_1 H_1) \cap \dots \cap \mathcal{N}_{M_2}(g_n H_n)).$$

Thus by (8-2) we have $d_S(aH, K) < M_2 + M'$. Since K is an elliptic subgroup, it has a finite diameter $D = D(K)$ and contains the identity. Hence every $aH \in \mathcal{A}$ intersects the finite ball of radius $D + M_2 + M'$ centered at the identity, which contradicts (8-1) if we choose $M_3 \geq D + M_2 + M'$.

Case 2 Suppose K is parabolic. By [Proposition 7.4](#), for each $H \in \mathcal{S}$ and each peripheral subgroup P , the peripheral cosets gP such that $H \cap gPg^{-1}$ is infinite lie in finitely many H -orbits. Let T be the maximum distance between H and any coset gP such that $H \cap gPg^{-1}$ is infinite. Equivalently, if $gHg^{-1} \cap P$ is infinite, then the distance between gH and P is at most T .

Since K is parabolic, it is contained in a conjugate of some peripheral subgroup P . Translating \mathcal{A} by an element of G , we can assume that K is a subgroup of P itself.

For each $i = 1, \dots, n$ the group $g_i H_i g_i^{-1} \cap P$ is infinite, since it contains the infinite group K . Therefore the distance between $g_i H_i$ and P is at most T . By assumption, the following intersection is nonempty:

$$\mathcal{N}_{M_2}(g_1 H_1) \cap \dots \cap \mathcal{N}_{M_2}(g_n H_n)$$

Thus by [Corollary 7.6](#) there is a constant $R_1 = R_1(\sigma, T + M_2, n)$ and a point

$$x \in \mathcal{N}_{R_1}(P) \cap \mathcal{N}_{R_1}(g_1 H_1) \cap \dots \cap \mathcal{N}_{R_1}(g_n H_n).$$

Translating \mathcal{A} by an element of P we can assume without loss of generality that x lies in the ball of radius R_1 centered at the identity. Consequently, the cosets $g_1 H_1, \dots, g_n H_n$ each intersect the ball of radius $2R_1$ centered at the identity. As in the elliptic case, there are a uniformly bounded number of choices for $g_1 H_1, \dots, g_n H_n$.

As before, [\(8-2\)](#) holds for some M' depending on $g_1 H_1, \dots, g_n H_n$ and M_2 . Since $K \leq P$, it follows from the same reasoning as above that $d_S(aH, P) \leq d_S(aH, K) < M_2 + M'$ for each $aH \in \mathcal{A}$. Furthermore, for each $aH, a'H' \in \mathcal{A}$, the pairwise distances between $aH, a'H'$ and P are at most $D + M_2 + M'$. Thus [Proposition 7.5\(2\)](#) gives a constant $R = R(\sigma, D + M_2 + M')$ such that for all $aH, a'H' \in \mathcal{A}$ the following intersection is nonempty:

$$\mathcal{N}_R(P) \cap \mathcal{N}_R(aH) \cap \mathcal{N}_R(a'H')$$

Recall that R depends on the cosets $g_1 H_1, \dots, g_n H_n$, which each intersect the ball of radius $2R_1$ centered at the identity. Maximizing over the finitely many possible choices for these cosets gives a uniform constant—which we again call R —that does not depend on our particular choice of \mathcal{A} .

In order to prove [\(1\)](#) and [\(2\)](#) it suffices to establish the uniqueness of P in the sense of [\(1\)](#) once M_3 is chosen sufficiently large. Suppose by way of contradiction that there were another peripheral coset $g'P' \neq P$ such that for all $aH \in \mathcal{A}$ the intersection $\mathcal{N}_R(g'P') \cap \mathcal{N}_R(aH)$ is nonempty. Then [Proposition 7.8](#) implies that the intersection $\bigcap_{aH \in \mathcal{A}} \mathcal{N}_{M_1}(aH)$ is nonempty for $M_1 = M_1(R, \sigma)$. If we choose M_3 larger than M_1 , this clearly contradicts [\(8-1\)](#).

We will now show that [\(2\)](#) implies [\(3\)](#) when M_3 is chosen sufficiently large. Indeed, suppose by way of contradiction that there exists $aH \in \mathcal{A}$ such that $\mathcal{N}_R(gP) \cap \mathcal{N}_R(aH)$ is bounded. After translating \mathcal{A} by an element of G , we can assume that gP and aH both intersect the ball of radius R in G . Thus there are a bounded number of choices for the cosets gP and aH , up to translation in G . Maximizing over all such choices gives a uniform upper bound S on the diameter of the intersection $I := \mathcal{N}_R(gP) \cap \mathcal{N}_R(aH)$. If $a'H'$ is any other element of \mathcal{A} , then [\(2\)](#) implies that $\mathcal{N}_R(a'H')$ intersects I .

Therefore every $a'H' \in \mathcal{A}$ intersects a fixed set of diameter $S + R$, which contradicts (8-1) if we choose M_3 larger than $S + R$.

Finally observe that (4) follows immediately from (3) using Lemma 4.5. □

Theorem 8.10 (Packing relatively quasiconvex subgroups) *Let (G, \mathbb{P}) be relatively hyperbolic, choose a finite generating set S for G , and let H be a σ -relatively quasiconvex subgroup of G . Suppose for each peripheral subgroup $P \in \mathbb{P}$ and each $g \in G$ the intersection $P \cap gHg^{-1}$ has bounded packing in P . Then H has bounded packing in G .*

Proof Let \mathcal{A} be any set of left cosets gH whose pairwise distances are at most D . Our goal is to show that \mathcal{A} is finite and to bound the cardinality of \mathcal{A} as a function of D .

Let R and M_3 be the constants given by Theorem 8.9. Suppose the following intersection is nonempty:

$$\bigcap_{gH \in \mathcal{A}} \mathcal{N}_{M_3}(gH)$$

Then there is a fixed ball of radius M_3 in G that intersects every $gH \in \mathcal{A}$. The cardinality of such a ball is an upper bound on the cardinality of \mathcal{A} .

Thus it suffices to assume that the following intersection is empty:

$$\bigcap_{gH \in \mathcal{A}} \mathcal{N}_{M_3}(gH)$$

In this case, Theorem 8.9 applies. After translating \mathcal{A} by an element of G , there exists a peripheral subgroup P such that for all $gH, g'H \in \mathcal{A}$ the following intersection is nonempty:

$$\mathcal{N}_R(P) \cap \mathcal{N}_R(gH) \cap \mathcal{N}_R(g'H)$$

For each $gH \in \mathcal{A}$, there is a point $p \in P$ such that $d_S(p, gH) < 2R$. Translating by p^{-1} , we see that the coset $p^{-1}gH$ intersects the ball of radius $2R$ centered at the identity. Therefore every coset in \mathcal{A} can be expressed as pgH for some $p \in P$ and $|g|_S < 2R$. The cosets $gH \in \mathcal{A}$ lie in a bounded number of orbits under the action of P ; therefore it is enough to bound the number of elements of \mathcal{A} lying in a single orbit $P(gH)$; in other words, a double coset.

Setting $Q := P \cap gHg^{-1}$ the map $pgH \mapsto pQ$ that takes a left coset of H in $P(gH)$ to the corresponding coset of Q in P is well-defined and injective; for $pgH = p'gH$ if and only if $p^{-1}p' \in P \cap gHg^{-1} = Q$ if and only if $pQ = p'Q$. By Lemma 4.5, we have

$$\mathcal{N}_R(P) \cap \mathcal{N}_R(gH) \subseteq \mathcal{N}_{M'}(Q)$$

for some M' depending on P , gH and R . Translating by p gives

$$\mathcal{N}_R(P) \cap \mathcal{N}_R(pgH) \subseteq \mathcal{N}_{M'}(pQ)$$

for each $p \in P$, as illustrated in Figure 2.




Figure 2: The coset pQ is coarsely the intersection of neighborhoods of P and pgH . The existence of a point close to P , pgH and $p'gH$ implies that pQ and $p'Q$ are close.

By hypothesis, for each $pgH, p'gH \in \mathcal{A}$ we have

$$\begin{aligned} \mathcal{N}_R(P) \cap \mathcal{N}_R(pgH) \cap \mathcal{N}_R(p'gH) \\ = (\mathcal{N}_R(P) \cap \mathcal{N}_R(pgH)) \cap (\mathcal{N}_R(P) \cap \mathcal{N}_R(p'gH)) \neq \emptyset. \end{aligned}$$

Therefore $\mathcal{N}_{M'}(pQ) \cap \mathcal{N}_{M'}(p'Q) \neq \emptyset$.

But $Q = P \cap gHg^{-1}$ has bounded packing in P so we have an upper bound on the number of cosets pQ in P that correspond to cosets $pgH \in \mathcal{A}$. □

Corollary 8.11 *Let (G, \mathbb{P}) be relatively hyperbolic and $H \leq G$ a relatively quasi-convex subgroup. Suppose $P \cap gHg^{-1}$ has bounded packing in P for each peripheral subgroup $P \in \mathbb{P}$ and each $g \in G$. Then $\text{width}_{(G, \mathbb{P})}(H)$ is finite.*

Proof If g_0H and g_1H are distinct cosets such that $g_0Hg_0^{-1} \cap g_1Hg_1^{-1}$ is loxodromic, then Corollary 8.5 implies that $d_S(g_0H, g_1H) < 2C$. The result follows immediately since H has bounded packing in G by Theorem 8.10. □

9 Quasiconvex and separable implies virtually malnormal

In this section we present a malnormality consequence of separability and finite height. The results are new even in the word-hyperbolic case.

Definition 9.1 A subgroup H of G is *malnormal* if $H \cap gHg^{-1}$ is trivial for each $g \in G - H$. In applications of malnormality it is usually sufficient to know that the intersection of conjugates is finite instead of trivial, and so we have elsewhere defined H to be *almost malnormal* if $H \cap gHg^{-1}$ is finite for each $g \in G - H$.

In light of our earlier definitions of width and height of subgroups of a relatively hyperbolic group G , it is natural to define a subgroup H of G to be *relatively malnormal* if $H \cap gHg^{-1}$ is either elliptic or parabolic for each $g \in G - H$.

Note that any malnormal subgroup and any parabolic or elliptic subgroup is immediately relatively malnormal.

Definition 9.2 A subgroup H of a group G is *separable* if H is an intersection of finite index subgroups of G .

Theorem 9.3 Let H be a separable, relatively quasiconvex subgroup of the relatively hyperbolic group G . Then there is a finite index subgroup K of G containing H such that H is relatively malnormal in K .

Proof By [Corollary 8.7](#), there are only finitely many double cosets Hg_1H, \dots, Hg_nH such that $H \cap g_iHg_i^{-1}$ is loxodromic. Since H is separable, there exists a finite index subgroup K of G such that $H \leq K$ but $g_i \notin K$ for $1 \leq i \leq n$. If $k \in K - H$ and $H \cap kHk^{-1}$ is loxodromic, then $kH = hg_iH$ for some g_i and some $h \in H$. Therefore $hg_iH \subset K$ and $H \leq K$ so $g_i \in K$, contradicting our choice of K . Consequently H is relatively malnormal in K . \square

Problem 9.4 Find a relatively hyperbolic group G with a relatively malnormal subgroup M that is not quasiconvex.

There is a related long-standing problem of constructing a malnormal subgroup of a word-hyperbolic group that is not quasiconvex. One expects that [Problem 9.4](#) won't be much easier, even if we relax the condition of malnormality to merely require finite width.

References

- [1] **J M Alonso, T Brady, D Cooper, V Ferlini, M Lustig, M Mihalik, H Short (editor)**, *Notes on word hyperbolic groups*, from: “Group theory from a geometrical viewpoint (Trieste, 1990)”, (E Ghys, A Haefliger, A Verjovsky, editors), World Sci. Publ., River Edge, NJ (1991) 3–63 [MR1170363](#)
- [2] **H-J Bandelt, M van de Vel**, *Superextensions and the depth of median graphs*, J. Combin. Theory Ser. A 57 (1991) 187–202 [MR1111556](#)
- [3] **B Bowditch**, *Relatively hyperbolic groups*, Univ. of Southampton Preprint Ser. (1999)[Http://www.warwick.ac.uk/masgak/papers/bhb-relhyp.pdf](http://www.warwick.ac.uk/masgak/papers/bhb-relhyp.pdf)
- [4] **V Chepoi**, *Graphs of some CAT(0) complexes*, Adv. in Appl. Math. 24 (2000) 125–179 [MR1748966](#)
- [5] **C Druţu, M Sapir**, *Tree-graded spaces and asymptotic cones of groups*, Topology 44 (2005) 959–1058 [MR2153979](#) With an appendix by D Osin and Sapir
- [6] **B Farb**, *Relatively hyperbolic groups*, Geom. Funct. Anal. 8 (1998) 810–840 [MR1650094](#)
- [7] **V N Gerasimov**, *Semi-splittings of groups and actions on cubings*, from: “Algebra, geometry, analysis and mathematical physics (Novosibirsk, 1996)”, (Y G Reshetnyak, L A Bokut’, S K Vodop’yanov, I A Taĭmanov, editors), Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk (1997) 91–109, 190 [MR1624115](#) In Russian; translated in Siberian Adv. Math. 8 (1998) 36–58
- [8] **R Gitik, M Mitra, E Rips, M Sageev**, *Widths of subgroups*, Trans. Amer. Math. Soc. 350 (1998) 321–329 [MR1389776](#)
- [9] **M Gromov**, *Hyperbolic groups*, from: “Essays in group theory”, (S M Gersten, editor), Math. Sci. Res. Inst. Publ. 8, Springer, New York (1987) 75–263 [MR919829](#)
- [10] **G C Hruska**, *Relative hyperbolicity and relative quasiconvexity for countable groups* [arXiv:0801.4596v1](#)
- [11] **G C Hruska, D Wise**, *Finiteness properties of cubulated groups*, in preparation
- [12] **E Martínez-Pedroza**, *Combination of quasiconvex subgroups of relatively hyperbolic groups*, Groups Geom. Dyn. 3 (2009) 317–342
- [13] **G A Niblo, L D Reeves**, *Coxeter groups act on CAT(0) cube complexes*, J. Group Theory 6 (2003) 399–413 [MR1983376](#)
- [14] **G A Niblo, M A Roller**, *Groups acting on cubes and Kazhdan’s property (T)*, Proc. Amer. Math. Soc. 126 (1998) 693–699 [MR1459140](#)
- [15] **D V Osin**, *Relatively hyperbolic groups: intrinsic geometry, algebraic properties, and algorithmic problems*, Mem. Amer. Math. Soc. 179 (2006) vi+100 [MR2182268](#)
- [16] **E Rips**, *Subgroups of small cancellation groups*, Bull. London Math. Soc. 14 (1982) 45–47 [MR642423](#)

- [17] **M Roller**, *Poc sets, median algebras and group actions. An extended study of Dunwoody's construction and Sageev's theorem*, Univ. of Southampton Preprint Ser. (1998)
- [18] **J H Rubinstein**, **M Sageev**, *Intersection patterns of essential surfaces in 3-manifolds*, Topology 38 (1999) 1281–1291 [MR1690158](#)
- [19] **J H Rubinstein**, **S Wang**, *π_1 -injective surfaces in graph manifolds*, Comment. Math. Helv. 73 (1998) 499–515 [MR1639876](#)
- [20] **M Sageev**, *Ends of group pairs and non-positively curved cube complexes*, Proc. London Math. Soc. (3) 71 (1995) 585–617 [MR1347406](#)
- [21] **M Sageev**, *Codimension-1 subgroups and splittings of groups*, J. Algebra 189 (1997) 377–389 [MR1438181](#)

*Department of Mathematical Sciences, University of Wisconsin–Milwaukee
PO Box 413, Milwaukee, WI 53201, USA*

*Department of Mathematics and Statistics, McGill University
Montreal, Quebec H3A 2K6, Canada*

chruska@uwm.edu, wise@math.mcgill.ca

<http://www.uwm.edu/~chruska>, <http://www.math.mcgill.ca/~wise>

Proposed: Martin Bridson

Received: 11 September 2006

Seconded: Walter Neumann, Benson Farb

Revised: 24 March 2009