

Notiz / Note

Palladium-Catalyzed Coupling of Vinylferrocene with Aromatic Halides – A Highly Efficient Route to (Ferrocenylvinyl)arenes

Burkhard König^{*a}, Harald Zieg^a, Peter Bubenitschek^a, and Peter G. Jones^bInstitut für Organische Chemie der Technischen Universität Braunschweig^a,
Hagenring 30, 38106 Braunschweig, GermanyInstitut für Anorganische und Analytische Chemie der Technischen Universität Braunschweig^b,
Hagenring 30, 38106 Braunschweig, Germany

Received May 11, 1994

Key Words: Vinylferrocene / Palladium-catalyzed coupling / (Ferrocenylvinyl)arenes

(Ferrocenylvinyl)arenes **3**, **5**, and **7** are obtained from vinylferrocene (**1**) and substituted aromatic and heteroaromatic halides by palladium-catalyzed Heck-type reactions. Up to three ferrocene units are introduced in one step by the multi-

fold reaction of 1,2-dibromo- (**4**) or 1,3,5-tribromobenzene (**6**) with **1**. The first crystal structure of a bis(ferrocenylvinyl)benzene chromophore (**5**) is reported.

There is considerable interest in the synthesis of new materials with large second-order optical non-linearities because of their potential use in optical devices for information processing^[1]. It is now well established that molecular structures that possess both differences between ground-state and excited-state dipole moments and large transition dipole moments will have large second-order non-linearities^[2]. Molecules with π donor-acceptor interactions are promising candidates to fulfil these requirements. But in comparison with the great efforts focussed on the synthesis of organic materials such properties^[3], organometallic compounds have received little attention until recently. The vinylferrocene moiety has now been used as a π electron donor in several compounds^[4] with high second harmonic generation^[5] (SHG) efficiencies.

However, most synthetic routes to (ferrocenylvinyl)arenes are based on the Wittig reaction of ferrocenecarboxaldehyde with ylides^[6a,4c] or (ferrocenylmethyl)triphenylphosphonium iodide with aldehydes^[6b]. Over the last decade palladium-catalyzed coupling reactions of vinyl and aryl halides with alkenes (Heck reactions^[7]) have become a useful synthetic method with numerous applications. In the course of our studies of an application of this powerful methodology to the synthesis of defined polynuclear metal complexes, we have observed the facile coupling of vinylferrocene with various aryl halides.


Iodobenzene reacts with vinylferrocene under the phase-transfer conditions described by Jeffery^[8] to yield (*E*)-styrylferrocene^[9] as the only product. The coupling reaction can be extended to electron-deficient or electron-rich aromatic rings with nitro or methoxy substituents. Heterocyclic examples^[10] of the procedure are the reactions of **1** with 2-bromo-6-methylpyridine (**2d**) and 2-bromo-pyrimidine (**2e**).

Two or three ferrocene units are introduced in a one-pot reaction by the multifold coupling of **1** with 1,2-di- (**4**) and 1,3,5-tribromobenzene (**6**)^[11].

Suitable crystals for an X-ray analysis of **5** have been obtained from dichloromethane. The crystal structure determination^[12] shows that only one of the two vinylferrocene moieties is essentially


parallel to the central aromatic ring (Figure 1). The interplanar angles to this ring are 17° for C9–13 and 60° for C9'–13', with corresponding torsion angles C1–7–8–9 179, C7–8–9–10 -7, C6–7'–8'–9' -175, and C7'–8'–9'–10' -158°.

Scheme 1. Coupling reactions of vinylferrocene (**1**) with aromatic halides. ^a Pd(OAc)₂, DMF, K₂CO₃, NBu₄Br




The palladium-catalyzed coupling of **1** with aryl halides is a simple and effective procedure for the synthesis of (ferrocenylvinyl)arenes with obvious advantages over stoichiometric reactions in terms of "atom economy"^[13]. Practical applications to the synthesis of ferrocene derivatives as electrochemically switchable non-linear optical materials^[14] and photodegradation protection units^[15] of conjugated chain molecules might be envisaged.

This work was supported by the *Fonds der Chemischen Industrie*. Generous gifts of chemicals by the *Degussa AG* are gratefully acknowledged. B. K. is grateful to the *Fonds der Chemischen Indu-*



a: 2 equiv. of **1**, Pd(OAc)₂ [5 mol-%], DMF, K₂CO₃, NBu₄Br.
Yield: 74 %.



a: 3 equiv. of **1**, Pd(OAc)₂ [5 mol-%], DMF, K₂CO₃, NBu₄Br.
Yield: 46 %.

strie for a Liebig stipend. We thank Prof. Dr. H. Hopf for his generous support.

Experimental

¹H NMR: Bruker AM 400; $\delta = 0$ for tetramethylsilane as internal standard, $\delta = 7.26$ for chloroform. – ¹³C NMR: AM 400; $\delta = 77.0$ for deuteriochloroform; the multiplicity of the ¹³C-NMR signals was determined by the DEPT technique and quoted as (+) for CH₃ and CH groups, (–) for CH₂- and (C_{quart}) for quaternary carbons. – IR: Nicolet 320 FT-IR. – MS: Finnigan MAT 8430. – UV/Vis: Hewlett Packard 8452A. – Melting points: Hot-plate microscope apparatus, not corrected. – Column chromatography (CC): Merck silica gel 60, mesh 70–230; PE (60/70) means petroleum ether with a boiling range of 60–70°C. – TLC: silica gel (Macherey-Nagel; Sil G/UV₂₅₄).

X-Ray Structure Determination of Compound 5^[12]: C₃₀H₂₆Fe₂, *M* = 498.21, monoclinic, space group *C2/c*, *a* = 2711.3(3), *b* = 722.2(1), *c* = 2337.2(3) pm, $\beta = 102.28(2)^\circ$, *V* = 4.4718(10) nm³, *Z* = 8, $\lambda(\text{Mo-K}\alpha) = 0.71073 \text{ \AA}$, $\mu = 1.3 \text{ mm}^{-1}$, *D*_x = 1.480 Mg m⁻³, *F*(000) = 2064, *T* = 143 K. A red prism with the dimensions 0.6 × 0.3 × 0.15 mm was mounted on a glass fiber in inert oil (type RS3000, donated by Fa. Riedel-de Haën) and transferred to the cold gas stream of a Stoe STADI-4 diffractometer with a Siemens LT-2 low-temperature attachment. Cell constants were refined from $\pm\infty$ values of 54 reflections in the 2 θ range 20–23°. A total 4051 intensities (3940 unique, *R*_{int} = 0.0208) were measured to 2 θ 50°. – The structure was solved by direct methods and refined anisotropically on *F*² (program SHELXL-93, G. M. Sheldrick, University of Göttingen). Hydrogen atoms were included with a riding model. The final *wR*(*F*²) for all reflections was 0.121, with a conventional *R*(*F*) of 0.046, for 289 parameters.

General Procedure (GP) for the Palladium-Catalyzed Coupling of Vinylferrocene (1): A mixture of 1.0 mmol of **1**, 10.0 mmol of potassium carbonate, 4.0 mmol of tetrabutylammonium bromide, the given amount of the appropriate halide, and 25 mg (0.11 mmol) of Pd(OAc)₂ in 20 ml of dimethylformamide (DMF) was heated in a capped pyrex tube under nitrogen for the time stated. The reaction mixture was filtered, diluted with 100 ml of dichloromethane and washed with six portions of 50 ml of water. The organic phase was dried with K₂CO₃, filtered, and the solvent was removed from the filtrate in vacuo. The crude products were purified by column chromatography (CC) on silica gel and subsequent recrystallization.

(*E*)-Styrylferrocene (**3a**)^[9]: **2a** (214 mg, 1.0 mmol) was allowed to react with **1** (204 mg, 1.0 mmol) according to the GP for 2 d at 70°C, and the crude product was chromatographed on silica gel (PE/CH₂Cl₂, 100:3) to yield 245 mg (85%) of pure **3a** (*R*_f = 0.17) as a red solid, m.p. 119°C^[9].

(*E*)-(4-Nitrostyryl)ferrocene (**3b**)^[4c]: **2b** (125 mg, 0.5 mmol) and **1** (107 mg, 0.5 mmol) were heated to 55°C for 2 d according to the GP. The crude product was chromatographed on silica gel (PE/CH₂Cl₂, 2:3) to yield 141 mg (84%) of **3b** (*R*_f = 0.44) as a dark red solid, m.p. 196°C^[4c].

(*E*)-(4-Methoxystyryl)ferrocene (**3c**)^[16]: **2c** (187 mg, 1.0 mmol) and **1** (214 mg, 1.0 mmol) were allowed to react according to the GP for 2 d at 80°C. The crude product was purified by CC on




Figure 1. Crystal structure of **5**. Selected bond lengths [pm]: C(7)–C(8) 133.2(5), C(1)–C(7) 146.7(6), C(8)–C(9) 146.1(6); bond angles [°]: C(8)–C(7)–C(1) 126.2(4), C(7)–C(8)–C(9) 127.1(4)

silica gel (PE/CH₂Cl₂, 7:3) to afford 142 mg (45%) of **3c** ($R_f = 0.27$) as orange plates, m.p. 128°C.

(*E*)-2-(2-Ferrocenylvinyl)-6-methylpyridine (**3d**): **2d** (170 mg, 1.0 mmol) and **1** (214 mg, 1.0 mmol) were allowed to react for 2 d at 80°C. By CC on silica gel (CH₂Cl₂/MeOH, 98:2) 112 mg (37%) of **3d** ($R_f = 0.38$) was obtained as an orange solid, m.p. 53°C. – IR (KBr): $\tilde{\nu} = 1636 \text{ cm}^{-1}$, 1565, 816, 782, 489. – UV (CH₃CN): λ_{max} (lg ϵ) = 194 nm (4.584), 260 (4.028), 320 (4.323), 414 (3.603). – ¹H NMR (CDCl₃): $\delta = 2.56$ (s, 3H), 4.13 (s, 5H), 4.39 (m, 2H), 4.50 (m, 2H), 6.77 (d, $J = 16.0$ Hz, 1H), 6.96 (d, $J = 7.6$ Hz, 1H), 7.17 (d, $J = 7.8$ Hz, 1H), 7.30 (d, $J = 16.0$ Hz, 1H), 7.49 (dd, $J = 7.7/7.6$ Hz, 1H). – ¹³C NMR (CDCl₃): $\delta = 24.64$ (+), 67.39 (+), 69.27 (+), 69.36 (+), 82.25 (C_{quart}), 117.46 (+), 120.78 (+), 126.14 (+), 131.26 (+), 136.49 (+), 156.30 (C_{quart}), 158.10 (C_{quart}). – MS (70 eV), m/z (%): 303 (50) [M⁺], 238 (100) [M⁺ – Cp]. – C₁₈H₁₇FeN: calcd. 303.0710; found 303.0710 (MS).

(*E*)-2-(2-Ferrocenylvinyl)pyrimidine (**3e**): **2e** (110 mg, 0.7 mmol) was allowed to react with **1** (200 mg, 0.9 mmol) for 16 h at 110°C according to the GP. CC on silica gel (CH₂Cl₂/MeOH, 97:3) yielded 27 mg (14%) of **3e** ($R_f = 0.64$) as a dark red solid, m.p. 143°C. – IR (KBr): $\tilde{\nu} = 1635 \text{ cm}^{-1}$, 1565, 1437. – UV (CH₃CN): λ_{max} (lg ϵ) = 208 nm (4.382), 264 (4.042), 310 (4.202), 464 (3.174). – ¹H NMR (CDCl₃): $\delta = 4.16$ (s, 5H), 4.37 (m, 2H), 4.56 (m, 2H), 6.82 (d, $J = 15.8$ Hz, 1H), 7.05 (t, $J = 4.9$ Hz, 1H), 7.80 (d, $J = 15.7$ Hz, 1H), 8.65 (d, $J = 4.8$ Hz, 2H). – ¹³C NMR (CDCl₃): $\delta = 68.08$ (+), 69.47 (+), 70.12 (+), 81.10 (C_{quart}), 117.70 (+), 124.51 (+), 138.31 (+), 157.00 (+), 160.30 (C_{quart}). – MS (70 eV), m/z (%): 290 (52) [M⁺], 225 (66) [M⁺ – Cp], 132 (100). – C₁₆H₁₄FeN₂: calcd. 290.0506; found 290.0502 (MS).

(*E,E*)-1,2-Bis(2-ferrocenylvinyl)benzene (**5**): **4** (188 mg, 0.8 mmol) and **1** (360 mg, 1.7 mmol) were heated for 3 d to 70°C according to the GP, and the crude product was chromatographed on silica gel (1. PE/CH₂Cl₂, 95:5; 2. CH₂Cl₂) to yield 313 mg (74%) of **5** ($R_f = 0.8$) as a red solid, m.p. 182°C. – IR (KBr): $\tilde{\nu} = 1626 \text{ cm}^{-1}$, 1104, 957, 833, 716, 487. – UV (CH₃CN): λ_{max} (lg ϵ) = 196 nm (4.770), 204 (4.905), 256 (4.389), 298 (4.473), 456 (3.383). – ¹H NMR (CDCl₃): $\delta = 4.18$ (s, 10H), 4.31 (m, 4H), 4.52 (m, 4H), 6.77 (d, $J = 16.0$ Hz, 2H), 7.04 (d, $J = 16$ Hz, 2H), 7.26 (m, 2H), 7.51 (m, 2H). – ¹³C NMR (CDCl₃): $\delta = 66.95$ (+), 69.10 (+), 69.27 (+), 83.60 (+), 124.19 (+), 126.31 (+), 126.99 (+), 129.17 (+), 135.71 (C_{quart}). – MS (70 eV), m/z (%): 498 (100) [M⁺], 433 (10) [M⁺ – Cp]. – C₃₀H₂₆Fe₂ (498.2): calcd. C 72.32, H 5.26; found C 72.29, H 5.22. – Mol. mass 498 (MS).

(*E,E,E*)-1,3,5-Tris(2-ferrocenylvinyl)benzene (**7**): 500 mg (0.236 mmol) of **1** and 189 mg (0.6 mmol) of 1,3,5-tribromobenzene (**6**) were allowed to react according to the GP for 2 d at 60°C. The crude product was chromatographed on silica gel (PE/CH₂Cl₂, 1:1) to yield 195 mg (46%) of **7** ($R_f = 0.47$) as a red solid, m.p. 172°C. – IR (KBr): $\tilde{\nu} = 1635 \text{ cm}^{-1}$, 491. – UV (CHCl₃): λ_{max} (lg ϵ) = 242 nm (4.328), 316 (4.681), 362 (3.995), 456 (3.552). – ¹H NMR (CDCl₃): $\delta = 4.16$ (s, 15H), 4.31 (m, 6H), 4.49 (m, 6H), 6.73 (d, $J = 16.1$ Hz, 3H), 6.95 (d, $J = 16.1$ Hz), 7.36 (s, 3H). – MS (70 eV), m/z (%): 708 (97) [M⁺], 44 (100). – C₄₂H₃₆Fe₃: calcd. 708.0868, found 708.0867 (MS).

- [1] [1a] D. J. Williams, *Angew. Chem.* **1984**, *96*, 637–651; *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 690–704. – [1b] R. W. Boyd, *Nonlinear Optics*, Academic Press, New York, **1992**. – [1c] D. F. Eaton, *Science* **1991**, *253*, 281–287. – [1d] S. R. Marder, D. N. Beratan, L.-T. Cheng, *Science* **1991**, *252*, 103–106.
- [2] [2a] J. L. Oudar, D. S. Chemla, *J. Chem. Phys.* **1977**, *66*, 2664–2668. – [2b] B. F. Levine, C. C. Betea, *J. Chem. Phys.* **1977**, *66*, 1070–1074.
- [3] For example: [3a] M. Barzoukas, M. Blanchard-Desce, D. Josse, J.-M. Lehn, J. Zyss, *Chem. Phys.* **1989**, *133*, 323–329. – [3b] R. A. Huijts, G. L. J. Hesselink, *Chem. Phys. Lett.* **1989**, *156*, 209–212.
- [4] [4a] M. L. H. Green, S. R. Marder, M. E. Thompson, J. A. Bandy, D. Bloor, P. V. Kolinsky, R. J. Jones, *Nature* **1987**, *330*, 360–362. – [4b] J. C. Calabrese, L.-T. Cheng, J. C. Green, S. R. Marder, W. Tam, *J. Am. Chem. Soc.* **1991**, *113*, 7227–7232. – [4c] H. E. Bunting, M. L. H. Green, S. R. Marder, M. E. Thompson, D. Bloor, P. V. Kolinsky, R. J. Jones, *Polyhedron* **1992**, *11*, 1489–1499. – [4d] S. R. Marder, J. W. Perry, B. G. Tiemann, W. P. Schaefer, *Organometallics* **1991**, *10*, 1896–1901. – [4e] S. Ghosal, M. Samoc, P. N. Prasad, J. J. Tufarillo, *J. Phys. Chem.* **1990**, *94*, 2847–2851. – [4f] Calculations: D. R. Kanis, M. A. Ratner, T. J. Marks, *J. Am. Chem. Soc.* **1992**, *114*, 10338–10357.
- [5] [5a] J. F. Nicoud, R. J. Twieg in *Nonlinear Optical Properties of Organic Molecules and Crystals* (Eds.: D. S. Chemla, J. Zyss), Academic Press, New York, **1987**. – [5b] The process of second harmonic generation (SHG) is one of the second-order nonlinear optical effects of practical importance. SHG is a process by which light of the frequency ω is converted to the frequency 2ω upon interaction with the non-linear medium.
- [6] [6a] Osgerby, P. L. Pauson, *J. Chem. Soc.* **1961**, 4604–4609. – [6b] P. L. Pauson, W. E. Watts, *J. Chem. Soc.* **1963**, 2990–2996. – [6c] For a McMurry-type coupling to (ferrocenylvinyl)arenes see: S. El-Tamany, F.-W. Raulfs, H. Hopf, *Angew. Chem.* **1983**, *95*, 631; *Angew. Chem. Int. Ed. Engl.* **1983**, *22*, 633.
- [7] [7a] R. F. Heck, *Palladium Reagents in Organic Synthesis*, Academic Press, London, **1985**. – [7b] R. F. Heck, G. Wu, W. Tao, A. L. Reingold in *Catalysis of Organic Reactions* (Ed.: D. W. Blackburn), Marcel Dekker, New York, **1989**. – [7c] H.-U. Reibig, *Nachr. Chem. Tech. Lab.* **1986**, *34*, 1066–1073.
- [8] [8a] T. Jeffery, *Tetrahedron Lett.* **1985**, *26*, 2667–2670. – [8b] T. Jeffery, *Tetrahedron Lett.* **1990**, *31*, 6641–6644.
- [9] G. Drefahl, G. Plötner, I. Winnefeld, *Chem. Ber.* **1962**, *95*, 2788–2791.
- [10] [10a] For the synthesis of 2-(ferrocenylvinyl)pyridine see: A. M. Osman, K. M. Hassan, Z. H. Khalil, V. D. Turin, *J. Appl. Chem. Biotechnol.* **1976**, *26*, 71–78. – [10b] Yields for **3d** and **3e** are not optimized.
- [11] The reaction of **1** with hexabromobenzene yielded a sixfold coupling product, as clearly indicated by mass spectroscopy. However, the compound was obtained as a mixture of conformers giving rise to complicated ¹H- and ¹³C-NMR spectra.
- [12] [12a] Full details of the crystal structure investigation are available on request from the Fachinformationszentrum Karlsruhe, Gesellschaft für wissenschaftlich-technische Information mbH, D-76344 Eggenstein-Leopoldshafen, on quoting the depository number CSD-400973, the names of the authors, and the journal citation. – [12b] For the structure of (Z)-(1,2-diphenylethynyl)-ferrocene see: Ch. J. Cardin, W. Crawford, W. E. Watts, B. J. Hathaway, *J. Chem. Soc., Dalton Trans.* **1979**, 970–974.
- [13] B. M. Trost, *Science* **1991**, *254*, 1471–1477.
- [14] [14a] K. L. Kott, R. J. McMahon, *J. Org. Chem.* **1992**, *57*, 3097–3101. – [14b] P. D. Beer, H. Sikanyika, *Polyhedron* **1990**, *9*, 1091–1094.
- [15] T. Arai, Y. Ogawa, H. Sakuragi, K. Tokumaru, *Chem. Phys. Lett.* **1992**, *196*, 145–149.
- [16] E. Solacianiova, S. Toma, T. Liptaj, *Collect. Czech. Chem. Commun.* **1986**, *51*, 670–676.

[174/94]