



Published in final edited form as:

Angew Chem Int Ed Engl. 2019 February 11; 58(7): 2144–2148. doi:10.1002/anie.201813699.

Palladium/Norbornene-Catalyzed Indenone Synthesis from Simple Aryl Iodides: Concise Syntheses of Pauciflorol F and Acredinone A

Feipeng Liu,

Department of Applied Chemistry, China Agricultural University, Beijing 100193, China,
Department of Chemistry, University of Chicago, Chicago, IL, 60637 (USA)

Zhe Dong,

Department of Chemistry, University of Chicago, Chicago, IL, 60637 (USA)

Jianchun Wang*, and

Department of Chemistry, University of Chicago, Chicago, IL, 60637 (USA)


Guangbin Dong*

Department of Chemistry, University of Chicago, Chicago, IL, 60637 (USA)

Abstract

To show the synthetic utility of palladium/norbornene (Pd/NBE) cooperative catalysis, here we report concise syntheses of indenone-based natural products, pauciflorol F and acredione A, which are enabled by direct annulation between aryl iodides and unsaturated carboxylic acid anhydrides. Compared to the previous indenone-preparation approaches, this method allows simple aryl iodides to be used as substrates with complete control of the regioselectivity. The total synthesis of acredione A features two different Pd/NBE-catalyzed *ortho* acylation reactions for constructing penta-substituted arene cores, including the development of a new *ortho* acylation/*ipso* borylation.

Graphical Abstract



A straightforward method for indenone synthesis was achieved via Pd/NBE-catalyzed annulation between simple aryl iodides and unsaturated carboxylic acid anhydrides. This method enabled streamlined syntheses of pauciflorol F and acredione A.

* jcwang1992@uchicago.edu (J.W.) gbdong@uchicago.edu (G.D.).

Keywords

acredininone A; pauciflorol F; indenone; norbornene; palladium

Indenones are commonly found in pharmaceuticals and bioactive natural products (Scheme 1a),^[1] and often serve as versatile intermediates to access polysubstituted indanones.^[2] While a number of methods have been developed for preparing indenones, conventional approaches usually take multiple steps and/or have limited scopes.^[3] Recently, the transition metal-catalyzed cyclization with alkynes as coupling partners offers a broad and useful strategy to form various indenones (Scheme 1b).^[4,5] However, it is not trivial to control regioselectivity when unsymmetrical alkynes are used.^[2] In addition, vicinal difunctionalized aryl substrates^[4] or those containing a directing group (DG)^[5] are needed. Thus, it remains attractive to realize a general and regioselective indenone synthesis from simple readily available starting materials.

On the other hand, palladium/norbornene (Pd/NBE) cooperative catalysis,^[6] originally discovered by Catellani,^[7] has become an increasingly useful approach for preparing polysubstituted arenes via simultaneous functionalization of both *ortho* and *ipso* positions of aryl halides.^[8] Notably, Lautens has pioneered in developing bifunctional reagents, which enabled forming various condensed rings via *ipso*-annulation reactions.^[8a,9] While numerous Pd/NBE catalysis methods have been developed, only a few have been successfully applied to the total synthesis of natural products.^[8p,10] Herein, we describe concise syntheses of pauciflorol F(**1**) and acredininone A (**2**), enabled by the development of a straightforward method to construct indenones from simple aryl iodides and unsaturated carboxylic anhydrides via Pd/NBE catalysis (Scheme 1c).

In 2015, we reported an initial study on the *ortho* acylation/*ipso* hydrogenation using a mixed anhydride.^[8i] Concurrently, the Liang^[8k] and Gu^[8j] groups independently developed the *ortho* acylation/*ipso* Heck reaction. Interestingly, when anhydrides derived from α,β -unsaturated carboxylic acids were employed, indenone products could be obtained (Scheme 1c). The reaction was likely initiated by oxidative addition of Pd(0) into the Ar–X bond, subsequent NBE migratory insertion and C–H metalation to generate an aryl-NBE-palladacycle (ANP) **I**,^[11] which would then react with the anhydride to afford intermediate **II**. After NBE extrusion, the resulting intermediate **III** could further undergo intramolecular cyclization to give the indenone product (*vide infra*, Scheme 3).

The reaction was further optimized using 2-iodotoluene (**3a**) as the substrate (see Supporting Information, Table S1). [Pd(allyl)Cl]₂ and tri(2-furyl)phosphine still proved to be the best metal/ligand combination, and the amide-substituted NBE (**N1**) was also found more efficient than simple NBE. Under the optimal conditions, the scope of the reaction was examined (Table 1). Aryl iodides contain both electron-rich and deficient substituents were suitable substrates, and different substitution patterns on arenes were tolerated. Note that sterically hindered aryl iodides, such as 2,5-disubstituted ones (**5q** and **5r**), were also competent substrates, which was the key requirement for the synthesis of acredininone A. Other unsaturated anhydrides were explored next as coupling partners. Anhydrides derived from alkyl (**5t**) and aryl-substituted acrylic acids (**5u-5ac**) could be successfully employed.

[12] In particular, when α - or α,β -disubstituted acrylic anhydrides were used, 2-substituted and 2,3-disubstituted indenones (**5aa-5ac**) were still afforded in good yields. Finally, a number of functional groups were found compatible, such as tertiary amine (**5c**, **5j** and **5k**), silyl ether (**5g**), tertiary alcohol (**5i**), MOM-protected alcohol (**5e**), Weinreb amide (**5n**), ester (**5f**, **5m** and **5p**), Boc-protected amine (**5l**), trifluoromethyl (**5w**), aryl bromide (**5v**) and thiophene (**5z**), making this method attractive for complex molecule synthesis.

Preliminary study showed that aryl bromides^[13] could also be used as the substrate (eq. 1, Scheme 2) albeit in a lower efficiency; the use of large bite-angle (DPEPhos) ligand^[13a] and additional CsI was important. Not surprisingly, *ortho*-unsubstituted aryl iodide **3ad** was not a suitable substrate using **N1** or simple NBE under the standard conditions due to the “*ortho* constraint”.^[14] However, with the recently developed bridgehead-substituted norbornene (**N2**),^[14b] the desired indenone product **5ad** was successfully isolated in 56% yield.

The mechanism regarding the conversion of intermediate **III** into indenone product was explored (Scheme 3). Two pathways are possible: **path a** involves activation of the vinyl β -C(sp²)-H bond of the enone intermediate followed by reductive elimination, while **path b** is through a Heck-like sequence: aryl migratory insertion into the olefin followed by β -H elimination. To differentiate these two pathways, both (*Z*) and (*E*)-anhydrides (**4aa**) were allowed to react with **3aa** in parallel under the standard conditions. Interestingly, they gave almost identical yields of **5aa** with a similar kinetic profile (see Supporting Information). Additionally, *Z/E* isomerization of the olefin moiety in **4aa** was not observed during the reaction. Taken together, the observations are consistent with the Heck-like pathway (**path b**), though other pathways including a Nazarov-type cyclization cannot be excluded at the current stage (see Supporting Information).

The utility of the method was first demonstrated in the formal synthesis of pauciflorol F (**1**) (Scheme 4). As a polyphenolic natural product derived from resveratrol, pauciflorol F has been an attractive synthetic target since its discovery in 2004.^[15] Several elegant routes have been developed to date.^[2,16] Our approach started with a two-step preparation of mixed anhydride **10** from commercially available aldehyde **8**.^[17] As expected, the subsequent Pd/**N1**-catalyzed annulation between anhydride **10** and aryl iodide **3t** provided indenone **11** in 82% yield, whose structure was confirmed by X-ray crystallography.^[18] Note that the prior synthesis of indenone **11** via an alkyne insertion strategy generated a 1:1 mixture of regioisomers.^[2] Demonstrated by Jeffery and Sarpong, indenone **11** can be converted to **1** in two steps;^[2] thus, the synthesis of pauciflorol F can now be accomplished in 5 total steps and a high overall yield (~50%).

We then focused on the total synthesis of acredinone A (**2**). Acredinone A was recently isolated in <5 mg (with $[\alpha]_D^{25}$: 0 in CHCl₃) from a marine-sponge-associated *acremonium* sp. Fungus, which represents the first nonpeptidic natural product that can inhibit the voltage-gated potassium channel.^[19] Such an activity holds potential for developing inhibitors for treating type-II diabetes.^[20] In addition, total synthesis of acredinone A has not been reported to date, and the structure-activity relationship (SAR) remains unknown. Given the pseudo-symmetry in **2**, it might be possible to establish an efficient biomimetic strategy. However, to enable rapid access to structurally diversified analogues for future SAR

studies, it could be more valuable to develop a modular and convergent approach. The *challenges* are how to construct the highly congested indenone core and how to prepare the two penta-substituted arenes efficiently. From a retrosynthetic viewpoint (Scheme 5), we envisioned that acredione A could be constructed via a late-stage Suzuki–Miyaura coupling of two fragments (**12** and **13**) with similar complexity. Each fragment would be synthesized via a unique Pd/NBE catalysis. Ultimately, the natural product would be assembled using four arene building blocks (**14–17**).

In a forward manner, aryl iodides **14** and **16** were synthesized from the same commercially available 3-methyl-4-anisaldehyde **18** in 4 steps (Scheme 6). The Pd/NBE-catalyzed indenone synthesis proceeded in a good yield on gram scales despite the bulkiness of the substrates. Bromination of indenone **19** with NBS afforded fragment **12**. Meanwhile, fragment **13** was proposed to be prepared via Pd/NBE-catalyzed *ortho* acylation/*ipso* borylation, which has been an unknown transformation. Encouraged by Ritter's *ortho* amination/*ipso* borylation,^[21] the three-component coupling of aryl iodide **16**, anhydride **17** and B₂pin₂ was carefully studied (Table 2). To our delight, the desired product **13a** was obtained in 21% yield in the first attempt using Pd(OAc)₂ as the catalyst, and the major side-reaction was the *ipso*-hydrogenation (**13a'**) (entry 1). A quick condition screening revealed that the use of Pd(dba)₂ could suppress the formation of side-product **13a'** (entries 1–4), though the exact reason is unclear. Further tuning the solvent and reaction temperature finally afforded the desired borylation product **13a** in 67% isolation yield (entries 5–7).

The Suzuki–Miyaura coupling between aryl bromide **12** and aryl boronic ester **13a** turned out to be extremely challenging due to the large steric hindrance in both coupling partners. The commonly employed monodentate or bidentate ligands, such as PPh₃, P^tBu₃, Buchwald's biarylphosphines, Hartwig's Q-Phos and NHC ligands, gave trace or no desired product. After extensive studies, gratifyingly, the bulky and electron-rich bidentate dtbpf ligand afforded the desired coupling product **20** albeit in a low yield.^[22] The major side-reaction was the protonation of boronic ester **13a**, suggesting low stability of this compound under the reaction conditions. Considering that Molander salts generally exhibit enhanced stability than the corresponding boronic esters in cross couplings,^[23] the corresponding potassium aryltrifluoroborate **13b** was then prepared. Indeed, the coupling between **12** and **13b** proceeded smoothly to afford product **20** in 68% yield, which was subsequently deprotected to give acredione A (**2**). The spectroscopic data of synthetic **2** matched those of the reported natural sample in all respects. Hence, the first total synthesis of acredione A was accomplished in eight steps in the longest linear sequence (LLS) from commercial chemicals.

In summary, a Pd/NBE-catalyzed annulation between aryl iodides and unsaturated carboxylic acid anhydrides has been developed, which provides a simple and straightforward approach for preparing indenones in a regio- and chemoselective manner. Using this method, concise syntheses of pauciflorol F and acredione A have been achieved. The utility of Pd/NBE catalysis for streamlined synthesis of tetra- and penta-substituted arenes could have broad implications beyond this work. Efforts on preparing analogues of acredione A for future SAR studies are ongoing.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial supports from the University of Chicago and NIGMS (1R01GM124414-01A1) are acknowledged. F.L. is supported by a CSC fellowship. Mr. Ki-Young Yoon is thanked for checking the experiment and for X-ray crystallography.


References

- [1]. (a)Morrell A, Placzek M, Parmley S, Grella B, Antony S, Pommier Y, Cushman M, J. Med. Chem 2007, 50, 4388; [PubMed: 17676830] (b)Ahn JH, Shin MS, Jung SH, Kang SK, Kim KR, Rhee SD, Jung WH, Yang SD, Kim SJ, Woo JR, Lee JH, Cheon HG, Kim SS, J. Med. Chem 2006, 49, 4781. [PubMed: 16854085]
- [2]. Jeffrey JL, Sarpong R, Org. Lett 2009, 11, 5450. [PubMed: 19902905]
- [3]. (a)Marvel CS, Hinman CW, J. Am. Chem. Soc 1954, 76, 5435;(b)Martens H, Hoornaer G, Tetrahedron 1974, 30, 3641;(c)Vasilyev AV, Walspurger S, Haouas M, Sommer J, Pale P, Rudenko AP, Org. Biomol. Chem 2004, 2, 3483; [PubMed: 15565241] (d)Vasilyev AV, Walspurger S, Pale P, Sommer J, Tetrahedron Lett 2004, 45, 3379;(e)Pan C, Huang B, Hu W, Feng X, Yu J, J. Org. Chem 2016, 81, 2087; [PubMed: 26840747] (f)Zhang X-S, Jiao J-Y, Zhang X-H, Hu B-L, Zhang X-G, J. Org. Chem 2016, 81, 5710; [PubMed: 27232156] (g)Suchand B, Satyanarayana G, J. Org. Chem 2017, 82, 372. [PubMed: 27935714]
- [4]. (a)Larock RC, Doty MJ, Cacchi S, J. Org. Chem 1993, 58, 4579;(b)Alonso DA, Najera C, Pacheco MC, Adv. Synth. Catal 2002, 344, 172;(c)Pletnev AA, Tian Q, Larock RC, J. Org. Chem 2002, 67, 9276; [PubMed: 12492329] (d)Miura T, Murakami M, Org. Lett 2005, 7, 3339; [PubMed: 16018655] (e)Harada Y, Nakanishi J, Fujihara H, Tobisu M, Fukumoto Y, Chatani N, J. Am. Chem. Soc 2007, 129, 5766; [PubMed: 17417848] (f)Tsukamoto H, Kondo Y, Org. Lett 2007, 9, 4227; [PubMed: 17867697] (g)Liu CC, Korivi RP, Cheng CH, Chem. Eur. J 2008, 14, 9503; [PubMed: 18810733] (h)Morimoto T, Yamasaki K, Hirano A, Tsutsumi K, Kagawa N, Kakiuchi K, Harada Y, Fukumoto Y, Chatani N, Nishioka T, Org. Lett 2009, 11, 1777; [PubMed: 19301863] (i)Yan X, Zou S, Zhao P, Xi C, Chem. Commun 2014, 50, 2775.
- [5]. (a)Li B-J, Wang H-Y, Zhu Q-L, Shi Z-J, Angew. Chem. Int. Ed 2012, 51, 3948; Angew. Chem. 2012, 124, 4014;(b)Chen S, Yu J, Jiang Y, Chen F, Cheng J, Org. Lett 2013, 15, 4754; [PubMed: 24011165] (c)Qi Z, Wang M, Li X, Org. Lett 2013, 15, 5440. [PubMed: 24117207]
- [6]. (a)Catellani M, Synlett 2003, 298–313;(b)Catellani M, Top. Organomet. Chem 2005, 14, 21; (c)Catellani M, Motti E, Della Ca' N, Acc. Chem. Res 2008, 41, 1512; [PubMed: 18680317] (d)Martins A, Mariampillai B, Lautens M, Top. Curr. Chem 2010, 292, 1; [PubMed: 21500401] (e)Ye J, Lautens M, Nat. Chem 2015, 7, 863; [PubMed: 26492005] (f)Della Ca' N, Fontana M, Motti E, Catellani M, Acc. Chem. Res 2016, 49, 1389. [PubMed: 27333299]
- [7]. Catellani M, Frignani F, Rangoni A, Angew. Chem. Int. Ed 1997, 36, 119; Angew. Chem. 1997, 109, 142.
- [8]. For selected examples of Pd/NBE catalysis: a)Lautens M, Piguel S, Angew. Chem. Int. Ed 2000, 39, 1045; Angew. Chem. 2000, 112, 1087;(b)Catellani M, Motti E, Baratta S, Org. Lett 2001, 3, 3611; [PubMed: 11700094] (c)Faccini F, Motti E, Catellani M, J. Am. Chem. Soc 2004, 126, 78; [PubMed: 14709068] (d)Bressy C, Alberico D, Lautens M, J. Am. Chem. Soc 2005, 127, 13148; [PubMed: 16173734] (e)Mariampillai B, Alberico D, Bidau V, Lautens M, J. Am. Chem. Soc 2006, 128, 14436; [PubMed: 17090008] (f)Maestri G, Motti E, Della Ca' N, Malacria M, Derat E, Catellani M, J. Am. Chem. Soc 2011, 133, 8574; [PubMed: 21563760] (g)Dong Z, Dong G, J. Am. Chem. Soc 2013, 135, 18350; [PubMed: 24256439] (h)Zhang H, Chen P, Liu G, Angew. Chem. Int. Ed 2014, 53, 10174; Angew. Chem. 2014, 126, 10338;(i)Dong Z, Wang J, Ren Z, Dong G, Angew. Chem. Int. Ed 2015, 54, 12664; Angew. Chem. 2015, 127, 12855;(j)Huang YZ, Zhu R, Zhao K, Gu ZH, Angew. Chem. Int. Ed 2015, 54, 12669; Angew. Chem. 2015, 127, 12860.(k)Zhou P-X, Ye Y-Y, Liu C, Zhao L-B, Hou J-Y, Chen D-Q, Tang Q, Wang A-Q, Zhang


J-Y, Huang Q-X, Xu P-F, Liang Y-M, ACS Catal 2015, 5, 4927;(l)Wang J, Zhang L, Dong Z, Dong G, Chem 2016, 1, 581;(m)Cheng H-G, Wu C, Chen H, Chen R, Qian G, Geng Z, Wei Q, Xia Y, Zhang J, Zhang Y, Zhou Q, Angew. Chem. Int. Ed 2018, 57, 3444; Angew. Chem. 2018, 130, 3502;(n)Li R, Dong G, Angew. Chem. Int. Ed 2018, 57, 1697; Angew. Chem. 2018, 130, 1713;(o)for Pd(II)-initiated Pd/NBE catalysis, see:Jiao L, Bach T, J. Am. Chem. Soc 2011, 133, 12990; [PubMed: 21806073] (p)Jiao L, Herdtweck E, Bach T, J. Am. Chem. Soc 2012, 134, 14563; [PubMed: 22913367] (q)Wang X-C, Gong W, Fang L-Z, Zhu R-Y, Li SH, Engle KM, Yu J-Q, Nature 2015, 519, 334; [PubMed: 25754328] (r)Dong Z, Wang J, Dong G, J. Am. Chem. Soc 2015, 137, 5887; [PubMed: 25909445] (s)Shen P-X, Wang X-C, Wang P, Zhu RY, Yu J-Q, J. Am. Chem. Soc 2015, 137, 11574; [PubMed: 26313012] (t)Shi H, Herron AN, Shao Y, Shao Q, Yu J-Q, Nature 2018, 558, 581; [PubMed: 29915312] (v)Zhang H, Wang H-Y, Luo Y, Chen C, Cao Y, Chen P, Guo Y-L, Lan Y, Liu G, ACS Catal 2018, 8, 2173.

- [9]. For the seminal use of bifunctional reagent in Pd/NBE catalysis, see ref 8a;for the fluorenone synthesis via Pd/NBE catalysis, see Zhao Y-B, Mariampillai B, Candito DA, Laleu B, Li M, Lautens M, Angew. Chem. Int. Ed 2009, 48, 1849; Angew. Chem. 2009, 121, 1881.
- [10]. (a)Sui X, Zhu R, Li G, Ma X, Gu Z, J. Am. Chem. Soc 2013, 135, 9318; [PubMed: 23758183] (b)Weinstabl H, Suhartono M, Qureshi Z, Lautens M, Angew. Chem. Int. Ed 2013, 52, 5305; Angew. Chem. 2013, 125, 5413(c)Liu Z-S, Qian G, Gao Q, Wang P, Cheng H-G, Wei Q, Liu Q, Zhou Q, ACS Catal 2018, 8, 4783.
- [11]. Catellani M, Chiusoli GP, J. Organomet. Chem 1988, 346, C27.
- [12]. The anhydride derived from unsubstituted acrylic acid did not yield the desired product, due to its instability under the reaction conditions.
- [13]. (a)Dong Z, Lu G, Wang JC, Liu P, Dong G, J. Am. Chem. Soc 2018, 140, 8551; [PubMed: 29906109] (b)Della Ca N, Casnati A, Fontana M, Coruzzi G, Aresta BM, Corriero N, Maggi R, Maestri G, Motti E, ChemCatChem 2018, 10, 4346;(c)Elsayed MSA, Griggs B, Cushman M, Org. Lett 2018, 20, 5228. [PubMed: 30102543]
- [14]. (a)Catellani M, Fagnola MC, Angew. Chem. Int. Ed. Engl 1994, 33, 2421; Angew. Chem. 1994, 106, 2559.(b)Wang J, Li R, Dong Z, Liu P, Dong G, Nat. Chem 2018, 10, 866. [PubMed: 29941906]
- [15]. Ito T, Tanaka T, Iinuma M, Nakaya K, Takahashi Y, Sawa R, Murata J, Darnaedi D, J. Nat. Prod 2004, 67, 932. [PubMed: 15217269]
- [16]. (a)Snyder SA, Zografos AL, Lin Y, Angew. Chem. Int. Ed 2007, 46, 8186; Angew. Chem. 2007, 119, 8334;(b)Lee BH, Choi YL, Shin S, Heo JN, J. Org. Chem 2011, 76, 6611; [PubMed: 21740065] (c)Yang Y, Philips D, Pan SF, J. Org. Chem 2011, 76, 1902; [PubMed: 21291249] (d)Kerr DJ, Miletic M, Manchala N, White JM, Flynn BL, Org. Lett 2013, 15, 4118. [PubMed: 23899350]
- [17]. Xiao C-F, Zou Y, Du J-L, Sun H-Y, Liu X-K, Synth. Commun 2012, 42, 1243.
- [18]. CCDC 1881705 (11) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre.
- [19]. Kim H, Yang I, Ryu SY, Won DH, Giri AG, Wang W, Choi H, Chin J, Hahn D, Kim E, Han C, Lee J, Nam SJ, Ho WK, Kang H, J. Nat. Prod 2015, 78, 363. [PubMed: 25689430]
- [20]. Wulff H, Castle NA, Pardo LA, Nat. Rev. Drug Discov 2009, 8, 982. [PubMed: 19949402]
- [21]. Shi H, Babinski DJ, Ritter T, J. Am. Chem. Soc 2015, 137, 3775. [PubMed: 25763682]
- [22]. Colacot TJ, Shea HA, Org. Lett 2004, 6, 3731. [PubMed: 15469335]
- [23]. Molander GA, Bernardi CR, J. Org. Chem 2002, 67, 8424. [PubMed: 12444620]


a) Indenone or indanone-containing natural products



b) Transition metal-catalyzed annulation approaches for indenone synthesis




c) Pd/NBE-catalyzed approach: this work




Scheme 1.


Indenone/indanone containing natural products and indenone synthesis




Scheme 2.
Further reaction exploration




Scheme 3.
Mechanistic exploration



Scheme 4.
Formal synthesis of pauciflorol F.




Scheme 5.
Retrosynthetic analysis of acedinone A.




Scheme 6.
Total synthesis of acredione A.

Table 1.

Substrate scope of Pd/NBE-catalyzed indenone synthesis.^[a]

[a] Unless otherwise noted, the reaction was run with **3** (0.3 mmol), **4** (0.6 mmol), [Pd(allyl)Cl]₂ (0.015 mmol), P(2-furyl)₃ (0.06 mmol), **N1** (0.15 mmol), Cs₂CO₃ (1.2 mmol) and NMe₄Cl (1.2 mmol) in dioxane (2.25 mL) and THF (0.75 mL) at 85 °C for 14 h. [b] **7** (0.6 mmol) was used instead of **4** and the reaction was run at 95 °C without NMe₄Cl and THF. [c] run for 3 h. [d] run for 4 h. [e] **4ac'** (0.6 mmol) was used instead of **4** and the reaction was run for 2 h.

Table 2.

Optimization for *ortho* acylation/*ipso* borylation.^[a]

entry	[Pd]	solvent	yield of 13a	yield of 13a'
1	Pd(OAc) ₂	dioxane	21	20
2	[Pd(allyl)Cl] ₂	dioxane	54	11
3	Pd ₂ (dba) ₃	dioxane	38	8
4	Pd(dba) ₂	dioxane	58	5
5	Pd(dba) ₂	toluene	56	4
6	Pd(dba) ₂	dioxane:toluene (4:1)	64	6
7 ^b	Pd(dba) ₂	dioxane:toluene (4:1)	67 ^c	5

[a] Conditions: **16** (0.1 mmol), **18** (0.12 mmol), B₂pin₂ (0.105 mmol), [Pd] (0.01 mmol), P(2-furyl)₃ (0.02 mmol), **N1** (0.05 mmol), Cs₂CO₃ (0.4 mmol), solvent (2 mL) at 100 °C for 24 h. Yields were determined by ¹H NMR using 1,3,5-trimethoxybenzene as the internal standard. [b] run at 120°C. [c] isolated yield.