

Palmitoylation-dependent Estrogen Receptor α Membrane Localization: Regulation by 17 β -Estradiol

Filippo Acconcia,* Paolo Ascenzi,*^{†‡} Alessio Bocedi,^{‡§} Enzo Spisni,^{||}
Vittorio Tomasi,^{||} Anna Trentalance,* Paolo Visca,*[‡] and Maria Marino*[¶]

*Department of Biology, University Roma Tre, I-00146 Rome, Italy; [†]Interdepartmental Laboratory for Electron Microscopy, University Roma Tre, I-00146 Rome, Italy; [‡]National Institute for Infectious Diseases Istituto di Ricovero e Cura a Carattere Scientifico Lazzaro Spallanzani, I-00149 Rome, Italy; [§]Department of Chemistry, Chemical Engineering, and Materials, University of L'Aquila, I-67100 L'Aquila, Italy; and ^{||}Department of Experimental Biology, University of Bologna, I-40126 Bologna, Italy

Submitted July 2, 2004; Revised September 28, 2004; Accepted October 4, 2004
Monitoring Editor: Keith Yamamoto

A fraction of the nuclear estrogen receptor α (ER α) is localized to the plasma membrane region of 17 β -estradiol (E2) target cells. We previously reported that ER α is a palmitoylated protein. To gain insight into the molecular mechanism of ER α residence at the plasma membrane, we tested both the role of palmitoylation and the impact of E2 stimulation on ER α membrane localization. The cancer cell lines expressing transfected or endogenous human ER α (HeLa and HepG2, respectively) or the ER α nonpalmitoylable Cys447Ala mutant transfected in HeLa cells were used as experimental models. We found that palmitoylation of ER α enacts ER α association with the plasma membrane, interaction with the membrane protein caveolin-1, and nongenomic activities, including activation of signaling pathways and cell proliferation (i.e., ERK and AKT activation, cyclin D₁ promoter activity, DNA synthesis). Moreover, E2 reduces both ER α palmitoylation and its interaction with caveolin-1, in a time- and dose-dependent manner. These data point to the physiological role of ER α palmitoylation in the receptor localization to the cell membrane and in the regulation of the E2-induced cell proliferation.

INTRODUCTION

The sex steroid 17 β -estradiol (E2) acts by binding to its nuclear receptors (i.e., ER α and ER β) that then transactivate target genes. In addition, E2 induces rapid, nongenomic actions involving plasma membrane-associated signaling that require a membrane ER (Coleman and Smith, 2001; Kelly and Levin, 2001; Jakacka *et al.*, 2002; Marino *et al.*, 2002). Although different structural and functional properties have been reported for the membrane-associated ER by comparison with nuclear ER α and ER β (Roper *et al.*, 2002; Toran-Allerand *et al.*, 2002; Deecher *et al.*, 2003), immunocytochemical studies revealed the presence of a significant fraction of nuclear ER also on the plasma membrane (Pappas *et al.*, 1995; Norfleet *et al.*, 1999; Dan *et al.*, 2003; Razandi *et al.*, 2003; Arvanitis *et al.*, 2004; Song *et al.*, 2004). In addition, a single mRNA originates a similarly sized nuclear and membrane ER in ER α -transfected Chinese hamster ovary and HeLa cells (Razandi *et al.*, 1999; Marino *et al.*, 2002, 2003). Thus, ER α localizes to both the nucleus and the plasma membrane. Moreover, the membrane ER α is emerging as the primary endogenous mediator of E2 rapid responses important in cell proliferation (Marino *et al.*, 1998, 2002; Castoria *et*

al., 1999, 2001; Razandi *et al.*, 1999, 2000; Lobenhofer *et al.*, 2000; Acconcia *et al.*, 2004a; Fernando and Wimalasena, 2004).

Debate is open regarding the structural bases and the mechanisms for ER α maintenance at and translocation to the plasma membrane. ER α does not display any intrinsic transmembrane domain (Song *et al.*, 2004); thus, ER α interaction with specific membrane proteins have been proposed to explain its membrane localization (Chambliss and Shaul, 2002; Chambliss *et al.*, 2002; Migliaccio *et al.*, 2002; Razandi *et al.*, 2002, 2003; Toran-Allerand *et al.*, 2002; Arvanitis *et al.*, 2004). In particular, the Ser522 residue has been reported as necessary for ER α interaction with caveolin-1. However, the expression of ER α -Ser522Ala mutant in Chinese hamster ovary cells decreased only 60% of ER α on the membrane, extracellular signal-regulated kinase (ERK) activation, and cyclin D₁ gene transcription (Razandi *et al.*, 2003). Thus, the direct ER α :caveolin-1 interaction could not be the sole molecular mechanism for ER α membrane association. Furthermore, because the protein:protein interaction occurs only after some minutes of E2 stimulation (5–15 min) (Song *et al.*, 2004), this mechanism does not justify the receptor presence at the plasma membrane of unstimulated target cells reported previously (Pappas *et al.*, 1995; Norfleet *et al.*, 1999; Razandi *et al.*, 1999; Levin, 2001; Dan *et al.*, 2003). Very recently, we reported that human wild-type ER α underwent palmitoyl acyl transferase (PAT)-dependent S-palmitoylation. Moreover, a point mutation of the ER α -Cys447 residue to Ala completely impaired receptor palmitoylation and the E2-induced rapid activation of the ERK/mitogen-activated protein kinase (MAPK) signaling pathway (Acconcia *et al.*, 2004b), although this mutant binds E2 with the same affinity than the wild-type ER α (Katzenellenbogen *et al.*,

Article published online ahead of print. Mol. Biol. Cell 10.1091/mbc.E04-07-0547. Article and publication date are available at www.molbiolcell.org/cgi/doi/10.1091/mbc.E04-07-0547.

[¶] Corresponding author. E-mail address: m.marino@uniroma3.it.

Abbreviations used: 2-Br, 2-bromohexadecanoic acid, 2-bromopalmitate; E2, 17 β -estradiol; ER, estrogen receptor; ERK, extracellular signal-regulated kinase; IGF-1, insulin-like growth factor-1; PAT, palmitoyl acyl transferase; PI3K, phosphoinositide-3-kinase.

1993; Neff *et al.*, 1994). This observation raises the intriguing hypothesis that the palmitoylation of ER α could be the major determinant for ER α residence at the plasma membrane.

Here, we examine the role of palmitoylation in localizing ER α to the plasma membrane and in promoting the ER α :caveolin-1 interaction, as well as the role of E2 in regulating ER α palmitoylation. Our results indicate that ER α palmitoylation is required for ER α :protein interaction (i.e., caveolin-1) and for the receptor localization to and maintenance at the plasma membrane. As a consequence, ER α palmitoylation enables rapid E2 signaling important for cell proliferation.

MATERIALS AND METHODS

Reagents

17 β -Estradiol, gentamicin, penicillin and other antibiotics, GenElute plasmid maxiprep kit, DMEM, RPMI 1640 medium (without phenol red), charcoal-stripped fetal calf serum, and the PAT inhibitor 2-bromohexadecanoic acid (2-bromo-palmitate; 2-Br) (IC₅₀ of ~4 μ M; Varner *et al.*, 2003) were purchased from Sigma-Aldrich (St. Louis, MO). The estrogen receptor inhibitor ICI 182,780 was obtained from Tocris Cookson (Ballwin, MO). 9,10-[³H]Palmitic acid (specific activity 57 Ci/mmol) was purchased from PerkinElmer Life and Analytical Sciences (Boston, MA). Methyl-1-[³H]thymidine (specific activity 81 Ci/mmol) was purchased from Amersham Biosciences UK (Little Chalfont, Buckinghamshire, United Kingdom). Lipofectamine reagent was obtained from Invitrogen (Carlsbad, CA). The luciferase kit was obtained from Promega (Madison, WI). Bradford protein assay was obtained from Bio-Rad (Hercules, CA). The monoclonal, anti-phospho-ERK, anti-ER α D12 (N terminus), anti-AKT, and anti- β -actin as well as the polyclonal anti-ERK, anti-caveolin-1, and anti-ER α MC20 (C terminus) antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). The monoclonal anti-ER α (full-length) antibody was purchased from Novocastra (Newcastle, United Kingdom). The polyclonal anti-phospho-AKT antibody was purchased from New England Biolabs (Beverly, MA). CDP-Star, chemiluminescence reagent for Western blot was obtained from PerkinElmer Life and Analytical Sciences (Boston, MA). All the other products were from Sigma-Aldrich. Analytical or reagent grade products, without further purification, were used.

Cell Culture and DNA Synthesis

HepG2 and HeLa cells were routinely grown in 5% CO₂ in modified, phenol red-free RPMI 1640 or DMEM media containing 10% (vol/vol) charcoal-stripped fetal calf serum, L-glutamine (2 mM), gentamicin (10 μ g/ml), and penicillin (100 U/ml). Cells were passaged every 2 d. DNA synthesis was assayed by incubating subconfluent cells (70–80%) with methyl-1-[³H]thymidine (final concentration, 1 μ Ci/ml). Cells were simultaneously treated with either E2 [final concentration, 10 nM in ethanol/phosphate-buffered saline (PBS) 1:10 (vol/vol)] or vehicle [ethanol/PBS 1:10 (vol/vol)]. Methyl-1-[³H]thymidine incorporation was assayed 1 h after E2 administration as reported previously (Marino *et al.*, 2001a). When indicated, the PAT inhibitor 2-Br was added 30 min before E2 administration (final concentration 10 μ M).

Plasmids

The reporter plasmid pXP2-D₁-2966-luciferase (pD₁), the empty vector (pCMV5), the expression vector pCR3.1- β -galactosidase, wild-type human ER α pSG5-HE0, pSG5-Cys447Ala (human ER α point mutant Cys447 residue to Ala), and pKCR2-HE14 (N-terminal deletion mutant of HE0 lacking A/B and DNA binding domains Δ 1–281) have been described previously (Herbert *et al.*, 1994; Marino *et al.*, 2001b, 2002; Acconcia *et al.*, 2004b). A luciferase dose-response curve showed that the maximum effect was obtained when 1.0 μ g of plasmids was transfected together with 1.0 μ g of pCR3.1- β -galactosidase to normalize for transfection efficiency (~55–65%). Plasmids were purified for transfection using the GenElute plasmid maxiprep kit according to the manufacturer's instructions.

Transfection and Luciferase Assay

HeLa cells were grown to ~70% confluence and then transfected using Lipofectamine reagent according to the manufacturer's instructions. Six hours after transfection, the medium was changed, and 24 h after the cells were stimulated with 10 nM E2 for 6 h. The cell lysis procedure as well as the subsequent measurement of luciferase gene expression was performed using the luciferase kit according to the manufacturer's instructions with a PerkinElmer Life and Analytical Sciences (Bad Wildbad, Germany) luminometer. When indicated, the PAT inhibitor 2-Br (final concentration 10 μ M) was added 30 min before E2 administration.

Cell Labeling with [³H]Palmitate and Immunoprecipitation

Forty-two hours after transfection with either plasmid containing empty, wild-type ER α or the ER α point mutant, HeLa cells and untransfected HepG2 cells were incubated with 0.5 mCi/ml [³H]palmitate at 37°C for different times ranging between 0 and 240 min. Where indicated, HepG2 and HeLa cells were stimulated with different concentrations of E2 (1, 10, and 100 nM) at different times (10, 60, and 240 min) in the presence of [³H]palmitate. Cells were then washed in ice-cold PBS, harvested by scraping, and lysed in 50 μ l of lysis buffer [10 mM Tris, pH 7.5, 1 mM EDTA, 0.5 mM EGTA, 10 mM NaCl, 1% (vol/vol) Triton X-100, and 1% (wt/vol) sodium cholate] containing protease inhibitors (0.5 mM phenylmethylsulfonyl fluoride, 1 μ g/ml leupeptin, and 5 μ g/ml aprotinin). The cell lysates were then clarified by centrifugation and immunoprecipitated as described previously (Han *et al.*, 2002). Briefly, equal amounts of soluble cell extracts were incubated with either 2 μ g of caveolin-1 antibody or different anti-ER α antibodies. The optimal signal was obtained using 1 μ g of anti-ER α D12 (N terminus) together with 1 μ g of either anti-ER α AER311 (region 495–595), anti-ER α MC20 (C terminus), or anti-ER α (full-length ER α). The lysates and antibodies were incubated for 90 min at 4°C, and then 20 μ l of protein A-agarose was added for 30 min at 4°C. After centrifugation (50,000 \times g for 15 min), the supernatant, and the immunoprecipitated proteins were separated in 7–10% SDS-PAGE. Proteins were electrophoretically transferred to nitrocellulose and then probed overnight at 4°C with anti-caveolin-1 or anti-ER α MC20 antibodies. The nitrocellulose was stripped and then probed with anti-caveolin-1 or anti-ER α MC20 antibodies. In some experiments, the radioactivity present in immunoprecipitated proteins and in the supernatant was monitored by counting with a Canberra Packard (Milan, Italy) liquid beta-counter.

Confocal Microscopy Studies

Wild-type ER α or ER α -Cys447Ala mutant was transiently expressed in HeLa cells grown on coverslips. To visualize only the cytosolic and the plasma membrane distribution of wild-type ER α and ER α -Cys447Ala mutant, HeLa cells were freeze-thawed before fixation in 4% (vol/vol) *p*-formaldehyde (Mardones and Gonzalez, 2003) and then incubated with MC-20 rabbit polyclonal antibody diluted 1:200 in PBS. After washing, cells were incubated with anti-rabbit fluorescein isothiocyanate-conjugated antibody (Sigma-Aldrich) diluted 1:50 in PBS containing 10 mg/ml bovine serum albumin (BSA). Finally, the slides were mounted in PBS medium containing 80% glycerol and 50 mg/ml 1,4-diazabicyclo[2.2.2]octane (Sigma-Aldrich). Evaluation of antibody specificity was carried out either by omitting the primary antibody or by using nonspecific sera. The confocal imaging was performed on a Radiance 2000 confocal laser scanning microscope (Bio-Rad), equipped with a Nikon 40 \times , 1.4 numerical aperture objective and with Krypton ion lasers.

Electrophoresis and Immunoblotting

After treatment, cells were lysed and solubilized in 0.125 M Tris, pH 6.8, containing 10% (wt/vol) SDS, 1 mM phenylmethylsulfonyl fluoride, and 5 μ g/ml leupeptin, and then the cell lysates were boiled for 2 min. Total proteins were quantified using the Bradford protein assay (Bradford, 1976). Solubilized proteins (20 μ g) were resolved by 10% SDS-PAGE at 100 V for 1 h at 24°C and then electrophoretically transferred to nitrocellulose for 45 min at 100 V and 4°C. The nitrocellulose was treated with 3% (wt/vol) BSA in 138 mM NaCl, 25 mM Tris, pH 8.0, at 24°C for 1 h and then probed overnight at 4°C with either anti-phospho-ERK or anti-phospho-AKT antibodies. The nitrocellulose was stripped by restore western blot stripping buffer (Pierce Chemical, Rockford, IL) for 10 min at room temperature and then probed with either anti-ERK or anti-AKT and anti- β -actin antibodies. Antibody reaction was visualized with chemiluminescence Western blotting detection reagent (Amersham Biosciences UK). Where indicated, the PAT inhibitor 2-Br (10 μ M) was added 30 min before E2 administration.

RESULTS

Palmitoylation Is Required for ER α Localization to the Plasma Membrane and Its Interaction with Caveolin-1

We previously demonstrated that ER α was a palmitoylated protein and that the ER α -Cys447 to Ala point mutation impaired palmitoylation of the receptor. This lipid modification was necessary for the induction of nongenomic ERK/MAPK signal transduction pathway (Acconcia *et al.*, 2004b). Based on these findings, we postulated that palmitoylation of ER α be important for the localization of this receptor to the membrane.

We first verified the occurrence of ER α palmitoylation in human cancer cells containing transfected (HeLa) or endogenous (HepG2) receptor. HeLa cells were transiently trans-




Figure 1. ER α palmitoylation. (a) [³H]Palmitate incorporation into the empty vector (control) or wild-type ER α or ER α -Cys447Ala mutant-transfected HeLa cells and into immunoprecipitated caveolin-1 (Cav-1) of untransfected HeLa cells has been evaluated after 4 h of palmitate labeling (top). Western blot analysis of the corresponding 67-kDa band of immunoprecipitated ER α or the 22-kDa band of immunoprecipitated caveolin-1 (bottom). (b) Time course of [³H]palmitate incorporation into immunoprecipitated ER α in ER α -transfected HeLa and in HepG2 cells. Data are the means of four independent experiments \pm SD.

fectured with either an empty vector, a wild-type ER α -encoding vector or the ER α missense mutant in which Cys447 was changed to Ala by site-directed mutagenesis (Acconcia *et al.*, 2004b), and then incubated with [³H]palmitate for 4 h at 37°C. After ER α immunoprecipitation, the radioactivity present in both the immunoprecipitate and the supernatant was determined. As expected, the cell transfected with the empty vector (control) did not immunoprecipitate ER α or incorporate [³H]palmitate (Figure 1a). ER α was successfully immunoprecipitated in wild-type ER α -transfected HeLa cells, as revealed by [³H]palmitate incorporation in the 67-kDa band corresponding to ER α (Figure 1a). The ER α -Cys447Ala mutant containing cells immunoprecipitate ER α but failed to incorporate [³H]palmitate (Figure 1a). ER α was not detected in the supernatant fractions (our unpublished data). In addition, the radioactivity present in the palmitoylated caveolin-1 (22-kDa band; Resh, 1999) immunoprecipitated from untransfected HeLa cells was detected as positive control (cav-1) (Figure 1a). ER α palmitoylation was essentially complete within 10 min and remained constant >240 min (Figure 1b). No difference in [³H]palmitate incorporation occurred between ER α -transfected HeLa cells and HepG2 cells (Figure 1b).

To visualize the cytosolic and/or plasma membrane localization of wild-type and ER α -Cys447Ala mutant, trans-




Figure 2. Subcellular localization of ER α and ER α -Cys447Ala mutant and ER α :caveolin-1 interaction. Confocal microscopy analysis of empty vector (a; control), wild-type ER α (b), and ER α -Cys447Ala mutant (c)-transfected HeLa cells by using anti-ER α MC20 antibody on freeze-thawed HeLa cells. White arrows indicate the membrane localization of wild-type ER α . In a, fluorescence signal is superimposed to laser pseudophase contrast images of the cells. Bar, 10 μ m. Empty vector (control) or wild-type ER α or ER α -Cys447Ala mutant-transfected HeLa cells were subjected to ER α immunoprecipitation with anti-ER α MC20 antibody followed by Western blot with anti-caveolin-1 antibody (d) or by Western blot with anti-ER α MC20 antibody (e).

fectured HeLa cells were studied by immunofluorescence staining (Figure 2, a–c). By using the anti-ER α MC20 antibody, a major population of membrane-associated ER α receptor was clearly present in HeLa cells expressing wild-type ER α , whereas a weak cytosolic staining and no membrane-bound receptor could be detected in HeLa cells expressing the ER α -Cys447Ala mutant (Figure 2, compare b with c). Moreover, no ER α was detected in HeLa cells that had been transfected with the empty vector (Figure 2a, control). We conclude that Cys447 is important for surface membrane localization.

The ability of wild-type ER α and of ER α -Cys447Ala mutant to bind the scaffolding plasma membrane protein caveolin-1 was examined by coimmunoprecipitation of caveolin-1 and ER α . This was accomplished in HeLa cells transiently transfected with empty, ER α , or ER α -Cys447Ala expression vectors. No association between caveolin-1 and ER α occurred in cells transfected with the empty vector (control), confirming the lack of any ER isoforms in HeLa cells (Figure 2, d and e, lane control). In ER α -expressing HeLa cells, a 22-kDa band, corresponding to caveolin-1, and a 67-kDa band, corresponding to ER α , were present (Figure 2, d and e, lane ER α). On the other hand, ER α -Cys447Ala mutant-expressing cells failed to interact with caveolin-1 because ER α could be immunoprecipitated, but coimmunoprecipitation of caveolin-1 could not be observed (Figure 2, d and e, lane Cys447Ala). These data are consistent with the role of palmitoylation in ER α localization to the plasma membrane and in the receptor association with caveolin-1.

ER α Palmitoylation Is Negatively Modulated by E2

To assess the ability of E2 to modulate ER α palmitoylation, HeLa cells transfected with the ER α expression vector and




Figure 3. E2 effect on ER α palmitoylation and ER α association with caveolin-1. Time course (a) and the corresponding Western blot analysis (b) of E2-stimulated ER α -transfected HeLa (open bars; 1) and HepG2 (filled bars; 2) cells. [3 H]palmitate incorporation in immunoprecipitated ER α was performed at 4 h. Data are the means of four independent experiments \pm SD. * $p < 0.001$, compared with unstimulated samples (0) were determined by using Student's *t* test. Wild-type ER α -transfected HeLa cells were subjected to caveolin-1 immunoprecipitation (c) or ER α immunoprecipitation (d) with anti-caveolin-1 or anti-ER α MC20 antibodies followed by Western blot with anti-caveolin-1 or with anti-ER α MC20 antibodies.

HepG2 cells were incubated with [3 H]palmitate for 4 h in the presence of different E2 concentrations. Physiological E2 concentration (10 nM) decreased by half the amount of [3 H]palmitate incorporated in wild-type ER α ; similar results were obtained at higher E2 concentrations (i.e., 100 nM), whereas a lower hormone concentration (i.e., 1 nM) was ineffective (our unpublished data). The time course of 10 nM E2 stimulation in cells containing transfected (HeLa) and endogenous (HepG2) ER α showed that 60 min of E2 stimulation induced the decrease of [3 H]palmitate incorporation (Figure 3a) without any change in the protein level (Figure 3b). The E2-induced reduction in [3 H]palmitate incorporation in both cell lines was completely prevented by pretreating the cells with the ER α inhibitor ICI 182,780 (1 μ M; our unpublished data). In addition to 48% reduction of [3 H]palmitate incorporation in ER α at 60 min, we found that E2 induced a >90% reduction in the ability of ER α to form a complex with caveolin-1 (Figure 3, c and d). It is important to note that although E2 reduces palmitoylation of ER α and complex formation of ER α with caveolin-1, it had no effect on the cellular protein level of either proteins (Figure 3, c and d). We further asked whether the A/B and C domains of ER α contributed in E2-induced down-regulation of both ER α palmitoylation and its association to caveolin-1. Surprisingly, from 60 min until 240 min, E2 induced a 40% increase in both receptor palmitoylation (Figure 4a) and receptor association with caveolin-1 (Figure 4, b and c) in HeLa cells transfected with ER α deletion mutant lacking the A/B and C domains (HE14).

ER α Palmitoylation Is Necessary for Nongenomic Activities

In HepG2 cells, the rapid E2-induced activation of both the ERK/MAPK and phosphoinositide-3-kinase (PI3K)/AKT





Figure 4. E2 effect on A/B and C domains deleted ER α mutant (HE14) palmitoylation and association with caveolin-1. (a) Time course of E2-stimulated ER α -HE14 mutant-transfected HeLa cells. [3 H]Palmitate incorporation in immunoprecipitated ER α -HE14 mutant was performed at 4 h in the absence (open bars) or presence (filled bars) of 10 nM E2. Data are the means of four independent experiments \pm SD. * $p < 0.001$, compared with unstimulated samples (open bars), were determined by using Student's *t* test. ER α -HE14 mutant-transfected HeLa cells were subjected to caveolin-1 immunoprecipitation (b) or ER α immunoprecipitation (c) with anti-caveolin-1 or anti-ER α MC20 antibodies followed by Western blot with anti-caveolin-1 or with anti-ER α MC20 antibodies.

pathways is sufficient and necessary for E2-induced cell cycle progression (i.e., DNA synthesis and transcription of the cyclin D₁ gene) (Marino *et al.*, 2002, 2003). We determined next whether ER α palmitoylation could impact on these rapid nongenomic ER α activities. E2 induced ERK (Figure 5, a and a') and AKT (Figure 5, b and b') phosphorylation in ER α -transfected HeLa and HepG2 cells. If the cells were pretreated with the PAT inhibitor 2-Br, the E2-induced activation of signaling kinases was completely blocked without affecting the basal level of phosphorylation. In agreement with our data mentioned above, E2 was ineffective in inducing ERK and AKT phosphorylation in ER α -Cys447Ala mutant-transfected HeLa cells. After reprobing the membranes using total ERK and AKT antibodies to recognize the nonphosphorylated forms of these proteins, we found that E2-induced phosphorylation of signaling proteins occurred in the absence of changes in their expression levels.

We have recently demonstrated that the ER-dependent cyclin D₁ transcription and DNA synthesis are some of the downstream targets of E2-induced ERK and AKT activation (Marino *et al.*, 2002, 2003). Accordingly, in ER α transfected cells, E2 induced cyclin D₁ promoter activity and [3 H]thymidine incorporation into DNA (Figure 6). In contrast, neither the cyclin D₁ promoter activity (Figure 6a) nor [3 H]thymidine incorporation into DNA (Figure 6b) was affected by E2 administration to the ER α -Cys447Ala mutant-transfected cells. In addition, the pretreatment of HeLa cells with the PAT inhibitor 2-Br prevented nongenomic E2-evoked effects (Figure 6). Notably, no changes in cyclin D₁ promoter basal transcription activity and DNA synthesis were detected after the treatment with the PAT inhibitor 2-Br alone. Thus, palmitoylated ER α mediates the E2-induced activation of ERK, AKT, cyclin D₁ promoter activity, and DNA synthesis. These findings demonstrate a critical role of palmitoylation in ER α -mediated cell proliferation.

Figure 5. Effect of ER α palmitoylation on E2-induced ERK and AKT phosphorylation. Western blot analysis of ERK phosphorylation in ER α (lanes 1–4) or ER α -Cys447Ala mutant (lanes 5 and 6)-transfected HeLa and HepG2 (lanes 7–9) cells were performed on unstimulated (–) and stimulated (+) cells for 10 min with E2 (10 nM) in the presence or absence of 10 μ M PAT inhibitor 2-Br (30-min pretreatment). The same filter was reprobed with anti-total ERK and anti-actin antibodies. (a) Typical Western blot. (a') Densitometric analysis of four different experiments. Data are the means \pm SD. $p < 0.001$, compared with unstimulated samples (*) or with E2-stimulated samples (°) were determined by using Student's *t* test. Western blot analysis of AKT phosphorylation in ER α (lanes 1–4) or ER α -Cys447Ala mutant (lanes 5 and 6)-transfected HeLa and HepG2 (lanes 7–9) cells were performed on unstimulated (–) and stimulated (+) cells for 30 min with E2 (10 nM) in the presence or absence of 10 μ M PAT inhibitor 2-Br (30-min pretreatment). The same filter was reprobed with anti-total AKT and anti-actin antibodies. (b) Typical Western blot. (b') Densitometric analysis of four different experiments. Data are the means \pm SD. $p < 0.001$, compared with unstimulated samples (*) or with E2-stimulated samples (°) were determined by using Student's *t* test.



DISCUSSION

The mechanism(s) underlying the mitogenic role played by E2 in different target tissues is now better understood based on the studies reported by different laboratories, including

our own. These studies designate a strict relationship between E2-induced nongenomic functions and cell proliferation (Marino *et al.*, 1998, 2001a,b, 2002, 2003; Castoria *et al.*, 1999, 2001; Razandi *et al.*, 1999, 2000; Lobenhofer *et al.*, 2000; Acconcia *et al.*, 2004a; Fernando and Wimalasena, 2004). The activation of ERK/MAPK, PI3K/AKT, and protein kinase C, rapidly generated after E2 binding to ER α , are all defined as necessary and sufficient for E2-induced cell cycle-regulating genes (e.g., cyclin D₁) and the G1-to-S phase progression in different cell lines. These actions are thought to require a plasma membrane ER α .

The idea that surface membrane ER is identical to the nuclear ER α seems now very plausible as deduced by using antibodies raised against multiple epitopes of the nuclear receptor (Pappas *et al.*, 1995; Norfleet *et al.*, 1999; Levin, 2001; Dan *et al.*, 2003). Moreover, the ER α presence on the human cell plasma membrane and the ER α role in the cellular effects of E2 are increasingly accepted (Razandi *et al.*, 1999; Simoncini *et al.*, 2000; Kousteni *et al.*, 2001; Wise *et al.*, 2001).

Despite clear progress, many questions are still unanswered such as the structural bases and the mechanism(s) for ER α localization to and maintenance at the plasma membrane. This issue has been addressed in a few papers that indicate that the ER α membrane localization could result from receptor interaction with specific membrane proteins and/or lipid modifications (Chambliss and Shaul, 2002; Chambliss *et al.*, 2002; Razandi *et al.*, 2002; Li *et al.*, 2003; Acconcia *et al.*, 2004b; Arvanitis *et al.*, 2004). In particular, in the region of plasma membrane, ER α can form a third-party protein (i.e., Shc/ER α /insulin-like growth factor-1 [IGF-1] receptor) and/or associates in a ternary complex to the membrane tyrosine kinase Src and p85 (the catalytic subunit of PI3K), causing Src/Ras/ERK and PI3K/AKT pathway activation and DNA synthesis (Castoria *et al.*, 2001; Migliaccio *et al.*, 2002; Song *et al.*, 2004). This association occurs only after some minutes of E2 stimulation (5–15 min), suggesting a recruitment of ER α from the cytosol, but does not clarify the receptor presence at the plasma membrane reported in several papers (Pappas *et al.*, 1995; Norfleet *et al.*, 1999; Razandi *et al.*, 1999; Levin, 2001; Dan *et al.*, 2003). Recently, the Ser522 residue within the E domain of ER α has been




Figure 6. Effect of ER α palmitoylation on E2-induced cyclin D₁ promoter activity and DNA synthesis. (a) Luciferase assay detection on HeLa cells cotransfected with pXP2-D₁-2966-luciferase and wild-type ER α or ER α -Cys447Ala mutant expression vectors. After transfection, cells were treated with vehicle (PBS) or E2 (10 nM) in the presence or absence of 10 μ M PAT inhibitor 2-Br (30-min pretreatment). (b) [³H]Thymidine incorporation into DNA in ER α or ER α -Cys447Ala mutant-transfected HeLa cells treated with vehicle (PBS) or E2 (10 nM) in the presence or in absence of 10 μ M PAT inhibitor 2-Br (30-min pretreatment). Data are the means \pm SD of four independent experiments. * $p < 0.001$, compared with respective control values (PBS); ° $p < 0.001$, compared with ER α -transfected cells stimulated with E2 values; and # $p < 0.001$, compared with respective E2 values, were determined by using Student's *t* test.

reported to be critical in linking the receptor to the cell membrane through an interaction with caveolin-1 (Razandi *et al.*, 2003). However, the expression of ER α -Ser522Ala mutant in Chinese hamster ovary cells decreased only 60% of ER α on the membrane, ERK activation, and cyclin D₁ gene transcription, suggesting that Ser522 is not the sole residue for ER α membrane association. We recently identified the ER α posttranslational modification with palmitate (Acconcia *et al.*, 2004b), and here, for the first time, we demonstrate that palmitoylation is a determinant for ER α residence at the plasma membrane caveolae. In fact, we prove that ER α localization to the membrane is dependent on ER α palmitoylation because the unpalmitoylable ER α -Cys477Ala mutant, which maintains an intact Ser522 residue, is unable to localize to the membrane, to interact with caveolin-1, and to generate the E2-induced rapid membrane-starting signal pathways important in cell proliferation. Thus, present data describe a new model in which Cys447 residue in the E domain of the ER α assures the receptor localization to and maintenance at the plasma membrane caveolae. Moreover, the A/B and C domains of ER α do not contribute to membrane ER localization; in fact, HE14 ER α deletion mutant, which contains only E and F domains, still undergoes to palmitoylation and associates to caveolin-1.

One of the main findings in this study is the time- and concentration-dependent negative regulation of wild-type ER α palmitoylation exerted by E2. Because palmitoylation is a rapid reversible chemical modification (Bijlmakers and Marsh, 2003), it is most likely that the membrane-bound full-length receptor undergoes to the E2-induced depalmitoylation. Moreover, the E2 binding to the soluble, cytosolic, receptor, enacting ER α structural modifications (Brzozowski *et al.*, 1997), could impair the PAT action (i.e., PAT:ER α recognition). In fact, it has been reported that the Cys447 residue is not yet able to react with the iodoacetic acid a cysteine-reacting reagent after E2 binding (Hegy *et al.*, 1996), suggesting that the Cys447 residue could be buried into the protein matrix. In addition, E2 also decreases the ER α :caveolin-1 complex further sustaining the pivotal role played by palmitoylation in ER α membrane localization. On the other hand, the deletion of A/B and C domains of the ER α may impair the E2-induced structural modifications of cytosolic ER α , thus enhancing its palmitoylation (40%). In addition these domains may facilitate the E2-dependent depalmitoylation and dissociation from caveolin-1 of the membrane bound ER α . As a consequence, the A/B and C domains render both soluble and membrane-bound receptor competent to E2 regulation.

The ER α palmitoylation is necessary for E2-induced ERK/MAPK and PI3K/AKT pathway activation as well as for cyclin D₁ transcription and DNA synthesis as demonstrated by the effect of the PAT inhibitor 2-Br and the nonresponsive behavior of the ER α -Cys447Ala mutant. The palmitoylation of ER α barely influences ER α genomic activities (i.e., estrogen responsive element-containing gene transcription) (Acconcia *et al.*, 2004b). Similar results were obtained in HepG2 cells, which express endogenous ER α . In fact, ER α palmitoylation was detected and regulated by E2 in the HepG2 cell line. Moreover, inhibition of the PAT activity by 2-Br prevents ERK/MAPK and PI3K/AKT activation. Our findings indicate a critical requirement of palmitoylation for the ER α plasma membrane localization and the initiation of the nongenomic events important to E2-dependent G1-S phase progression (i.e., cyclin D₁ transcription and DNA synthesis).

In conclusion, palmitoylation enables ER α to reside at the plasma membrane and to interact with caveolin-1. On E2

binding, the ER α may undergo depalmitoylation and dissociate from caveolin-1. Thus, ER α could be relocated by docking to other partner proteins (i.e., Shc/IGF-1 receptor; Src/p85) (Castoria *et al.*, 2001; Migliaccio *et al.*, 2002; Song *et al.*, 2002, 2004). As a consequence, the nongenomic signals could be generated (i.e., ERK/MAPK, PI3K/AKT). Thus, a cycle of inactive/active receptor may occur in the proximity of the plasma membrane as reported for the ER α nuclear pool (Reid *et al.*, 2003). In the future, electron microscopic strategies (e.g., double labeling of ER α and caveolin-1, association/release of ER α from membrane in the presence of E2) will substantiate this action mechanism. However, from now on palmitoylation can be regarded as a physiological regulatory device enabling ER α to initiate E2-induced cell proliferation and provides a new potential target in the treatment of E2-related cancers.

ACKNOWLEDGMENTS

We thank Prof. Rakesh Kumar (Department of Molecular and Cellular Oncology, M.D. Anderson Cancer Center, University of Texas, Houston, TX) and Prof. Hynda Kleinman (Craniofacial and Developmental Biology, National Institutes of Health, Bethesda, MD) for the helpful and critical discussions. The generous gift of antiserum anti-ER α AR311 from Dr. Ciro Abbondanza (Dipartimento di Patologia Generale, Seconda Università degli Studi di Napoli, Napoli, Italy) and of human ER α -H14 mutant expression vector from Prof. Pierre Chambon (Institut de Génétique et Biologie Moléculaire et Cellulaire, Strasbourg, France) are gratefully acknowledged. We express gratitude to Drs. Spartaco Santi and Massimo Riccio (Consiglio Nazionale delle Ricerche, Bologna, Italy) for helpful advices in performing confocal microscopy analysis. The editorial assistance of Peter DeMuro also is acknowledged. This work was supported by grants from 2004 Università "Roma Tre" and Fondo per gli Investimenti della Ricerca di Base 2001 to M.M. V.T. was supported by a grant from Istituto Superiore di Sanità (1%), Rome, Italy.

REFERENCES

- Acconcia, F., Ascenzi, P., Fabozzi, G., Visca, P., and Marino, M. (2004b). S-Palmitoylation modulates human estrogen receptor- α functions. *Biochem. Biophys. Res. Commun.* 316, 878–883.
- Acconcia, F., Totta, P., Ogawa, S., Cardillo, I., Inoue, S., Leone, S., Trentalance, A., Muramatsu, M., and Marino, M. (2004a). Survival versus apoptotic 17 β -estradiol effect: role of ER α and ER β activated non-genomic signalling. *J. Cell. Physiol.* Published online at www.interscience.wiley.com/DOI10.1002/5CP.20219.
- Arvanitis, D. N., Wang, H., Bagshaw, R. D., Callahan, J. W., and Boggs, J. M. (2004). Membrane-associated estrogen receptor and caveolin-1 are present in central nervous system myelin and oligodendrocyte plasma membranes. *J. Neurosci. Res.* 75, 603–613.
- Bijlmakers, M. J., and Marsh, M. (2003). The on-off story of protein palmitoylation. *Trends Cell Biol.* 13, 32–42.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.
- Brzozowski, A. M., Pike, A.C.W., Dauter, Z., Hubbard, R. E., Bonn, T., Engstrom, O., Homan, L., Greene, G. F., Gustafsson, J.-Å., and Carlquist, M. (1997). Molecular basis of agonism and antagonism in the oestrogen receptor. *Nature* 389, 753–758.
- Castoria, G., Barone, M. V., Di Domenico, M., Bilancio, A., Ametrano, D., Migliaccio, A., and Auricchio, F. (1999). Non-transcriptional action of oestradiol and progestin triggers DNA synthesis. *EMBO J.* 18, 2500–2510.
- Castoria, G., Migliaccio, A., Bilancio, A., Di Domenico, M., de Falco, A., Lombardi, M., Fiorentino, R., Varricchio, L., Barone, M. V., and Auricchio, F. (2001). PI3-kinase in concert with Src promotes the S-phase entry of oestradiol-stimulated MCF-7 cells. *EMBO J.* 20, 6050–6059.
- Chambliss, K. L., and Shaul, P. W. (2002). Rapid activation of endothelial NO synthase by estrogen: evidence for a steroid receptor fast-action complex (SRFC) in caveolae. *Steroids* 67, 413–419.
- Chambliss, K. L., Yuhanna, I. S., Anderson, R. G., Mendelsohn, M. E., and Shaul, P. W. (2002). ER β has nongenomic action in caveolae. *Mol. Endocrinol.* 16, 938–946.

- Coleman, K. M., and Smith, C. L. (2001). Intracellular signaling pathways: nongenomic actions of estrogens and ligand-independent activation of estrogen receptors. *Front. Biosci.* 6, D1379–D1391.
- Dan, P., Cheung, J.C., Scriven, D. R., and Moore, E. D. (2003). Epitope-dependent localization of estrogen receptor- α , but not - β , in en face arterial endothelium. *Am. J. Physiol.* 284, H1295–H1306.
- Deecher, D. C., Swiggard, P., Frail, D. E., and O'Connor, L. T. (2003). Characterization of a membrane-associated estrogen receptor in a rat hypothalamic cell line (D12). *Endocrine* 22, 211–223.
- Fernando, R. I., and Wimalasena, J. (2004). Estradiol abrogates apoptosis in breast cancer cells through inactivation of BAD: Ras dependent non-genomic pathways requiring signaling through ERK and AKT. *Mol. Biol. Cell* 15, 3266–3284.
- Han, J. M., Kim, Y., Lee, J. S., Lee, C. S., Lee, B. D., Ohba, M., Kuroki, T., Suh, P.-G., and Ryu, S. H. (2002). Localization of phospholipase D1 to caveolin-enriched membrane via palmitoylation: implications for epidermal growth factor signalling. *Mol. Biol. Cell* 13, 3976–3988.
- Hegy, G. B., Shackleton, C. H., Carlquist, M., Bonn, T., Engstrom, O., Sjöholm, P., and Witkowska, H. E. (1996). Carboxymethylation of the human estrogen receptor ligand-binding domain-estradiol complex: HPLC/ESMS peptide mapping shows that cysteine 447 does not react with iodoacetic acid. *Steroids* 61, 367–373.
- Herbert, B., Truss, M., Beato, M., and Müller, R. (1994). Inducible regulatory elements in the human cyclin D₁ promoter. *Oncogene* 9, 1295–1304.
- Jakacka, M., Ito, M., Martinson, F., Ishikawa, T., Lee, E. J., and Jameson, J. L. (2002). An estrogen receptor ER α deoxyribonucleic acid-binding domain knock-in mutation provides evidence for nonclassical ER pathway signaling *in vivo*. *Mol. Endocrinol.* 16, 2188–2201.
- Katzenellenbogen, B. S., Bhardwaj, B., Fang, H., Ince, B. A., Pakdel, F., Reese, J. C., Schodin, D. and Wrenn, C. K. (1993) Hormone binding and transcription activation by estrogen receptors: analyses using mammalian and yeast systems. *J. Steroid. Biochem. Mol. Biol.* 47, 39–48.
- Kelly, M. J., and Levin, E. R. (2001). Rapid actions of plasma membrane estrogen receptors. *Trends Endocrinol. Metab.* 12, 152–156.
- Kousteni, S., *et al.* (2001). Nongenotropic, sex-nonspecific signaling through the estrogen or androgen receptors: dissociation from transcriptional activity. *Cell* 104, 719–790.
- Levin, E. R. (2001). Cell localization, physiology, and nongenomic actions of estrogen receptors. *J. Appl. Physiol.* 91, 1860–1867.
- Li, L., Haynes, M. P., and Bender, J. R. (2003). Plasma membrane localization and function of the estrogen receptor α variant (ER46) in human endothelial cells. *Proc. Natl. Acad. Sci. USA* 100, 4807–4812.
- Lobenhofer, E. K., Huper, G., Iglehart, J. D., and Marks, J. R. (2000). Inhibition of mitogen-activated protein kinase and phosphatidylinositol 3-kinase activity in MCF-7 cells prevents estrogen-induced mitogenesis. *Cell Growth Differ.* 11, 99–110.
- Mardones, G., and Gonzalez, A. (2003). Selective plasma membrane permeabilization by freeze-thawing and immunofluorescence epitope access to determine the topology of intracellular membrane proteins. *J. Immunol. Methods* 275, 169–177.
- Marino, M., Pallottini, V., and Trentalance, A. (1998). Estrogens cause rapid activation of IP₃-PKC- α signal transduction pathway in HEPG2 cells. *Biochem. Biophys. Res. Commun.* 245, 254–258.
- Marino, M., Distefano, E., Pallottini, V., Caporali, S., Ceracchi, G., and Trentalance, A. (2001a). β -Estradiol stimulation of DNA synthesis requires different PKC isoforms in HepG2 and MCF7 cells. *J. Cell Physiol.* 188, 170–177.
- Marino, M., Distefano, E., Trentalance, A., and Smith, C. L. (2001b). Estradiol-induced IP₃ mediate the estrogen receptor activity expressed in human cells. *Mol. Cell. Endocrinol.* 182, 19–26.
- Marino, M., Acconcia, F., Bresciani, F., Weisz, A., and Trentalance, A. (2002). Distinct non-genomic signal transduction pathways controlled by 17 β -estradiol regulate DNA synthesis and cyclin D₁ gene transcription in HepG2 cells. *Mol. Biol. Cell* 13, 3720–3729.
- Marino, M., Acconcia, F., and Trentalance, A. (2003). Biphasic estradiol-induced AKT-phosphorylation is modulated by PTEN via MAP kinase in HepG2 cells. *Mol. Biol. Cell* 14, 2583–2591.
- Migliaccio, A., Castoria, G., Di Domenico, M., de Falco, A., Bilancio, A., and Auricchio, F. (2002). Src is an initial target of sex steroid hormone action. *Ann. N.Y. Acad. Sci.* 963, 185–190.
- Neff, S., Sadowski, C., and Miksicek, R. J. (1994). Mutational analysis of cysteine residues within the hormone-binding domain of the human estrogen receptor identifies mutants that are defective in both DNA-binding and subcellular distribution. *Mol. Endocrinol.* 8, 1215–1223.
- Norfleet, A. M., Thomas, M. L., Gametchu, B., and Watson, C. S. (1999). Estrogen receptor- α detected on the plasma membrane of aldehyde-fixed GH3/B6/F10 rat pituitary tumor cells by enzyme-linked immunocytochemistry. *Endocrinology* 140, 3805–3814.
- Pappas, T. C., Gametchu, B., and Watson, C. S. (1995). Membrane estrogen receptors identified by multiple antibody labeling and impeded-ligand binding. *FASEB J.* 9, 404–410.
- Razandi, M., Pedram, A., Greene, G. L., and Levin, E. R. (1999). Cell membrane and nuclear estrogen receptors (ERs) originate from a single transcript: studies of ER α and ER β expressed in Chinese hamster ovary cells. *Mol. Endocrinol.* 13, 307–319.
- Razandi, M., Pedram, A., and Levin, E. R. (2000). Plasma membrane estrogen receptors signal to antiapoptosis in breast cancer. *Mol. Endocrinol.* 14, 1434–1447.
- Razandi, M., Oh, P., Pedram, A., Schnitzer, J., and Levin, E. R. (2002). ERs associate with and regulate the production of caveolin: implications for signaling and cellular actions. *Mol. Endocrinol.* 16, 100–115.
- Razandi, M., Alton, G., Pedram, A., Ghonshani, S., Webb, P., and Levin, E. R. (2003). Identification of a structural determinant necessary for the localization and function of estrogen receptor α at the plasma membrane. *Mol. Cell Biol.* 23, 1633–1646.
- Reid, G., Hubner, M. R., Metivier, R., Brand, H., Denger, S., Manu, D., Beaudouin, J., Ellenberg, J., and Gannon, F. (2003). Cyclic, proteasome-mediated turnover of unliganded and liganded ER α on responsive promoters is an integral feature of estrogen signaling. *Mol. Cell* 11, 695–707.
- Resh, M. D. (1999). Fatty acylation of proteins: new insights into membrane targeting of myristoylated and palmitoylated proteins. *Biochim. Biophys. Acta* 1451, 1–16.
- Ropero, A. B., Soria, B., and Nadal, A. (2002). A nonclassical estrogen membrane receptor triggers rapid differential actions in the endocrine pancreas. *Mol. Endocrinol.* 16, 497–505.
- Simoncini, T., Hafezi-Moghadam, A., Brazil, D. P., Ley, K., Chin, W. W., and Liao, J. K. (2000). Interaction of oestrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase. *Nature* 407, 538–541.
- Song, R. X., McPherson, R. A., Adam, L., Bao, Y., Shupnik, M., Kumar, R., and Santen, R. J. (2002). Linkage of rapid estrogen action to MAPK activation by ER α -Shc association and Shc pathway activation. *Mol. Endocrinol.* 16, 116–127.
- Song, R. X., Barnes, C. J., Zhang, Z., Bao, Y., Kumar, R., and Santen, R. J. (2004). The role of Shc and insulin-like growth factor 1 receptor in mediating the translocation of estrogen receptor α to the plasma membrane. *Proc. Natl. Acad. Sci. USA* 101, 2076–2081.
- Toran-Allerand, C. D., Guan, X., MacLusky, N. J., Horvath, T. L., Diano, S., Singh, M., Connolly, E. S., Jr., Nethrapalli, I. S., and Tinnikov, A. A. (2002). ER-X: a novel, plasma membrane-associated, putative estrogen receptor that is regulated during development and after ischemic brain injury. *J. Neurosci.* 22, 8391–8401.
- Varner, A. S., Ducker, C. E., Xia, Z., Zhuang, Y., De Vos, M. L., and Smith, C. D. (2003). Characterization of human palmitoyl-acyl transferase activity using peptides that mimic distinct palmitoylation motifs. *Biochem. J.* 373, 91–99.
- Wise, P. M., Dubal, D. B., Wilson, M. E., Rau, S. W., and Bottner, M. (2001). Neuroprotective effects of estrogen-new insights into mechanisms of action. *Endocrinology* 142, 969–973.