

Parallel adaptive radiations in two major clades of placental mammals

Ole Madsen^{1,2}, Mark Scally^{2,3,4}, Christophe J. Douady^{3,4}, Diana J. Kao³, Ronald W. DeBry⁵, Ronald Adkins⁶, Heather M. Amrine^{3,7}, Michael J. Stanhope^{4,8}, Wilfried W. de Jong^{9,1} & Mark S. Springer^{3,7}

1. Department of Biochemistry, University of Nijmegen, PO Box 9101, 6500 HB Nijmegen, The Netherlands
2. Department of Biology, University of California, Riverside, California 92521, USA
3. Queen's University of Belfast, Biology and Biochemistry, 97 Lisburn Road, Belfast, BT9 7BL, United Kingdom
4. Department of Biological Sciences, Box 210006, University of Cincinnati, Cincinnati, Ohio 45221 USA
5. Biology Department, University of Massachusetts, Amherst, Massachusetts, 01003 USA
6. Graduate Group in Genetics, University of California, Riverside, California 92521 USA
7. Bioinformatics, SmithKline Beecham Pharmaceuticals, 1250 South Collegeville Road, UP1345, Collegeville, Pennsylvania 19426 USA
8. Institute for Biodiversity and Ecosystem Dynamics, 1090 GT Amsterdam, The Netherlands
9. These authors contributed equally to this work

Correspondence to: Mark S. Springer^{3,7} Correspondence and requests for materials should be addressed to M.S.S. (e-mail: Email: mark.springer@ucr.edu).

Top of page

Abstract

Higher level relationships among placental mammals, as well as the historical biogeography and morphological diversification of this group, remain unclear^{1, 2, 3}. Here we analyse independent molecular data sets, having aligned lengths of DNA of 5,708 and 2,947 base pairs, respectively, for all orders of placental mammals. Phylogenetic analyses resolve placental orders into four groups: Xenarthra, Afrotheria, Laurasiatheria, and Euarchonta plus Glires. The first three groups are consistently monophyletic with different methods of analysis. Euarchonta plus Glires is monophyletic or paraphyletic depending on the phylogenetic method. A unique nine-base-pair deletion in exon 11 of the *BRCA1* gene provides additional support for the monophyly of Afrotheria, which includes proboscideans, sirenians, hyracoids, tubulidentates, macroscelideans, chrysochlorids and tenrecids. Laurasiatheria contains cetartiodactyls, perissodactyls, carnivores, pangolins, bats and eulipotyphlan insectivores. Parallel adaptive radiations have occurred within Laurasiatheria and Afrotheria. In each group, there are aquatic, ungulate and insectivore-like forms.

<http://www.nature.com/nature/journal/v409/n6820/full/409610a0.html>

