

**WHOI-87-17**

*Copy 2*

Joint Program:  
Woods Hole Oceanographic Institution, USA  
University of Kiel, FRG

NORDIC SEAS SEDIMENTATION  
DATA FILE, Vol. 1, 1987

**Particle Fluxes,  
Northeastern Nordic Seas  
1983 - 1986**



**DOCUMENT  
LIBRARY**  
Woods Hole Oceanographic  
Institution

Sponsored by  
Office of Naval Research  
Arlington, Virginia USA

Particle Fluxes, North-Eastern Nordic Seas: 1983-1986

(Nordic Seas Sedimentation Data File, Vol. 1)

by

Susumu Honjo, Steven J. Manganini, Amy Karowe, Bonnie L. Woodward

Woods Hole Oceanographic Institution  
Woods Hole, Massachusetts 02543

April, 1987

Technical Report

Funding was provided by the Office of Naval Research  
under Grant Number N00014--85-C-0001.

Reproduction in whole or in part is permitted for any purpose of  
The United States Government. This report should be cited as:  
Woods Hole Oceanog. Inst. Tech. Rept. WHOI-87-17.

Approved for publication; distribution unlimited

Approved for Distribution:

  
\_\_\_\_\_  
David A. Ross, Chairman  
Geology and Geophysics Department



Joint Program:

Woods Hole Oceanographic Institution  
and  
University of Kiel, University of Bremen, FRG

NORDIC SEAS SEDIMENTATION

DATA FILE, Vol. 1

PARTICLE FLUXES,

NORTH-EASTERN NORDIC SEAS:

1983 - 1986

Sponsored by  
Office of Naval Research  
Arlington, Virginia USA

## Table of Contents

	Page
Abstract . . . . .	1
Introduction . . . . .	1
Field Program . . . . .	3
Laboratory Analysis . . . . .	3
Results . . . . .	4
Acknowledgments . . . . .	5
References . . . . .	6
Table 1. Sediment trap moorings, 1983-1987 . . . . .	7
Table 2. Average mass fluxes in the northern Nordic Seas, 1983-1986 .	10
Table 3. Comparison of mass fluxes between stations in the northern Nordic Seas, 1983-1986 . . . . .	11
Figure 1. Location of map of sediment trap moorings . . . . .	12
Figure 2. PARFLUX Mark 5 and 6 sediment traps . . . . .	13
Figure 3. Analytical procedure . . . . .	14
Flux Data Files:	
Norwegian-Atlantic Current Area . . . . .	15
LB-1     East Lofoten Basin . . . . .	16
BI-1     Bear Island-West of Starfjord . . . . .	26
NA-1     Aegir Ridge . . . . .	36
NB-1     East of Jan Mayen . . . . .	46
East Greenland/Fram Strait Area . . . . .	56
FS-1     Central Fram Strait . . . . .	57
GB-2.2K   Greenland Basin, 1,966 m . . . . .	67
GB-2.3K   Greenland Basin, 2,817 m . . . . .	75

Nordic Seas Sedimentation Data File, Volume 1

Particle Fluxes, North-Eastern Nordic Seas: 1983 - 1986

by

Susumu Honjo, Steven J. Manganini, Amy Karowe, Bonnie L. Woodward

Woods Hole Oceanographic Institution  
Woods Hole, Massachusetts 02543

April, 1987

**Abstract**

Seventy-nine particle flux samples were collected from 1983 to 1986 using 7 automated time-series sediment traps at 6 stations distributed in the northern and eastern portion of the Nordic Seas as part of a German/U.S. joint program on arctic sedimentation studies. Each sample represents either one month or two weeks of sedimentation at approximately 400 m above the sea floor. In this data file the results of laboratory analysis conducted at the Woods Hole Oceanographic Institution, U.S.A. of the main sedimentological criteria: total mass, carbonate, opal, combustible, organic carbon, nitrogen, and lithogenic mass are presented in both tabular and histogram form. Results from the southern and western portion of the Nordic Seas will be published as they become available.

**Introduction**

Supported by the United States Office of Naval Research, the Woods Hole Oceanographic Institution (WHOI), with the cooperation of the University of Kiel and the University of Bremen, Federal Republic of Germany, has conducted a basin-wide sedimentological research program in the Nordic Sea since the summer of 1983. One of the major field experiments was deployment of 16 sets of sediment trap-current meter moorings for a period of about one year each throughout the basin. During the first half of the program we deployed 6 year-round moorings between August 1983 and August 1986 in the Fram Strait and Norwegian Basin. Details of mooring positions, depths, duration of deployment are summarized in Table 1. During the second part of the program, sediment trap mooring deployments and laboratory analyses of incoming samples will continue around Iceland, coastal Greenland and selected stations in cooperation with the Marine Research Institute, Reykjavik. The

University of Hamburg maintains 3 sediment trap mooring stations in the southern North Sea and we cooperate with their program on some of the laboratory analyses (Fig. 1).

The Nordic Sea is a basin, approximately 2.5 million square kilometers, defined by the east coast of Greenland to the west, Iceland to the south, the Norwegian coast to the east, and Spitsbergen to the north. It connects to the Arctic Ocean via the Fram Strait and to the North Atlantic via the Faeroe and Denmark straits (Hurdle, 1986). In short, the Nordic Sea is the bridge between the Arctic Ocean and the North Atlantic Ocean, and therefore is of global significance in regard to the Atlantic environment.

Most of the Nordic Sea lies north of the Arctic Circle. The net solar energy input is strongly limited in this basin due to low angle insolation during the summer and day-long darkness in the winter. Three longitudinal zones of ocean characteristics can be distinguished in this basin: 1) a zone along the east coast of Greenland which is covered by southerly flowing ice packs and floes in the East Greenland Current combined with fast-ice conditions on the immediate coast (Vinje, 1977). The surface temperature in this zone is 0°C throughout the year; 2) a zone on the east side of the basin where the warm, saline northward-flowing Norwegian-Atlantic Current prevails (Gathman, 1986); and 3) a zone in the central gyre which is often associated with mixed ice conditions where the other two zones meet in the middle of the basin (Wadhams, 1986; Swift, 1986). This unique arrangement of currents form several ocean fronts (Johannessen, 1986) and strong contrasts of oceanic conditions are seen within this relatively small basin. For example, the summer surface temperature difference between the east and west side of the basin along the 70th latitude (off Tromso, Norway to Scoresby Sound, Greenland, which are only about 1,000 km apart) is as great as 10°C in some years (Detrich, 1969). Thus the Nordic Sea embodies highly diversified specific environments within the basin boundary.

Very little is known about particle sedimentation and recycling schemes in the North Sea environment. Ocean particles in the Nordic Basin also involve specific origins, flux and processes which reflect varied oceanic characteristics. Questions include: how much of the particulate carbon and other biogenic particles settle down to the sea floor and how do they compare with surface production which is produced under severely limiting Arctic conditions? What is the sedimentary mechanism of lithogenic particles in the Arctic open ocean environment? How are these sedimentary particle processes related to ice coverage and mixed ice zone conditions? This research aims to answer these questions and, optimally, to draw a realistic model of particle flux and sedimentation in relation to other critical high latitude ocean environmental factors.

## Field Program

Experimental logistics in the Nordic Sea are generally very difficult compared to lower latitude oceanographic endeavors; winter storms and ice coverage hinder deployment and recovery of large bottom tethered mooring arrays. Because of strong seasonality, flux measurements in high latitudes must cover at least a one-year cycle of seasons. We have used automated time-series sediment traps left unattended for about one year. A sediment trap used in this environment requires a large opening in order to collect enough volume of sample during the winter months when the flux is estimated to be extremely small. We used a PARFLUX Mark 5 and Mark 6 whose apertures are 1.2 and 0.5 m<sup>2</sup> with 12 and 13 sampling increments, respectively (Honjo and Doherty, 1987, in press) (Table 1, Fig. 2). The sediment traps were deployed at approximately 400 m above the sea floor at most mooring sites. The exception was a mooring with two sediment traps deployed along a taut line which was set in the Greenland Basin. One to three current meters were deployed with each sediment trap mooring. A transmissometer was deployed with two Fram Strait moorings for one year, 1984-1985. The results from the current meter and transmissometer experiments will be published elsewhere. The deployment/recovery procedure for sediment trap mooring arrays was described in a separate paper (Honjo and Doherty, 1987, in press). We used sodium azide as a preservative (Honjo, 1980).

## Laboratory Analysis

Recovered samples were refrigerated throughout the transportation and storage period. Each sample was equally shared with Dr. Gerold Wefer's laboratory (University of Bremen). Our responsibility at WHOI was to clarify the nature of the sediment trap collected samples with regard to basic sedimentological criteria. Dr. Wefer's group is investigating stable isotopes in planktonic foraminiferal tests and some biocoenosis composition in the samples.

Upon arrival at WHOI, each sample was sieved through a 1mm Nylon mesh. This was necessary to maintain precise sample splitting. Particles smaller than 1 mm were further split into smaller aliquots by a precision wet sample splitter (Honjo, 1980). The split aliquots were further sieved through a 62 micron mesh for the LB-1, FS-1, and BI-1 samples in order to separate foraminiferal tests and radiolarian shells in this size category more efficiently. We analyzed individually samples in each size category for the following criteria. All results were normalized to flux values in mg m<sup>2</sup>day (Honjo, 1980).

Total mass  
Carbonate mass  
Combustible mass

Noncombustible mass  
Opal mass  
Lithogenic mass  
Organic carbon, nitrogen, and hydrogen mass

A detailed description of the analytical methods applied to this research will be published elsewhere. In summary as illustrated in Fig. 3, the total mass flux was obtained as the average of dry mass weight of the three 16th aliquots. The carbonate content was obtained from the dry weight difference before and after decalcification by 1N acetic acid at room temperature. A decalcified aliquot was combusted for 3 hours at 500°C to obtain the mass of combustible organic matter as the difference between a decalcified sample and ash weight. Biogenic silica, or opal, content was analyzed by the sodium carbonate leaching method modified from Eggiman *et al.*, 1980, on decalcified aliquots. Lithogenic particle flux, mostly clay and fine rock-forming detritus, was gained by subtracting the opal flux from the noncombustible flux. Organic carbon, nitrogen, and hydrogen content were analyzed using a Perkin-Elmer Elemental Analyzer, type 240C. We used at least 100 mg of decalcified samples (Fig. 3).

Total flux, therefore, is equal to the sum of carbonate, noncombustible, and combustible fluxes. The sum of biogenic opal and lithogenic fluxes should be the noncombustible flux. Insignificant discrepancies appear in some total flux values in this data file due to the rounding out processes during calculation. We regard the combustible portion of the flux as organic matter flux (Honjo, 1980). Combustible flux consists of organic carbon, nitrogen, and hydrogen balanced with oxygen and other unidentifiable ignition loss. The amount of organic nitrogen in the GB-1 sample was too small to analyze within our level of confidence. The opal content in the GB-2, 1966 trap sample was also too small to analyze with the leaching method at the time but we are making an effort to bring up significant numbers.

The phosphorus flux from this area will be published in a separate file. The results of analysis of 15 trace elements from all time-series sediment trap samples treated in the present data file (total of 1,185 analyses) will be published in a separate volume.

#### Results

The purpose of this data file is to publish a summary of available data on the flux in the north-eastern Nordic Sea for public use. Scientific interpretations and models will not be included in this publication.

The annual averages in two major areas, Norwegian-Atlantic current area and the East Greenland current area (sea ice prevailed) based upon fluxes from 6 stations presented in this report, is given in Table 2. The annual fluxes of sedimentary components from 6 stations are tabulated in Table 3 for comparison. At the beginning of each data file for individual stations are given the sample identification numbers, opening and closing dates, length of collection period and mid-point date during which the samples were collected. On subsequent pages are given the percentages of total flux of three size categories: particles which passed through 62 micrometer mesh ( $< 63 \mu\text{m}$ ), particles retained in a 1 mm mesh ( $> 1 \text{ mm}$ ), and particles in between ( $63 \mu\text{m} - 1 \text{ mm}$ ). In the rightmost column of the table, the total flux of size categories combined is given. The columns of each histogram are labeled according to mid-point day of the sampling period. The six flux categories listed in the previous section are included in each data set.

#### Acknowledgments

Without the encouragement and support of Dr. G. Leonard Johnson, Office of Naval Research, this first entire ocean basin sedimentation study applying the flux concept would never have been started. We sincerely thank him for his insight and strong commitment to excellent science.

The Nordic Sea is one of the most difficult oceans with regard to experimental logistics. We have received a large amount of good will support from international colleagues; a large part of our success is due to them and even the unusually long acknowledgment in this paper may cover only a portion. In particular, the Alfred Wegener Institution of Polar and Marine Research, Bremerhaven provided us with vital shiptime on board R/V Polarstern for this experiment. Dr. Jörn Thiede, Chief Scientist of the 1984 and 1985 legs, took every possible opportunity to help us with his professional competence and personal care during this experiment. We also thank the R/V Meteor (old) and the Deutche Hydrographische Institute, Hamburg, which supported us in a difficult mission to recover a malfunctioned mooring system and to deploy a large array in the Greenland Sea during the summer of 1985. We also thank the R/V Meteor (new) and R/V Valdivia, University of Hamburg, for their high quality support of the mooring experiments in 1986.

The Nordic Sea program has been carried out under the mutual cooperation among the University of Bremen, University of Kiel and Woods Hole Oceanographic Institution. Dr. Gerold Wefer, our partner, has provided many useful suggestions in research and has been very helpful in providing vital logistic support. We owe him our sincere gratitude. We thank for their dedication and imagination: Dr. Vernon L. Asper,

University of Southern Mississippi, and Dorinda Ostermann, WHOI, who made it possible to deploy and recover the first 4 mooring arrays in the northern Nordic Sea in 1983 and 1984; Peter Clay and Thomas Crook who provided vital assistance in recovering a stranded GB-1 mooring in the summer of 1985; Emily Evans who took care of communication traffic and data editing during this program.

#### References

- Detrich, G., 1969. Atlas of the Hydrography of the Northern North Atlantic, International Council for the Exploration of the Sea. Copenhagen.
- Eggemann, D.W., Manheim, F.T. and Betzer, P.R., 1980. Dissolution and Analysis of Amorphous Silica in Marine Sediments. Journal of Sedimentary Petrology, 50(1): 215-225.
- Gathman, S.G., 1986. Climatology. In: The Nordic Seas (Hurdle, B.G., ed.), 1-18, Springer-Verlag, New York, 777 pp.
- Hurdle, B.G. (ed.), 1986. The Nordic Seas. Springer-Verlag, New York, 777 pp.
- Honjo, S., 1980. Material Fluxes and Modes of Sedimentation in the Mesopelagic and Bathypelagic zones. Journal of Marine Research, 38: 53-97.
- Honjo, S. and Doherty, K.W., 1987 (in press). Large Aperture Time-Series Oceanic Sediment Traps; Design Objectives, Construction and Applications. Deep-Sea Research.
- Johannessen, O., 1986. Brief overview of Physical Oceanography. In: The Nordic Seas (Hurdle, B.G., ed.), 103-127, Springer-Verlag, New York, 777 pp.
- Swift, J.H., 1986. The Arctic Waters. In: The Nordic Seas (Hurdle, B.G., ed.), 129-153, Springer-Verlag, New York, 777 pp.
- Vinje, T.E., 1977. Sea Ice Conditions in the European Sector of the Marginal Sea of the Arctic, 1966-75. Norsk Polarinstitutt Arbok, 1975: 163-174.
- Wadhams, P., 1986. The Ice Cover. In: The Nordic Seas (Hurdle, B.G., ed.), 21-84, Springer-Verlag, New York, 777 pp.

Table 1a

SEDIMENT TRAP MOORINGS IN THE GREENLAND BASIN/NORWEGIAN SEA, 1983-1987.  
SUPPORTED BY ONR (WHOI), DFG/GMIS (KIEL UNIV.), GMIS (HAMBURG UNIV.)  
COORDINATED BY WHOI/ONR

March 20, 1985  
Revised: September 26, 1985  
Revised: August 26, 1986  
Revised: February 24, 1987  
Revised: April 13, 1987

Table 1b

Rev: March 20, 1985  
 Rev: September 26, 1985  
 Rev: February 24, 1987  
 Rev: April 13, 1987

No.	Year	ID No.	Location	Latitude/ Longitude	Water depth (m)	Mooring height (m)	No. 501b spheres	Trap type and number/mooring (off floor)	Trap depth, m	Cups/ duration dys:hrs	Extra Instrs.	Deploy/ Recov:	Ship: dep/rec:
7	1985	GB-2	Greenland Basin	74°35'N 06°43'W	3,445	2,008	34	Mark 5 (2)	2,823 (622) 881 (2564)	25/ 14:0	None	8/02/85 8/23/86	Meteor Valdivia
<b>All equipment worked.</b>													
8	1985	FS-2	Central Fram Str.	79°00'N 04°55.0'E	2,430	1,862	32	Mark 6 (2)	1,929 (501) 1,000 (1,500)	13/ 27:0	1 CM 1 TMM	7/29/85 8/17/86	Pstern Valdivia
<b>Only 5 cups worked. Transmissometer flooded. Current record OK.</b>													
9	1985	NB-1	W. Norwegian Sea	70°00'N 01°58'W	3,296	2,773	15	Mark 6 (1)	2,749 (520)	13/ 30:0	2 DEP	8/18/85 7/15/86 Meteor (new)	Pstern
<b>New 25 cup system; 17 samples recovered (U. Bremen, G. Wefer, P.I.).</b>													
10	1985	NA-1	Norway Abyssal Plain	65°31'N 00°64'E	3,058	2,558	15	Mark 6 (1)	2,630 (428)	13/ 30:0	2 DEP	8/19/85 7/18/86 Meteor (new)	Pstern
<b>New electronics. 13 samples recovered.</b>													
11	1985	LB-2 (Kiel)	Lofoten Basin	69°30'N 10°00'E	3,160	432	25	Mark 5 (1)	2,760 (400) (15?) (25?) 30/	12/ 25? 30/	2 CM 2 DEP	8/9/85 8/86	Pstern Valdivia
<b>Permanent station for Bremen group to be funded by RG-95, DFG, after 1986 (U. Bremen, G. Wefer, P.I.).</b>													

Table 1c

March 20, 1985  
 Rev: September 26, 1985  
 Rev: February 24, 1987  
 Rev: April 13, 1987

No.	Year	ID No.	Location	Latitude/ Longitude	Water depth (m)	Mooring height (m)	No. 501b spheres	Trap type and number/mooring (off floor)	Trap depth, m	Cups/ dys:hrs	Extra duration Instrs.	Deploy/ depth m	Ship: dep/rec
□ 12	1986	IP-1	Iceland Plateau	68°01'N 12°39'W	1,884	458	10	Mark 6 (1)	1,454 (430)	13/ 30:0	1 CM (1458)	10/20/86 10/2/87	Smdsn Smdsn
□ 13	1986	MRI-1	South Iceland	62°58'N 21°32'W	1,004	459	10	Mark 5 (1)	574 (430)	13/ 30:0	1 CM (577)	10/3/86 10/2/87	Smdsn Smdsn
□ 14	1986	NDS-1	N. Denmark Strait	69°30'N 21°13'W	446	156	20	Mark 6 (1)	136 (100)	13/ 30:0	1 CM (350)	10/14/86 9/2/87	Smdsn Smdsn

Very high-tension mooring.

ONR: Office of Naval Research, Code 1125AR  
 WHOI: Woods Hole Oceanographic Institution  
 DFG: Deutsche Forschungsgemeinschaft  
 GMIS: German Ministry of Industry and Science  
 DHI: Deutsche Hydrographische Institute  
 MRI: Marine Research Institute, Iceland

ID No: ONR/WHOI mooring identification number  
 Ship, deployment and recovery:  
 Pstern: R/V Polarstern, Bremerhaven  
 Meteor: R/V Meteor, Hamburg  
 Valdivia: R/V Valdivia, Hamburg  
 Smdsn: R/V Bjarni Saemundsson, Reykjavik

CM: Current meter  
 TMM: Transmissometer  
 DEP: Dissolution experiment package

Table 2. Average Mass Fluxes; Northern Nordic Seas, 1983-1986.  
 Average Fluxes and (standard deviation) mg m<sup>-2</sup>day<sup>-1</sup>.

	Norwegian-Atlantic Current Area:	E. Greenland and Fram Strait Area:
Moorings:	LB-1, BI-1, NA-1, NB-1	FS-1, GB-21, GB-23
Total Flux	21.31 (5.39)	8.45 (1.81)
Carbonate Flux	9.03 (1.96)	2.42 (0.95)
Noncombustible Flux	9.14 (4.87)	4.56 (1.05)
Combustible Flux	3.24 (1.55)	1.55 (0.83)
Biogenic Opal Flux	1.55 (0.36)	--- *
Lithogenic Flux	7.55 (4.69)	--- *
Organic Carbon Flux	1.34 (1.00)	0.58 (0.31)
Nitrogen Flux	0.16 (0.11)	0.09 (0.06)

\* Not detectable

Trap Station Codes:

- LB-1: East Lofoten Basin
- FS-1: Central Fram Strait
- BI-1: Bear Island - west of Storfjord
- NA-1: Aegir Ridge
- NB-1: East of Jan Mayen
- GB-21: Greenland Basin (shallow)
- GB-23: Greenland Basin (deep)

Table 3. Comparison of mass fluxes between 6 stations in the Northern Nordic Seas, 1983-1986.

Area:	Norwegian-Atlantic Current				East Greenland/Fram Strait		
Trap Station:	LB-1	BI-1	NA-1	NB-1	FS-1	GB-21	GB-23
Latitude	69°30N	75°51N	65°31N	70°00N	78°52N	74°35N	75°35N
Longitude	10°00E	11°28E	00°64E	01°58W	01°22E	06°43W	06°43W
Trap Depth	2,760m	1,700m	2,630m	2,749m	2,440m	1,966m	2,871m
Total Flux**	22.80	28.40	17.36	16.79	7.20	8.79	10.21
Carbonate Flux	11.40	6.61	9.18	8.93	1.40	2.59	3.28
Noncombustible Flux	8.07	16.31	5.94	6.24	4.26	3.65	5.73
Combustible Flux	3.37	5.35	2.31	1.90	0.92	2.50	1.23
Biogenic Opal Flux	1.12	1.96	1.68	1.44	0.60	---	2.61
Lithogenic Flux	6.95	14.35	4.26	4.65	4.00	---	3.12
Organic Carbon Flux	1.37	2.85	0.59	0.53	0.41	0.94	0.40
Nitrogen Flux	0.18	0.30	0.08	0.08	0.06	0.16	0.06

\*\* Flux is in mg m<sup>2</sup>day

Trap Station Codes:

- LB-1: East Lofoten Basin
- FS-1: Central Fram Strait
- BI-1: Bear Island - west of Storfjord
- NA-1: Aegir Ridge
- NB-1: East of Jan Mayen
- GB-21: Greenland Basin (shallow)
- GB-23: Greenland Basin (deep)





Figure 1. Approximate positions of sediment trap-current meter moorings in the Nordic Seas, 1983-1987.



Mark 6-13 ( $0.5 \text{ m}^2$  aperture with 13 sampling bottles).

Mark 5-12 ( $1.2 \text{ m}^2$  aperture with 12 sampling bottles).





Figure 2. PARFLUX Mark 5 and 6 sediment traps (from Honjo and Doherty, 1987, Fig. 2).






Figure 3. Sample process flow diagram for sedimentological analysis of Nordic Sea flux samples.

NORWEGIAN-ATLANTIC CURRENT AREA

LB-1  
 EAST LOFOTEN BASIN  
 69°30' N, 10°00'E  
 Trap depth: 2,760m Water depth: 3,160m


Annual Fluxes. (g/m/yr):

Total.....	22.80
Carbonate.....	11.40
Noncombustible.....	8.07
Combustible.....	3.37
Opal.....	1.12
Lithogenic.....	6.95
Organic C.....	1.37
N.....	0.18

PARFLUX Mark 5-12

Sample ID	Opening Date	Closing Date	Span	Mid. Date
1 LB1-2600-1	15-AUG-83	13-SEP-83	29.33	30-AUG-83
2 LB1-2600-2	13-SEP-83	12-OCT-83	29.33	28-SEP-83
3 LB1-2600-3	12-OCT-83	11-NOV-83	29.33	27-OCT-83
4 LB1-2600-4	11-NOV-83	10-DEC-83	29.33	26-NOV-83
5 LB1-2600-5	10-DEC-83	08-JAN-84	29.33	25-DEC-83
6 LB1-2600-6	08-JAN-84	07-FEB-84	29.33	23-JAN-84
7 LB1-2600-7	07-FEB-84	07-MAR-84	29.33	22-FEB-84
8 LB1-2600-8	07-MAR-84	05-APR-84	29.33	22-MAR-84
9 LB1-2600-9	05-APR-84	05-MAY-84	29.33	20-APR-84
10 LB1-2600-10	05-MAY-84	03-JUN-84	29.33	20-MAY-84
11 LB1-2600-11	03-JUN-84	02-JUL-84	29.33	18-JUN-84
12 LB1-2600-12	02-JUL-84	01-AUG-84	29.33	17-JUL-84

Total Flux at Lofoten Basin (LB-1), 2600m, 1983-1984



Lofoten Basin I had 12 cups each open 29.33 days.


Mark 5 trap open from August 15 1983 to August 1 1984 at 2600 meters.

TOTAL FLUX (mg / m<sup>2</sup> / day)

Ttl is total Flux in all size classes

Cup #	< 63um			63um - 1			> 1mm			TOTAL		
	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	% of Ttl	FLUX	
1	70.74	72.41	17.90	18.32	11.36	11.63	100.00	102.36				
2	60.82	98.24	26.73	43.17	12.45	20.11	100.00	161.52				
3	64.75	91.36	20.72	29.24	14.53	20.51	100.00	141.10				
4	41.90	18.37	17.84	7.82	40.26	17.65	100.00	43.84				
5	55.73	10.74	13.75	2.65	30.50	5.88	100.00	19.27				
6	37.26	2.28	6.70	.41	56.20	3.44	100.00	6.12				
7	40.92	3.43	6.56	.55	52.51	4.40	100.00	8.38				
8	58.69	2.84	8.89	.43	32.28	1.56	100.00	4.84				
9	60.09	2.68	7.62	.34	32.26	1.44	100.00	4.46				
10	85.00	48.28	8.26	4.69	6.75	3.84	100.00	56.80				
11	88.39	64.87	8.61	6.32	3.00	2.20	100.00	73.39				
12	89.13	114.52	9.23	11.86	1.64	2.11	100.00	128.49				

Carbonate Flux at Lofoten Basin (LB-1), 2600m, 1983-1984



Lofoten Basin I had 12 cups each open 29.33 days.


Mark 5 trap open from August 15 1983 to August 1 1984 at 2600 meters.

Carbonate Flux

Ttl is total Flux in all size classes

Cup #	< 63um		63um - 1		> 1mm		TOTAL	
	% of Ttl	FLUX						
1	32.92	33.69	15.18	15.54	6.81	6.97	54.91	56.20
2	25.46	41.13	22.92	37.01	9.02	14.56	57.40	92.71
3	26.19	36.95	17.96	25.34	10.58	14.93	54.73	77.23
4	19.56	8.58	15.52	6.80	19.38	8.50	54.46	23.88
5	22.72	4.38	7.57	1.46	14.13	2.72	44.43	8.56
6	14.62	.89	3.22	.20	18.57	1.14	36.41	2.23
7	17.43	1.46	3.16	.26	23.57	1.98	44.16	3.70
8	24.67	1.19	4.83	.23	17.51	.85	47.02	2.28
9	23.20	1.03	3.55	.16	10.75	.48	37.50	1.67
10	32.73	18.59	3.31	1.88	2.64	1.50	38.69	21.97
11	35.03	25.71	4.57	3.35	1.25	.92	40.85	29.98
12	35.70	45.87	4.89	6.28	.84	1.08	41.43	53.23

Noncombustible Flux at Lofoten Basin (LB-1), 2600 m, 1983-84



Lofoten Basin I had 12 cups each open 29.33 days.


Mark 5 trap open from August 15 1983 to August 1 1984 at 2600 meters.

NON COMBUSTIBLE FLUX (mg / m<sup>2</sup> / day)

'Ttl is total Flux in all size classes

Cup #	< 63um		63um - 1		> 1mm		TOTAL	
	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX
1	29.76	30.46	1.15	1.18	.14	.15	31.06	31.79
2	27.47	44.37	2.29	3.69	.30	.48	30.05	48.54
3	30.15	42.55	1.45	2.05	.32	.45	31.93	45.05
4	15.00	6.57	1.35	.59	.58	.26	16.93	7.42
5	23.62	4.55	3.72	.72	1.78	.34	29.12	5.61
6	14.30	.87	2.03	.12	.83	.05	17.16	1.05
7	18.51	1.55	2.14	.18	.44	.04	21.10	1.77
8	24.04	1.16	2.63	.13	.00	.00	26.68	1.29
9	27.97	1.25	2.11	.09	4.28	.19	34.36	1.53
10	41.79	23.74	3.33	1.89	2.44	1.38	47.55	27.01
11	43.15	31.66	2.67	1.96	1.28	.94	47.10	34.57
12	43.49	55.88	2.89	3.71	.05	.06	46.43	59.65

Combustible Flux at Lofoten Basin (LB-1), 2600 m, 1983-84



Lofoten Basin I had 12 cups each open 29.33 days.


Mark 5 trap open from August 15 1983 to August 1 1984 at 2600 meters.

Combustible Flux

Ttl is total Flux in all size classes

Cup #	< 63um			63um - 1			> 1mm			TOTAL		
	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	% of Ttl	FLUX	
1	8.07	8.26	1.56	1.60	4.41	4.51	14.04	14.37				
2	7.89	12.74	1.52	2.46	3.14	5.07	12.55	20.27				
3	8.41	11.86	1.31	1.84	3.63	5.12	13.34	18.83				
4	7.34	3.22	.97	.42	20.30	8.90	28.61	12.54				
5	9.39	1.81	2.46	.47	14.59	2.81	26.43	5.09				
6	8.34	.51	1.45	.09	36.80	2.25	46.59	2.85				
7	4.98	.42	1.26	.11	28.50	2.39	34.74	2.91				
8	9.97	.48	1.42	.07	0.00	0.00	11.39	.55				
9	8.92	.40	1.97	.09	17.23	.77	28.12	1.25				
10	10.48	5.95	1.62	.92	1.68	.95	13.77	7.82				
11	10.21	7.50	1.37	1.00	.46	.34	12.05	8.84				
12	9.94	12.77	1.43	1.87	.76	.98	12.15	15.61				


Biogenic Silica flux at Lofoten Basin (LB-1), 2600m, 1983-1984



Sample ID#	OPAL <63	OPAL % tot.<63	OPAL 63-1	OPAL % tot.63-1	OPAL >1	OPAL % tot.>1	OPAL total	OPAL % of total
1 LB1-2600-1	3.87	3.78	0.38	0.37	0.03	0.03	4.28	4.18
2 LB1-2600-2	6.40	3.96	0.99	0.61	0.01	0.01	7.40	4.58
3 LB1-2600-3	3.82	2.71	0.60	0.43	0.16	0.11	4.57	3.24
4 LB1-2600-4	0.82	1.87	0.13	0.30	0.15	0.35	1.11	2.53
5 LB1-2600-5	0.51	2.65	0.05	0.26	0.10	0.54	0.66	3.45
6 LB1-2600-6*	0.09	1.47	0.09	0.02	0.00	0.03	0.09	1.47
7 LB1-2600-7	0.17	2.03	0.03	0.32	0.19	2.27	0.39	4.55
8 LB1-2600-8*	0.16	2.61	0.25	0.64	0.06	1.23	0.25	5.17
9 LB1-2600-9*	0.26	4.25	0.25	0.67	0.15	3.31	0.44	9.87
10 LB1-2600-10	3.42	6.02	0.53	0.93	0.30	0.53	4.25	7.47
11 LB1-2600-11	3.75	5.11	0.52	0.71	0.18	0.24	4.45	6.05
12 LB1-2600-12	8.12	6.32	1.04	0.81	0.05	0.04	9.21	7.17

Flux is in mg/m<sup>2</sup>/day.


Lithogenic Flux at Lofoten Basin (LB-1) 2600m, 1983-84



Sample I.D.	LITH <63	LITH<63 %tot.	LITH 63-1	LITH63-1 %tot.	LITH >1	LITH %tot.	LITH total	LITH% total
1 LB1-2600-1	26.59	25.98	0.80	0.78	0.12	0.12	27.51	26.88
2 LB1-2600-2	37.97	23.51	2.71	1.68	0.47	0.29	41.15	25.48
3 LB1-2600-3	38.73	27.45	1.45	1.03	0.29	0.21	40.47	28.68
4 LB1-2600-4	5.75	13.11	0.46	1.05	0.11	0.25	6.32	14.42
5 LB1-2600-5	4.04	20.96	0.67	3.48	0.24	1.25	4.95	25.69
6 LB1-2600-6	0.78	12.75	0.43	7.03	0.05	0.82	1.26	20.59
7 LB1-2600-7	1.38	16.46	0.18	2.15	0.04	0.48	1.60	19.09
8 LB1-2600-8	1.01	20.87	0.99	20.46	0.04	0.83	2.04	42.16
9 LB1-2600-9	0.99	22.20	0.99	22.20	0.04	0.90	2.02	45.29
10 LB1-2600-10	20.31	35.76	1.36	2.39	1.08	1.90	22.75	40.05
11 LB1-2600-11	27.91	38.03	1.43	1.95	0.76	1.04	30.10	41.01
12 LB1-2600-12	47.76	37.17	2.87	2.23	0.01	0.01	50.64	39.41

Flux is in mg/m²/day

Carbon Flux at Lofoten Basin (LB-1), 2600 m, 1983-84




Sample I.D.	NTGN .63	NTGN-63 %cmbf.	NTGN 63-1	NTGN63-1 %cmbf.	NTGN .1	NTGN1 %cmbf.	NTGN total	NTGNTot. %cmbf.
1 LB1-2600-1	0.40	4.84	0.16	1.11	0.21	1.46	0.76	5.29
2 LB1-2600-2	0.76	3.75	0.19	0.94	0.36	1.78	1.31	6.46
3 LB1-2600-3	0.66	3.51	0.10	0.53	0.30	1.59	1.06	5.63
4 LB1-2600-4	0.11	0.88	0.01	0.08	0.44	3.51	0.56	4.47
5 LB1-2600-5	0.10	1.96	0.03	0.59	0.21	4.13	0.34	6.68
6 LB1-2600-6*	0.04	1.27	0.00	0.02	0.13	4.56	0.17	5.85
7 LB1-2600-7*	0.02	0.69	0.01	0.34	0.13	4.47	0.16	5.50
8 LB1-2600-8*	0.03	1.42	0.01	0.47	0.10	4.50	0.04	5.90
9 LB1-2600-9*	0.02	1.60	0.00	0.00	0.06	4.80	0.08	5.40
10 LB1-2600-10	0.25	3.20	0.06	0.77	0.08	1.02	0.39	4.99
11 LB1-2600-11	0.25	2.83	0.04	0.45	0.03	0.34	0.32	3.82
12 LB1-2600-12	0.60	3.84	0.14	0.90	0.07	0.45	0.81	5.19

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Nitrogen Flux at Lofoten Basin (LB-1), 2600 m, 1983-84



Lofoten Basin I had 12 cups each open 29.33 days.


Mark 5 trap open from August 15 1983 to August 1 1984 at 2600 meters.

NITROGEN FLUX (mg / m<sup>2</sup> / day)

Ttl is Total Flux in all size classes.

Cup #	< 63um			63um - 1			> 1mm			TOTAL		
	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	FLUX	% of Ttl	% of Ttl	FLUX	
1	.39	.40	.15	.16	.20	.21	.75	.76				
2	.47	.76	.12	.19	.22	.36	.81	1.31				
3	.47	.66	.07	.10	.21	.30	.75	1.06				
4	.24	.11	.02	.01	1.01	.44	1.27	.56				
5	.53	.10	.17	.03	1.08	.21	1.78	.34				
6	.60	.04	.10	.01	2.09	.13	2.79	.17				
7	.27	.02	.09	.01	1.58	.13	1.94	.16				
8	.69	.03	.12	.01	.00	.00	.81	.04				
9	.43	.02	.11	.00	1.28	.06	1.82	.08				
10	.44	.25	.11	.06	.14	.08	.68	.39				
11	.35	.25	.06	.04	.04	.03	.44	.32				
12	.46	.60	.11	.14	.06	.07	.63	.81				

Hydrogen Flux at Lofoten Basin (LB-1), 2600 m, 1983-84



Sample I.D.	HYDC <sub>&lt;63</sub>	HYDC <sub>&lt;63</sub> %cmbf.	HYDC <sub>63-1</sub>	HYDC <sub>63-1</sub> %cmbf.	HYDC <sub>&gt;1</sub>	HYDC <sub>&gt;1</sub> %cmbf.	HYDC total	HYDC <sub>tot.</sub> %cmbf.
1 LB1-2600-1	0.59	4.11	0.09	0.63	0.40	3.78	1.08	7.51
2 LB1-2600-2	0.77	3.80	0.15	0.74	0.35	1.73	1.26	6.22
3 LB1-2600-3	0.85	4.51	0.07	0.37	0.35	1.86	1.28	6.80
4 LB1-2600-4	0.14	1.12	0.01	0.08	0.79	6.30	0.94	7.50
5 LB1-2600-5	0.12	2.36	0.03	0.59	0.26	5.11	0.41	8.06
6 LB1-2600-6*	0.04	1.40	0.01	0.35	0.20	7.02	0.24	9.42
7 LB1-2600-7*	0.03	1.03	0.03	0.92	0.22	7.56	0.25	9.59
8 LB1-2600-8*	0.03	1.42	0.03	1.56	0.06	2.93	0.04	1.90
9 LB1-2600-9*	0.03	2.40	0.01	0.80	0.08	6.40	0.11	8.90
10 LB1-2600-10	0.42	5.37	0.05	0.64	0.12	1.53	0.59	7.54
11 LB1-2600-11	0.37	4.19	0.05	0.57	0.03	0.34	0.45	5.09
12 LB1-2600-12	0.95	6.09	0.10	0.64	0.09	0.58	1.15	7.37

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

BI-1  
BEAR ISLAND - WEST OF STORFJORD  
75°51'N, 11°28'E  
Trap depth: 1,700m Water depth: 2,123m


Annual Fluxes. (g/m<sup>2</sup>/yr):

Total.....28.30  
Carbonate.....6.61  
Noncombustible.....16.31  
Combustible.....5.38  
Biogenic Opal.....1.96  
Lithogenic.....14.35  
Organic C.....2.85  
N.....0.30

PARFLUX Mark 5-13

Sample ID	Opening Date	Closing Date	Span	Mid. Date
26 BI1-1700-1	12-AUG-84	11-SEP-84	30.17	27-AUG-84
27 BI1-1700-2	11-SEP-84	11-OCT-84	30.17	26-SEP-84
28 BI1-1700-3	11-OCT-84	11-NOV-84	30.17	27-OCT-84
29 BI1-1700-4	11-NOV-84	11-DEC-84	30.17	26-NOV-84
30 BI1-1700-5	11-DEC-84	12-JAN-85	30.17	27-DEC-84
31 BI1-1700-6	12-JAN-85	11-FEB-85	30.17	27-JAN-85
32 BI1-1700-7	11-FEB-85	12-MAR-85	30.17	26-FEB-85
33 BI1-1700-8	12-MAR-85	11-APR-85	30.17	27-MAR-85
34 BI1-1700-9	11-APR-85	12-MAY-85	30.17	27-APR-85
35 BI1-1700-10	12-MAY-85	11-JUN-85	30.17	27-MAY-85
36 BI1-1700-11	11-JUN-85	11-JUL-85	30.17	26-JUN-85
37 BI1-1700-12	11-JUL-85	10-AUG-85	30.17	26-JUL-85


Total Flux at Bear Island (BI-1), 1700m, 1984-1985



Sample I.D.	TTLF <1	<1 % of total		TTLF >1	>1 % of total		TTLF total
		TTLF <1	<1 % of total		TTLF >1	>1 % of total	
26 BI1-1700-1	42.41	79.97	10.62	20.03	53.03		
27 BI1-1700-2	100.12	82.55	21.17	17.45	121.29		
28 BI1-1700-3	89.79	82.63	18.88	17.37	108.67		
29 BI1-1700-4	65.74	81.60	14.82	18.40	80.56		
30 BI1-1700-5	118.82	92.68	9.38	7.32	128.20		
31 BI1-1700-6	125.34	92.12	10.72	7.88	136.06		
32 BI1-1700-7	57.81	80.41	14.08	19.59	71.89		
33 BI1-1700-8	35.88	85.41	6.13	14.59	42.01		
34 BI1-1700-9	21.65	89.65	2.50	10.35	24.15		
35 BI1-1700-10	47.31	95.42	2.27	4.58	49.58		
36 BI1-1700-11	13.22	56.98	9.98	43.02	23.20		
37 BI1-1700-12	91.78	94.35	5.50	5.65	97.28		

Flux is in mg/m<sup>2</sup>/day.


Carbonate Flux at Bear Island (BI-1). 1700m. 1984-1985



Sample I.D.	CRTA <1	CRTA % tot.<1	CRTA >1	CRTA % tot.>1	CRTA total	CRTA % total
26 BI1-1700-1	9.80	18.48	1.77	3.34	11.56	21.30
27 BI1-1700-2	22.83	18.82	11.61	9.57	34.44	28.39
28 BI1-1700-3	23.67	21.78	5.66	5.21	29.55	27.19
29 BI1-1700-4	15.80	19.61	4.75	5.90	20.73	25.73
30 BI1-1700-5	21.95	17.12	2.42	1.89	24.52	19.13
31 BI1-1700-6	22.88	16.82	4.38	3.22	27.31	20.07
32 BI1-1700-7	11.63	16.18	3.47	4.83	15.26	21.23
33 BI1-1700-8	7.04	16.76	2.41	5.74	9.62	22.90
34 BI1-1700-9	4.18	17.31	1.25	5.18	5.56	23.02
35 BI1-1700-10	8.74	17.63	0.72	1.45	9.52	19.20
36 BI1-1700-11	2.56	11.03	4.20	18.10	6.81	29.35
37 BI1-1700-12	20.38	20.95	2.00	2.06	22.56	23.19

Flux is in mg/m²/day.


Noncombustible Flux at Bear Island 1, 1700m, 1984-85



Sample ID#	NONC <1	NONC % tot.<1	NONC >1	NONC % tot.>1	NONC total	NONC % total
26 BI1-1700-1*	22.26	41.98	4.25	8.01	26.52	50.01
27 BI1-1700-2*	59.87	49.36	0.56	0.46	60.43	49.82
28 BI1-1700-3*	54.04	49.73	2.59	2.38	56.63	52.11
29 BI1-1700-4*	39.17	48.62	0.39	0.48	39.56	49.11
30 BI1-1700-5*	80.57	62.85	1.55	1.21	82.11	64.05
31 BI1-1700-6*	88.35	64.93	0.24	0.18	88.59	65.11
32 BI1-1700-7*	39.29	54.65	2.14	2.98	41.43	57.63
33 BI1-1700-8*	23.82	56.70	0.46	1.09	24.28	57.80
34 BI1-1700-9*	14.51	60.08	0.13	0.54	14.63	60.58
35 BI1-1700-10*	32.45	65.45	0.62	1.25	33.08	66.72
36 BI1-1700-11*	7.86	33.88	0.58	2.50	8.44	36.38
37 BI1-1700-12*	59.40	61.06	1.21	1.24	60.61	62.30

Flux is in mg/m<sup>2</sup>/day.


**Combustible Flux at Bear Island (BI-1), 1700m, 1984-85**



Sample ID#	COMB <1	COMB % tot.<1	COMB >1	COMB % tot.>1	COMB TOTAL	COMB % total
26 BI1-1700-1*	10.03	18.91	4.60	8.67	14.64	27.81
27 BI1-1700-2*	17.33	14.29	9.00	7.42	26.33	21.71
28 BI1-1700-3*	11.29	10.39	10.62	9.77	21.91	20.16
29 BI1-1700-4*	10.03	12.45	9.67	12.00	19.71	24.47
30 BI1-1700-5*	15.48	12.07	5.42	4.23	20.90	16.30
31 BI1-1700-6*	13.78	10.13	6.10	4.48	19.87	14.60
32 BI1-1700-7*	6.08	8.46	8.47	11.78	14.55	20.24
33 BI1-1700-8*	4.14	9.85	3.27	7.78	7.41	17.64
34 BI1-1700-9*	2.31	9.57	1.12	4.64	3.43	14.20
35 BI1-1700-10*	5.80	11.70	0.93	1.88	6.73	13.57
36 BI1-1700-11*	2.58	11.12	5.20	22.41	7.78	33.53
37 BI1-1700-12*	11.22	11.53	2.29	2.35	13.51	13.69

Flux is in mg/m<sup>2</sup>/day.

Biogenic Silica Flux at Bear Island 1, 1700m, 1984-85




Sample ID#	OPAL <1	OPAL<1 %Ncmb.	OPAL >1	OPAL>1 %Ncmb.	OPAL total	OPAL %Ncmb.	OPAL %total
26 BI1-1700-1	6.51	24.56	0.959340	3.62	7.47	28.17	14.09
27 BI1-1700-2	11.62	19.22	0.061184	0.10	11.68	19.32	9.63
28 BI1-1700-3	9.41	16.62	0.520868	0.92	9.93	17.54	9.14
29 BI1-1700-4	4.94	12.49	0.123738	0.31	5.07	12.81	6.29
30 BI1-1700-5	7.12	8.67	0.191400	0.23	7.31	8.90	5.70
31 BI1-1700-6	5.75	6.50	0.065302	0.07	5.82	6.57	4.28
32 BI1-1700-7	2.69	6.50	0.595221	1.44	3.29	7.93	4.57
33 BI1-1700-8	1.53	6.28	0.119040	0.49	1.64	6.77	3.91
34 BI1-1700-9	1.04	7.08	0.020000	0.14	1.06	7.21	4.37
35 BI1-1700-10	2.56	7.74	0.136090	0.41	2.70	8.15	5.44
36 BI1-1700-11	0.90	10.66	0.115022	1.36	1.01	12.02	4.37
37 BI1-1700-12	7.27	12.00	0.186550	0.31	7.46	12.31	7.67

Flux is in mg/m<sup>2</sup>/day.

%Ncmb. is "% noncombustible flux".

Lithogenic Flux at Bear Island 1, 1700m, 1984-85




Sample ID#	LITH <1	LITH<1 %Ncmb.	LITH >1	LITH>1 %Ncmb.	LITH total	LITH %Ncmb.
26 BI1-1700-1	15.75	59.38	3.29	12.41	19.04	71.79
27 BI1-1700-2	48.25	79.85	0.50	0.83	48.75	80.68
28 BI1-1700-3	44.63	78.81	2.07	3.65	46.70	82.46
29 BI1-1700-4	34.23	86.52	0.27	0.67	34.49	87.19
30 BI1-1700-5	73.45	89.45	1.36	1.65	74.81	91.11
31 BI1-1700-6	82.60	93.23	0.17	0.20	82.77	93.43
32 BI1-1700-7	36.60	88.34	1.54	3.73	38.14	92.07
33 BI1-1700-8	22.29	91.82	0.34	1.40	22.64	93.23
34 BI1-1700-9	13.47	92.10	0.11	0.75	13.58	92.86
35 BI1-1700-10	29.89	90.35	0.48	1.46	30.37	91.82
36 BI1-1700-11	6.96	82.47	0.46	5.51	7.43	87.98
37 BI1-1700-12	52.13	86.01	1.02	1.69	53.15	87.69

Flux is in mg/m<sup>2</sup>/day.

%Ncmb. is "% of noncombustible flux".

Carbon Flux at Bear Island 1, 1700m, 1984-85




Sample I.D.	CRNC <1	CRNC<1 %cmbf.	CRNC >1	CRNC>1 %cmbf.	CRNC total	CRNCtot. %cmbf.
26 BI1-1700-1*	4.65	31.76	2.54	17.35	7.19	49.11
27 BI1-1700-2*	7.66	29.09	5.06	19.20	12.72	43.29
28 BI1-1700-3*	12.31	56.18	6.35	28.98	18.66	85.17
29 BI1-1700-4*	4.72	23.96	5.82	29.55	10.55	53.51
30 BI1-1700-5*	7.04	33.68	3.07	14.69	10.11	48.37
31 BI1-1700-6*	5.50	27.67	3.61	18.15	9.11	45.82
32 BI1-1700-7*	2.48	17.04	4.97	34.16	7.45	51.20
33 BI1-1700-8*	1.83	24.68	1.96	26.46	3.79	51.15
34 BI1-1700-9*	0.88	25.66	0.59	17.20	1.47	42.86
35 BI1-1700-10*	2.54	37.71	0.47	7.02	3.01	44.73
36 BI1-1700-11*	1.23	15.81	2.88	37.02	4.11	52.83
37 BI1-1700-12*	4.57	33.84	1.16	8.61	5.74	42.45

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux"

Nitrogen Flux at Bear Island 1, 1700m, 1984-1985




Sample I.D.	NTGN <1	NTGN<1 %cmbf.	NTGN >1	NTGN>1 %cmbf.	NTGN total	NTGNtot. %cmbf.
26 BI1-1700-1*	0.64	4.37	0.30	2.05	0.94	6.42
27 BI1-1700-2*	1.07	4.05	0.76	2.87	1.82	6.92
28 BI1-1700-3*	0.62	2.83	0.60	2.74	1.22	5.57
29 BI1-1700-4*	0.63	3.21	0.53	2.66	1.16	5.87
30 BI1-1700-5*	0.93	4.45	0.31	1.48	1.24	5.93
31 BI1-1700-6*	0.74	3.71	0.41	2.04	1.14	5.75
32 BI1-1700-7*	0.32	2.20	0.33	2.27	0.65	4.47
33 BI1-1700-8*	0.24	3.29	0.20	2.73	0.45	6.01
34 BI1-1700-9*	0.12	3.50	0.09	2.62	0.21	6.12
35 BI1-1700-10*	0.40	5.95	0.08	1.21	0.48	7.16
36 BI1-1700-11*	0.21	2.70	0.41	5.27	0.62	7.97
37 BI1-1700-12*	0.71	5.26	0.18	1.35	0.89	6.59

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Hydrogen Flux at Bear Island 1, 1700m, 1984-85



Sample I.D.	HYDC <1	HYDC<1 %cmbf.	HYDC >1	HYDC>1 %cmbf.	HYDC total	HYDCtot. %cmbf.
26 BI1-1700-1*	0.87	5.94	0.40	2.73	1.27	8.67
27 BI1-1700-2*	1.55	5.90	0.80	3.04	2.35	8.94
28 BI1-1700-3*	1.94	8.85	1.06	4.84	3.00	13.69
29 BI1-1700-4*	0.95	4.80	0.93	4.71	1.87	9.50
30 BI1-1700-5*	1.54	7.37	0.51	2.44	2.05	9.81
31 BI1-1700-6*	1.35	6.80	0.56	2.80	1.91	9.60
32 BI1-1700-7*	0.57	3.92	0.82	5.64	1.39	9.55
33 BI1-1700-8*	0.40	5.38	0.33	4.43	0.73	9.81
34 BI1-1700-9*	0.21	6.12	0.10	2.92	0.31	9.04
35 BI1-1700-10*	0.56	8.35	0.08	1.20	0.64	9.55
36 BI1-1700-11*	0.23	2.96	0.48	6.17	0.71	9.13
37 BI1-1700-12*	0.86	6.38	0.20	1.47	1.06	7.95

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux"

NA-1  
 AEGIR RIDGE  
 65°31'N, 00°64'E  
 Trap depth: 2,630m Water depth: 3,058m


Annual Fluxes ( $\text{g/m}^2/\text{yr}$ ):

Total.....	17.36
Carbonate.....	9.18
Noncombustible.....	5.94
Combustible.....	2.31
Lithogenic.....	4.26
Biogenic Opal.....	1.68
Organic C.....	0.59
N .....	0.08

PARFLUX Mark 6-13

Sample ID	Opening Date	Closing Date	Span	Mid. Date
57 NA1-3058-1	21-AUG-85	13-SEP-85	23	02-SEP-85
58 NA1-3058-2	13-SEP-85	06-OCT-85	23	25-SEP-85
59 NA1-3058-3	06-OCT-85	29-OCT-85	23	18-OCT-85
60 NA1-3058-4	29-OCT-85	21-NOV-85	23	10-NOV-85
61 NA1-3058-5	21-NOV-85	14-DEC-85	23	03-DEC-85
62 NA1-3058-6	14-DEC-85	06-JAN-86	23	26-DEC-85
63 NA1-3058-7	06-JAN-86	29-JAN-86	23	18-JAN-86
64 NA1-3058-8	29-JAN-86	21-FEB-86	23	10-FEB-86
65 NA1-3058-9	21-FEB-86	16-MAR-86	23	05-MAR-86
66 NA1-3058-10	16-MAR-86	08-APR-86	23	28-MAR-86
67 NA1-3058-11	08-APR-86	01-MAY-86	23	20-APR-86
68 NA1-3058-12	01-MAY-86	24-MAY-86	23	13-MAY-86
69 NA1-3058-13	24-MAY-86	16-JUN-86	23	05-JUN-86


Total Flux at Aegir Ridge (NA1), 3058m, 1985-1986



Sample I.D.	TTLF <1	<1 % of total	TTLF >1	>1 % of total	TTLF total
<hr/>					
57 NA1-3058-1	47.97	89.03	5.91	10.97	53.89
58 NA1-3058-2	62.76	91.27	6.01	8.73	68.76
59 NA1-3058-3	62.53	90.83	6.32	9.17	68.84
60 NA1-3058-4	63.93	94.15	3.97	5.85	67.90
61 NA1-3058-5	89.06	97.02	2.73	2.98	91.79
62 NA1-3058-6	52.31	95.90	2.23	4.10	54.54
63 NA1-3058-7	45.37	99.78	0.10	0.22	45.47
64 NA1-3058-8	27.87	97.70	0.65	2.30	28.52
65 NA1-3058-9	21.54	99.56	0.10	0.44	21.63
66 NA1-3058-10	6.89	98.73	0.09	1.27	6.98
67 NA1-3058-11	5.44	85.35	0.93	14.65	6.38
68 NA1-3058-12	38.72	89.64	4.48	10.36	43.20
69 NA1-3058-13	53.18	89.73	6.09	10.27	59.27

Flux is in mg/m<sup>2</sup>/day.


Carbonate Flux at Aegir Ridge 1 (NA-1), 3058 m, 1985-1986



Sample I.D.	CRTA	CRTA %	CRTA	CRTA %	CRTA	CRTA %
	<1	tot.<1	>1	tot.>1	total	total
57 NA1-3058-1	21.54	39.87	1.55	2.88	23.09	42.85
58 NA1-3058-2	35.54	51.86	7.37	10.72	42.91	62.40
59 NA1-3058-3	36.44	52.83	3.37	4.90	39.81	57.93
60 NA1-3058-4	35.29	51.97	1.80	2.69	37.09	54.82
61 NA1-3058-5	46.88	51.07	1.09	1.19	47.97	52.16
62 NA1-3058-6	28.00	47.87	0.91	1.67	28.91	49.34
63 NA1-3058-7	21.76	47.86	0.02	0.04	21.78	47.80
64 NA1-3058-8	15.31	53.68	0.34	1.19	16.65	54.87
65 NA1-3058-9	10.52	49.09	0.00	0.00	10.52	49.09
66 NA1-3058-10	4.70	67.36	0.00	0.00	4.70	67.36
67 NA1-3058-11	3.26	51.12	0.68	10.68	3.94	61.79
68 NA1-3058-12	20.23	46.93	1.88	4.35	22.11	51.18
69 NA1-3058-13	28.81	46.61	1.95	3.29	30.76	51.30

Flux is in mg/m<sup>2</sup>/day


Noncombustible Flux at Aegir Ridge (NA-1), 3058m, 1985-1986



Sample ID#	NONC <1	NONC % tot,<1	NONC >1	NONC % tot,>1	NONC total	NONC % total
57 NA1-3058-1	20.46	37.96	0.24	0.45	20.70	38.42
58 NA1-3058-2	5.99	8.71	0.25	0.37	6.24	9.08
59 NA1-3058-3	21.05	30.58	0.73	1.06	21.79	31.65
60 NA1-3058-4	23.74	34.96	0.63	0.93	24.38	35.90
61 NA1-3058-5	35.60	38.78	0.11	0.12	35.71	38.91
62 NA1-3058-6	22.42	41.10	0.13	0.24	22.55	41.33
63 NA1-3058-7	19.74	43.41	0.00	0.01	19.74	43.42
64 NA1-3058-8	11.34	39.77	0.03	0.10	11.37	39.87
65 NA1-3058-9	9.04	41.80	0.02	0.07	9.06	41.87
66 NA1-3058-10	1.84	26.34	0.04	0.50	1.87	26.84
67 NA1-3058-11	1.91	29.91	0.01	0.23	1.92	30.14
68 NA1-3058-12	14.81	34.28	0.92	2.13	15.73	35.41
69 NA1-3058-13	19.06	32.15	1.60	2.70	20.66	34.86

Flux is in mg/m<sup>2</sup>/day.


Combustible Flux at Aegir Ridge (NA-1), 3058m, 1985-86



Sample ID#	COMB <1	COMB % tot.<1	COMB >1	COMB % tot.>1	COMB TOTAL	COMB % total	
						COMB <1	COMB % tot.<1
57 NA1-3058-1	5.97	11.08	4.12	7.64	10.09	18.72	
58 NA1-3058-2	21.23	30.88	0.39	0.56	21.62	31.44	
59 NA1-3058-3	5.04	7.31	4.22	6.13	9.25	13.44	
60 NA1-3058-4	4.90	7.21	1.54	2.26	6.43	9.47	
61 NA1-3058-5	6.58	7.17	1.53	1.66	8.11	8.83	
62 NA1-3058-6	3.89	7.14	1.19	2.18	5.08	9.32	
63 NA1-3058-7	3.87	8.52	0.08	0.17	3.95	8.68	
64 NA1-3058-8	1.22	4.27	0.48	1.68	1.70	5.96	
65 NA1-3058-9	1.88	8.68	0.08	0.39	1.96	9.07	
66 NA1-3058-10	0.35	5.05	0.05	0.79	0.41	5.84	
67 NA1-3058-11	0.27	4.27	0.24	3.69	0.51	7.97	
68 NA1-3058-12	3.68	8.52	1.68	3.89	5.36	12.40	
69 NA1-3058-13	5.31	8.96	2.54	4.28	7.85	13.25	

Flux is in mg/m<sup>2</sup>/day.

Biogenic Silica Flux at Aegir Ridge (NA-1), 3058m, 1985-86




Sample ID#	OPAL <1	OPAL % Ncf.<1	OPAL % tot.<1
57 NA1-3058-1	5.88	28.40	10.91
58 NA1-3058-2	5.56	89.07	8.09
59 NA1-3058-3	5.55	25.47	8.06
60 NA1-3058-4	7.17	29.41	10.56
61 NA1-3058-5	9.67	27.08	10.53
62 NA1-3058-6	4.95	21.96	9.08
63 NA1-3058-7	5.12	25.93	11.26
64 NA1-3058-8	2.55	22.42	8.94
65 NA1-3058-9	3.59	39.63	16.59
66 NA1-3058-10	0.58	30.98	8.31
67 NA1-3058-11	0.58	30.18	9.10
68 NA1-3058-12	3.76	23.90	8.70
69 NA1-3058-13	4.87	23.57	8.22

Flux is in  $\text{mg}/\text{m}^2/\text{day}$ .

Not enough sample in >1 mm fraction to analyze for Opal.

"%Ncf." = "% of noncombustible flux".

Lithogenic Flux at Aegir Ridge (NA-1), 3058 m., 1985-6




Sample I.D.	LITH <1	LITH<1 %Ncmb.	LITH<1 %tot.
57 NA1-3058-1	14.58	70.42	27.05
58 NA1-3058-2	0.43	6.86	0.62
59 NA1-3058-3	15.50	71.16	22.52
60 NA1-3058-4	16.57	67.99	24.41
61 NA1-3058-5	25.93	72.61	28.25
62 NA1-3058-6	17.47	77.47	32.02
63 NA1-3058-7	14.62	74.05	32.15
64 NA1-3058-8	8.79	77.31	30.82
65 NA1-3058-9	5.45	60.19	25.20
66 NA1-3058-10	1.26	67.15	18.02
67 NA1-3058-11	1.33	69.07	20.82
68 NA1-3058-12	11.05	70.25	25.58
69 NA1-3058-13	14.19	68.67	23.94

Flux is in mg/m<sup>2</sup>/day.

%Ncmb. = "% of noncombustible flux".

Not enough >1 mm fraction to do analysis.

Carbon Flux at Aegir Ridge (NA-1), 3058m, 1985-86




Sample I.D.	CRNC <1	CRNC<1 %cmbf.	CRNC >1	CRNC>1 %cmbf.	CRNC total	CRNCtot. %cmbf.
57 NA1-3058-1	2.55	25.25	2.16	21.36	4.70	46.62
58 NA1-3058-2	3.10	14.33	0.26	1.19	3.36	15.52
59 NA1-3058-3	2.43	26.28	1.54	16.62	3.97	42.90
60 NA1-3058-4	1.50	23.37	0.58	8.97	2.08	32.34
61 NA1-3058-5	3.26	40.17	0.80	9.90	4.06	50.06
62 NA1-3058-6	1.34	26.29	0.52	10.25	1.86	36.54
63 NA1-3058-7	1.10	27.99	0.04	0.89	1.14	28.88
64 NA1-3058-8	0.95	56.04	0.22	13.16	1.18	69.21
65 NA1-3058-9	0.84	42.58	0.04	2.23	0.88	44.81
66 NA1-3058-10	0.14	33.53	0.04	9.68	0.18	43.21
67 NA1-3058-11	0.15	29.44	0.11	21.58	0.26	51.02
68 NA1-3058-12	2.09	39.02	1.14	21.27	3.23	60.30
69 NA1-3058-13	1.55	19.77	1.79	22.74	3.34	42.51

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux"

Nitrogen Flux at Aegir Ridge (NA-1), 3058m, 1985-86




Sample I.D.	NTGN <1	NTGN<1 %cmbf.	NTGN >1	NTGN>1 %cmbf.	NTGN total	NTGNtot. %cmbf.
57 NA1-3058-1	0.33	3.25	0.27	2.71	0.60	5.95
58 NA1-3058-2	0.42	1.93	0.03	0.16	0.45	2.08
59 NA1-3058-3	0.31	3.33	0.21	2.32	0.52	5.64
60 NA1-3058-4	0.18	2.80	0.09	1.37	0.27	4.17
61 NA1-3058-5	0.47	5.78	0.11	1.30	0.57	7.07
62 NA1-3058-6	0.18	3.62	0.06	1.18	0.24	4.81
63 NA1-3058-7	0.14	3.65	0.00	0.09	0.15	3.74
64 NA1-3058-8	0.11	6.36	0.02	1.38	0.13	7.74
65 NA1-3058-9	0.10	5.12	0.00	0.23	0.11	5.36
66 NA1-3058-10	0.02	3.98	0.00	1.02	0.02	4.99
67 NA1-3058-11	0.02	4.12	0.01	2.27	0.03	6.39
68 NA1-3058-12	0.29	5.45	0.12	2.24	0.41	7.69
69 NA1-3058-13	0.20	2.55	0.12	1.51	0.32	4.05

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux"

Hydrogen Flux at Aegir Ridge (NA-1), 3058m, 1985-86



Sample I.D.	HYDC <1	HYDC<1 %cmbf.	HYDC >1	HYDC>1 %cmbf.	HYDC total	HYDCtot. %cmbf.
57 NA1-3058-1	0.44	4.40	0.30	2.94	0.74	7.34
58 NA1-3058-2	0.44	2.05	0.03	0.16	0.48	2.21
59 NA1-3058-3	0.36	3.92	0.19	2.09	0.56	5.01
60 NA1-3058-4	0.41	6.37	0.07	1.08	0.48	7.44
61 NA1-3058-5	0.56	6.92	0.11	1.38	0.67	8.30
62 NA1-3058-6	0.21	4.19	0.08	1.51	0.29	5.70
63 NA1-3058-7	0.26	6.64	0.00	0.12	0.27	6.76
64 NA1-3058-8	0.13	7.54	0.03	1.82	0.16	9.36
65 NA1-3058-9	0.14	7.24	0.01	0.31	0.15	7.54
66 NA1-3058-10	0.03	6.50	0.01	1.34	0.03	7.84
67 NA1-3058-11	0.03	5.45	0.02	2.99	0.04	8.44
68 NA1-3058-12	0.28	5.21	0.16	2.94	0.44	8.15
69 NA1-3058-13	0.22	2.82	0.23	2.95	0.45	5.78

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux"

NB-1  
 EAST OF JAN MAYEN  
 (NB-1) 70°00'N, 01°58'W  
 Trap depth: 2,749m Water depth: 2,773m


Annual Fluxes ( $\text{g/m}^2/\text{yr}$ ):

Total.....	16.78
Carbonate.....	8.93
Noncombustible.....	6.24
Lithogenic.....	4.65
Combustible.....	1.90
Biogenic Opal.....	1.44
Organic C.....	0.53
N .....	0.08

PARFLUX Mark 5-25

Sample ID	Opening Date	Closing Date	Span	Mid. Date
70 NB1-2815-1	20-AUG-85	01-SEP-85	12	26-AUG-85
71 NB1-2815-2	01-SEP-85	13-SEP-85	12	07-SEP-85
72 NB1-2815-3	13-SEP-85	25-SEP-85	12	19-SEP-85
73 NB1-2815-4	25-SEP-85	07-OCT-85	12	01-OCT-85
74 NB1-2815-5	07-OCT-85	19-OCT-85	12	13-OCT-85
75 NB1-2815-6	19-OCT-85	31-OCT-85	12	25-OCT-85
76 NB1-2815-7	31-OCT-85	12-NOV-85	12	06-NOV-85
77 NB1-2815-8	12-NOV-85	24-NOV-85	12	18-NOV-85
78 NB1-2815-9	24-NOV-85	06-DEC-85	12	30-NOV-85
79 NB1-2815-10	06-DEC-85	18-DEC-85	12	12-DEC-85
80 NB1-2815-11	18-DEC-85	30-DEC-85	12	24-DEC-85
81 NB1-2815-12	30-DEC-85	30-DEC-85	12	05-JAN-86
82 NB1-2815-13	30-DEC-85	11-JAN-86	12	05-JAN-86
83 NB1-2815-14	11-JAN-86	23-JAN-86	12	17-JAN-86
84 NB1-2815-15	23-JAN-86	04-FEB-86	12	29-JAN-86
85 NB1-2815-16	04-FEB-86	16-FEB-86	12	10-FEB-86
86 NB1-2815-17	16-FEB-86	28-FEB-86	12	22-FEB-86
87 NB1-2815-18	28-FEB-86	12-MAR-86		06-MAR-86
88 NB1-2815-19	12-MAR-86	24-MAR-86		18-MAR-86


Total Flux at Jan Mayen (NB1), 2815m, 1985-1986



Sample I.D.	TTLF <1	<1 % of total		TTLF >1	>1 % of total		TTLF total
		<1	>1		<1	>1	
70 NB1-2815-1	72.19	91.14		7.01	8.86		79.21
71 NB1-2815-2	91.93	89.06		11.29	10.94		103.22
72 NB1-2815-3	65.98	89.10		8.07	10.90		74.05
73 NB1-2815-4	71.94	93.86		4.71	6.14		76.65
74 NB1-2815-5	52.90	97.22		1.51	2.78		54.41
75 NB1-2815-6	45.13	97.55		1.13	2.45		46.26
76 NB1-2815-7	48.36	76.51		14.85	23.49		63.22
77 NB1-2815-8	52.95	98.41		0.85	1.59		53.80
78 NB1-2815-9	40.77	97.33		1.12	2.67		41.89
79 NB1-2815-10	36.70	97.73		0.85	2.27		37.55
80 NB1-2815-11	46.52	98.56		0.68	1.44		47.20
81 NB1-2815-12	21.13	97.30		0.59	2.70		21.72
82 NB1-2815-13	26.87	98.79		0.33	1.21		27.20
83 NB1-2815-14	38.43	99.13		0.34	0.87		38.76
84 NB1-2815-15	4.70	98.10		0.09	1.90		4.79
85 NB1-2815-16	2.72	97.75		0.06	2.25		2.79
86 NB1-2815-17	8.03	90.12		0.88	9.88		8.91

Flux is in  $\text{mg}/\text{m}^2/\text{day}$ .


Carbonate Flux at Jan Mayen (NB-1), 2815m, 1985-1986



Sample I.D.	CRTA <1	CRTA % tot:<1	CRTA >1	CRTA % tot.>1	CRTA total	CRTA % total
70 NB1-2815-1	33.84	42.72	6.66	8.41	40.50	51.13
71 NB1-2815-2	47.19	45.72	10.35	10.03	57.54	55.74
72 NB1-2815-3	44.21	59.70	7.59	10.25	51.80	69.95
73 NB1-2815-4	36.25	47.29	3.49	4.55	39.74	51.85
74 NB1-2815-5	25.33	46.55	0.89	1.64	26.22	48.19
75 NB1-2815-6	24.04	51.96	0.32	0.69	24.36	52.66
76 NB1-2815-7	24.50	38.76	5.67	8.97	30.17	47.72
77 NB1-2815-8	28.01	52.06	0.13	0.24	28.14	52.30
78 NB1-2815-9	22.43	53.54	0.15	0.36	22.58	53.90
79 NB1-2815-10	20.74	55.24	0.07	0.19	20.81	55.42
80 NB1-2815-11	24.52	51.95	0.07	0.15	24.59	52.10
81 NB1-2815-12	11.80	54.34	0.12	0.00	11.80	54.34
82 NB1-2815-13	14.59	53.64	0.10	0.37	14.69	54.00
83 NB1-2815-14	19.14	49.38	0.05	0.13	19.19	49.51
84 NB1-2815-15	1.55	32.38	0.06	1.25	1.61	33.64
85 NB1-2815-16	1.23	44.14	0.04	1.44	1.27	45.58
86 NB1-2815-17	0.80	9.01	0.14	1.57	0.94	10.54

Flux is in mg/m<sup>2</sup>/day.


Noncombustible Flux at Jan Mayen (NB-1), 2815m, 1985-1986



Sample ID#	NONC <1	NONC % tot.<1	NONC >1	NONC % tot.>1	NONC total	NONC % total
70 NB1-2815-1	30.37	38.35	0.14	0.17	30.51	38.52
71 NB1-2815-2	35.84	34.72	1.42	1.37	37.25	36.09
72 NB1-2815-3	17.49	23.61	1.60	2.16	19.09	25.77
73 NB1-2815-4	29.94	39.07	0.81	1.06	30.75	40.12
74 NB1-2815-5	22.22	40.84	0.18	0.34	22.40	41.18
75 NB1-2815-6	17.48	37.79	0.03	0.05	17.51	37.85
76 NB1-2815-7	19.30	30.53	2.34	3.71	21.65	34.24
77 NB1-2815-8	21.70	40.33	0.07	0.13	21.77	40.46
78 NB1-2815-9	15.42	36.82	0.24	0.57	15.66	37.38
79 NB1-2815-10	13.68	36.43	0.09	0.24	13.77	36.67
80 NB1-2815-11	18.57	39.34	0.05	0.10	18.61	39.44
81 NB1-2815-12	8.05	37.08	0.05	0.23	8.10	37.31
82 NB1-2815-13	10.13	37.24	0.01	0.04	10.14	37.28
83 NB1-2815-14	16.17	41.70	0.02	0.06	16.19	41.76
84 NB1-2815-15	2.31	48.17	0.00	0.02	2.31	48.19
85 NB1-2815-16	1.49	53.48	0.03	1.21	1.52	54.63
86 NB1-2815-17	3.43	38.43	0.04	0.49	3.47	38.92

Flux is in mg/m<sup>2</sup>/day.


Combustible Flux at Jan Mayen (NB-1), 2815m, 1985-86



Sample ID#	COMB <1	COMB % tot.<1	COMB >1	COMB % tot.>1	COMB TOTAL	COMB % total
70 NB1-2815-1	7.98	10.07	0.21	0.27	8.19	10.34
71 NB1-2815-2	8.90	8.63	2.50	2.43	11.41	11.05
72 NB1-2815-3	4.18	5.65	3.10	4.19	7.28	9.84
73 NB1-2815-4	5.75	7.50	2.10	2.74	7.85	10.24
74 NB1-2815-5	5.35	9.83	0.24	0.44	5.59	10.27
75 NB1-2815-6	3.61	7.80	0.19	0.42	3.80	8.22
76 NB1-2815-7	4.56	7.21	12.49	19.75	17.04	26.96
77 NB1-2815-8	3.24	6.03	0.44	0.82	3.68	6.84
78 NB1-2815-9	2.92	6.96	0.88	2.11	3.80	9.07
79 NB1-2815-10	2.28	6.08	0.76	2.02	3.04	8.10
80 NB1-2815-11	3.43	7.27	0.56	1.20	4.00	8.47
81 NB1-2815-12	1.28	5.89	0.54	2.48	1.82	8.37
82 NB1-2815-13	2.15	7.90	0.22	0.80	2.37	8.71
83 NB1-2815-14	3.12	8.06	0.27	0.69	3.39	8.75
84 NB1-2815-15	0.84	17.64	0.01	0.19	0.85	17.83
85 NB1-2815-16	0.00	0.00	0.01	0.22	0.01	0.22
86 NB1-2815-17	3.80	42.64	0.70	7.81	4.50	50.45

Flux is in mg/m<sup>2</sup>/day.

Biogenic Silica Flux at Jan Mayen (NB-1), 2815m, 1986




Sample ID#	OPAL <1	OPAL % Ncf. <1	OPAL % tot. <1
70 NB1-2815-1	9.44	30.94	11.92
71 NB1-2815-2	12.99	34.87	12.58
72 NB1-2815-3	3.91	20.49	5.28
73 NB1-2815-4	8.17	26.57	10.66
74 NB1-2815-5	4.70	20.98	8.64
75 NB1-2815-6	4.30	24.56	9.29
76 NB1-2815-7	4.08	18.85	6.45
77 NB1-2815-8	3.69	16.95	6.86
78 NB1-2815-9	3.03	19.35	7.23
79 NB1-2815-10	2.28	16.56	6.07
80 NB1-2815-11	3.23	17.35	6.84
81 NB1-2815-12	1.22	15.06	5.62
82 NB1-2815-13	1.79	17.67	6.58
83 NB1-2815-14	2.80	17.32	7.22
84 NB1-2815-15	0.46	19.91	9.60
85 NB1-2815-16	0.22	14.77	7.89
86 NB1-2815-17	0.92	26.82	10.33

Flux is in mg/m<sup>2</sup>/day.

Not enough sample in >1 mm fraction to analyze for Opal.

"%Ncf." = "% of noncombustible flux".

Lithogenic Flux at Jan Mayen (NB-1), 2815, 1986




SAMPLE I.D.	LITH <1	LITH<1 %Ncmb.	LITH<1 %total
NB1-2815-1	20.93	68.61	26.43
NB1-2815-2	22.85	61.33	22.13
NB1-2815-3	13.58	71.14	18.33
NB1-2815-4	21.77	70.80	28.41
NB1-2815-5	17.52	78.21	32.20
NB1-2815-6	13.18	75.30	28.50
NB1-2815-7	15.22	70.33	24.08
NB1-2815-8	18.01	82.72	33.47
NB1-2815-9	12.39	79.14	29.58
NB1-2815-10	11.40	82.79	30.36
NB1-2815-11	15.34	82.40	32.50
NB1-2815-12	6.83	84.32	31.46
NB1-2815-13	8.34	82.24	30.66
NB1-2815-14	13.37	82.56	34.48
NB1-2815-15	1.85	80.02	38.56
NB1-2815-16	1.27	83.35	45.58
NB1-2815-17	2.51	72.22	28.11

Flux is in mg/m<sup>2</sup>/day.

Insufficient material to analyze >1mm fraction.

Carbon Flux at Jan Mayen (NB-1), 2815m, 1985-86




Sample I.D.	CRNC <1	CRNC<1 %cmbf.
70 NB1-2815-1	3.41	41.63
71 NB1-2815-2	4.13	36.16
72 NB1-2815-3	1.69	23.20
73 NB1-2815-4	1.91	24.29
74 NB1-2815-5	1.41	25.17
75 NB1-2815-6	1.70	44.72
76 NB1-2815-7	2.35	13.77
77 NB1-2815-8	1.77	47.97
78 NB1-2815-9	1.41	37.03
79 NB1-2815-10	1.12	36.93
80 NB1-2815-11	1.56	39.03
81 NB1-2815-12	0.72	39.43
82 NB1-2815-13	0.05	2.23
83 NB1-2815-14	0.04	1.31
84 NB1-2815-15	0.02	1.99

Flux is in  $\text{mg}/\text{m}^2/\text{day}$ .

"%cmbf" = "% of combustible flux".

Not enough >1 mm fraction to do analysis.

Nitrogen Flux at Jan Mayen (NB-1), 2815m, 1985-86




Sample I.D.	NTGN <1	NTGN<1 %cmbf.
70 NB1-2815-1	0.48	5.90
71 NB1-2815-2	0.60	5.29
72 NB1-2815-3	0.24	3.24
73 NB1-2815-4	0.25	3.23
74 NB1-2815-5	0.22	3.85
75 NB1-2815-6	0.25	6.55
76 NB1-2815-7	0.31	1.82
77 NB1-2815-8	0.24	6.64
78 NB1-2815-9	0.20	5.21
79 NB1-2815-10	0.15	5.04
80 NB1-2815-11	0.23	5.67
81 NB1-2815-12	0.09	5.08
82 NB1-2815-13	0.01	0.36
83 NB1-2815-14	0.01	0.17
84 NB1-2815-15	0.00	0.00

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Not enough >1 mm fraction to do analysis.

Hydrogen Flux at Jan Mayen (NB-1), 2815m, 1985-86



Sample I.D.	HYDC <1	HYDC<1 %cmbf.
70 NB1-2815-1	0.51	6.28
71 NB1-2815-2	0.76	6.63
72 NB1-2815-3	0.34	4.67
73 NB1-2815-4	0.45	5.69
74 NB1-2815-5	0.33	5.92
75 NB1-2815-6	0.25	6.66
76 NB1-2815-7	0.36	2.11
77 NB1-2815-8	0.32	8.74
78 NB1-2815-9	0.24	6.42
79 NB1-2815-10	0.19	6.14
80 NB1-2815-11	0.26	6.55
81 NB1-2815-12	0.13	6.88
82 NB1-2815-13	0.03	1.35
83 NB1-2815-14	0.02	0.63

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Not enough >1 mm fraction to do analysis.

EAST GREENLAND/FRAM STRAIT AREA


FS-1  
CENTRAL FRAM STRAIT  
78°52' N, 01°22'E  
Trap depth: 2,440m Water depth: 2,527m

Annual Fluxes ( $\text{g/m}^2/\text{yr}$ ):  
Total.....6.61  
Carbonate.....1.40  
Noncombustible.....4.26  
Combustible.....0.92  
Lithogenic.....4.00  
Biogenic Opal.....0.60  
Organic C.....0.41  
N.....0.06

PARFLUX Mark S-13

Sample ID	Opening Date	Closing Date	Span	Mid. Date
13 FS1-2000-1	20-AUG-84	17-SEP-84	27.5	03-SEP-84
14 FS1-2000-2	17-SEP-84	15-OCT-84	27.5	01-OCT-84
15 FS1-2000-3	15-OCT-84	11-NOV-84	27.5	29-OCT-84
16 FS1-2000-4	11-NOV-84	09-DEC-84	27.5	25-NOV-84
17 FS1-2000-5	09-DEC-84	07-JAN-85	27.5	24-DEC-84
18 FS1-2000-6	07-JAN-85	03-FEB-85	27.5	21-JAN-85
19 FS1-2000-7	03-FEB-85	03-MAR-85	27.5	17-FEB-85
20 FS1-2000-8	03-MAR-85	30-MAR-85	27.5	17-MAR-85
21 FS1-2000-9	30-MAR-85	27-APR-85	27.5	13-APR-85
22 FS1-2000-10	27-APR-85	25-MAY-85	27.5	11-MAY-85
23 FS1-2000-11	25-MAY-85	21-JUN-85	27.5	08-JUN-85
24 FS1-2000-12	21-JUN-85	19-JUL-85	27.5	05-JUL-85
25 FS1-2000-13	19-JUL-85	15-AUG-85	27.5	02-AUG-85

Total Flux at Fram Strait (FS-1), 2000m, 1984-1985




FRAM STRAIT 1 POISONED WITH HG CL2 359 DAYS  
 Mark 5 trap open from AUGUST 20 1984 to AUGUST 15 1985 at 2000 meters.

TOTAL FLUX (mg / m<sup>2</sup> / day)

Ttl is total Flux in all size classes

Cup #	< 63um		63um - 1		> 1mm		TOTAL	
	% of Ttl	FLUX						
1	57.18	12.30	23.52	5.06	19.29	4.15	100.00	21.51
2	63.02	25.34	17.56	7.06	19.42	7.81	100.00	40.21
3	64.12	29.17	14.99	6.82	20.88	9.50	100.00	45.49
4	50.03	9.05	29.41	5.32	20.56	3.72	100.00	18.09
5	63.83	6.46	17.89	1.81	18.28	1.85	100.00	10.12
6	52.92	3.90	34.60	2.55	12.48	.92	100.00	7.37
7	59.75	2.91	12.73	.62	27.52	1.34	100.00	4.87
8	51.98	3.80	15.18	1.11	32.83	2.40	100.00	7.31
9	24.86	14.00	27.73	15.62	47.41	26.70	100.00	56.32
10	36.46	3.81	45.45	4.75	18.09	1.89	100.00	10.45
11	72.32	4.39	19.60	1.19	8.07	.49	100.00	6.07
12	31.80	5.31	16.11	2.69	52.10	8.70	100.00	16.70
13	31.56	2.13	12.00	.81	56.44	3.81	100.00	6.75


Carbonate Flux at Fram Strait (FS-1), 2000m, 1984-1985



Sample I.D.	CRTA <1	CRTA % tot.<1	CRTA >1	CRTA % tot.>1	CRTA total	CRTA % total
13 FS1-2000-1	3.51	14.52	0.81	3.77	4.32	3.35
14 FS1-2000-2	6.84	17.13	2.30	5.72	9.14	5.76
15 FS1-2000-3	7.72	18.16	2.05	4.51	9.77	4.82
16 FS1-2000-4	3.31	18.33	2.78	15.37	6.09	15.39
17 FS1-2000-5	1.64	18.10	0.54	5.34	2.18	5.96
18 FS1-2000-6	0.78	14.44	0.39	3.85	1.17	7.22
19 FS1-2000-7	0.81	15.52	0.72	9.78	1.53	13.79
20 FS1-2000-8	0.98	13.44	0.98	13.41	1.96	13.44
21 FS1-2000-9	0.71	1.60	1.57	2.79	2.28	3.53
22 FS1-2000-10	0.82	7.27	0.33	3.16	1.15	2.93
23 FS1-2000-11	0.72	11.98	0.10	1.65	0.82	1.66
24 FS1-2000-12	1.21	7.79	1.64	9.80	2.85	10.56
25 FS1-2000-13	0.50	7.76	2.21	32.74	2.71	34.32

Flux is in mg/m<sup>2</sup>/day.


Noncombustible Flux at Fram Strait 1, 2000 m, 1984-85



Sample ID#	NONC <1	NONC % tot.<1	NONC >1	NONC % tot.>1	NONC total	NONC % total
13 FS1-2000-1*	13.22	54.67	2.14	8.85	15.36	63.52
14 FS1-2000-2*	20.86	52.23	2.64	6.61	23.50	58.84
15 FS1-2000-3*	21.77	51.21	4.02	9.46	25.79	60.67
16 FS1-2000-4*	9.16	50.72	0.45	2.49	9.61	53.21
17 FS1-2000-5*	4.56	50.33	0.45	4.97	5.01	55.30
18 FS1-2000-6*	2.87	53.15	0.13	2.41	3.00	55.56
19 FS1-2000-7*	2.06	39.46	0.13	2.49	2.29	43.87
20 FS1-2000-8*	2.75	37.72	0.47	6.45	3.22	44.17
21 FS1-2000-9*	15.74	35.43	24.30	54.69	40.04	90.12
22 FS1-2000-10*	7.67	68.00	1.23	10.90	8.90	78.90
23 FS1-2000-11*	3.67	61.06	0.29	4.83	3.96	65.89
24 FS1-2000-12*	4.09	26.34	4.92	31.68	9.01	58.02
25 FS1-2000-13*	1.35	20.96	0.59	9.16	1.94	30.12

Flux is in  $\text{mg}/\text{m}^2/\text{day}$ .


**Combustible Flux at Fram Strait 1, 2000 m, 1984-85**



Sample ID#	COMB <1	COMB % tot.<1	COMB >1	COMB % tot.>1	COMB TOTAL	COMB % total
13 FS1-2000-1*	3.28	13.56	1.21	5.00	4.49	18.57
14 FS1-2000-2*	4.32	10.82	2.87	7.19	7.19	18.00
15 FS1-2000-3*	3.51	8.26	3.43	8.07	6.94	16.33
16 FS1-2000-4*	1.61	8.91	0.49	2.71	2.10	11.63
17 FS1-2000-5*	0.85	9.38	0.86	9.49	1.71	18.87
18 FS1-2000-6*	0.43	7.96	0.39	7.22	0.82	15.19
19 FS1-2000-7*	0.31	5.94	0.49	9.39	0.80	15.33
20 FS1-2000-8*	0.45	6.17	0.95	13.03	1.40	19.20
21 FS1-2000-9*	0.58	1.31	0.83	1.87	1.41	3.17
22 FS1-2000-10*	0.55	4.88	0.33	2.93	0.88	7.80
23 FS1-2000-11*	0.68	11.31	0.10	1.66	0.78	12.98
24 FS1-2000-12*	0.85	5.47	2.13	13.72	2.98	19.19
25 FS1-2000-13*	0.27	4.19	1.01	15.68	1.28	19.88

Flux is in mg/m<sup>2</sup>/day.

Biogenic Opal Flux at Fram Strait 1, 2000m, 1984-85




Sample ID#	OPAL <1	OPAL<1 %Ncmb.	OPAL<1 %Total
13 FS1-2000-1	2.81	18.28	11.61
14 FS1-2000-2	5.58	23.73	13.96
15 FS1-2000-3	5.41	20.98	12.73
16 FS1-2000-4	1.58	16.46	8.76
17 FS1-2000-5	0.71	14.23	7.87
18 FS1-2000-6	0.52	17.33	9.63
19 FS1-2000-7	0.33	14.47	6.35
20 FS1-2000-8	1.32	40.91	18.07
21 FS1-2000-9	0.78	1.95	1.75
22 FS1-2000-10	0.41	4.63	3.65
23 FS1-2000-11	1.02	25.73	16.96
24 FS1-2000-12	0.53	5.93	3.44
25 FS1-2000-13	0.27	14.14	4.26

Flux is in  $\text{mg}/\text{m}^2/\text{day}$ .

%Ncmb. is "% noncombustible flux".

Not enough >1 mm fraction to do analysis.

Lithogenic Flux at Fram Strait 1, 2000m, 1984-85




Sample ID#	LITH <1	LITH<1 %Ncmb.
13 FS1-2000-1	10.41	67.79
14 FS1-2000-2	15.28	65.03
15 FS1-2000-3	16.36	63.44
16 FS1-2000-4	7.58	78.86
17 FS1-2000-5	3.85	76.79
18 FS1-2000-6	2.35	78.34
19 FS1-2000-7	1.73	75.49
20 FS1-2000-8	1.43	44.49
21 FS1-2000-9	14.96	37.36
22 FS1-2000-10	7.26	81.55
23 FS1-2000-11	2.65	66.94
24 FS1-2000-12	3.56	39.46
25 FS1-2000-13	1.08	55.45

Flux is in mg/m<sup>2</sup>/day.

%Ncmb. is "% noncombustible flux".

Not enough >1 mm fraction to do analysis.

Carbon Flux at Fram Strait 1, 2000m, 1984-85




Sample I.D.	CRNC <1	CRNC<1 %cmbf.	CRNC >1	CRNC>1 %cmbf.	CRNC total	CRNCTot. %cmbf.
13 FS1-2000-1*	1.39	30.89	0.58	12.92	1.97	43.81
14 FS1-2000-2*	2.13	29.62	1.41	19.61	3.54	49.23
15 FS1-2000-3*	1.75	25.25	2.04	29.39	3.79	54.65
16 FS1-2000-4*	0.79	37.66	0.26	12.38	1.05	50.04
17 FS1-2000-5*	0.48	27.80	0.36	21.05	0.84	48.85
18 FS1-2000-6*	0.34	41.45	0.19	22.71	0.53	64.16
19 FS1-2000-7*	0.22	27.77	0.17	21.25	0.39	49.02
20 FS1-2000-8*	0.18	12.86	0.51	36.48	0.69	49.29
21 FS1-2000-9*	0.13	9.08	0.51	36.17	0.64	45.25
22 FS1-2000-10*	0.29	32.92	0.14	16.22	0.43	49.14
23 FS1-2000-11*	0.43	55.10	0.06	7.69	0.49	62.79
24 FS1-2000-12*	0.35	11.81	1.15	38.59	1.50	50.40
25 FS1-2000-13*	0.22	17.39	0.26	20.31	0.48	37.70

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Nitrogen Flux at Fram Strait, 2000m, 1984-85




Sample I.D.	NTGN <1	NTGN<1 %cmbf.	NTGN >1	NTGN>1 %cmbf.	NTGN total	NTGNtot. %cmbf.
13 FS1-2000-1*	0.19	4.17	0.08	1.78	0.27	5.95
14 FS1-2000-2*	0.30	4.24	0.15	2.09	0.45	6.33
15 FS1-2000-3*	0.23	3.36	0.17	2.45	0.40	5.81
16 FS1-2000-4*	0.11	5.09	0.04	1.90	0.15	6.99
17 FS1-2000-5*	0.07	3.86	0.03	1.75	0.10	5.61
18 FS1-2000-6*	0.04	5.21	0.02	2.77	0.07	7.98
19 FS1-2000-7*	0.02	3.11	0.01	1.25	0.03	4.36
20 FS1-2000-8*	0.14	9.87	0.06	4.35	0.20	14.22
21 FS1-2000-9*	0.01	0.95	0.03	2.13	0.04	3.08
22 FS1-2000-10*	0.04	4.09	0.01	1.65	0.05	5.74
23 FS1-2000-11*	0.05	6.13	0.01	1.28	0.06	7.41
24 FS1-2000-12*	0.05	1.54	0.12	4.03	0.17	5.57
25 FS1-2000-13*	0.03	2.14	0.03	2.34	0.06	4.48

Flux is in mg/m<sup>2</sup>/day.

"/%cmbf" = "% of combustible flux"

Hydrogen Flux at Fram Strait, 2000m, 1984-1985



Sample I.D.	HYDC <1	HYDC<1 %cmbf.	HYDC >1	HYDC>1 %cmbf.	HYDC total	HYDCtot. %cmbf.
13 FS1-2000-1*	0.27	6.06	0.10	2.23	0.37	8.29
14 FS1-2000-2*	0.40	5.52	0.23	3.20	0.63	8.72
15 FS1-2000-3*	0.45	6.47	0.35	5.04	0.80	11.51
16 FS1-2000-4*	0.17	7.95	0.04	1.90	0.21	9.86
17 FS1-2000-5*	0.11	6.28	0.06	3.52	0.17	9.80
18 FS1-2000-6*	0.08	9.64	0.03	3.23	0.11	12.87
19 FS1-2000-7*	0.04	5.54	0.03	3.87	0.08	9.40
20 FS1-2000-8*	0.05	32.14	0.08	5.75	0.13	21.13
21 FS1-2000-9*	0.05	3.52	0.12	8.51	0.17	12.03
22 FS1-2000-10*	0.06	7.30	0.05	5.35	0.11	12.66
23 FS1-2000-11*	0.08	10.76	0.01	1.51	0.10	12.27
24 FS1-2000-12*	0.07	2.39	0.21	7.05	0.28	9.43
25 FS1-2000-13*	0.04	3.01	0.05	3.77	0.09	6.78

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

GB-2 (1,900m)  
 GREENLAND BASIN  
 74°35'N, 06°43'W  
 Trap depth: 881m Water depth: 3,445m


Annual Fluxes (g/m<sup>2</sup>/yr):

Total.....	8.79
Carbonate.....	2.59
Noncombustible.....	3.69
Combustible.....	2.50
Organic C.....	0.94
N .....	0.16

PARFLUX Mark 5-12

Sample ID	Opening Date	Closing Date	Span	Mid. Date
89 GB2-1000-1	06-AUG-85	05-SEP-85	30	21-AUG-85
90 GB2-1000-2	05-SEP-85	05-OCT-85	30	20-SEP-85
91 GB2-1000-3	05-OCT-85	04-NOV-85	30	20-OCT-85
92 GB2-1000-4	04-NOV-85	04-DEC-85	30	19-NOV-85
93 GB2-1000-5	04-DEC-85	03-JAN-86	30	19-DEC-85
94 GB2-1000-6	03-JAN-86	02-FEB-86	30	18-JAN-86
95 GB2-1000-7	02-FEB-86	04-MAR-86	30	17-FEB-86
96 GB2-1000-8	04-MAR-86	03-APR-86	30	19-MAR-86
97 GB2-1000-9	03-APR-86	03-MAY-86	30	18-APR-86
98 GB2-1000-10	03-MAY-86	02-JUN-86	30	18-MAY-86
99 GB2-1000-11	02-JUN-86	02-JUL-86	30	17-JUN-86
100 GB2-1000-12	02-JUL-86	01-AUG-86	30	17-JUL-86

Total Flux at Greenland Basin 2, 1900m, 1985-86




Sample I.D.	TTLF <1	<1 % of total	TTLF >1	>1 % of total	TTLF total
89 GB2-1900-1	30.25	82.01	6.63	17.99	36.88
90 GB2-1900-2	23.44	78.53	6.41	21.47	29.86
91 GB2-1900-3	30.75	86.77	4.69	13.23	35.44
92 GB2-1900-4	21.21	92.38	1.75	7.62	22.96
93 GB2-1900-5	16.44	92.42	1.35	7.58	17.79
94 GB2-1900-6	13.42	92.27	1.12	7.73	14.54
95 GB2-1900-7	9.46	86.77	1.44	13.23	10.91
96 GB2-1900-8	0.10	100.00			0.10
97 GB2-1900-9	0.10	100.00			0.10
98 GB2-1900-10	0.08	100.00			0.08
99 GB2-1900-11					
100 GB2-1900-12					

Flux is in mg/m<sup>2</sup>/day.

Trap malfunctioned beginning at cup #8.


Carbonate Flux at Greenland Basin 2, 1900m, 1985-86



Sample I.D.	CRTA <1	CRTA % tot.<1	CRTA >1	CRTA % tot.>1	CRTA total	CRTA % total
89 GB2-1900-1	7.14	19.35	0.78	2.11	7.92	21.47
90 GB2-1900-2	7.03	23.57	1.30	4.36	8.33	27.93
91 GB2-1900-3	12.76	36.57	1.04	2.95	13.81	39.51
92 GB2-1900-4	7.78	33.91	0.35	1.53	8.14	35.44
93 GB2-1900-5	4.14	23.29	0.23	1.27	4.37	24.56
94 GB2-1900-6	3.72	25.51	0.29	1.97	4.00	27.48
95 GB2-1900-7	2.58	23.66	0.26	2.36	2.84	26.02
96 GB2-1900-8						
97 GB2-1900-9						
98 GB2-1900-10						
99 GB2-1900-11						
100 GB2-1900-12						

Flux is in m/m<sup>2</sup>/day  
Trap malfunctioned beginning at cup #8.

Noncombustible Flux at Greenland Basin 2, 1900m, 1985-86




Sample ID#	NONC <1	NONC % tot.<1	NONC >1	NONC % tot.>1	NONC total	NONC % total
89 GB2-1900-1	14.12	38.29	3.91	10.59	18.03	48.88
90 GB2-1900-2	9.84	32.95	3.18	10.66	13.02	43.62
91 GB2-1900-3	10.06	28.38	2.36	6.66	12.42	35.04
92 GB2-1900-4	8.63	37.60	1.07	4.64	9.70	42.25
93 GB2-1900-5	6.53	36.69	0.99	5.54	7.51	42.22
94 GB2-1900-6	5.60	38.49	0.32	2.19	5.92	40.68
95 GB2-1900-7	3.61	33.06	0.55	5.01	4.15	38.07
96 GB2-1900-8						
97 GB2-1900-9						
98 GB2-1900-10						
99 GB2-1900-11						
100 GB2-1900-12						

Flux is in mg/m<sup>2</sup>/day.

Trap malfunctioned beginning at cup #8.

Combustible Flux at Greenland Basin 2, 1900m, 1985-86




Sample ID#	COMB <1	COMB % tot.<1	COMB >1	COMB % tot.>1	COMB TOTAL	COMB % total
89 GB2-1900-1	8.99	24.37	1.95	5.28	10.94	29.65
90 GB2-1900-2	6.57	22.01	1.93	6.46	8.50	28.47
91 GB2-1900-3	7.93	22.38	1.29	3.63	9.22	26.01
92 GB2-1900-4	4.79	20.87	0.33	1.45	5.12	22.32
93 GB2-1900-5	5.77	32.44	0.14	0.78	5.91	33.22
94 GB2-1900-6	4.10	28.22	0.52	3.57	4.62	31.79
95 GB2-1900-7	3.27	30.02	0.64	5.86	3.91	35.93
96 GB2-1900-8						
97 GB2-1900-9						
98 GB2-1900-10						
99 GB2-1900-11						
100 GB2-1900-12						

Flux is in mg/m<sup>2</sup>/day.

Trap malfunctioned beginning at cup #8.


Carbon Flux at Greenland Basin 2, 1900m, 1985-86



Sample I.D.	CRNC <1	CRNC<1 %cmbf.	CRNC >1	CRNC>1 %cmbf.	CRNC total	CRNCtot. %cmbf.
89 GB2-1900-1	4.21	38.53	0.82	7.46	5.03	45.98
90 GB2-1900-2	3.10	36.43	0.87	10.21	3.97	46.71
91 GB2-1900-3	3.78	41.02	0.69	7.49	4.47	48.48
92 GB2-1900-4	2.24	43.70	0.19	3.65	2.43	47.46
93 GB2-1900-5	2.56	43.32	0.36	6.15	2.92	49.41
94 GB2-1900-6	1.48	32.00	0.15	3.28	1.63	35.28
95 GB2-1900-7	0.68	17.49	0.57	14.68	1.25	31.97
96 GB2-1900-8						
97 GB2-1900-9						
98 GB2-1900-10						
99 GB2-1900-11						
100 GB2-1900-12						

Flux is in mg/m<sup>2</sup>/day.

Nitrogen Flux at Greenland Basin 2, 1900m, 1985-86




Sample I.D.	NTGN <1	NTGN<1 %cmbf.	NTGN >1	NTGN>1 %cmbf.	NTGN total	NTGNtot. %cmbf.
89 GB2-1900-1	0.68	6.19	0.13	1.17	0.81	7.36
90 GB2-1900-2	0.49	5.75	0.14	1.61	0.63	7.36
91 GB2-1900-3	0.56	6.05	0.10	1.11	0.66	7.16
92 GB2-1900-4	0.37	7.23	0.03	0.66	0.40	7.39
93 GB2-1900-5	0.45	7.66	0.05	0.85	0.50	8.50
94 GB2-1900-6	0.27	5.92	0.02	0.54	0.30	6.45
95 GB2-1900-7	0.33	8.44	0.09	2.22	0.42	10.66
96 GB2-1900-8						
97 GB2-1900-9						
98 GB2-1900-10						
99 GB2-1900-11						
100 GB2-1900-12						

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Hydrogen Flux at Greenland Basin 2. 1900m. 1985-86



Sample I.D.	HYDC <1	HYDC<1 %cmbf.	HYDC >1	HYDC>1 %cmbf.	HYDC total	HYDCTot. %cmbf.
89 GB2-1900-1	0.59	5.37	0.11	0.98	0.69	6.35
90 GB2-1900-2	0.41	4.87	0.11	1.27	0.52	6.14
91 GB2-1900-3	0.47	5.07	0.09	0.98	0.56	6.05
92 GB2-1900-4	0.30	5.76	0.02	0.45	0.32	6.21
93 GB2-1900-5	0.35	5.87	0.05	0.82	0.40	6.69
94 GB2-1900-6	0.20	4.43	0.02	0.47	0.23	4.39
95 GB2-1900-7	0.14	3.52	0.08	1.92	0.21	5.44
96 GB2-1900-8						
97 GB2-1900-9						
98 GB2-1900-10						
99 GB2-1900-11						
100 GB2-1900-12						

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".


GB-2 3,000m  
GREENLAND BASIN  
74°35'N, 06°43'W  
Trap depth: 2,823m Water depth: 3,445m

Annual Fluxes (g/m<sup>2</sup>/yr):  
Total.....10.21  
Carbonate.....3.28  
Noncombustible.....5.73  
Combustible.....1.23  
Biogenic Opal.....2.61  
Lithogenic.....3.12  
Organic C.....0.40  
N .....0.06

PARFLUX Mark 6-13

Sample ID	Opening Date	Closing Date	Span	Mid. Date
101 GB2-3000-1	04-AUG-85	01-SEP-85	28	18-AUG-85
102 GB2-3000-2	01-SEP-85	29-SEP-85	28	15-SEP-85
103 GB2-3000-3	29-SEP-85	27-OCT-85	28	13-OCT-85
104 GB2-3000-4	27-OCT-85	24-NOV-85	28	10-NOV-85
105 GB2-3000-5	24-NOV-85	22-DEC-85	28	08-DEC-85
106 GB2-3000-6	22-DEC-85	19-JAN-86	28	05-JAN-86
107 GB2-3000-7	19-JAN-86	16-FEB-86	28	02-FEB-86
108 GB2-3000-8	16-FEB-86	16-MAR-86	28	02-MAR-86
109 GB2-3000-9	16-MAR-86	13-APR-86	28	30-MAR-86
110 GB2-3000-10	13-APR-86	11-MAY-86	28	27-APR-86
111 GB2-3000-11	11-MAY-86	08-JUN-86	28	25-MAY-86
112 GB2-3000-12	08-JUN-86	06-JUL-86	28	22-JUN-86
113 GB2-3000-13	06-JUL-86	03-AUG-86	28	20-JUL-86


Total Flux at Greenland Basin 2, 3000m, 1985-86



Sample I.D.	TTLF <1	<1 % of total	TTLF >1	>1 % of total	TTLF total
<hr/>					
101 GB2-3000-1	19.29	99.48	0.10	0.52	19.39
102 GB2-3000-2	30.23	99.43	0.17	0.57	30.40
103 GB2-3000-3	28.68	98.94	0.31	1.06	28.99
104 GB2-3000-4	31.92	98.34	0.54	1.66	32.46
105 GB2-3000-5	18.15	99.37	0.11	0.63	18.27
106 GB2-3000-6	25.71	93.48	1.79	6.52	27.51
107 GB2-3000-7	26.64	92.41	2.19	7.59	28.83
108 GB2-3000-8	20.24	89.10	2.48	10.90	22.71
109 GB2-3000-9	17.57	99.26	0.13	0.74	17.70
110 GB2-3000-10	26.59	98.12	0.51	1.88	27.10
111 GB2-3000-11	23.64	100.00			23.64
112 GB2-3000-12	25.28	97.96	0.53	2.04	25.80
113 GB2-3000-13	58.24	96.00	2.43	4.00	60.67

Flux is in mg/m<sup>2</sup>/day.


Carbonate Flux at Greenland Basin 2, 3000m, 1985-86



Sample I.D.	CRTA <1	CRTA % tot.<1	CRTA >1	CRTA % tot.>1	CRTA total	CRTA % total
101 GB2-3000-1	6.19	31.93	0.00	31.93	6.20	31.95
102 GB2-3000-2	10.01	32.91	0.04	32.91	10.04	33.03
103 GB2-3000-3	8.20	28.30	0.06	28.30	8.27	28.52
104 GB2-3000-4	9.70	29.90	0.17	29.90	9.87	30.41
105 GB2-3000-5	5.81	31.80	0.00	31.80	5.81	31.83
106 GB2-3000-6	8.20	29.82	0.26	29.82	8.46	30.76
107 GB2-3000-7	8.42	29.20	0.56	29.20	8.98	31.14
108 GB2-3000-8	7.33	32.25	1.44	32.25	8.77	38.59
109 GB2-3000-9	5.90	33.35	0.07	33.35	5.98	33.76
110 GB2-3000-10	8.80	32.48	0.10	32.48	8.90	32.85
111 GB2-3000-11	7.94	33.59	0.00	33.59	7.94	33.59
112 GB2-3000	7.89	30.56	0.13	30.56	8.02	31.07
113 GB2-3000	17.30	28.51	0.42	28.51	17.72	29.20

Flux is in mg/m<sup>2</sup>/day.


Noncombustible Flux at Greenland Basin 2, 3000m, 1985-6



Sample ID#	NONC <1	NONC % tot.<1	NONC >1	NONC % tot.>1	NONC total	NONC % total
101 GB2-3000-1	10.19	52.55	0.04	0.20	10.23	52.75
102 GB2-3000-2	15.96	52.48	0.07	0.23	16.03	52.71
103 GB2-3000-3	16.61	57.29	0.08	0.28	16.69	57.57
104 GB2-3000-4	18.04	55.58	0.28	0.86	18.32	56.44
105 GB2-3000-5	10.16	55.61	0.06	0.30	10.21	55.91
106 GB2-3000-6	13.66	49.66	1.15	4.18	14.81	53.84
107 GB2-3000-7	15.04	52.15	1.62	5.63	16.66	57.78
108 GB2-3000-8	12.86	56.62	0.49	2.18	13.35	58.79
109 GB2-3000-9	9.44	53.32	0.06	0.33	9.50	53.65
110 GB2-3000-10	14.50	53.50	0.35	1.29	14.85	54.79
111 GB2-3000-11	13.14	55.58			13.14	55.58
112 GB2-3000-12	14.66	56.81	0.32	1.25	14.98	58.07
113 GB2-3000-13	34.76	57.30	0.38	0.62	35.14	57.92

Flux is in mg/m<sup>2</sup>/day.


Combustible Flux at Greenland Basin 2, 3000m, 1985-86



Sample ID#	COMB <1	COMB % tot.<1	COMB >1	COMB % tot.>1	COMB TOTAL	COMB % total
101 GB2-3000-1	2.91	15.00	0.06	0.30	2.97	15.30
102 GB2-3000-2	4.27	14.04	0.07	0.23	4.34	14.26
103 GB2-3000-3	3.87	13.35	0.16	0.56	4.03	13.92
104 GB2-3000-4	4.18	12.87	0.09	0.28	4.27	13.15
105 GB2-3000-5	2.18	11.96	0.06	0.30	2.24	12.26
106 GB2-3000-6	3.85	14.01	0.39	1.40	4.24	15.41
107 GB2-3000-7	3.19	11.06	0.01	0.02	3.20	11.08
108 GB2-3000-8	0.05	0.23	0.54	2.39	0.59	2.61
109 GB2-3000-9	2.23	12.59	0.00	0.00	2.23	12.59
110 GB2-3000-10	3.29	12.14	0.06	0.22	3.35	12.36
111 GB2-3000-11	2.56	10.82	0.00	0.00	2.56	10.82
112 GB2-3000-12	2.73	10.58	0.07	0.28	2.80	10.86
113 GB2-3000-13	6.18	10.19	1.63	2.69	7.81	12.88

Flux is in mg/m<sup>2</sup>/day.

Biogenic Silica Flux at Greenland Basin 2, 3000m, 1985-86




Sample ID#	OPAL <1	OPAL % Ncf.<1	OPAL % tot.<1
<hr/>			
101 GB2-3000-1	6.18	60.40	31.86
102 GB2-3000-2	7.99	49.86	26.28
103 GB2-3000-3	8.34	49.97	28.77
104 GB2-3000-4	8.81	48.09	27.14
105 GB2-3000-5	4.58	44.84	25.07
106 GB2-3000-6	6.25	42.21	22.72
107 GB2-3000-7	5.99	35.96	20.78
108 GB2-3000-8	4.17	31.23	18.36
109 GB2-3000-9	3.78	39.81	21.36
110 GB2-3000-10	5.23	35.23	19.30
111 GB2-3000-11	4.60	35.01	19.46
112 GB2-3000-12	7.12	47.52	27.59
113 GB2-3000-13	19.83	56.43	32.69

Flux is in mg/m<sup>2</sup>/day.

"%Ncf." = "% of noncombustible flux".

Not enough >1 mm fraction to do analysis.

Lithogenic Flux at Greenland Basin 2, 3000 m, 1985-86




Sample I.D.	LITH <1	LITH<1 %Ncmb.	LITH<1 %tot.
<hr/>			
101 GB2-3000-1	4.01	39.22	20.69
102 GB2-3000-2	7.97	49.71	26.20
103 GB2-3000-3	8.27	49.55	28.52
104 GB2-3000-4	9.23	50.39	28.44
105 GB2-3000-5	5.58	54.62	30.54
106 GB2-3000-6	7.41	50.03	26.93
107 GB2-3000-7	9.05	54.30	31.37
108 GB2-3000-8	8.69	78.70	38.26
109 GB2-3000-9	5.66	59.58	31.96
110 GB2-3000-10	9.27	62.41	34.20
111 GB2-3000-11	8.54	64.99	36.12
112 GB2-3000-12	7.54	50.32	29.22
113 GB2-3000-13	14.93	42.49	24.61

Flux is in mg/m<sup>2</sup>/day.

%Ncmb. = "% of noncombustible flux".

Not enough >1 mm fraction to do analysis.

Carbon Flux at Greenland Basin 2, 3000m, 1985-86




Sample I.D.	CRNC <1	CRNC<1 %cmbf.
101 GB2-3000-1	1.04	35.06
102 GB2-3000-2	1.52	35.12
103 GB2-3000-3	0.36	8.99
104 GB2-3000-4	1.38	32.43
105 GB2-3000-5	0.70	31.47
106 GB2-3000-6	1.07	25.21
107 GB2-3000-7	1.09	34.22
108 GB2-3000-8	1.08	37.24
109 GB2-3000-9	1.07	47.91
110 GB2-3000-10	1.10	32.72
111 GB2-3000-11	1.08	42.25
112 GB2-3000-12	0.44	15.77
113 GB2-3000-13	2.48	31.70

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Not enough >1 mm fraction to do analysis.

Nitrogen Flux at Greenland Basin 2, 3000m, 1985-86




Sample I.D.	NTGN <1	NTGN<1 %cmbf.
101 GB2-3000-1	0.15	5.21
102 GB2-3000-2	0.22	4.99
103 GB2-3000-3	0.05	1.27
104 GB2-3000-4	0.21	4.89
105 GB2-3000-5	0.10	4.57
106 GB2-3000-6	0.17	3.93
107 GB2-3000-7	0.14	4.51
108 GB2-3000-8	0.16	5.52
109 GB2-3000-9	0.12	5.24
110 GB2-3000-10	0.14	4.14
111 GB2-3000-11	0.15	5.95
112 GB2-3000-12	0.05	1.61
113 GB2-3000-13	0.36	4.66

Flux is in  $\text{mg}/\text{m}^2/\text{day}$ .

"%cmbf" = "% of combustible flux".

Not enough >1 mm fraction to do analysis.

Hydrogen Flux at Greenland Basin 2, 3000m, 1985-86



Sample I.D.	HYDC <1	HYDC<1 %cmbf.
101 GB2-3000-1	0.16	5.39
102 GB2-3000-2	0.24	5.60
103 GB2-3000-3	0.05	1.12
104 GB2-3000-4	0.22	5.10
105 GB2-3000-5	0.11	4.85
106 GB2-3000-6	0.16	3.68
107 GB2-3000-7	0.20	6.10
108 GB2-3000-8	0.16	5.52
109 GB2-3000-9	0.17	7.43
110 GB2-3000-10	0.17	5.10
111 GB2-3000-11	0.17	6.56
112 GB2-3000-12	0.06	2.05
113 GB2-3000-13	0.41	5.29

Flux is in mg/m<sup>2</sup>/day.

"%cmbf" = "% of combustible flux".

Not enough >1 mm fraction to do analysis.

<b>REPORT DOCUMENTATION PAGE</b>		1. REPORT NO. WHOI-87-17	2.	3. Recipient's Accession No.
4. Title and Subtitle  Particle Fluxes, North-Eastern Nordic Seas: 1983-1986 (Nordic Seas Sedimentation Data File, Vol. 1)		5. Report Date April 1987		6.
7. Author(s) Susumu Honjo, Steven J. Manganini, Amy Karowe, Bonnie L. Woodward		8. Performing Organization Rept. No. WHOI-87-17		10. Project/Task/Work Unit No.
9. Performing Organization Name and Address  Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543		11. Contract(C) or Grant(G) No. (C) (G) N00014-85-C-0001		13. Type of Report & Period Covered  Technical
12. Sponsoring Organization Name and Address  Office of Naval Research		14.		
15. Supplementary Notes  This report should be cited as: Woods Hole Oceanogr. Inst. Tech. Rept. WHOI-87-17				
16. Abstract (Limit: 200 words)  Seventy-nine particle flux samples were collected from 1983 to 1986 using 7 automated time-series sediment traps at 6 stations distributed in the northern and eastern portion of the Nordic Seas as part of a German/U.S. joint program on arctic sedimentation studies. Each sample represents either one month or two weeks of sedimentation at approximately 400 m above the sea floor. In this data file the results of laboratory analysis conducted at the Woods Hole Oceanographic Institution, U.S.A. of the main sedimentological criteria: total mass, carbonate, opal, combustible, organic carbon, nitrogen, and lithogenic mass are presented in both tabular and histogram form. Results from the southern and western portion of the Nordic Seas will be published as they become available.				
17. Document Analysis a. Descriptors  1. Nordic Seas 2. Material flux 3. Sediment trap				
b. Identifiers/Open-Ended Terms				
c. COSATI Field/Group				
18. Availability Statement:  Approved for publication; distribution unlimited		19. Security Class (This Report) UNCLASSIFIED	21. No. of Pages 84	20. Security Class (This Page)
		22. Price		

# DOCUMENT LIBRARY

November 21, 1986

## *Distribution List for Technical Report Exchange*

Institute of Marine Sciences Library  
University of Alaska  
O'Neill Building  
905 Koyukuk Ave., North  
Fairbanks, AK

Attn: Stella Sanchez-Wade  
Documents Section  
Scripps Institution of Oceanography  
Library, Mail Code C-075C  
La Jolla, CA 92093

Hancock Library of Biology & Oceanography  
Alan Hancock Laboratory  
University of Southern California  
University Park  
Los Angeles, CA 90089-0371

Gifts & Exchanges  
Library  
Bedford Institute of Oceanography  
P.O. Box 1006  
Dartmouth, NS, B2Y 4A2, CANADA

Office of the International  
Ice Patrol  
c/o Coast Guard R & D Center  
Avery Point  
Groton, CT 06340

Library  
Physical Oceanographic Laboratory  
Nova University  
8000 N. Ocean Drive  
Dania, FL 33304

NOAA/EDIS Miami Library Center  
4301 Rickenbacker Causeway  
Miami, FL 33149

Library  
Skidaway Institute of Oceanography  
P.O. Box 13687  
Savannah, GA 31416

Institute of Geophysics  
University of Hawaii  
Library Room 252  
2525 Correa Road  
Honolulu, HI 96822

Library  
Chesapeake Bay Institute  
4800 Atwell Road  
Shady Side, MD 20876

MIT Libraries  
Serial Journal Room 14E-210  
Cambridge, MA 02139

Director, Ralph M. Parsons Laboratory  
Room 48-311  
MIT  
Cambridge, MA 02139

Marine Resources Information Center  
Bldg. E38-320  
MIT  
Cambridge, MA 02139

Library  
Lamont-Doherty Geological Observatory  
Colombia University  
Palisades, NY 10964

Library  
Serials Department  
Oregon State University  
Corvallis, OR 97331

Pell Marine Science Library  
University of Rhode Island  
Narragansett Bay Campus  
Narragansett, RI 02882

Working Collection  
Texas A&M University  
Dept. of Oceanography  
College Station, TX 77843

Library  
Virginia Institute of Marine Science  
Gloucester Point, VA 23062

Fisheries-Oceanography Library  
151 Oceanography Teaching Bldg.  
University of Washington  
Seattle, WA 98195

Library  
R.S.M.A.S.  
University of Miami  
4600 Rickenbacker Causeway  
Miami, FL 33149

Maury Oceanographic Library  
Naval Oceanographic Office  
Bay St. Louis  
NSTL, MS 39522-5001  
ATTN: Code 4601