

## Path and Association Analysis and Stress Indices for Salinity Tolerance Traits in Promising Rice (*Oryza sativa* L.) Genotypes

S.L. KRISHNAMURTHY<sup>1\*</sup>, S.K. SHARMA<sup>1</sup>, R.K. GAUTAM<sup>2</sup> and V. KUMAR<sup>1</sup>

<sup>1</sup>Central Soil Salinity Research Institute, Karnal – 132001, India

<sup>2</sup>Central Agricultural Research Institute for Port Blair, Andaman and Nicobar Islands – 744101, India

(Received 17 July 2013; Accepted 15 August 2013;

Communicated by A. Anioł)

Effects of salinity on correlation, path and stress indices, yield and its components were studied in a set of 34 promising rice genotypes collected from various national and international organizations. These genotypes were evaluated in a randomized complete block design with three replications during the wet seasons (*khariif*) of 2009 and 2010 in normal ( $EC_{iw} \sim 1.2$  dS/m) and salinity stress ( $EC_{iw} \sim 10$  dS/m) environments in micro plots at Central Soil Salinity Research Institute (CSSRI), Karnal, India. Grain yield per plant showed positive significant association with plant height, total tillers, productive tillers, panicle length, and biological yield per plant and harvest index under normal environment, whereas grain yield showed positive significant association with biological yield and harvest index under salinity stress. These results clearly indicate that selection of high yielding genotypes would be entirely different under normal and saline environments. The stress susceptibility index (SSI) values for grain yield ranged from 0.35 (HKR 127) to 1.55 (TR-2000-008), whereas the stress tolerance index (STI) values for grain yield ranged from 0.07 (PR 118) to 1.09 (HKR 120). The genotypes HKR 120, HKR 47 and CSR-RIL-197 exhibited higher values of stress tolerance index (STI) in salinity. Under salinity, negative and significant association was shown by SSI and grain yield in contrast to positive and significant association shown by STI and grain yield. These associations could be useful in identifying salt tolerant and sensitive high yielding genotypes. The stress susceptible and stress tolerance indices suggest that the genotypes developed for salinity tolerance could exhibit higher tolerance, adaptability and suitability. Harvest index and biological yield traits emerged as the ideal traits for improvement through selection and could be used to increase the rice productivity under saline stress environments.

**Keywords:** rice, salinity, correlation, path coefficient, SSI and STI

### Introduction

Rice is the staple food and most important cereal crop which contributes more to human food requirements annually and accounts for about 43 per cent of total food grain production in India. At the current rate of population growth (1.8 per cent), rice requirement by 2020, would be around 125 million tons (Mishra 2005). The need and importance of rice is

\* Corresponding author; E-mail: krishnagene@gmail.com

increasing day by day due to the increase in global population. Therefore, improving the productivity of rice is crucial for food security and economic development. Higher population growth rate and the conversion of some highly productive rice cultivation lands for industrial and residential purposes, has pushed rice cultivation to less productive area such as saline, drought and flood prone areas. In India, salt affected area accounts for 6.73 million hectares of land. Though most of the cultivated rice varieties are susceptible to salinity but variability for salt tolerance traits has also been reported (Sharma 1986, 2010; Flowers et al. 2000). The detrimental effects of salt stress are through alterations in plant metabolism, reduced water potential, ion imbalances, toxicity and reduction of crop yield and severe salt stress may lead to total failure of crop. The effects of salinity are also related to the stage of plant development at which salinity occurs, concentration and nature of the salt, and the duration of salinization (Pearson et al. 1966; Aslam et al. 1993; Zeng et al. 2001). In rice, plant height, total number of tillers, panicle length, grain weight per panicle, 1000-seed weight and quality and quantity of grains decreased progressively with increase in salinity. Involvement of some undesirable characters in the traditional varieties and complex nature of salinity tolerance have presented challenges in efforts to improve salt tolerance of crops (Yamaguchi and Blumwald 2005). Breeding of saline tolerant varieties necessitates understanding of salinity, its effects on various plant characters and including selection for one or several characters in artificially generated populations or natural populations (Simmonds 1983; Arshadullah et al. 2011). Success of any plant breeding programme depends on the magnitude of variation for yield and yield components present in the germplasm and the nature of association among these components. To improve grain yield, evaluation of germplasm is the most important aspect. Being a complex trait, grain yield is influenced by various genetic and environmental factors; therefore information on the extent of association between traits is important and should be given importance in selection. With these points in view, the present investigation was carried out to understand the effects of salinity on genetic variability, correlation, direct and indirect influences and stress indices of grain yield with important yield components in rice.

## Materials and Methods

### *Plant materials*

The experimental materials comprised of 34 promising rice genotypes collected from various national and international organizations (Table S1\*). Genotypes included popular high yielding varieties, high yielding salt tolerant varieties in India and recombinant inbred lines (derived from salt tolerant and salt sensitive parents) developed by CSSRI, Karnal and elite salt tolerant breeding lines developed by IRRI, Philippines. These genotypes were evaluated in randomized complete block design with three replications during *kharif*2009 and *kharif*2010 in control ( $EC_{iw} \sim 1.2$  dS/m) and salinity ( $EC_{iw} \sim 10$  dS/m) environments in micro plots at Central Soil Salinity Research Institute, Karnal, Haryana, In-

\* Further details about the Electronic Supplementary Material (ESM) can be found at the end of the article.

dia, situated at 29.43° N latitude and 76.58° N longitude and 245 m above sea level. These micro plots were covered with polycarbonate sheet sheets to protect against rains. Thirty-five days old seedlings from the wet bed nurseries were transplanted two seedlings per hill with a spacing of 15 × 20 cm. Basal fertilizers for the main crop were 120–60–60 kg of NPK/ha and the recommended package of practices was followed to grow a healthy crop. One month after transplanting, salinity stress was imposed by irrigating with saline water created using 7 NaCl: 1 Na<sub>2</sub>SO<sub>4</sub>: 2 CaCl<sub>2</sub> salts on equivalent basis. Subsequently, salinity was recorded weekly and maintained around ( $EC_{iw} \sim 10 \text{ dSm}^{-1}$ ) throughout the cropping season.

#### *Evaluation for yield and its yield components and stress indices analysis*

Five plants from each genotype were randomly tagged in each replication and data recorded for plant height, total tillers per plant, number of productive tillers per plant, panicle length, grain yield per plant, biological yield per plant and harvest index (%). The stress susceptibility indices (SSIs) and stress tolerance indices (STIs) for each genotype were calculated for grain yield as per Fischer and Maurer (1978) and Fernandez (1993).

#### *Statistical analysis*

Homogeneity of error variance across the two seasons was tested by the F test (Gomez and Gomez 1983) and the combined analyses of variances for genotypes were performed. Significant levels were determined (McIntosh 1983) for the combined analysis. Analysis of variance (ANOVA) for all the traits was estimated (Panse and Sukhatme 1967). The genotypic and phenotypic correlations were estimated (Al-Jibouri 1958) and the genotypic correlation coefficients were partitioned into direct and indirect effects (Dewey and Lu 1959).

## **Results**

#### *Analysis of variance*

Homogeneity of error variance across the two seasons was tested by the F test and none of the error mean squares were found significant for any of the traits. Therefore, combined analyses of variances for genotypes between the two seasons were determined by comparing the genotypes × seasons interaction for each trait. Significant levels were also determined for combined analysis. As significant differences among genotypes, seasons interaction were not observed (Table S2), data of both the seasons were therefore combined for further analysis. ANOVA revealed highly significant variation among the treatments for all the seven traits in both the control and saline environments thus suggesting the presence of sufficient variation among the genotypes along with ample scope for improving salinity tolerance in rice.

*Effect of salinity on grain yield and yield attributing traits*

Grain yield per plant was the trait most affected by salinity as expressed by 58 per cent reduction in saline environment compared to control non-saline environment (Table S3) and was followed by reductions in biological yield (52%), productive tillers per plant (28%), plant height (26%), total tillers per plant (24%), panicle length (18%) and harvest index (14%), respectively.

*Correlation and path analysis*

Grain yield per plant showed significant and positive association with plant height (0.48 and 0.60), total tillers per plant (0.26 and 0.18), productive tillers per plant (0.25 and 0.20), panicle length (0.66 and 0.71), biological yield per plant (0.68 and 0.75) and harvest index (0.85 and 0.65) at genotypic and phenotypic levels, respectively under normal environment (Table 1). The direct effects of plant height and panicle length were positive but lower in magnitude. Direct positive effects of these traits on grain yield indicated their im-

Table 1. Genotypic (G) and phenotypic (P) correlations for yield and yield contributing traits in rice in control ( $EC_{iw} \sim 1.2$  dS/m) and saline ( $EC_{iw} \sim 10$  dS/m) environments

| Characters                     | Environments | Total tillers | Productive tillers | Panicle length | Biological yield | Harvest index | Grain yield |        |
|--------------------------------|--------------|---------------|--------------------|----------------|------------------|---------------|-------------|--------|
| Plant height (cm)              | G            | Normal        | 0.68**             | 0.64**         | 0.66**           | 0.64**        | 0.25**      | 0.48** |
|                                |              | Saline        | -0.09              | 0.14*          | 0.37**           | 0.35**        | 0.03        | 0.25** |
|                                | P            | Normal        | 0.51**             | 0.51**         | 0.70**           | 0.71**        | 0.19*       | 0.60** |
|                                |              | Saline        | -0.15*             | -0.14*         | 0.38**           | 0.07          | -0.1        | -0.01  |
| Total tillers per plant        | G            | Normal        | 1                  | 0.96**         | 0.46**           | 0.47**        | 0.17*       | 0.26** |
|                                |              | Saline        | 1                  | 0.49**         | 0.07             | 0.26**        | 0.05        | 0.14*  |
|                                | P            | Normal        | 1                  | 0.96**         | 0.43**           | 0.35**        | -0.01       | 0.18** |
|                                |              | Saline        | 1                  | 0.73**         | -0.01            | 0.12          | 0.02        | 0.08   |
| Productive tillers per plant   | G            | Normal        | 1                  | 1              | 0.46**           | 0.46**        | 0.18*       | 0.25** |
|                                |              | Saline        | 1                  | 1              | 0.16*            | 0.022         | 0.20**      | 0.07   |
|                                | P            | Normal        | 1                  | 1              | 0.45**           | 0.37**        | 0.001       | 0.20** |
|                                |              | Saline        | 1                  | 1              | -0.01            | 0.027         | 0.15*       | 0.1    |
| Panicle length (cm)            | G            | Normal        | 1                  | 1              | 0.24**           | 0.77**        | 0.66**      |        |
|                                |              | Saline        | 1                  | 1              | -0.11            | 0.40**        | 0.16*       |        |
|                                | P            | Normal        | 1                  | 1              | 0.53**           | 0.47**        | 0.71**      |        |
|                                |              | Saline        | 1                  | 1              | -0.063           | 0.20**        | 0.11        |        |
| Biological yield per plant (g) | G            | Normal        | 1                  | 1              | 1                | 0.24**        | 0.68**      |        |
|                                |              | Saline        | 1                  | 1              | 1                | 0.04**        | 0.76**      |        |
|                                | P            | Normal        | 1                  | 1              | 1                | 0.03          | 0.75**      |        |
|                                |              | Saline        | 1                  | 1              | 1                | 0.18**        | 0.75**      |        |
| Harvest index (%)              | G            | Normal        | 1                  | 1              | 1                | 1             | 0.85**      |        |
|                                |              | Saline        | 1                  | 1              | 1                | 1             | 0.67**      |        |
|                                | P            | Normal        | 1                  | 1              | 1                | 1             | 0.65**      |        |
|                                |              | Saline        | 1                  | 1              | 1                | 1             | 0.77**      |        |

\* significant at  $p = 0.05$ ; \*\* significant at  $p = 0.01$

portance in determining the complex character and, therefore, should be given priority while making selections for the improving grain yield. Under salinity stress, biological yield per plant had the highest positive direct effect on grain yield (0.76) and was followed by harvest index (0.65) (Table 2).

Table 2. Direct (diagonal) and indirect (off-diagonal) effects of different traits in rice on grain yield in control ( $EC_{iw} \sim 1.2$  dS/m) and saline ( $EC_{iw} \sim 10$  dS/m) environments

| Characters                     | Environ-ments | Plant height (cm) | Total tillers per plant | Productive tillers per plant | Panicle length (cm) | Biological yield per plant (g) | Harvest index (%) | Genotypic correlation |
|--------------------------------|---------------|-------------------|-------------------------|------------------------------|---------------------|--------------------------------|-------------------|-----------------------|
| Plant height (cm)              | Normal        | <b>0.03</b>       | 0.018                   | 0.02                         | 0.018               | 0.017                          | 0.007             | 0.48**                |
|                                | Saline        | <b>-0.04</b>      | 0.003                   | -0.005                       | -0.015              | -0.01                          | -0.001            | 0.25**                |
| Total tillers per plant        | Normal        | -0.07             | <b>-0.01</b>            | -0.009                       | -0.004              | -0.004                         | -0.002            | 0.26**                |
|                                | Saline        | 0.007             | <b>-0.07</b>            | -0.04                        | -0.005              | -0.019                         | -0.003            | 0.14*                 |
| Productive tillers per plant   | Normal        | -0.10             | -0.15                   | <b>-0.16</b>                 | -0.07               | -0.072                         | -0.028            | 0.25**                |
|                                | Saline        | -0.01             | -0.02                   | <b>-0.03</b>                 | -0.005              | -0.001                         | -0.006            | 0.07                  |
| Panicle length (cm)            | Normal        | 0.02              | 0.01                    | 0.01                         | <b>0.02</b>         | 0.005                          | 0.018             | 0.66**                |
|                                | Saline        | 0.01              | 0.001                   | 0.002                        | <b>0.01</b>         | -0.002                         | 0.001             | 0.17*                 |
| Biological yield per plant (g) | Normal        | 0.36              | 0.27                    | 0.26                         | 0.14                | <b>0.57</b>                    | 0.14              | 0.69**                |
|                                | Saline        | 0.27              | 0.12                    | 0.017                        | -0.08               | <b>0.76</b>                    | 0.03              | 0.76**                |
| Harvest index (%)              | Normal        | 0.18              | 0.12                    | 0.12                         | 0.55                | 0.17                           | <b>0.72</b>       | 0.85**                |
|                                | Saline        | 0.02              | 0.03                    | 0.13                         | 0.26                | 0.02                           | <b>0.65</b>       | 0.67**                |

Residual effects under normal and saline environments are 0.11 and 0.08, respectively.

\* significant at  $p = 0.05$ ; \*\* significant at  $p = 0.01$

### Stress indices

The stress susceptibility index (SSI) values for grain yield ranged from 0.35 (HKR 127) to 1.55 (TR-2000-008), whereas indices of other genotypes with higher grain yield fell in between these values under high salinity (Table 3). The association between the SSI and grain yield in salinity revealed negative correlation ( $r = -0.43$ ). The lower values of SSI indicated lower differences in yield across stress and normal environments and hence more stability and indicate genotypes performing well under stress with sufficient plasticity to respond to the potential environment.

The stress tolerance index (STI) values for grain yield ranged from 0.07 (PR 118) to 1.09 (HKR 120), with the genotypes HKR 120, HKR 47 and CSR-RIL-197 exhibiting higher values of stress tolerance index (Table 3). The association between the STI and grain yield under salinity revealed a strong positive correlation ( $r = 0.81$ \*\*) thus indicating better utility of STI in identifying higher yielding genotypes across environments and prediction of stress tolerance as compared to SSI ( $r = -0.43$ \*\*) in these materials. Overall results indicate better utility of STI in identifying higher yielding genotypes across environments and prediction of stress tolerance in such materials as compared to SSI.

Table 3. Salinity stress susceptibility (SSI) and salinity stress tolerance (STI) indices for yield and its component traits in rice genotypes

| S. No. | Genotypes                  | SSI          |               |               |             | STI          |               |               |             |
|--------|----------------------------|--------------|---------------|---------------|-------------|--------------|---------------|---------------|-------------|
|        |                            | Plant height | Prod. tillers | Biolog. yield | Grain yield | Plant height | Prod. tillers | Biolog. yield | Grain yield |
| 1      | IR77674-3B-8-1-3-13-2-AJY2 | 1.12         | 0.90          | 1.42          | 1.31        | 0.63         | 0.83          | 0.43          | 0.61        |
| 2      | IR77674-3B-8-1-3-13-4-AJY2 | 0.89         | 1.20          | 1.06          | 1.12        | 0.66         | 0.66          | 0.26          | 0.44        |
| 3      | IR77674-3B-8-1-3-14-2-AJY3 | 0.93         | 0.71          | 1.00          | 1.11        | 0.63         | 0.68          | 0.32          | 0.60        |
| 4      | IR77674-3B-8-1-3-14-2-AJY4 | 0.87         | 0.74          | 1.15          | 1.22        | 0.72         | 0.87          | 0.26          | 0.43        |
| 5      | IR78806-B-B-16-1-2-2-AJY1  | 0.96         | 1.12          | 0.87          | 0.74        | 0.70         | 0.66          | 0.36          | 0.48        |
| 6      | IR60997-16-2-3-2-2R        | 0.73         | 0.47          | 0.95          | 1.00        | 0.68         | 0.74          | 0.44          | 0.66        |
| 7      | BCW 56                     | 1.49         | 1.57          | 1.30          | 1.38        | 1.01         | 0.52          | 1.05          | 0.59        |
| 8      | CSR-RIL(CSR 27/MI 48)-50   | 1.04         | 0.59          | 0.61          | 0.48        | 0.70         | 0.83          | 0.43          | 0.53        |
| 9      | CSR-RIL(CSR 27/MI 48)-197  | 1.01         | 0.57          | 0.80          | 0.71        | 0.81         | 0.77          | 0.60          | 0.79        |
| 10     | CSR-RIL(CSR 27/MI 48)-75   | 0.98         | 1.63          | 0.47          | 0.61        | 0.77         | 0.73          | 0.46          | 0.51        |
| 11     | CSR-RIL(CSR 27/MI 48)-192  | 0.33         | 1.29          | 0.64          | 0.37        | 0.99         | 0.84          | 0.29          | 0.25        |
| 12     | CSR-RIL(CSR 27/MI 48)-169  | 1.13         | 0.71          | 1.14          | 1.04        | 0.79         | 0.79          | 0.71          | 0.64        |
| 13     | CSR-RIL(CSR 27/MI 48)-170  | 1.28         | 0.83          | 0.44          | 0.92        | 0.59         | 0.74          | 0.27          | 0.19        |
| 14     | TR-2000-008                | 1.43         | 1.77          | 1.60          | 1.55        | 1.31         | 1.04          | 0.64          | 0.17        |
| 15     | NDR-359                    | 1.26         | 0.86          | 0.50          | 0.80        | 0.75         | 0.56          | 0.31          | 0.24        |
| 16     | PR 113                     | 0.85         | 0.96          | 0.97          | 1.06        | 0.64         | 0.71          | 0.24          | 0.09        |
| 17     | PR 114                     | 0.66         | 1.13          | 0.96          | 0.68        | 0.62         | 0.70          | 0.36          | 0.16        |
| 18     | PR 115                     | 1.11         | 0.74          | 0.94          | 0.56        | 0.55         | 0.68          | 0.35          | 0.24        |
| 19     | PR 116                     | 0.65         | 0.86          | 1.01          | 0.78        | 0.68         | 0.57          | 0.49          | 0.28        |
| 20     | PR 118                     | 0.29         | 0.96          | 0.71          | 0.87        | 0.59         | 0.62          | 0.29          | 0.07        |
| 21     | PAU 201                    | 0.83         | 0.77          | 0.94          | 0.69        | 0.57         | 0.71          | 0.36          | 0.27        |
| 22     | PR 120                     | 1.27         | 1.33          | 0.95          | 0.96        | 0.60         | 0.68          | 0.53          | 0.35        |
| 23     | PUSA 44                    | 1.12         | 0.90          | 1.34          | 1.24        | 0.55         | 0.64          | 0.27          | 0.07        |
| 24     | HKR 46                     | 1.21         | 0.80          | 0.82          | 0.52        | 0.60         | 0.61          | 0.37          | 0.26        |
| 25     | HKR 47                     | 1.24         | 0.51          | 0.81          | 1.09        | 0.74         | 0.71          | 0.79          | 0.84        |
| 26     | HKR 120                    | 0.68         | 0.90          | 1.04          | 0.95        | 0.76         | 0.61          | 1.00          | 1.09        |
| 27     | HKR 127                    | 1.18         | 0.55          | 0.68          | 0.35        | 0.73         | 0.62          | 0.57          | 0.48        |
| 28     | CSR 10                     | 1.18         | 0.83          | 0.59          | 0.45        | 0.53         | 0.91          | 0.31          | 0.15        |
| 29     | CSR 13                     | 1.09         | 0.78          | 0.95          | 0.67        | 0.66         | 0.79          | 0.56          | 0.27        |
| 30     | CSR 23                     | 0.51         | 1.23          | 0.96          | 0.54        | 0.98         | 1.05          | 0.76          | 0.56        |
| 31     | CSR 27                     | 1.07         | 1.30          | 1.08          | 0.59        | 0.83         | 0.54          | 0.52          | 0.58        |
| 32     | CSR 36                     | 1.04         | 0.95          | 0.93          | 0.50        | 0.78         | 0.81          | 0.96          | 0.63        |
| 33     | MI 48                      | 1.31         | 1.18          | 0.83          | 0.80        | 0.74         | 0.69          | 0.42          | 0.42        |
| 34     | VSR 156                    | 0.78         | 1.53          | 1.39          | 1.49        | 1.43         | 0.67          | 0.53          | 0.35        |
|        | Mean                       | 0.99         | 0.98          | 0.94          | 0.92        | 0.75         | 0.72          | 0.49          | 0.43        |
|        | Range                      | 0.29–1.49    | 0.44–1.60     | 0.53–1.43     | 0.24–1.05   |              |               |               |             |
|        |                            |              | 0.47–1.77     | 0.35–1.55     | 0.52–1.05   |              |               | 0.07–1.09     |             |

## Discussion

### *Effect of salinity on grain yield and yield attributing traits*

Grain yield per plant was the trait most affected by salinity. Reduction in grain yield and its component traits occurs through osmotic effects reducing the ability of plants to take up

water and causing reduced growth. These results are in agreement with the earlier reported findings of Mohammadi-Nejad et al. (2008), Yasseen et al. (2010), and Joseph and Jini (2010). In rice, plant height, total number of tillers, panicle length, grain weight per panicle, 1000-seed weight and quantity of grains decreased progressively with increase in salinity levels (Abdullah et al. 2001). Therefore, identification of plant genotypes with tolerance to salt and incorporation of desirable traits into economically useful crop plants will help overcome the effects of salinity on crop productivity. The phenotypic coefficient of variation (PCV) for all the traits indicated influence of environment on the manifestation of these traits. High magnitude (>15%) of genotypic as well as phenotypic coefficients of variation were observed for biological yield per plant, grain yield per plant and harvest index under both the normal and saline environments. Higher estimates of genetic advance (>25%) and heritability were observed for harvest index and biological yield under normal and saline environments. This suggests that the evaluated materials might provide higher response to selection for these traits exhibiting higher heritability along with higher genetic advance owing to their high transmissibility and variability.

#### *Correlation and path analysis*

The grain yield or economic yield in almost all crops is referred to as a complex character resulting from multiplicative interactions of several other characters termed as yield contributing components. Thus genetic architecture of grain yield in rice as well as other crops is based on the balance or overall net effect produced by various yield components directly or indirectly by interacting with one another. Therefore, identification of important yield components and information about their associations is very useful for developing efficient breeding strategies for evolving high yielding varieties. Grain yield per plant showed positive significant associations with plant height, total tillers per plant, productive tillers, panicle length, and biological yield and harvest index in normal environment. Under salinity stress, grain yield showed positive significant associations with biological yield and harvest index. These results clearly indicate that selection of genotype under normal and saline conditions should be entirely different. These results are in agreement with the findings of Raju et al. (2003), Ganapathy et al. (2006), Sabesan et al. (2009) and Chakraborty et al. (2010) in rice. The lower estimates of phenotypic correlation coefficient indicate that the relations were affected by environment at the phenotypic level and are in conformity of the findings of Jayasudha and Sharma (2010) and Anbanandan et al. (2009). The differential responses exhibited by the traits under salinity stress might be due to the complexity of the traits under stress. As grain yield is a complex trait governed by many traits with large environmental influence which along with salinity tolerance also being a complex trait governed by multi genes makes improvement of yield under salinity stress conditions a tough challenge for scientists.

A positive association between two traits warrants simultaneous improvement of both the traits while restricting selection to any one of the associated traits. Relative importance of the attributes should therefore be decided on the basis of higher correlation of the trait with grain yield. However, the correlation coefficient alone is inadequate to interpret the cause and effect relationships among the traits and ultimately with yield. Path analysis

technique furnishes a method portioning the correlation coefficients into direct and indirect effects providing information on the actual contribution of a trait on the yield. Direct positive effects of these traits on grain yield indicate their importance in determining complex characters and, therefore, should be accorded priority while making selections for improvements in grain yield. Panicle length exerted low but positive direct effect on grain yield, while plant height, total tillers per plant and productive tillers per plant exhibited negative direct effects on grain yield (Sajjad 1990).

#### *Stress indices*

Lower values of SSI indicate lower differences in yield across stress and normal environments and hence more stability thereby indicating better performing genotypes under stress and also possessing sufficient plasticity to respond to the potential environment. This suggested that the genotypes developed for salinity tolerance could exhibit higher adaptability and suitability, particularly under saline conditions. Porch (2006) has reported that several genotypes of common beans were superior for heat tolerance based on the stress indices and consistency of their reactions across environments. White and Singh (1991) reported low yield potential and low SSI scores of drought stress in bean indicated that selection for low SSI could identify stress tolerant lines but with low yield potential under non-stress conditions.

Association between the STI and grain yield in salinity revealed a strong positive correlation thus indicating better utility of STI in identifying higher yielding genotypes across environments and prediction of stress tolerance as compared to SSI in these materials. Overall results indicate better utility of STI in identifying higher yielding genotypes across environments and prediction of stress tolerance as compared to SSI in such materials. Genotypes experiencing minimum yield reduction under stress compared to normal conditions are distinguished by stress susceptibility index (Fischer and Maurer 1978). Lower values of stress susceptibility index (SSI) indicated lower differences in yield across stress and normal environments and hence more stability. This indicates genotypes performing well under stress and also have sufficient plasticity to respond to the potential environment. On the other hand, the higher values of stress tolerance index (STI) indicate superiority of genotypes having higher yield potential and stress tolerance as well (Fernandez 1993), thus making it a more desirable and helpful criterion in identifying and developing genotypes having both higher yield and stress tolerance.

The genotypes CSR-RIL-197, CSR 36, HKR 127 and HKR 120 showed better performance in normal and stress environments and could be useful in increasing productivity under saline stress environments. Correlations among grain yield and its contributing traits, direct and indirect effects of different traits on yield are useful in identification of the trait(s) strongly associated with yield under normal and salinity stress. The traits harvest index and biological yield, which are associated with yield emerged as the ideal traits for improvement through selection under saline stress environments and could be used to increase rice productivity under saline stress environments. The negative and significant association between the SSI and grain yield; positive and significant association between the STI and grain yield at salinity could be utilized by rice breeders in identifying higher



yielding genotypes having salt tolerance and sensitivity. The stress susceptible and tolerance indices suggest that genotypes developed for salinity tolerance could exhibit higher tolerance, adaptability and suitability.

### Acknowledgements

Authors express sincere thanks to the Director, Central Soil Salinity Research, Karnal, and the Indian Council of Agricultural Research, New Delhi, India and NPTC for encouragement and providing the facilities.

### References

- Abdullah, Z., Khan, M.A., Flowers, T.Z. 2001. Causes of sterility in seed set of rice under salinity stress. *Agron. J. Crop Sci.* **167**:25–32.
- Al-Jibouri, H.A., Miller, P.A., Robinson, H.F. 1958. Genotypic and environmental variance in an upland cotton crosses of inter-specific origin. *Agron. J.* **50**:633–637.
- Anbanandan, V., Saravanan, K., Sabesan, T. 2009. Variability, heritability and genetic advance in rice (*Oryza sativa* L.). *Int. J. Plant Sci.* **3**:61–63.
- Arshadullah, M., Rasheed, M., Zaidi, S.A.R. 2011. Salt tolerance of different rice cultivars for their salt tolerance under salt-affected soils. *Int. Res. J. Agric. Sci. Soil Sci.* **1**:183–184.
- Aslam, M., Qureshi, R.H., Mahmood, I.A., Perveen, S. 1993. Determination of critical toxicity levels of sodium and chloride in rice. *Pak. J. Soil Sci.* **8**:13–18.
- Chakraborty, S., Das, P.K., Guha, B., Kalyan, K.S., Barman, B. 2010. Quantitative genetic analysis for yield and yield components in boro rice (*Oryza sativa* L.). *Nat. Sci. Biol.* **2**:117–120.
- Dewey, D.R., Lu, K.H. 1959. A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron. J.* **51**:515–518.
- Fernandez, C.G.J. 1993. Effective selection criteria for assessing plant stress tolerance. In: Kuo, C.G. (ed.), *Adaptation of Food Crops to Temperature and Water Stress*. AVRDC, Shanhua, Taiwan, pp. 257–270.
- Fischer, R.A., Maurer, R. 1978. Drought resistance in spring wheat cultivars. I. Grain yield responses. *Aust. J. Agric. Res.* **29**:897–912.
- Flowers, T.J., Koyama, M.L., Flowers, S.A., Sudhakar, C., Singh, K.P., Yeo, A.R. 2000. QTL: Their place in engineering tolerance of rice to salinity. *J. Exp. Bot.* **51**:99–106.
- Ganapathy, S., Ganesh, S.K., Vivekanandan, P., Chandra Babu, R., Shanmugasundaram, P. 2006. Genetic variability and association analysis for drought tolerance, yield and its contributing traits in rice. In: *Plant Breeding in Post Genomics Era*. Second National Plant Breeding Congress. TNAU, Coimbatore, India, 1–3 March, p. 111.
- Gomez, K.A., Gomez, A.A. 1983. *Statistical Procedures for Agricultural Research*. John Wiley & Sons, New York, USA, 680 pp.
- Jayasudha, S., Sharma, D. 2010. Genetic parameters of variability, correlation and path-coefficient for grain yield and physiological traits in rice (*Oryza sativa* L.) under shallow lowland situation. *Electronic J. Plant Breed.* **1**:33–38.
- Joseph, B., Jini, D. 2010. Salinity induced programmed cell death in plants: Challenges and opportunities for salt-tolerant plants. *J. Plant Sci.* **5**:376–390.
- McIntosh, M.S. 1983. Analysis of combined experiments. *Agron. J.* **75**:153–155.
- Mishra, B. 2005. *More Crop per Drop*. Survey of Indian Agriculture. The Hindu Kasthuri Publishers, Chennai, India, pp. 41–46.
- Mohammadi-Nejad, G., Arzani, A., Rezai, A.M., Singh, R.K., Gregorio, G.B. 2008. Assessment of rice genotypes for salt tolerance using microsatellite markers associated with the *saltol* QTL. *African J. Biotech.* **7**:730–736.
- Panse, V.G., Sukhatme, P.V. 1967. *Statistical Methods for Agricultural Workers*. ICAR, New Delhi, India.

- Pearson, G.A., Ayers, A.D., Eberhard, D.L. 1966. Relative salt tolerance of rice during germination and early seedling development. *Soil Sci.* **102**:151–156.
- Porch, T.G. 2006. Application of stress indices for heat tolerance screening of common bean. *J. Agro. Crop Sci.* **192**:390–394.
- Raju, C.H.S., Rao, M.V.B., Sudarshanan, A. 2003. Association of physiological growth parameters in rice hybrids. *Madras Agric. J.* **90**:621–624.
- Sabesan, T., Suresh, R., Saravanan, K. 2009. Genetic variability and correlation for yield and grain quality characters of rice grown in coastal saline low land of Tamilnadu. *Electronic J. Plant Breed.* **1**:56–59.
- Sajjad, M.S. 1990. Correlations and path coefficient analysis of rice under controlled saline environment. *Pak. J. Agric. Res.* **11**:164–168.
- Sharma, S.K. 1986. Mechanisms of tolerance in rice varieties differing in sodicity tolerance. *Plant and Soil* **93**:141–145.
- Sharma, S.K. 2010. Salt tolerant crop varieties developed at CSSRI Karnal, India: A success story. <http://www.PlantStress.com>, 14 pp.
- Simmonds, N.W. 1983. Plant breeding: The state of the art. In: Genetic Engineering of Plants. An Agricultural Stress Susceptibility Index. New York, London, pp. 5–26.
- White, J.W., Singh, S.P. 1991. Breeding for adaptation to drought. In: van Schoonhoven, A., Voysest, O. (eds), Common Beans: Research for Crop Improvement. CAB Int. Wallingford, U.K. and CIAT, Cali, Colombia, pp. 501–560.
- Yamaguchi, T., Blumwald, E. 2005. Developing salt-tolerant crop plants: Challenges and opportunities. *Trends Plant Sci.* **10**:615–620.
- Yasseen, B.T., Abu-Al-Basal, M.A., Alhadi, F.A. 2010. An analysis of leaf growth under osmotic stress. *J. Plant Sci.* **5**:391–401.
- Zeng, L., Shannon, M.C., Lesch, S.M. 2001. Timing of salinity stress affects rice growth and yield components. *Agri. Water Manage.* **48**:191–206.

### Electronic Supplementary Material (ESM)

Electronic Supplementary Material (ESM) associated with this article can be found at the website of CRC at <http://www.akademai.com/content/120427/>

Electronic Supplementary *Table S1*. List of 34 rice genotypes and their pedigree/geographical location

Electronic Supplementary *Table S2*. Pooled analysis of variance (ANOVA) for grain yield and its attributing traits in control and saline ( $EC_{iw} \sim 10$  dS/m) environments in rice

Electronic Supplementary *Table S3*. Range, mean, coefficient of variation, heritability and genetic advance for grain yield and other attributes in rice in control and saline conditions