



Patterns and distribution of tobacco consumption in India: cross sectional multilevel evidence from the 1998-9 national family health survey

Citation

Subramanian, S V. 2004. "Patterns and Distribution of Tobacco Consumption in India: Cross Sectional Multilevel Evidence from the 1998-9 National Family Health Survey." *BMJ* 328 (7443) (April 3): 801-806. doi:10.1136/bmj.328.7443.801.

Published Version

doi:10.1136/bmj.328.7443.801

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:27002629>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Patterns and distribution of tobacco consumption in India: cross sectional multilevel evidence from the 1998-9 national family health survey

S V Subramanian, Shailen Nandy, Michelle Kelly, Dave Gordon, George Davey Smith

Abstract

Objective To investigate the demographic, socioeconomic, and geographical distribution of tobacco consumption in India.

Design Multilevel cross sectional analysis of the 1998-9 Indian national family health survey of 301 984 individuals in 92 447 households in 3215 villages in 440 districts in 26 states.

Setting Indian states.

Participants 301 984 adults (≥ 18 years).

Main outcome measures Dichotomous variable for smoking and chewing tobacco for each respondent (1 if yes, 0 if no) as well as a combined measure of whether an individual smokes, chews tobacco, or both.

Results Smoking and chewing tobacco are systematically associated with socioeconomic markers at the individual and household level. Individuals with no education are 2.69 times more likely to smoke and chew tobacco than those with postgraduate education. Households belonging to the lowest fifth of a standard of living index were 2.54 times more likely to consume tobacco than those in the highest fifth. Scheduled tribes (odds ratio 1.23, 95% confidence interval 1.18 to 1.29) and scheduled castes (1.19, 1.16 to 1.23) were more likely to consume tobacco than other caste groups. The socioeconomic differences are more marked for smoking than for chewing tobacco. Socioeconomic markers and demographic characteristics of individuals and households do not account fully for the differences at the level of state, district, and village in smoking and chewing tobacco, with state accounting for the bulk of the variation in tobacco consumption.

Conclusion The distribution of tobacco consumption is likely to maintain, and perhaps increase, the current considerable socioeconomic differentials in health in India. Interventions aimed at influencing change in tobacco consumption should consider the socioeconomic and geographical determinants of people's susceptibility to consume tobacco.

Introduction

Consumption of tobacco is a major risk factor for mortality.¹ Recent shifts in global tobacco consumption indicate that an estimated 930 million of the world's 1.1 billion smokers live in developing countries,² with

182 million in India alone.³ By 2020 tobacco consumption has been projected to account for 13% of all deaths in India.^{1 4}

Smoking is not only associated with lung cancer⁵ but is also linked to cardiovascular diseases, tuberculosis, and chronic respiratory diseases.⁶ Although 20% of total tobacco consumption in India is through cigarette smoking,¹ bidis (handrolled cigarettes that contain unprocessed tobacco) and hookahs are alternatives, with bidi smoking accounting for 40% of total tobacco consumption.^{1 3} Tobacco is also consumed, especially in India and South East Asian countries, through chewing (for example, paan masala, gutka, and mishri).^{7 8} Chewing tobacco is a risk factor for oral cancers.⁷ The annual incidence of oral cancer in men in India is estimated to be 10 per 100 000.⁹ Regardless of how tobacco is consumed, its adverse influence on disease and mortality among individuals and populations is clear.

Importantly, however, the distribution of tobacco consumption is not uniform. Tobacco consumption is often found to be disproportionately higher among lower socioeconomic groups.¹⁰ However, barring a few local studies,¹¹ little systematic investigation has been done into how tobacco consumption is socioeconomically and geographically distributed in India. The gaps in tobacco consumption need to be examined to see which people are most likely to consume tobacco and which areas are more likely to have higher tobacco consumption. Such analyses are critical for designing policies and interventions aimed at achieving overall reductions in tobacco consumption at the population level and at reducing the inequalities in susceptibility to consume tobacco.

We investigate how tobacco consumption (in its smoking and smokeless form) is distributed across a range of demographic and socioeconomic markers at individual and household level in India. Conditional on this distribution we also estimate the extent to which the prevalence of tobacco consumption varies between localities, districts, and states.

Methods

Sources of data

The analysis was based on a representative, cross sectional 1998-9 national family health survey of

See Editorial by Samarasinghe and Goonaratna

Department of Society, Human Development and Health, Harvard School of Public Health, 677 Huntington Avenue, KRESGE 7th floor, Boston MA 02115-6096, USA

S V Subramanian
assistant professor

School of Policy Studies, University of Bristol, Bristol Shailen Nandy
PhD student

Dave Gordon
professor of social justice

Social Science Research Unit, Institute of Education, University of London, London Michelle Kelly
PhD student

Department of Social Medicine, University of Bristol, Bristol George Davey Smith
professor of clinical epidemiology

Correspondence to: S V Subramanian svsubram@hsph.harvard.edu

BMJ 2004;328:801-6

Table 1 Descriptive information on the individual sample considered for the analytical multilevel models from the 1998-9 Indian national family health survey, showing the frequency of different exposure variables along with the counts and prevalence of tobacco smoking, chewing, and smoking and chewing across different exposure variables. Values are numbers (%) of study participants for each variable

Variable	Participants	Smoking	Chewing	Smoking and chewing
Living environment				
Large city	38 930 (12.9)	5 340 (13.7)	6 101 (15.7)	9 744 (25.0)
Small city	20 189 (6.7)	2 524 (12.5)	3 029 (15.0)	4 817 (23.9)
Town	42 304 (14.0)	6 380 (15.1)	7 633 (18.0)	11 985 (28.3)
Village	200 561 (66.4)	41 448 (20.7)	46 743 (23.3)	72 950 (36.4)
Sex				
Female	149 939 (49.7)	5 091 (3.4)	19 472 (13.0)	23 170 (15.5)
Male	152 045 (50.3)	50 601 (33.3)	44 034 (29.0)	76 326 (50.2)
Marital status				
Married or cohabiting	221 391 (73.3)	46 792 (21.1)	49 629 (22.4)	79 958 (36.1)
Single	53 416 (17.7)	4 876 (9.1)	6 857 (12.8)	9 799 (18.3)
Widowed	24 121 (8.0)	3 490 (14.5)	6 178 (25.6)	8 556 (35.5)
Divorced or separated	3 056 (1.0)	534 (17.5)	842 (27.6)	1 183 (38.7)
Religion				
Hindu	231 498 (76.7)	41 755 (18.0)	48 316 (20.9)	75 321 (32.5)
Muslim	35 304 (11.7)	7 537 (21.3)	6 601 (18.7)	12 104 (34.3)
Christian	19 125 (6.3)	4 798 (25.1)	5 718 (29.9)	8 238 (43.1)
Other	15 762 (5.2)	1 545 (9.8)	2 795 (17.7)	3 727 (23.6)
Missing	295 (0.1)	57 (19.3)	76 (25.8)	106 (35.9)
Social caste				
Other caste	118 978 (39.4)	18 863 (15.9)	18 578 (15.6)	32 227 (27.1)
Scheduled caste	48 350 (16.0)	10 399 (21.5)	11 174 (23.1)	17 862 (36.9)
Scheduled tribe	36 735 (12.2)	9 124 (24.8)	13 075 (35.6)	17 857 (48.6)
Other backward caste	83 598 (27.7)	14 510 (17.4)	17 612 (21.1)	26 688 (31.9)
No caste or missing	14 323 (4.7)	2 796 (19.5)	3 067 (21.4)	4 862 (33.9)
Education				
Illiterate	115 704 (38.3)	22 635 (19.6)	27 740 (24.0)	42 412 (36.7)
Primary school	50 456 (16.7)	12 241 (24.3)	13 093 (25.9)	20 764 (41.2)
Secondary school	87 649 (29.0)	16 059 (18.3)	16 893 (19.3)	27 232 (31.1)
Higher school	21 741 (7.2)	2 353 (10.8)	2 894 (13.3)	4 471 (20.6)
College	19 952 (6.6)	1 866 (9.4)	2 255 (11.3)	3 555 (17.8)
Postgraduate	6 482 (2.1)	538 (8.3)	631 (9.7)	1 062 (16.4)
Household standard of living index				
Lowest fifth	55 003 (18.2)	13 615 (24.8)	17 102 (31.1)	25 038 (45.5)
Second fifth	56 821 (18.8)	13 225 (23.3)	15 011 (26.4)	23 094 (40.6)
Third fifth	59 569 (19.7)	12 150 (20.4)	13 421 (22.5)	21 352 (35.8)
Fourth fifth	62 409 (20.7)	9 945 (15.9)	10 715 (17.2)	17 710 (28.4)
Highest fifth	68 182 (22.6)	6 757 (9.9)	7 257 (10.6)	12 302 (18.0)
Total	301 984 (100.0)	55 692 (18.4)	63 506 (21.0)	99 496 (32.9)

301 984 adults aged 18 and older, from 92 447 households from 26 Indian states.¹² The household data, obtained through an interview based structured questionnaire and answered by an available adult household member, provide a range of demographic and socioeconomic markers on all the members of the household, including information on smoking and chewing tobacco.¹² All households are geographically referenced to the primary sampling unit, district, and the state to which they belong. The primary sampling units (hereafter termed: local areas) are villages or groups of villages for rural areas, and wards or municipal localities for urban areas. The response rate to the survey ranged from 89% to 100%; in 24 of the 26 states it exceeded 94%.

Outcome measures

Our analysis used two dichotomous outcomes, based on the responses to the questions: "Does 'household member' chew paan masala or tobacco?" and "Does 'household member' smoke?" In addition, a combined measure of participants who smoke and chew tobacco

was constructed in order to assess the distribution of consuming any tobacco. In our sample the overall prevalence for smoking was 18.4% and for chewing 21.0%; the combined prevalence was 32.9%. Table 1 provides the descriptive characteristics of the outcome and exposure measures in the sample population considered for analysis.

Exposure measures

At the individual level we considered age (treated as continuous variable centred at its mean), sex, marital status, and educational attainment. At the household level we considered caste, religion, and a standard of living index based on material possessions. Caste status was based on the following mutually exclusive classification: scheduled caste, scheduled tribe, other backward caste; other caste; and no caste. "Scheduled caste" and "scheduled tribe" represents population groups identified by India's constitution as being marginal to the mainstream socioeconomic and political processes and, since 1951, are eligible for affirmative action. "Other backward caste" is another grouping of populations that are identified as socially and educationally backward. Since 1990 the other backward castes, while not sharing the constitutional affirmative action rights of scheduled castes and tribes are legally defined and covered by other legislative measures. "Other caste" is a residual category of people who are not scheduled caste or tribe, or other backward caste; "no caste" represents population groups for whom caste is not applicable (Muslims, Christians, or Buddhists, for example) and participants who did not report any caste affiliation in the survey. In general, the "other caste" category is considered to have higher social status, with the government of India designating the scheduled caste or tribe and other backward caste as socially and economically disadvantaged.¹² We divided religious affiliation into four categories: Hindu, Muslim, Christian, and other. We used the consumption and material possessions based on assets to create the standard of living index for households, weighted for the proportion of each possession at the all India level.¹³ Since the standard of living index is a constructed measure it does not have an absolute interpretation. Consequently, it is more appropriate to use this measure in categorical, hierarchically ordered fashion. We followed the convention of dividing the population into every fifth of the standard of living index for our analysis. Households were also characterised by whether they were located in a large city (a population of 1 million or more), a small city (population of 100 000 or more but less than 1 million), a town (population of less than 100 000), or villages and rural areas.

Statistical approach

We applied multilevel statistical procedures¹⁴ to model the variation in tobacco consumption according to the different analytical levels.¹⁵ Specifically we estimated the effect of the demographic and socioeconomic markers on tobacco consumption ("fixed parameters") and the variations in tobacco consumption in local areas, districts, and states that are not accounted for by individual and household demographic and socioeconomic markers. We calibrated a five level binary logistic model with a nested structure: 301 984 individuals (level 1) in 92 447 households (level 2) in

3215 local areas (level 3) in 440 districts (level 4) in 26 states (level 5). To calibrate models, we used the marginal quasi-likelihood (MQL) approximation with first order Taylor linearisation procedure. Model estimates are maximum likelihood based, derived by using the iterative generalised least squares algorithm, as implemented within the *MLwiN* program, version 1.10.0006.¹⁶

Results

Table 2 presents the odds ratios along with the 95% confidence interval derived from the fixed part of a multiple multilevel regression model calibrated for tobacco smoking, chewing, and smoking and chewing.

Smoking and demographic and socioeconomic markers

Age was positively associated with the probability of smoking; for a 10 year change in age the odds ratio related to smoking increased by 1.16. Men were considerably more likely to smoke than women (odds ratio 19.69). Marital status was also predictive of smoking: single, widowed, and divorced or separated people were less likely to smoke (odds ratios 0.32, 0.88, and 0.93, respectively), although the association was weak and imprecisely estimated for divorced and separated people. We also observed religion-based differences: Christians and the residual category of "other religion" were less likely to smoke than Muslims or Hindus. Caste status was also associated with smoking; in comparison to the other caste (the reference category), the scheduled tribe and scheduled caste were more likely to smoke. We observed a strong gradient between education and smoking, with the odds of being a smoker approximately three times higher in the educationally worst off group (illiterate people) than in the educationally best off group (people with postgraduate education). We observed a similar gradient between household standard of living and smoking, with individuals in the lowest fifth having an odds ratio of 2.5 of being smokers (compared with the highest fifth). The prevalence of smoking was greater in rural areas (odds ratio 1.19) and towns (odds ratio 1.14) than in large cities.

Chewing and demographic and socioeconomic markers

Age was positively associated with chewing (odds ratio 1.14 for a 10 year change), and men were more likely to chew than women (odds ratio 3.27). Single people were less likely to chew, but both widowed and separated or divorced people were more likely to chew than married people. Muslims were more likely to chew (odds ratio 1.15) and Christians and other religions less likely to chew (odds ratios 0.76 and 0.85, respectively) compared with Hindus. Scheduled caste and scheduled tribe groups were more likely to chew (odds ratios 1.15 and 1.11, respectively) than other caste. We observed a strong gradient between education and chewing; the odds of chewing in the educationally worst off group was 1.84 times that of people with postgraduate education. A similar household standard of living gradient became apparent for chewing; the odds of chewing in the lowest fifth was nearly twice that of the highest fifth. Chewing prevalence did not differ substantially between different types of urban and rural areas.

The pattern in demographic and socioeconomic inequalities for smoking and chewing combined was similar to those observed for the separate analysis of smoking and chewing. In general the socioeconomic gap in combined tobacco consumption was smaller than for smoking alone and larger than for chewing alone.

Distribution of tobacco consumption across local areas, districts, and states

Table 3 provides the variance estimates for each of the levels for two models—the first does not account for age, sex, marital status, education, religion, caste, household standard of living and urban or rural status; the second does account for these demographic and socioeconomic markers. The variance estimates in table 3 show that socioeconomic markers at the individual and household level do not entirely explain the differences in the prevalence of smoking, chewing, and their combined prevalence by local areas, districts

Table 2 Odds ratios with 95% confidence intervals from the fixed part of a multivariable five level binomial logit model that is calibrated for tobacco smoking, chewing, and smoking and chewing, conditional on random effects at the level of state, district, local area, and household

	Smoking	Chewing	Smoking and chewing
Age (years)	1.16 (1.15 to 1.16)	1.14 (1.13 to 1.15)	1.23 (1.22 to 1.23)
Living environment			
Large city*	1.00	1.00	1.00
Small city	1.06 (0.94 to 1.19)	0.91 (0.81 to 1.02)	1.02 (0.92 to 1.14)
Town	1.14 (1.03 to 1.26)	0.97 (0.88 to 1.07)	1.12 (1.02 to 1.22)
Village	1.19 (1.08 to 1.31)	0.92 (0.84 to 1.01)	1.09 (1.00 to 1.18)
Sex			
Female*	1.00	1.00	1.00
Male	19.69 (19.20 to 20.19)	3.27 (3.21 to 3.32)	8.53 (8.37 to 8.70)
Marital status			
Married or cohabiting*	1.00	1.00	1.00
Single	0.32 (0.31 to 0.33)	0.55 (0.53 to 0.56)	0.36 (0.36 to 0.37)
Widow	0.88 (0.85 to 0.92)	1.20 (1.16 to 1.24)	1.01 (0.98 to 1.05)
Divorced or separated	0.93 (0.84 to 1.03)	1.23 (1.14 to 1.32)	1.20 (1.11 to 1.30)
Religion			
Hindu*	1.00	1.00	1.00
Muslim	1.03 (0.98 to 1.08)	1.15 (1.11 to 1.20)	1.15 (1.10 to 1.20)
Christian	0.88 (0.82 to 0.94)	0.76 (0.72 to 0.81)	0.76 (0.72 to 0.81)
Other	0.54 (0.50 to 0.58)	0.85 (0.80 to 0.90)	0.66 (0.63 to 0.70)
Missing	0.77 (0.55 to 1.08)	0.95 (0.71 to 1.27)	0.80 (0.60 to 1.08)
Social caste			
Other caste*	1.00	1.00	1.00
Scheduled caste	1.16 (1.12 to 1.20)	1.15 (1.11 to 1.18)	1.19 (1.16 to 1.23)
Scheduled tribe	1.27 (1.20 to 1.33)	1.11 (1.06 to 1.16)	1.23 (1.18 to 1.29)
Other backward caste	1.03 (0.99 to 1.06)	1.00 (0.98 to 1.03)	1.01 (0.98 to 1.04)
No caste or missing	1.07 (1.00 to 1.14)	0.96 (0.91 to 1.01)	0.98 (0.92 to 1.03)
Education			
Postgraduate*	1.00	1.00	1.00
College	1.22 (1.12 to 1.33)	1.16 (1.08 to 1.24)	1.18 (1.10 to 1.26)
Higher school	1.40 (1.29 to 1.53)	1.30 (1.21 to 1.40)	1.38 (1.29 to 1.48)
Secondary school	1.99 (1.84 to 2.15)	1.54 (1.44 to 1.65)	1.85 (1.74 to 1.98)
Primary school	2.56 (2.36 to 2.77)	1.73 (1.62 to 1.85)	2.37 (2.21 to 2.53)
Illiterate	2.94 (2.71 to 3.19)	1.84 (1.72 to 1.97)	2.69 (2.51 to 2.87)
Household standard of living index			
Highest fifth*	1.00	1.00	1.00
Fourth fifth	1.43 (1.37 to 1.48)	1.30 (1.25 to 1.34)	1.43 (1.38 to 1.47)
Third fifth	1.78 (1.71 to 1.85)	1.49 (1.44 to 1.55)	1.76 (1.70 to 1.82)
Second fifth	2.14 (2.05 to 2.24)	1.67 (1.61 to 1.74)	2.10 (2.02 to 2.18)
Lowest fifth	2.43 (2.32 to 2.55)	1.90 (1.82 to 1.97)	2.54 (2.44 to 2.64)

*Reference categories in each of the categorical exposure variable. The reference category is a 40 year old, postgraduate, married Hindu woman, belonging to the "other caste" group and to the highest fifth (quintile) of household standard of living index and who lives in a large city. For this group the predicted probability of smoking and chewing was 0.8% and 6.4%, respectively, and the combined probability 4.4%.

Table 3 Variance estimates from a five level binomial logit model, before and after adjusting for demographic and socioeconomic markers of individuals or households, at the level of the state, district, local areas, and household, for tobacco smoking, chewing, and smoking and chewing

Levels	Smoking		Chewing		Smoking and chewing	
	Before*	After†	Before*	After†	Before*	After†
Level 5: states	0.243	0.452	0.726	1.010	0.326	0.613
Level 4: districts	0.063	0.102	0.148	0.146	0.057	0.082
Level 3: local areas	0.151	0.132	0.151	0.128	0.124	0.106
Level 2: households	0.045	0.839	0.480	0.652	0.158	0.560
Dispersion	0.859	0.667	0.725	0.665	0.839	0.772

Note: All variance variables were significant at the 0.001 level.

*Gives the variance estimates at different levels before accounting for demographic and socioeconomic markers of individuals or households.

†Gives the variance estimates at different levels after accounting for demographic and socioeconomic markers of individuals or households.

and states. The differences between those geographical areas (especially at state level) increased once account was taken of the composition of the population residing in them. After accounting for individual or household demographic and socioeconomic markers, the bulk of the remaining variation lies at the household and state level. The other spatial levels of local areas and districts seem to matter less. Although the patterning of the multilevel variation was not substantially different across smoking, chewing, and smoking and chewing, the exact magnitude varied: the variation between states was largest for chewing, followed by smoking and chewing, and smoking.

Table 4 Odds ratios for smoking, chewing, and smoking and chewing for the different Indian states, based on the state level residuals estimated from a five level multiple binomial logit model

States	Sample size	Smoking		Chewing		Smoking and chewing	
		Odds ratio	Rank	Odds ratio	Rank	Odds ratio	Rank
Andhra Pradesh	11 668	0.90	15	0.47	6	0.59	6
Assam	10 531	0.80	12	2.62	23	1.71	22
Bihar	21 260	0.61	6	1.42	18	0.78	13
Goa	5 397	0.67	9	0.52	10	0.54	3
Gujarat	12 514	0.63	7	0.74	13	0.66	9
Haryana	9 317	1.72	21	0.36	3	0.70	11
Himachal Pradesh	10 460	1.43	20	0.35	2	0.59	5
Jammu	10 595	2.26	23	0.31	1	0.76	12
Karnataka	14 580	0.61	5	0.65	12	0.63	8
Kerala	10 055	0.79	11	0.51	9	0.59	4
Madhya Pradesh	21 853	0.66	8	1.26	17	0.91	16
Maharashtra	19 488	0.35	1	1.43	19	0.87	15
Manipur	5 547	2.04	22	2.33	22	2.28	23
Meghalaya	3 519	2.44	24	1.26	16	2.61	24
Mizoram	4 335	8.11	26	29.83	26	18.43	26
Nagaland	2 961	0.94	16	1.83	20	1.01	18
Orissa	15 072	0.46	2	2.70	24	1.44	20
Punjab	10 232	0.54	3	0.39	4	0.35	1
Rajasthan	21 815	0.97	17	0.49	7	0.61	7
Sikkim	4 074	0.76	10	1.95	21	1.45	21
Tamil Nadu	15 118	0.57	4	0.50	8	0.48	2
West Bengal	14 891	1.09	18	0.82	14	0.94	17
Uttar Pradesh	29 489	0.88	14	1.05	15	0.86	14
New Delhi	9 222	1.13	19	0.54	11	0.70	10
Arunachal Pradesh	4 045	0.85	13	4.09	25	2.72	25
Tripura	3 946	2.45	25	0.43	5	1.05	19

The model additionally included the demographic and socioeconomic markers of individuals or households in the fixed part and variance component at the level of households, local areas, districts, and states, in addition to the dispersion variable.

The reference point for the calculation of the state odds ratios is the all India log of the odds of smoking, chewing, and smoking and chewing.

Table 4 presents the predicted odds ratio for the different states after taking demographic and socioeconomic markers into account, using the all India prevalence as a reference. For smoking, Maharashtra has the lowest odds ratio (0.35); for chewing, Jammu (0.31); and Punjab for smoking and chewing (0.35). Mizoram has the highest odds ratio for smoking, chewing, and smoking and chewing. The burden of tobacco consumption in India falls disproportionately on Mizoram¹⁷ and the other northeastern states.

Figures 1 and 2 map the unconditional (crude) and conditional (model based) prevalence for combined tobacco consumption. In both maps a “natural break algorithm” was used to divide the states of India into four groups.¹⁸ A strong geographical pattern is evident in the prevalence for tobacco consumption both before (fig 1) and after (fig 2) controlling for demographic and socioeconomic markers at the individual level. In the northeastern states a greater proportion of the adult population smoke and chew tobacco than in the southern and western states of India. As is evident from figure 2, much but not all of the state variation in tobacco consumption observed in figure 1 is accounted for by the differences in the socioeconomic circumstances of the people who live in them and by other variations attributable to households, local areas, and districts.

Discussion

Tobacco consumption in India has a distinct socioeconomic and spatial distribution; the worse off population groups are at greater risk of consuming tobacco. Our analyses show the extent to which tobacco consumption is distributed across population subgroups and across states in India.

Differential socioeconomic and geographical susceptibilities to tobacco consumption

Men are more likely to consume tobacco than women. A strong gradient with regards to education and standard of living is apparent. Higher levels of education and standard of living are inversely related to the probability of smoking and chewing; the gradient is stronger for smoking. The relation between these socioeconomic markers and tobacco consumption is similar to relations observed in developed countries.¹⁰ Further, the caste based differences in tobacco consumption show the persistent effect of caste as a key axis along which health and other wellbeing outcomes are stratified, over and above the adverse effects of low education and an index of material standard of living. In addition, the distribution of tobacco consumption by marital status is contrary to what has been observed in developed countries, where marriage is seen to have a protective effect.¹⁰ Importantly, the large differences observed between states in tobacco consumption, even after controlling for the demographic and socioeconomic markers at individual or household level, highlight the potential importance of the state context in influencing this behaviour.¹⁹

Limitations of the study

In addition to the general limits to drawing causal inferences based on cross sectional, observational data, one caveat that is pertinent to our analysis relates to the extent to which the observed magnitude of socioeconomic gap reflects “actual” gaps compared with

“reporting” gaps, especially since formal and informal social conventions related to tobacco consumption can influence reporting patterns.

Moreover, data were available only on overall tobacco smoking. Differences in the socioeconomic gradient in the use of manufactured cigarettes (higher among people with more education) and bidis (higher among people with less education) have been shown to




Fig 1 Crude prevalence of adults aged 18 and above who smoke or chew tobacco in 1998-9, by Indian state. The term “crude” means unadjusted prevalence and is computed as number of individuals who smoke and chew tobacco divided by the total number of individuals, in each state, and expressed as percentages




Fig 2 Model based predicted proportion of adults aged 18 and above who smoke or chew tobacco in 1998-9 by Indian state after controlling for demographic and socioeconomic markers at the individual level and for variation in tobacco consumption between households, local areas, and districts. The term “model based” means conditional prevalence and is based on model based, residual, state level differences in smoking and chewing after accounting for between-individual differences in tobacco consumption that are due to age, sex, marital status, caste, religion, education, standard of living, and urban and rural differences, and after taking account of within-state variation attributable to the level of households, local areas, and districts, and expressed as percentages

What is already known on this topic

Tobacco consumption is a key adverse health influence in South Asia in general and India in particular

Little is known about how the consumption of tobacco is distributed in India

What this study adds

Tobacco consumption in India is heterogeneous, with higher prevalence rates observed for population groups with low social caste, education, and standard of living

The prevalence of smoking and tobacco chewing shows marked geographical differences (at the level of villages, districts, and states), even after controlling for the individual and household demographic and socioeconomic markers

States account for the bulk of spatial variation in tobacco consumption

exist.¹¹ Smoking bidis is more strongly related to lung and oral cancer than smoking manufactured cigarettes,^{20, 21} and the preponderance of bidi smoking among less advantaged socioeconomic groups will tend to exacerbate health inequalities.

Conclusion

The presence of socioeconomic and demographic gradients in tobacco consumption in India counter the perspective of “poor versus non-poor” that is often adopted when examining health differentials in societies that are labelled as poor or low income. Furthermore, the variations in the prevalence of these risk behaviours by states, even after accounting for the type of individuals and households that reside in these states, show the potential importance of the state context in influencing tobacco consumption. The state of Maharashtra took the first legislative steps to discourage tobacco consumption through prohibitions on use and sale,²² and this state has the lowest current overall smoking (table 4).

The current large socioeconomic and geographical gap in tobacco consumption in India is likely to feed into substantial, and perhaps increasing, socioeconomic differentials in the health of adults over the future decades. A need exists to document and monitor such inequalities in tobacco consumption in India systematically, to understand their determinants better, and to provide an evidence base for public health interventions that takes account of differences at state level as well as between population groups, in tobacco consumption.

The research from which this paper borrows part of its data was commissioned and funded by the UK Department for International Development (DFID). The views expressed in this paper do not represent the official position of UK DFID. We acknowledge the support of Macro International (www.measuredhs.com) for providing us access to use the data of the 1998-9 Indian national family health survey. We are also grateful for the extremely helpful comments by the BMJ editorial committee and an independent reviewer.

Contributors: SVS conceived the study, analysed and interpreted the data, wrote and edited the manuscript, and is guarantor. SN, MK, and DG contributed to data preparation, interpretation of the results and to the editing of the manuscript. GDS contributed to the interpretation of the results and to the editing of the manuscript.

Funding: None.

Competing interests: None declared.

Ethical approval: This research is based on a secondary analysis of a public dataset, with the survey being conducted by international agencies under a cooperative agreement with the national governments. Ethical approval has been granted by the ethics committee at the School for Policy Studies at the University of Bristol, which conforms to guidance of the Social Research Association (SRA) for secondary research of this nature.

- 1 World Health Organization. *Tobacco or health: a global status report*. Geneva: WHO, 1997.
- 2 Jha P, Ranson MK, Nguyen SN, Yach D. Estimates of global and regional smoking prevalence in 1995 by age and sex. *Am J Public Health* 2002;92:1002-6.
- 3 Shinkhada R, Peabody JW. Tobacco control in India. *Bull WHO* 2003;81:48-52.
- 4 Kumar S. India steps up anti-tobacco measures. *Lancet* 2000;356:1089.
- 5 International Agency for Research on Cancer. *Tobacco smoking: monographs on the evaluation of carcinogenic risk of chemicals to humans*. Lyons: IARC, 1985.
- 6 Gajalakshmi V, Peto R, Kanaka TS, Jha P. Smoking and mortality from tuberculosis and other diseases in India: retrospective study of 43000 adult male deaths and 35000 controls. *Lancet* 2003;362:507-15.
- 7 Critchley JA, Unal B. Health effects associated with smokeless tobacco: a systematic review. *Thorax* 2003;58:435-43.
- 8 Hirayama T. An epidemiological study of oral and pharyngeal cancers in central and south east Asia. *Bull WHO* 1966;34:41-69.
- 9 Moore SR, Johnson NW, Pierce AM, Wilson D. The epidemiology of mouth cancer: a review of global incidence. *Oral Dis* 2000;6:65-74.

- 10 Giovino GA, Henningfield JE, Tomar SL. Epidemiology of tobacco use and dependence. *Epidemiol Rev* 1995;17:48-65.
- 11 Gupta PC. Survey of sociodemographic characteristics of tobacco use among 99,598 individuals in Bombay, India using handheld computers. *Tobacco Control* 1996;5:114-120.
- 12 International Institute of Population Sciences. *National family health survey 1998-99*. Mumbai: IIPS, 2000.
- 13 Davey Smith G, Gordon D, Kelly M, Nandy S, Subramanian SV. Inequalities in health in India: the methodological construction of indices and measures: report submitted to UK Department for International Development, 2004.
- 14 Goldstein H. *Multilevel statistical models*. 3rd ed. London: Arnold, 2003.
- 15 Subramanian SV, Jones K, Duncan C. Multilevel methods for public health research. In: Kawachi I, Berkman LF, eds. *Neighborhoods and health*. New York: Oxford University Press, 2003:65-111.
- 16 Rasbash J, Browne W, Goldstein H, Yang M, Plewis I, Healy M, et al. *A user's guide to MLwiN, Version 2.1*. London: Multilevel Models Project, Institute of Education, University of London, 2000.
- 17 Chaturvedi HK, Phukan RK, Zoramtharga K, Hazarika NC, Mahanta J. Tobacco use in Mizoram, India: sociodemographic differences in pattern. *Southeast Asian J Trop Med Public Health* 1998;29:66-70.
- 18 Jenks GF, Caspall FC. Error in choroplethic maps: definition, measurement, reduction. *Ann Assoc Am Geograph* 1971;61:217-44.
- 19 Joseph DT. Antitobacco campaign in Maharashtra, India: achievements and perspectives. In: Gupta PC, Hammer JEI, Murti PR, eds. *Control of tobacco-related cancers and other diseases. Proceedings of an international symposium, TIFR, Bombay, January 15-19, 1990*. Bombay: Oxford University Press, 1992:345-7.
- 20 Gajalakshmi V, Hung RJ, Mathew A, Varghese C, Brennan P, Boffetta P. Tobacco smoking and chewing, alcohol drinking and lung cancer risk among men in southern India. *Int J Cancer* 2003;107:441-7.
- 21 Znaor A, Brennan P, Gajalakshmi V, Mathew A, Shanta V, Varghese C, et al. Independent and combined effects of tobacco smoking, chewing and alcohol drinking on the risk of oral, pharyngeal and esophageal cancer in Indian men. *Int J Cancer* 2003;105:681-6.
- 22 Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology* 2003;8:419-31.

(Accepted 10 February 2004)

A memorable call

A house call in Nepal's Himalayas

The young man came into the Himalayan Rescue Association aid post in Manang at midday. He asked me to go and see a man in a nearby village who had been sick since the day before and was unable to move. I stuffed my medical gear into a backpack and followed the man to the horse that he had brought along, and, after a two hour ride, we finally arrived at his village.

We walked through the narrow alleys to the patient's house, where a cow greeted us on the ground floor, and climbed the wooden staircase to the terrace to find a middle aged man lying still on a mattress. Though in agony, he tried to smile on seeing me. The history of his ailment and subsequent physical examination made me think that he had either acute cholecystitis or liver abscess. He looked pretty sick, so I started treatment with intravenous fluids and antibiotics, followed by pethidine, which made him doze off for a while.

He needed an abdominal scan, and the nearest place where it was available was four days' walk away. Helicopter evacuation seemed unfeasible as this is too expensive for most Nepalese. The only viable option was to fly him out from the local seasonal airfield, which was two hours' walk away. After telephone calls to the city, we found out there was a plane flying in tomorrow. This was our best bet. By now it was apparent that my patient needed medical supervision until he could be flown out to the city hospital, and I sent the young man back to the aid post to get extra drug supplies.



STEVE MCCURRY/MAGNUM

Strong winds started in the evening, and it snowed late into the night. This had stopped by the morning, but the whole valley was now covered with thick fog, and it seemed unlikely that the flight would come in. After a few hours, however, the fog started to clear, so we put the patient in a large basket, his legs dangling from two holes in its side, and started carrying him to the airport. The flight came in when we were halfway along the trail. Two of us rushed ahead to ask the pilots to wait, and the party finally made it just as the pilots, impatient to leave, were preparing to take off. To save a few precious minutes, the sick man was lifted over the perimeter fence of the airfield.

A few days later, I got the news that my patient had had liver abscess and was recuperating in a hospital in Kathmandu.

Puncho Gurung *volunteer physician, Himalayan Rescue Association, Nepal*

We welcome articles up to 600 words on topics such as *A memorable patient, A paper that changed my practice, My most unfortunate mistake*, or any other piece conveying instruction, pathos, or humour. Please submit the article on <http://submit.bmj.com> Permission is needed from the patient or a relative if an identifiable patient is referred to. We also welcome contributions for "Endpieces," consisting of quotations of up to 80 words (but most are considerably shorter) from any source, ancient or modern, which have appealed to the reader.