



Published in final edited form as:

J Am Chem Soc. 2013 May 22; 135(20): 7567–7571. doi:10.1021/ja400659s.

Pd(II)-Catalyzed *Ortho*- or *Meta*-C–H Olefination of Phenol Derivatives

Hui-Xiong Dai[†], Gang Li[†], Xing-Guo Zhang[†], Antonia F. Stepan[‡], and Jin-Quan Yu[†]

Jin-Quan Yu: yu200@scripps.edu

[†]Department of Chemistry, The Scripps Research Institute, 10550 N. Torrey Pines Road, La Jolla, California 92037, United States;

[‡]Neuroscience Medicinal Chemistry, Pfizer Global Research and Development, Cambridge, Massachusetts 02139, United States

Abstract

A combination of weakly coordinating auxiliaries and ligand acceleration allows for the development of both *ortho*- and *meta*-selective C–H olefination of phenol derivatives. These reactions demonstrate the feasibility of directing C–H functionalizations when functional groups are distal to target C–H bonds. The *meta*-C–H functionalization of electron-rich phenol derivatives is unprecedented and orthogonal to previous electrophilic substitution of phenols in terms of regioselectivity. These methods are also applied to functionalize α -phenoxyacetic acids, a fibrate class of drug scaffolds.

1. Introduction

Directed C–H activation via 5-, 6-membered cyclopalladation has recently emerged as a versatile tool for developing synthetic transformations.¹ However, activation of C–H bonds that are more than five bonds away from the chelating atom are still rare due to the difficulty of forming larger membered palladacycles.² Alternative strategies relying on the proximity between remote C–H bonds and highly reactive species such as radical or meta-oxo species have been reported.³ In contrast to the directed metalation processes, these strategies do not produce systematic and predictable structural patterns at this stage of development. On the other hand, directed metalation of C–H bonds at certain geometric positions that are resistant to the assembly of cyclic transition states due to ring strain has limited success, as exemplified by the lack of examples of directed *meta*-C–H activation of arenes until recently.^{2b} Development of *meta*-C–H activation of broad range of synthetically useful substrates will significantly enhance the versatility of directed C–H activation reactions in synthetic applications.

Phenol and their derivatives have drawn intensive efforts from the community of C–H functionalizations due to their broad synthetic utility. Early studies by Rawal and Miura used the phenol to direct arylation of biphenyls with ArI and Pd(0)/PPh₃ catalysts.⁴ Bedford and others succeeded in using catalytic amount of phosphites as directing groups to promote Rh(I)-catalyzed *ortho*-arylation of phenols (Fig 1).⁵ Hartwig reported Ir(I)-catalyzed borylation of hydrosilyl ethers of phenols.⁶ Recent emergence of C–H activation reactions with Pd(II),⁷ Rh(III),⁸ Ru(II)⁹ catalysts using weakly coordinating carbonyls have led to a

Correspondence to: Jin-Quan Yu, yu200@scripps.edu.

Supporting Information Available. Detailed experimental procedures, characterization of new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

range of *ortho*-C–H functionalization reactions of phenol derivatives (Fig 1). Due to the strong electron-donating ability of the phenoxy group, *meta*-selective C–H activation is especially challenging and yet synthetically enabling.

Herein we report two approaches for achieving *ortho*- and *meta*-C–H olefination of phenol derivatives respectively through remote C–H activation promoted by weak coordination and ligand acceleration (Fig 2). Removal of the acetic acid auxiliaries affords synthetically valuable *ortho*- or *meta*-olefinated phenols. This protocol was successfully applied to the parent α -phenoxyacetic acids of drug molecules fenofibrate, clofibrate and etofibrate (Fig 3).

2. Results and Discussion

We initially aimed to develop *ortho*-C–H functionalization methods for late-stage diversification of α -phenoxyacetic acids, important pharmacophores found in the fibrate class of lipid-lowering agents (Figure 3).¹⁰ Despite a number of elegant directing groups previously developed for the *ortho*-C–H functionalizations of phenols,^{5–9} the distance between the C–H bonds and the functional groups in α -phenoxyacetic acids presents a challenge. Although we have developed various *ortho*-C–H functionalizations of phenylacetic acids via a relatively distal directing group, the *ortho*- C–H bonds and the chelating atom of the carboxyl group in α -phenoxyacetic acids are further apart (six bonds away) as shown in ciprofibrate (Figure 3). The presence of the α -oxygen atom in the side chain could also be problematic as Pd(II) could form a bis-chelation with this oxygen atom and the directing group, which could prevent Pd center from aligning with C–H bonds with a relatively small dihedral angle, a requisite for facile C–H activation.

Encouraged by our recently observed ligand acceleration,¹¹ we began to identify ligands that can cooperate with the weak coordination of the distal carbonyl in COOK moiety to promote the C–H activation reactivity. Guided by our previous C–H olefination of phenyl acetic acids¹¹ and early olefination reactions,¹² we established conditions for the *ortho*-olefination of α -phenoxyacetic acids. Thus, substrate **1a** was reacted with 2 equiv of ethyl acrylate under 1 atm of O₂ in the presence of 5 mol% of Pd(OAc)₂, 10 mol% Boc-Val-OH, 2 equiv of KHCO₃ in *t*-amyl alcohol at 90 °C for 24 hours to give 90% of the mono-olefinated product **3a** (Table 1). The high mono-selectivity can be attributed to the steric buttress imparted by the α,α -dimethyl groups. Interestingly, the absence of the α,α -dimethyl groups resulted in a significant loss of reactivity leading to poor yields, presumably due to the loss of favorable Thrope-Ingle effect (less than 10%, see SI). The replacement of O₂ with air dropped the yield to 34%.

This reaction tolerates electron-donating substituents such as methyl and methoxyl (**3b–d**). The naphthalene ring in example **3i** is selectively olefinated at the 2-position. Olefination also proceeds smoothly with a wide range of electron deficient arenes, including those with fluoro (**3e–g**), chloro (**3h**), trifluoromethyl (**3i**, **3j**) and keto functionality (**3k**). This method can be used to directly functionalize medicinally useful α -phenoxyacetic acids or prepare *ortho*-substituted phenols after removal of the acetic acid directing groups (Scheme 1). Notably, substrates **1h** and **1k** are derived from drug molecules clofibrate and fenofibrate respectively.

Prompted by our recent development of *meta*-selective C–H olefination reactions of hydrocinnamic acids,^{2b} we wondered if the end-on coordinating nitrile template can be modified to direct *meta*-C–H olefination of α -phenoxyacetic acids. We were pleased to find that olefination of **4a** proceeded with high *meta*-selectivity under our previous conditions to give the mono- and di-olefinated products in 60% and 29% yields respectively (Table 2).

Meta-substitutions of phenol derivatives complement the *ortho*- and *para*- electrophilic substitutions, thus providing a new strategy to construct densely substituted arenes. This reaction is compatible with electron-donating groups (**5b–f**), as well as electron-withdrawing groups (**5g–m**). High mono-selectivities were obtained with *ortho*-substituted substrates (**5b**, **5e**, **5g**, **5j**, **5l**). Importantly the template overrides the electronic influence of substituents on the arenes. The regioselectivity observed with **5l** is most striking as olefination occurs at the position that is *meta* to alkoxy and *para* to trifluoromethyl group. The use of a directing group to govern *meta*-selectivity is fundamentally a different approach compared to previously developed *meta*-C–H functionalizations where the electronic or steric bias of arene substrates plays a decisive role.^{13–17}

To further demonstrate the synthetic utility of this method, we performed *meta*-olefination of an *ortho*-brominated substrate **5p** (Table 3). We were pleased to find that a wide range of olefins including disubstituted olefins are reactive. Coupling of **4p** with *trans*-2-butenolate gave the desired product **5s** stereospecifically. It is worth noting that di-substituted olefins are generally not compatible with directed C–H olefination reactions; only rare examples have been shown to date.^{2b} Additionally, the bromine functionality is a highly versatile handle that allows for subsequent transformations. Finally, the nitrile template can be removed by hydrolysis^{2b} to give the medicinally useful *meta*-substituted α -phenoxyacetic acids. The acetic acid moiety attached to the phenoxy can then be removed by treatment with DPPA to afford the *meta*-substituted phenol (Scheme 1).

Extensive studies on C–H olefination have shown that electron-deficient olefins are in general more effective coupling partners while electron-rich olefins tend to react with Pd(II) catalyst via the Wacker oxidation pathway with rare exceptional examples.¹⁸ To examine the scope of this *meta*-selective olefination reaction with respect to the olefins, we tested alkenes and styrenes as the coupling partners and found only those styrenes with an electron-withdrawing group attached to the arenes reacted to some extent to give the desired product in moderate yields (Table 4). Further improvement of the C–H olefination with electron-rich olefins such as simple styrene and hexane will require the modification of the ligand.

While the detailed mechanism and origin of the *meta*-selectivity remains to be elucidated by computational, kinetic as well as structural studies, We have also observed a significant isotope effect ($k_H/k_D = 3.8$) of the nitrile-directed *meta*-selective C–H olefination with analogous substrates (see supporting information), suggesting that the C–H cleavage may be involved in the rate-determining step. A tentative catalytic cycle can also be proposed based on our previous studies (Fig 4).

3. Conclusion

In summary, we have developed two new approaches for *ortho*- and *meta*-C–H functionalization of phenol derivatives. These C–H activation reactions feature rare cyclometalation processes of C–H bonds directed by distal chelating atoms (six bonds away). The selective functionalizations of phenol derivatives at the *meta*-positions are unprecedented and especially useful as the regioselectivity is orthogonal to the electrophilic substitution. The methods are also applied to functionalize α -phenoxyacetic acids, a fibrates class of drug scaffolds.

4. Experimental Section

4.1. General Information

All commercial reagents were purchased from Sigma-Aldrich, Fluka, Alfa Aesar, TCI, Oakwood, and Acros of the highest purity grade. They were used without further purification unless specified. Palladium acetate, silver acetate was purchased from Sigma-Aldrich. The amino acid ligands were bought from Novabiochem, Bachem and Sigma-Aldrich. 2,2'-azanediylidibenzonitrile was prepared by literature methods.¹⁹ ¹H and ¹³C NMR spectra were recorded on Bruker AV 400, Varian Inova 400 (400 MHz and 100 MHz, respectively), Bruker DRX 500 (500 MHz and 125 MHz, respectively) and Bruker DRX 600 (600 MHz and 150 MHz, respectively) instruments. The peaks were internally referenced to TMS (0.00 ppm) or residual undeuterated solvent signal. The following abbreviations were used to explain multiplicities: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, and br = broad. High resolution mass spectra were recorded at the Center for Mass Spectrometry, The Scripps Research Institute.

4.2 General procedure for Pd(II)-catalyzed *ortho*-C–H olefination of α -phenoxyacetic acids (1a–1l)

A 50 mL Schlenk-type tube (with a Teflon high pressure valve and side arm) equipped with a magnetic stir bar was charged with acid **1** (0.20 mmol, 1.0 equiv.), Pd(OAc)₂ (2.3 mg, 0.010 mmol, 5 mol %), Boc-Val-OH (4.3 mg, 0.020 mmol, 10 mol %), KHCO₃ (40 mg, 0.40 mmol, 2.0 equiv.), ethyl acrylate **2a** (0.40 mmol, 2.0 equiv.), and *t*-AmylOH (2.0 mL). The reaction tube was evacuated and backfilled with O₂ (5-times, balloon) and heated to 90 °C for 24 hours under vigorous stirring. The reaction vessel was then cooled to 0 °C in an ice bath. A 2.0 N HCl solution (5 mL) was then added, and the mixture was extracted with EtOAc (3 × 20 mL). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The resulting residue was purified by preparative TLC using hexanes/EtOAc as the eluent to afford the product (**3a–3l**).

4.3 General procedure for Pd(II)-catalyzed *meta*-C–H olefination of phenol derivative (4a–4p)

A 35 mL sealed tube (with a Teflon cap) equipped with a magnetic stir bar was charged with amide **4** (0.10 mmol, 1.0 equiv.), Pd(OAc)₂ (2.3 mg, 0.010 mmol, 10 mol %), Ac-Gly-OH (2.4 mg, 0.020 mmol, 20 mol %) and AgOAc (50 mg, 0.30 mmol, 3.0 equiv.). HFIP (0.60 mL) was added to the mixture, followed by ethyl acrylate **2a** (2.0 equiv.) and then another 1.0 mL of HFIP. The tube was then capped and submerged into a pre-heated 90 °C oil bath. The reaction was stirred for 24 h and cooled down to room temperature. The crude reaction mixture was diluted with EtOAc (2 mL) and filtered through a short pad of Celite. The sealed tube and Celite pad were washed with an additional 15 mL of EtOAc. The filtrate was concentrated *in vacuo*, and the resulting residue was purified by preparative TLC using hexanes/EtOAc as the eluent. The positional selectivity was determined by ¹H NMR analysis of the unpurified reaction mixture.

4.4 Procedure for removal of template²⁰

To a 50 mL of flask, **3a** (111.2 mg, 0.4 mmol) was dissolved in anhydrous toluene (5 mL) and DMF (0.5 mL), then Et₃N (46.4 mg, 0.4 mmol) and DPPA (110 mg, 0.4 mmol) were added and the mixture was refluxed for 3h. 30 mL of water was added and continue to reflux for 2h. After the mixture cool to room temperature, the solution was acidified with 2N HCl solution (5 mL) and extracted with EtOAc (3 × 5 mL). The combined organic phases were washed with brine, dried over anhydrous MgSO₄ and evaporated. The residue was purified via column chromatography on silica with hexane and EtOAc (10:1) as eluents to afford

product **A**²¹ in 71% yield (54.3 mg). ¹H NMR (400 MHz, CDCl₃) δ: 8.05 (d, *J* = 16.2 Hz, 1H), 7.47 (d, *J* = 8.0 Hz, 1H), 7.26–7.21 (m, 1H), 6.93–6.85 (m, 2H), 6.80 (br, 1H), 6.65 (d, *J* = 16.2 Hz, 1H), 4.30 (q, *J* = 7.2 Hz, 2H), 1.35 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (600 MHz, CDCl₃) δ: 169.3, 158.3, 141.5, 132.2, 130.1, 122.6, 121.5, 119.2, 117.3, 61.6, 15.2.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge The Scripps Research Institute, the National Institutes of Health (NIGMS 1 R01 GM102265-01), and Eli Lilly for financial support. X.-G.Z. is a visiting scholar from Wenzhou University and was sponsored by the China Scholarship Council and the National Natural Science Foundation of China (21002070). We thank Li Wan for experimental assistance.

References

- (a) Chen X, Engle KM, Wang DH, Yu JQ. *Angew Chem, Int Ed.* 2009; 48:5094.(b) Daugulis O, Do HQ, Shabashov D. *Acc Chem Res.* 2009; 42:1074. [PubMed: 19552413] (c) Lyons TW, Sanford MS. *Chem Rev.* 2010; 110:1147. [PubMed: 20078038] (d) Engle KM, Mei TS, Wasa M, Yu JQ. *Acc Chem Res.* 2012; 45:788. [PubMed: 22166158]
- (a) Li JJ, Giri R, Yu JQ. *Tetrahedron.* 2008; 64:6979.(b) Leow D, Li G, Mei TS, Yu JQ. *Nature.* 2012; 486:518. [PubMed: 22739317]
- For early discussion and other approaches for remote C–H activation: Breslow R. *Acc Chem Res.* 1980; 13:170.Schwarz H. *Acc Chem Res.* 1989; 22:282.Das S, Incarvito CD, Crabtree RH, Brudvig GW. *Science.* 2006; 312:1941. [PubMed: 16809537]
- (a) Hennings DD, Iwasa S, Rawal VH. *J Org Chem.* 1997; 62:2. [PubMed: 11671356] (b) Satoh T, Kawamura Y, Miura M, Nomura M. *Angew Chem, Int Ed.* 1997; 36:1740.(c) Miura M, Tsuda T, Satoh T, Nomura M. *Chem Lett.* 1997:1103.
- (a) Bedford RB, Coles SJ, Hursthouse MB, Limmert ME. *Angew Chem Int Ed.* 2003; 42:112.(b) Oi S, Watanabe S, Fukita S, Inoue Y. *Tetrahedron Lett.* 2003; 44:8665.(c) Bedford RB, Limmert ME. *J Org Chem.* 2003; 68:8669. [PubMed: 14575501]
- Ir(I)-catalyzed C–H activation of hydrosilyl ether of Phenols: Boebel TA, Hartwig JF. *J Am Chem Soc.* 2008; 130:7534. [PubMed: 18494474]
- Pd(II)-catalyzed C–H functionalizations of Phenol derivatives: Huang C, Gevorgyan V. *J Am Chem Soc.* 2009; 131:10844. [PubMed: 19722665] Zhao X, Yeung CS, Dong VM. *J Am Chem Soc.* 2010; 132:5837. [PubMed: 20359220] Xiao B, Fu Y, Xu J, Gong TJ, Dai JJ, Yi J, Liu L. *J Am Chem Soc.* 2010; 132:468. [PubMed: 20020760] Huang C, Chattopadhyay B, Gevorgyan V. *J Am Chem Soc.* 2011; 133:12406. [PubMed: 21766826] Huang C, Ghavtadze N, Chattopadhyay B, Gevorgyan v. *J Am Chem Soc.* 2011; 133:17630. [PubMed: 21999512] Xiao B, Gong TJ, Liu ZJ, Liu JH, Luo DF, Xu J, Liu L. *J Am Chem Soc.* 2011; 133:9250. [PubMed: 21609019] John A, Nicholas KM. *J Org Chem.* 2012; 77:5600. [PubMed: 22663545]
- Rh(III)-catalyzed C–H activation of Phenol derivatives: Gong TJ, Xiao B, Liu ZJ, Wan J, Xu J, Luo DF, Fu Y, Liu L. *Org Lett.* 2011; 13:3235. [PubMed: 21604790]
- Ru(II)-catalyzed C–H activation of Phenol derivatives: Ackermann L, Diers E, Manvar A. *Org Lett.* 2012; 14:1154. [PubMed: 22313092]
- (a) Razavi B, Song W, Cooper WJ, Greaves J, Jeong J. *J Phys Chem.* 2009; 113:1287.(b) Chuang S, Velkov T, Horne J, Wielens J, Chalmers DK, Porter CJH, Scanlon MJ. *J Med Chem.* 2009; 52:5344. [PubMed: 19663428]
- (a) Engle KM, Wang DH, Yu JQ. *J Am Chem Soc.* 2010; 132:14137. [PubMed: 20853838] (b) Engle KM, Wang DH, Yu JQ. *Angew Chem Int Ed.* 2010; 49:6169.
- (a) Moritan I, Fujiwara Y. *Tetrahedron Lett.* 1967; 8:1119.(b) Fujiwara Y, Noritani I, Danno S, Asano R, Teranishi S. *J Am Chem Soc.* 1969; 91:7166.(c) Jia C, Kitamura T, Fujiwara Y. *Acc Chem Res.* 2001; 34:63.

13. Cho J-Y, Tse MK, Holmes D, Maleczka RE Jr, Smith MR III. *Science*. 2002; 295:305. [PubMed: 11719693]
14. Ishiyama T, Takagi J, Ishida K, Miyaura N, Anastasi NR, Hartwig JF. *J Am Chem Soc*. 2002; 124:390. [PubMed: 11792205]
15. Zhang YH, Shi BF, Yu JQ. *J Am Chem Soc*. 2009; 131:5072. [PubMed: 19296661]
16. Phipps RJ, Gaunt MJ. *Science*. 2009; 323:1593. [PubMed: 19299616]
17. Saidi O, Marae J, Ledger AEW, Liu PM, Mahon MF, Kociok-Köhn G, Whittlesey MK, Frost CG. *J Am Chem Soc*. 2011; 133:19298. [PubMed: 22047022]
18. Wang DH, Engle KM, Shi BF, Yu JQ. *Science*. 2010; 327:315. [PubMed: 19965380]
19. Gorvin JH. *J Chem Soc, Perkin Trans1*. 1988:1331.
20. Mirk D, Waldvogel SR. *Tetrahedron Lett*. 2004; 45:7911.
21. Park JO, Youn SW. *Org Lett*. 2010; 12:2258. [PubMed: 20429502]




Figure 1.
Previous *ortho*-C–H Functionalizations of Phenol




Figure 2.
Remote *ortho*- and *meta*-C-H Functionalizations




Figure 3.
Drug Molecules Based on α -Phenoxyacetic Acids






Figure 4.
Catalytic Cycle of *meta*-C-H Olefination

**Scheme 1.**

The removal of Directing Group^a

^a Reaction conditions: **3a** (0.4 mmol), **DPPA** (0.4 mmol), Et₃N (0.4 mmol), toluene (5 mL), DMF (0.5 mL), refluxed 3h. **DPPA**: diphenylphosphoryl azide. Yield: 71%.

Table 1


Pd(II)-Catalyzed *ortho*-Olefination of Phenol Derivatives^{a,b}

^aReaction conditions: **1a-1l** (0.1 mmol), **2a** (0.2 mmol), Pd(OAc)₂ (5 mol %), Boc-Val-OH (10 mol %), KHCO₃ (0.2 mmol), O₂ (1 atm), *t*-AmylOH (1 mL), 90 °C, 24h.

^bIsolated yield.

^cPd(OAc)₂ (10 mol %), Boc-Val-OH (20 mol %).

Table 2


Pd(II)-Catalyzed *meta*-Olefination of Phenol Derivatives^{a,b,c}

^aReaction conditions: **4a-4o** (0.1 mmol), **2a** (0.2 mmol), Pd(OAc)₂ (10 mol %), Ac-Gly-OH (20 mol %), AgOAc (0.3 mmol), HFIP (1 mL), 90 °C, 24h.

^bIsolated yield.

^cRegioselectivity were determined by ¹H NMR analysis of the crude product and confirmed by one-dimensional selective NOESY experiments; the variance is estimated to be within 5%.

Table 3


meta-Olefination of 2-Bromo-Phenol Derivatives^{a,b,c}

^aReaction conditions: **4p** (0.1 mmol), **2a–2f** (0.2 mmol), Pd(OAc)₂ (10 mol %), Ac-Gly-OH (20 mol %), AgOAc (0.3 mmol), HFIP (1 mL), 90 °C, 24h.

^bIsolated yield.

^cRegioselectivity were determined by ¹H NMR analysis of the crude product and confirmed by one-dimensional selective NOESY experiments; the variance is estimated to be within 5%.

Table 4

meta-Olefination with Styrenes^{a,b,c}

^aReaction conditions: **4b** (0.1 mmol), **7a-7c** (0.2 mmol), $\text{Pd}(\text{OAc})_2$ (10 mol %), Ac-Gly-OH (30 mol %), AgOAc (0.3 mmol), HFIP (1 mL), 90 °C, 24h.

^bIsolated yield.

^cRegioselectivity were determined by ^1H NMR analysis of the crude product and confirmed by one-dimensional selective NOESY experiments; the variance is estimated to be within 5%.

^dAc-Gly-OH (20 mol%).

^e30 h.

^fOnly one major regioisomer was observed.

^gNMR yield.