

Pectinase Production with Waste Materials by Local Water Isolate of *Aspergillus fumigatus* Strain 2101 and Purification and Characterization of the Enzyme

Yerel Su İzolatı *Aspergillus Fumigatus* 2101 Suşu Tarafından Atık Malzemeler ile Pektinaz Üretimi ve Enzimin Saflaştırılması ve Karakterizasyonu

Research Article

Hacer Yamancı¹, Kader Poturcu¹, İsmail Özmen^{1*}, Hacı Halil Bıyık²

¹Department of Chemistry, Suleyman Demirel University, Isparta, Turkey.

²Department of Biology, Adnan Menderes University, Aydın, Turkey.

ABSTRACT

In this study, pectin lyase (PL) was obtained from *Aspergillus fumigatus* strain 2101 which was isolated from local water sources. PL was purified using ammonium sulphate precipitation, gel filtration and ion-exchange chromatography. After these purification steps, 12.4 purification fold was obtained. The optimum pH and temperature of PL were 8.0 and 40°C at 70 min, respectively. The molecular weight of PL was 11.5 kDa. Maximum PL activity was obtained with fruit pulps in submerged culture medium. Stability assays showed that the PL was stable till 20 months at 4°C. The highest PL activity was obtained using citrus pectin as carbon source. PL was strongly activated by FeSO₄, FeCl₃ and ascorbic acid.

Key Words

Pectin lyase, *Aspergillus*, chromatography, inhibition, ions.

ÖZET

Bu çalışmada, pektin liyaz (PL) yerel su kaynaklarından izole edilen *Aspergillus fumigatus* 2101 suşundan elde edildi. PL, amonyum sülfat çöktürmesi, jel filtrasyon ve iyon değişim kromatografisi kullanılarak saflaştırıldı. Bu saflaştırma basamaklarından sonra, 12.4 saflaştırma katsayısı elde edildi. PL'nin optimum pH ve sıcaklığı sırasıyla 8.0 ve 40°C'de 70 dakikadır. PL'nin molekül ağırlığı 11.5 kDa'dır. Sıvı besi ortamında maksimum PL aktivitesi meyve pulpları ile elde edilmiştir. Stabilité deneyleri, PL'nin 4°C'de 20 aya kadar kararlı olduğunu göstermiştir. En yüksek PL aktivitesi karbon kaynağı olarak sitrus pectin kullanıldığında elde edilmiştir. PL, FeSO₄, FeCl₃ ve askorbik asit tarafından güçlü bir şekilde aktive edilmiştir.

Anahtar Kelimeler

Pektin liyaz, *Aspergillus*, kromatografi, inhibisyon, iyonlar.

Article History: Received: Jan 8, 2016; Revised: Mar 16, 2016; Accepted: Mar 20, 2016; Available Online: Apr 01, 2016.

DOI: 10.15671/HJBC.20164417567

Correspondence to: İ. Özmen, Department of Chemistry, Suleyman Demirel University, Isparta, Turkey.

Tel: +90 246 211 3811

Fax: +90 246 237 1106

E-Mail: ismailozmen@sdu.edu.tr

INTRODUCTION

Pectin is complex colloidal acid polysaccharide presents in plant primary cell wall and middle lamella [1]. Pectin includes homogalacturonan (HG), xylogalacturonan (XGA) rhamnogalacturonan I (RG I), rhamnogalacturonan II (RG II) parts. HG is the smooth region of the pectin, it consists α -1,4-D-galacturonic acid residues. XGA includes β -1,3-D-xylose residues which are linked to galacturonyl residues of HG. α -1,2-rhamnosyl and galactosyluronic acid residues are present in RG I parts of pectin. RG II includes approximately nine α -1,4-D-galactosyluronic acid residues. XGA, RG I, RG II form hairy region of the pectin [2-4].

Pectinases are upcoming enzymes and among the first enzyme groups used in homes [5]. Pectinases are produced by many microorganisms such as bacteria, yeast, fungi and plants [6]. *Aspergillus* and *Penicillium* genus are preferred for industrial applications [7]. Pectinase is general name used for enzymes that catalyze hydrolysis of glycosidic bonds in pectin polysaccharides [8]. Pectinases can be divided 3 major groups: protopectinase, pectinesterases, depolymerizing enzymes [5].

Protopectinase solubilizes protopectin and constitutes highly polymerized soluble pectin. Pectinesterases catalyze removal of methoxyl group from pectin [5]. Depolymerizing enzymes are divided 2 sub-groups: hydrolases and lyases. Hydrolases consist of endo- and exo-polygalacturonase. Lyases include endo-, exo-pectate lyase and endo-pectin lyase (endo-PL) [9].

The objective of this study is to produce of endo-pectin lyase in submerged culture medium from *Aspergillus fumigatus* strain 2101 isolated from local water sources. Effects of waste materials, for maximum PL production in submerged culture were investigated. Purification studies with ammonium sulphate, chromatographic techniques and characterization studies of PL were also performed. Effects of metal ions and several chemicals were determined. Stability tests of PL were carried out at 4°C.

MATERIALS AND METHOD

Chemicals and Reagents

Citrus pectin, yeast extract, DEAE-cellulose, Sephadex G-100 and Coomassie Brilliant Blue were purchased from Sigma-Aldrich. Ethyl alcohol, Na_2HPO_4 , NaCl, HCl, $(\text{NH}_4)_2\text{SO}_4$ were purchased from Merck. All chemicals were analytical grade. *Aspergillus fumigatus* strain 2101 was procured Adnan Menderes University Arts and Science Faculty, Department of Biology, Section of Molecular Biology.

Identification of Fungi

In our study, DNA isolation was made by classical phenol chloroform method. Purity and concentration measurements were made by nonodrop spectrophotometer (Thermo). PCR reactions were carried out at 95°C 5 min, 35 cycle with 94°C 1 min, 55°C 1 min and 72°C 1 min, and then 72°C 7 min. 5X Firepol Master Mix was used in reactions with ITS primers. Electrophoresis was made in agarose gel (1.2%). For sequencing, samples were sent to MacroGen Europe. Sequence data was processed with Blastn. Sequence alignment and neighbour joining tree construction was made by using MEGA 6.06 software. For aligning sequences ClustalW was used which was in MEGA software.

Culture Conditions of Fungi

Aspergillus fumigatus strain 2101 was isolated from local water sources in Aydın, Turkey and it was used for PL production. First of all, *Aspergillus fumigatus* strain 2101 was grown on potato dextrose agar and incubated at 35°C for 7 days. It was stored at -80°C for further usage. Submerged culture for PL production was prepared according to Sandri method with some modifications [10].

Maximum Enzyme Production with Waste Materials

For maximum enzyme production, different waste materials were used as carbon sources. Apricot, apple, peach, sour cherry were used as fruit pulps. Fruit pulps (I), wheat bran (II), fruit pulps+wheat bran (III) and citrus pectin (IV) were added different flask cultures and incubation

was performed in rotating shaker. Crude enzyme was taken every flask, centrifuged and enzyme activity was measured. The best carbon source and optimum time for enzyme production in submerged culture were determined between 6-168 hours.

Enzyme Purification Steps

Flask was taken from rotating shaker when incubation time ended. Centrifugation was carried out at 8000 rpm for 25 min. After centrifugation, crude enzyme was partially purified with $(\text{NH}_4)_2\text{SO}_4$. Precipitation was carried out between 0-95% ranges. Then, gel filtration column (1x15 cm) was pre-equilibrated with phosphate buffer (0.05 M, pH 8.0) and 0-95% precipitates were loaded on to column. Elution was performed with the same buffer and enzyme was pooled. Ion-exchange column was pre-equilibrated and pooled enzyme taken from gel filtration column was loaded on to ion-exchange column (1x15 cm) with 1.5 and 2.0 M NaCl gradient. Elution was performed with phosphate buffer (0.05 M, pH 8.0) and fractions were collected and pooled. After every purification steps, enzyme activity and protein amounts of PL were measured.

PL Activity Tests

PL activity was performed according to Albersheim methods with some modifications [11]. Optical density of 4,5-unsaturated oligogalacturonides were measured at 235 nm with UV-vis spectrophotometers (Shimadzu, UV-mini-1240 Spectrophotometers). Enzyme activity was defined in terms of μmole unsaturated 4,5-oligogalacturonides released per min and molar extinction coefficient value of the product was used as $5550 \text{ M}^{-1}\text{cm}^{-1}$. All experiments were carried out triplicate.

Molecular Weight Determination of PL

Molecular weight of PL and its purity were determined with SDS-PAGE gel electrophoresis method. 12% resolving gel and 5% stacking gel were used. The electrophoresis was run 30 min at 50 V and 200 min at 100 V. After running, methanol (50%, (v/v)), Coomassie Brilliant Blue

R-250 (0.1% (w/v)) and acetic acid (50%, (v/v)) were used for staining. Destaining was performed with the same solution used for staining without Coomassie Brilliant Blue [12,13].

Protein Determination

Protein concentration was determined according to Bradford method, bovine serum albumin (BSA) was used as the standard [14]. All experiments were carried out triplicate.

Determination of Optimum pH and Temperature

Citrate-phosphate buffer (pH 4.0-6.0), phosphate buffer (pH 6.0-8.0) and carbonate buffer (pH 8.0-9.0) were used with 0.05 M concentration. Optimum pH for PL was determined with these buffer solutions.

To determine optimum temperature and time for PL, the experiments were carried out between 40-100°C with different time intervals (to 100 from 10 min). All experiments were carried out triplicate.

Effects of Chemical Substances on PL Activity

Effects of chemical substances on PL such as BaCl_2 , HgCl_2 , ZnSO_4 , FeCl_3 , FeSO_4 , EDTA, CaCl_2 , CoCl_2 , LiCl_3 , MgCl_2 , MgSO_4 , NaF, $\text{K}_4[\text{Fe}(\text{CN})_6]$, KCl, KMnO_4 , NaN_3 , NaCl, ascorbic acid, glycine, SDS, L-Cysteine, citric acid monohydrate, NaH_2AsO_4 , Triton X-100, $(\text{NH}_4)_2\text{SO}_4$, L-Arginine monohydrate, L-Asparagine were determined. All experiments were carried out triplicate.

Storage Tests of PL

Purified enzyme was stored in refrigerator at 4°C for 98 days. 500 μL PL was taken at special time ranges and enzyme activity was measured.

Substrate Specificity Tests of PL

To perform substrate specificity of PL against citrus pectin, peptone and PDA; substrate solutions were prepared. Enzyme was incubated with these substrates at 40°C 40 min. Substrate preference of PL was determined.

AATTCACTCTCCTCCGGCATCGATCGAGGTCACCTTAGAAAATAAAGTTGGGTGTCGGCT
 GGCGCCGGCCGGGCCTACAGAGCAGGTGACAAAGCCCCATACGCTCGAGGACCGGACGC
 GGTGCCGCCGCTGCCTTTCGGGGCCCGTCCCCGGGAGAGGGGGACGGGGGCCAACACA
 CAAGCCGTGCTTGAGGGCAGCAATGACGCTCGGACAGGCATGCCCCCGGAATACCAGG
 GGGCGCAATGTGCGTTCAAAGACTCGATGATTCACTGAATTCTGCAATTCACATTACTTAT
 CGCATTTTCGCTGCGTTCTTCATCGATGCCGGAACCAAGAGATCCGTTGTTGAAAGTTTAA
 CTGATTACGATAATCAACTCAGACTGCATACTTTCAGAACAGCGTTCATGTTGGGGTCTTC
 GGCGGGCGCGGGCCCGGGGGCGCAAGGCCTCCCCGGCGGCCGTCGAAACGGCGGGCCCG
 CCGAAGCAACAAGGTACGATAGACACGGGTGGGAGGTTGGACCCAGAGGGCCCTCACTC
 GGTAATGATCCTTCCGCAGGTCCCCTAACGGAAGGGTCATTCCCGAG

Figure 1. Nucleotide sequence of *Aspergillus fumigatus* strain 2101.

RESULTS AND DISCUSSION

Nucleotide Sequence of *Aspergillus fumigatus* Strain 2101

According to phylogenetic tree of our microorganism, fungus studied in this report was defined as *Aspergillus fumigatus* strain 2101 and Figure 1 shows that nucleotide sequence of *Aspergillus fumigatus* strain 2101.

Maximum Enzyme Production with Waste Materials

To recycle and reuse of waste materials, fruit pulps (I), wheat bran (II), fruit pulps+wheat bran (III) and citrus pectin (IV) were used as a carbon source in submerged culture media for enzyme production. The weight of fruit pulps in all submerged culture media was taken in equal amounts. PL showed maximum activity in submerged culture which included fruit pulps whereas submerged culture medium contained citrus pectin gave the lowest enzyme activity values. So, fruit pulps (I) were the best carbon sources in submerged culture media for PL production. After this submerged culture media optimization with carbon sources, we used fruit pulps in all submerged cultures (Figure 2).

Enzyme Purification

PL was purified with three steps. It was partially purified with $(\text{NH}_4)_2\text{SO}_4$ fractionation. Then, PL was purified with gel filtration and ion-exchange

chromatography. It was obtained 12.4 purification fold, after 3 steps purification. Purification steps of PL are shown in Table 1.

Molecular Weight Determination of PL

The homogeneity of PL was validated by SDS-PAGE electrophoresis. Molecular weight of PL was determined by plotting log of molecular weight of markers versus their Rf values. Molecular weight of PL was found 11.5 kDa (Figure 3).

In previous studies, molecular mass of PL secreted by *Penicillium italicum* was 22 kDa [21]. Molecular weight of PL from *Bacillus sp.* DT7 was 106 kDa [21,22]. Molecular weight of PL secreted by *Aspergillus flavus* was 38 kDa [23]. PL-overproducing hybrid was obtained by protoplast fusion between mutant pectinolytic *Aspergillus flavipes* and *Aspergillus niveus* CH-Y-1043 had 30 kDa molecular weight [19].

Optimum pH and Temperature

PL showed activity between pH 4.0-9.0 with different buffers. PL had maximum activity at pH 8.0 with phosphate buffer. PL also gave high activity values at pH 6.0 and pH 7.0 with phosphate buffer and their results were similar with pH 8.0. PL gave low activity values at pH 4.0 and 8.0 with citrate-phosphate and carbonate buffers, respectively. PL produced in our laboratory can be used at industrial scale with wide pH ranges




Figure 2. Effects of waste materials on PL activity in submerged culture medium (fruit pulps + wheat bran (I), wheat bran (II), fruit pulps (III) and citrus pectin (IV)).

Table 1. Purification steps of PL secreted by *Aspergillus fumigatus* strain 2101.

Purification steps	Total activity (EU/ml)	Total protein	Specific Activity (EU/mg)	Purification fold
Crude Extract	0.07	0.98	0.071	1
Ammonium sulphate precipitation (0-95%)	0.81	6.30	0.13	1.83
Gel filtration chromatography	0.52	2.26	0.23	3.24
Ion-exchange chromatography	0.39	0.44	0.88	12.4

(pH 4.0-9.0). Optimum pH results of PL are demonstrated in Figure 4. PL lost its activity in citrate-phosphate buffer for pH 4.0; 5.0; 6.0; 41.10%, 9.53 %, 15.96%; in phosphate buffer for pH 6.0; 7.0; 5.51%; 9.53%; in carbonate buffer for pH 8.0; 9.0; 48.31%; 4.24%, respectively.

PL showed activity between 40 to 90°C and had maximum activity at 40°C for 70 min, at 50°C for 10 min, at 60°C for 30 min, at 70°C for 30 min, at 80°C for 10 min, at 90°C for 10 min. The enzyme obtained in this study showed activity at 90°C till 100 min and didn't lose its activity largely as far as 100 min at 80°C. It also showed high activity values at 50°C. Optimum temperature results of PL are shown in Figure 5.

In the literature, PL secreted by *Debaryomycesflavus nepalensis* had maximum activity at pH 6.4 [24]. Pectinase from *Streptomyces sp.-QG-11-3* was optimally active at pH 3.0 in citrate-phosphate buffer and it retained more than 85% its activity between pH 2.0 to 9.0 [25].

In the previous studies, maximum PL activity was obtained at 35°C from *Debaryomycesflavus nepalensis* [24]. PL from *Penicillium italicum* completely inactivated after 10 min at 80°C [26]. *Paenibacillus amylolyticus* endophytic strain had maximum PL activity at 40°C and pH 7.9 [20]. PL secreted by *Aspergillus sp.* CH-Y-1043 showed maximum activity at pH and temperature values 8.5-8.8 and 40-45°C, respectively [2]. Optimum




Figure 3. SDS-PAGE analysis of purified PL. Lane 1 marker, Lane 2, 3 pooled PL after ion-exchange chromatography.




Figure 4. pH results of PL between pH 4.0-9.0.




Figure 5. PL activity at different temperatures between 10 to 100 minutes.




Figure 6. Storage results of PL at 4°C.




Figure 7. Substrate specificity results of PL.

pH and temperature for recombinant PL from a *Penicillium purpurogenum* was 6.0 (McIlvaine buffer (0.1 M citric acid/0.2 M disodium hydrogen phosphate) and 50°C, respectively [27].

Storage Results of PL

To determine storage stability of PL at 4°C, it was stored in refrigerator. PL lost 32% of its starting activity at 98 days. At the end of 20 months, the PL conserved 60% of its activity. Storage results of PL are shown in Figure 6.

Substrate Specificity Results of PL

Substrate specificity results of PL against citrus pectin, peptone and PDA were graphed in Figure 7. PL gave the best result against citrus pectin as a substrate.

In the literature, PL from *Paenibacillus amylolyticus* was used for substrate specificity tests. Citrus pectin, apple pectin, polygalacturonic

acid were used as a substrate and PL was more active on citrus pectin than other substrates [20].

Effects of Chemical Substances on PL Activity

Effects of metal ions BaCl₂, HgCl, ZnSO₄, FeCl₃·6H₂O, FeSO₄, CaCl₂, CoCl₂, LiCl₃, MgCl₂, MgSO₄, NaF, K₄[Fe(CN)₆], KCl, KMnO₄, NaN₃, NaCl; amino acids and the other chemicals were summarized in Table 2. PL activity was reached up 306.192% at 6.0×10⁻³ M concentration with BaCl₂, 655.39% at 1.33×10⁻³ M concentration with HgCl, 1108.68% at 6.0×10⁻³ M concentration with FeCl₃, 840.10% at 4.0×10⁻³ M concentration with FeSO₄, 488.91 % at 6.0×10⁻³ M concentration with K₄[Fe(CN)₆], 407.2% at 5.33×10⁻³ M concentration with ascorbic acid; 707.90% at 1.33×10⁻³ M concentration with glycine, 172.43% at 1.33×10⁻³ M concentration with Triton X-100. Activation and inhibition effects of metal ions and the other chemicals on PL activity are demonstrated in Table 2 and Figure 8.




Figure 8. Effects of metal ions, amino acids and several chemicals on PL activity A) ascorbic acid B) HgCl₂ C) FeCl₃ D) FeSO₄ E) EDTA F) glycine G) CaCl₂ H) CoCl₂ I) K₄[Fe(CN)₆] J) KCl K) KMnO₄ L) citric acid monohydrate M) NaCl N) Triton X-100.

Table 2. Effects of chemical substances on PL activity.

Chemicals	In this study	Relative Activity (%)	Concentration (M)	Previous studies
Metal ions				
BaCl ₂	activation	249.02	up to 5.33	
HgCl	activation	623.09	up to 0.67	
ZnSO ₄	no effect		-	Inhibition [20,23,28,] Activation [29]
FeCl ₃ .6H ₂ O	activation	1059.41	up to 2.67	
FeSO ₄	activation		up to 1.33	
CaCl ₂	inhibition	703.16	up to 0.333	Inhibition [20,23]
CoCl ₂	inhibition	88.30	up to 0.67	Inhibition [20,25,28]
LiCl ₃	no effect	89.84	-	
MgCl ₂	no effect		-	
MgSO ₄	no effect		-	
NaF	no effect		-	
K ₄ [Fe(CN) ₆]	activation	198.82	up to 0.333	Inhibition [20,23]
KCl	inhibition	57.51	up to 2.67	Inhibition [20,25,28]
KMnO ₄	inhibition	60.52	up to 0.133	
NaN ₃	no effect		-	
NaCl	inhibition	75.48	up to 0.333	
<i>other chemicals</i>				
Ascorbic acid	activation	162.90	up to 1.33	Inhibition [25]
Glycine	activation	171.29	up to 0.133	
SDS	inhibition	97.03	up to 0.333	
L-Cysteine	no effect		-	
Citric acid monohydrate	inhibition	75.57	up to 4.0	Inhibition [23,25,28,29,30]
NaH ₂ AsO ₄	no effect		-	
Triton X-100	activation	136.38	up to 0.67	
EDTA	inhibition	86.36	up to 0.333	No effect [31] Activation [20]
(NH ₄) ₂ SO ₄	no effect		Inhibition [25]	
L-Arginine monohydrate	no effect	-		
L-Asparagine	no effect	-		

References

1. J.N. Be Miller, An introduction to pectins: Structure and properties, In Fishman, M.L., & Jem, J (Eds.), Chemistry and Functions of Pectins, ACS Symposium Series 310. American Chemical Society, Washington, DC, (1986).
2. L. Delgado, B.A. Trejo, C. Huitrón, G. Aguilar, Pectin lyase from *Aspergillus* sp. CH-Y-1043, Applied Microbiology Biotechnology, 39 (1992) 515-519.
3. R.P. De Vries, and J. Visser, *Aspergillus* enzymes involved in degradation of plant cell wall polysaccharides, Microbiology and Molecular Biology Reviews, 65 (2001) 497-552.
4. A. Martínez-Trujillo, J.S. Aranda, C. Gómez-Sánchez, B. Trejo-Aguilar, G. Aguilar-Osorio, Constitutive and inducible pectinolytic enzymes from *Aspergillus flavipes* FP-500 and their modulation by pH and carbon source, Brazilian Journal of Microbiology, 40 (2009) 40-47.
5. D.R. Kashyap, P.K. Vohra, S. Chopra, R. Tewari, Applications of pectinases in the commercial sector: a review, Bioresource Technology, 77 (2001) 215-227.
6. R.S.S. Jayani, S. Saxena, R. Gupta, Microbial pectinolytic enzymes: A review. Process Biochemistry, 40 (2005) 2931-2944.
7. S.R. Nair, S.K. Rakshit, T. Panda, Effect of carbon sources on the synthesis of pectinase by *Aspergillus*, Bioprocess Engineering, 13 (1995) 37-40.
8. I.Reid, and M. Ricard, Pectinase in papermaking: solving retention problems in mechanical pulps bleached with hydrogen peroxide, Enzyme and Microbial Technology, 26 (2000) 115-123.
9. S. Yadav, P.K. Yadav, D. Yadav, K.D.S. Yadav, Pectin lyase: A review, Process Biochemistry, 44 (2009) 1-10.
10. I.G. Sandri, R.C. Fontana, D.M. Barfknecht, M.M. Silveira, Clarification of fruit juices by fungal pectinases, LWT-Food Science and Technology, 44 (2011) 2217-2222.
11. P. Albersheim, Pectin lyase from fungi, Methods in Enzymology, 8 (1966) 628-631.
12. U.K. Laemmli, Cleavage of structural proteins during the assembly of the head of Bacteriophage T4, Nature, 15 (1970) 680-685.
13. G. Temizkan, S. Yilmazer, M. Ozturk, S. Arı, H. Ertan, A. Topal-Sarıkaya, N. Arda, Moleküler Biyolojide Kullanılan Yöntemler. In: G Temizkan & N Arda (Eds.), Proteinlerin İzolasyonu, Analizi, Saflaştırılması, (2008) (pp. 161-273). İstanbul, Turkey.
14. M.M. Bradford, A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, Analytical Biochemistry, 72 (1976) 248-254.
15. D. Silva, E.S. Martins, R. Silva, E. Gomes, Pectinase production by *Penicillium viridicatum* RFC3 by solid state fermentation using agricultural wastes and agro-industrial by-products, Brazilian Journal of Microbiology, 33 (2002) 318-324.
16. N. Martin, S.R. Souza, R. Silva, E. Gomes, Pectinase production by fungal strains in solid-state fermentation using agro-industrial bioproduct, Brazilian Archives of Biology and Technology, 47 (2004) 813-819.
17. A. Alaña, A. Gabilondo, F. Hernando, M.D. Moragues, J.B. Domínguez, M.J. Llana, J.L. Serra, Pectin Lyase Production by a *Penicillium italicum* strain, Applied and Environmental Microbiology, 55 (1989) 1612-1616.
18. S. Koser, Z. Anwar, Z. Iqbal, A. Anjum, T. Aqil, S. Mehmood, M. Irshada, Utilization of *Aspergillus oryzae* to produce pectin lyase from various agro-industrial residues, J of Radiation Research and Applied Sciences, 7 (2014) 327-332.
19. S. Solís, J. Loeza, G. Segura, J. Tello, N. Reyes, P. Lappe, L. Guitérrez, F. Ríos, C. Huitrón, Hydrolysis of orange peel by a pectin lyase-overproducing hybrid obtained by protoplast fusion between mutant pectinolytic *Aspergillus flavipes* and *Aspergillus niveus* CH-Y-1043, Enzyme and Microbial Technology, 44 (2009) 123-128.
20. C.C.H. Sakiyama, E.M. Paula, P.C. Pereira, A.C. Borges, D.O. Silva, Characterization of pectin lyase produced by an endophytic strain isolated from coffee cherries, Letters in Applied Microbiology, 33 (2001) 117-121.
21. A. Alaña, I. Alkorta, J.B. Domínguez, M.J. Llana, J.L. Serra, Pectin lyase activity in a *Penicillium italicum* strain, Applied Environmental Microbiology, 56 (1990) 3755-3759.
22. D.R. Kashyap, S. Chandra, A. Kaul, R. Tewari, Production, purification and characterization of pectinase from a *Bacillus* sp. DT7, World Journal of Microbiology and Biotechnology, 16 (2000) 277-282.
23. S. Yadav, P.K. Yadav, D. Yadav, K.D.S. Yadav, Purification and characterization of an alkaline pectin lyase from *Aspergillus flavus*, Process Biochemistry, 43 (2008) 547-552.
24. S.N. Gummadi, and D.S. Kumar, Optimization of chemical and physical parameters affecting the activity of pectin lyase and pectate lyase from *Debaryomyces flavus nepalensis*: A statistical approach, Biochemical Engineering Journal, 30 (2006) 130-137.
25. Q.K. Beg, B. Bhushan, M. Kapoor, G.S. Hoondal, Production and characterization of thermostable xylanase and pectinase from *Streptomyces* sp. QG-11-3, Journal of Industrial Microbiology & Biotechnology, 24 (2000) 396-402.
26. A. Alaña, M.J. Llana, J.L. Serra, Purification and some properties of the pectin lyase from *Penicillium italicum*, FEBS Letters, 280 (1991) 335-340.
27. C. Perez-Fuentes, M.C. Ravanal, J.H. Eyzaguirre, Heterologous expression of a *Penicillium purpurogenum* pectin lyase in *Pichia pastoris* and its characterization, Fungal Biology, 118 (2014) 507-515.
28. S.H. Hamdy, Purification and characterization of the pectin lyase produced by *Rhizopus oryzae* grown on orange peels, Annals of Microbiology, 55 (2005). 205-211.

29. S. Yadav, A.K. Dubey, G. Anand, R. Kumar, D. Yadav, Purification and biochemical characterization of an alkaline pectin lyase from *Fusarium decemcellulare* MTCC 2079 suitable for *Crotalaria juncea* fiber retting, *Journal of Basic Microbiology*, 53 (2013) 1-9.
30. A. Losonczi, E. Csisza' r, G. Szaka' cs, L. Bezur, Role of the EDTA chelating agent in bioscouring of cotton, *Textile Research Journal*, 75 (2005). 411-417.
31. W.C. Chen, E.C. Hsieh, T.C. Tseng, Purification and characterization of a pectin lyase from *Pythium splendens* infected cucumber fruits, *Botanical Bulletin Academia Sinica*, 39 (1998) 81-186.

