



Peculiarities and applications of aryl-alcohol oxidases from fungi

Vlada B. Urlacher¹ · Katja Koschorreck¹

Received: 5 March 2021 / Revised: 3 May 2021 / Accepted: 6 May 2021 / Published online: 17 May 2021
© The Author(s) 2021

Abstract

Aryl-alcohol oxidases (AAOs) are FAD-containing enzymes that oxidize a broad range of aromatic as well as aliphatic allylic alcohols to aldehydes. Their broad substrate spectrum accompanied by the only need for molecular oxygen as cosubstrate and production of hydrogen peroxide as sole by-product makes these enzymes very promising biocatalysts. AAOs were used in the synthesis of flavors, fragrances, and other high-value-added compounds and building blocks as well as in dye decolorization and pulp biobleaching. Furthermore, AAOs offer a huge potential as efficient suppliers of hydrogen peroxide for peroxidase- and peroxxygenase-catalyzed reactions. A prerequisite for application as biocatalysts at larger scale is the production of AAOs in sufficient amounts. Heterologous expression of these predominantly fungal enzymes is, however, quite challenging. This review summarizes different approaches aiming at enhancing heterologous expression of AAOs and gives an update on substrates accepted by these promising enzymes as well as potential fields of their application.

Key points

- Aryl-alcohol oxidases (AAOs) supply ligninolytic peroxidases with H_2O_2 .
- AAOs accept a broad spectrum of aromatic and aliphatic allylic alcohols.
- AAOs are potential biocatalysts for the production of high-value-added bio-based chemicals.

Keywords Aryl-alcohol oxidase · GMC family · Lignin degradation · Aryl alcohols · Aliphatic allylic alcohols · Biocatalysis · Bio-based precursors · Flavors

Introduction

Since decades, oxidoreductases have been recognized as valuable tools for synthetic chemistry, because they catalyze reactions that are often difficult or impossible to achieve with classical chemical catalysts (Hall 2020). Oxidases, reductases, oxygenases, and dehydrogenases that enable selective synthesis of chiral alcohols, ketones, aldehydes, and carbon acids have been successfully integrated into biocatalytic and chemoenzymatic processes. Oxidoreductases are of vital interest also for developing the bioeconomy via efficient conversion of renewable feedstocks like lignocellulose to fuels and high-value-added chemicals. White-rot fungi secrete a bunch of oxidative enzymes participating in the degradation of lignocellulosic biomass as a central step in carbon recycling. Peroxidases

responsible for lignin degradation require sources of H_2O_2 (Hammel 1997). Thus, fungi produce oxidases such as aryl-alcohol oxidases that generate H_2O_2 required by the peroxidases (Fig. 1). Aryl-alcohol oxidases (AAOs, EC 1.1.3.7), also known as veratryl-alcohol oxidases, aromatic alcohol oxidases, or benzyl-alcohol oxidases, are flavin-adenine-dinucleotide (FAD)-containing enzymes that catalyze the oxidation of aromatic and aliphatic allylic primary alcohols to the corresponding aldehydes while reducing molecular oxygen to H_2O_2 (Guillen et al. 1992). Some other flavin-containing oxidases like vanillyl alcohol oxidase (EC 1.1.3.38) or 4-hydroxymandelate oxidase (decarboxylating; EC 1.1.3.19) that can also oxidize aromatic or phenolic compounds (Bhat and Vaidyanathan 1976; de Jong et al. 1992b; Ewing et al. 2020; Martin et al. 2020) will not be considered herein.

Within this review, we will focus on AAOs from fungi. Since their discovery in 1960, significant knowledge has been gained with regard to structural, spectral, biochemical, and catalytic properties of fungal AAOs. All these achievements are a subject of several excellent reviews (Hernandez-Ortega et al. 2012a; Serrano et al. 2020) and therefore will only be

✉ Katja Koschorreck
Katja.Koschorreck@hhu.de

¹ Institute of Biochemistry, Heinrich-Heine-University Düsseldorf, Universitätsstraße 1, 40225 Düsseldorf, Germany




Fig. 1 Involvement of AAO in lignin degradation. VP versatile peroxidase, MnP manganese peroxidase, LiP lignin peroxidase, AADH aryl-alcohol dehydrogenase

touched briefly here. Like all oxidases, AAOs are independent of the costly nicotinamide cofactors required by oxygenases, and rely only on molecular oxygen as final electron acceptor. These features combined with the broad substrate spectrum and quite high stability make these enzymes potential biocatalysts for the production of bio-based precursors for plastics, bioactive compounds, and fragrances. Indeed, a number of publications, particularly those which have appeared in the last years, are dedicated to the use of AAOs in biocatalysis. Recently, progress was also made in protein engineering of AAOs for improved expression, enhanced activity, and increased selectivity. Broader exploitation and protein engineering of AAOs are however challenged by their difficult accessibility due to low concentrations in original fungi and insufficient expression in recombinant microorganisms. Whereas we will not specifically focus on the methods of enzyme engineering of AAOs described in a recent review (Vina-Gonzalez and Alcalde 2020), we will include a number of examples focusing on synthetic and biotechnological applications of wild-type AAOs and engineered variants thereof, and discuss various heterologous hosts and strategies aiming to enhance heterologous expression of these enzymes.

Occurrence and GMC superfamily

Since the first report in 1960 by Farmer et al. (1960), who identified aromatic alcohol oxidase activity in the growth medium of the basidiomycete *Polystictus (Trametes) versicolor*, AAO activity has been found in cultures of many other basidiomycetes like *Pleurotus* species (Bourbonnais and Paice 1988; Galperin et al. 2016; Guillen et al. 1990; Sannia et al. 1991; Varela et al. 2000), *Bjerkandera adusta* (Muheim et al. 1990b; Romero et al. 2009), *Bjerkandera* sp. BOS55 (de Jong et al. 1992a), *Coprinopsis cinerea* (Tamaru et al. 2018),

Phanerochaete chrysosporium (Asada et al. 1995), *Ustilago maydis* (Couturier et al. 2016), and ascomycetous fungi such as *Aspergillus terreus* (Chakraborty et al. 2014), *Botrytis cinerea* (Goetghebeur et al. 1992), *Fusarium proliferatum* (Regalado et al. 1999), *Geotrichum candidum* (Kim et al. 2001), and *Thermothelemyces thermophilus* (Kadowaki et al. 2020). Although predominantly found in fungi, enzymes with AAO activities have also been reported in bacteria *Sphingobacterium* sp. (Tamboli et al. 2011), *Streptomyces* sp. (Riyadi et al. 2020), and *Methylovorus* sp. (Dijkman and Fraaije 2014), in the phytopathogenic insects *Chrysomela populi* and *Phratora vitellinae* (Brückmann et al. 2002), and in gastropods such as *Arion ater*, *Helix aspersa*, and *Limax flavus* (Large and Connock 1993; Mann et al. 1989).

According to their sequence and structure similarity, most flavin-containing oxidases have been classified in six families (Dijkman et al. 2013; Martin et al. 2020). AAOs belong to one of these families, the glucose-methanol-choline (GMC) oxidoreductase family, defined by Cavener in 1992 (Cavener 1992). GMC oxidoreductases were identified in ascomycetous and basidiomycetous fungi, plants, insects, and bacteria (Cavener 1992; Iida et al. 2007; Martin et al. 2020). Recently, hydroxymethylfurfural oxidases (HMFOs; EC 1.1.3.47) have been discovered which also belong to the GMC oxidoreductase family and oxidize aromatic primary alcohols, aldehydes, and thiols (Dijkman and Fraaije 2014; Ewing et al. 2014; Vinambres et al. 2020). While AAOs are secreted by fungi, HMFOs have been found in bacteria like *Methylovorus* sp. strain MP688 and *Pseudomonas nitroreducens*. HMFO from *Methylovorus* sp. strain MP688 is, for example, an intracellular bacterial enzyme with low sequence identity to AAOs (Dijkman and Fraaije 2014).

In 2013, the GMC oxidoreductases were assigned to the “Auxiliary Activities 3” (AA3) family in the Carbohydrate-Active enZyme (CAZy) database (Levasseur et al. 2013). The

“auxiliary” enzymes help the carbohydrate-active enzymes like glycoside hydrolases, polysaccharide lyases, and carbohydrate esterases to gain a better access to carbohydrates in plant cell wall. Four subfamilies belong to the AA3 family: AA3_1 includes for the most part cellobiose dehydrogenases (CDHs, EC 1.1.99.18); AAOs together with glucose 1-oxidases (GOx, EC 1.1.3.4), glucose-1-dehydrogenases (GDHs, EC 1.1.5.9), and pyranose dehydrogenases (PDHs, EC 1.1.99.29) belong to the subfamily AA3_2; AA3_3 contains alcohol oxidases (AOx, EC 1.1.3.13); and AA3_4 comprises pyranose 2-oxidase (POx, EC 1.1.3.10).

The spectrum of substrates accepted by GMC oxidoreductases covers a broad range of sugars and alcohols. Depending on the type of the final electron acceptor, the GMC family members can be divided into two groups: oxidases, that transfer electrons to molecular oxygen, and dehydrogenases that transfer electrons to quinones, phenoxy radicals, or metal ions and show no or negligible oxygen reactivity. Fungal aryl-alcohol:quinone oxidoreductases (AAQOs) possess a similar substrate spectrum as AAOs but utilize quinones as final electron acceptor rather than molecular oxygen (Mathieu et al. 2016). Based on sequence similarities of ~10,000 putative fungal GMC oxidoreductases, these enzymes were classified in five clusters: AOx, CDH, POx, GOx–GDH, and AAO–PDH (Sützl et al. 2019). Phylogenetic and sequence studies of the individual clusters shed light into the sequence-structure-function relationship of the GMC family (Sützl et al. 2019). According to this analysis, the AAO-PDH cluster showed a split into three clades: AAO, AAO-like, and PDH. AAOs and AAQOs did not appear in separate clades, indicating that only minor amino acid changes determine the specificity for the final electron acceptor.

Functions in nature

As mentioned above, in nature, AAOs are involved in degradation of lignocellulosic biomass. Lignocellulose is the most abundant renewable feedstock that provides a rich source for biofuels and value-added chemicals. Thus, effective bioconversion of lignocellulosic biomass has attracted significant attention from both academic and industrial points of view (Becker and Wittmann 2019). The most recalcitrant component of lignocellulose is lignin that can be decomposed by fungi and some bacterial species (Bugg et al. 2011; de Gonzalo et al. 2016). Wood-decaying white-rot fungi degrade lignin by using a large set of enzymes including lignin-modifying oxidative enzymes and “auxiliary” enzymes that do not degrade lignin on their own but assist other enzymes during lignin decomposition. Fungal oxidative enzymes involved in lignin degradation are laccases and the class II peroxidases lignin peroxidase (LiP), manganese peroxidase (MnP), and versatile peroxidase (VP) (Fig. 1). “Auxiliary”

enzymes involved in lignin degradation include H₂O₂-generating AAO, glyoxal oxidase (GLOX, EC 1.2.3.5), and pyranose 2-oxidase but also CDH, GDH, and others. The well-studied lignin-degrading white-rot fungus *Pycnoporus cinnabarinus* contains, for example, 9 class II peroxidase encoding sequences, 5 putative laccase genes, 7 putative glyoxal oxidase genes, and 3 AAO encoding sequences (Levasseur et al. 2013). In comparison, brown-rot fungi lack class II peroxidases and instead produce enzymes that are involved in extracellular formation of Fenton’s reagent (Levasseur et al. 2014) (Fig. 1). Fe(II) reduces H₂O₂ to hydroxyl free radicals which cause cellulose degradation and methanol release from lignin by demethylation (Martinez et al. 2011). Methanol is used by methanol oxidase (MOX) to produce H₂O₂. MOX is the most abundant GMC oxidoreductase and the main oxidase involved in wood decay by brown-rot fungi (Daniel et al. 2007; Ferreira et al. 2015a). Thus, white-rot fungi rely on H₂O₂-producing enzymes like AAOs for ligninolytic peroxidases while in brown-rot fungi H₂O₂ released by MOX is utilized for Fenton’s reagent formation. This is in agreement with the higher number of AAOs found in peroxidase-producing white-rot fungi compared to brown-rot fungi (Ferreira et al. 2015a; Levasseur et al. 2013). Furthermore, extracellular production of AAO along with LiP by the white-rot fungus *B. adusta* (Muheim et al. 1990a) and AAO together with LiP and MnP by the white-rot fungi *P. ostreatus* and *T. pubescens* (Casieri et al. 2008) supports the hypothesis that AAOs are involved in lignin decomposition. Notably, the thermophilic cellulolytic ascomycete *Thermothelomyces thermophilus* M77 produces AAO but no lignin or manganese peroxidase (Berka et al. 2011), which suggests that AAO might be also involved in Fenton’s reagent formation.

To supply class II peroxidases with H₂O₂ for breakdown of lignin, wood-decaying fungi produce small aromatic aldehydes like anisaldehyde or veratraldehyde that are reduced by NADPH-dependent aryl-alcohol dehydrogenases to the corresponding alcohols which are accepted by AAOs as substrates (Guillen and Evans 1994). AAOs complete the redox cycle by transferring the electrons from aryl alcohols to molecular oxygen thus producing H₂O₂ and aldehydes. Lignin-derived aryl alcohols and aldehydes may also be used as substrates by AAOs for H₂O₂ production. Besides oxidizing benzylic alcohols produced by fungi or released in course of lignin decomposition, AAOs accept a broad range of other substrates, which are described in more detail in the following.

Structure and mechanism

Members of the GMC family contain FAD as cofactor and share a similar topologic fold: They are composed of a highly conserved N-terminal domain involved in FAD binding, and a


less conserved C-terminal domain involved in substrate binding (Martin et al. 2020). The N-terminal domain has the Rossmann super-secondary structure with an ADP-binding $\beta\alpha\beta$ fold. The variability of the C-terminal domain reflects the diversity of substrates accepted by the members of this family (Sützl et al. 2018). Common to all GMC members is the presence of a highly conserved catalytic histidine residue involved in substrate oxidation and FAD reoxidation (Martin et al. 2020).

For many years, only the crystal structure of *PeAAO* from *P. eryngii* with and without inhibitor was available (Carro et al. 2017; Fernandez et al. 2009). The enzyme was produced as selenomethionine derivative in *E. coli* and refolded in vitro. Structure analysis revealed that this AAO is a monomer that is, similar to other GMC oxidoreductases, composed of two domains, an FAD-binding domain and a substrate-binding domain. Non-covalently bound FAD cofactor is stabilized by H bonds formed by the residues of the conserved $\beta\alpha\beta$ fold in the N-terminal domain. In this enzyme, the isoalloxazine ring of FAD is buried inside the protein. The substrate-binding domain is formed by a six-stranded antiparallel β -sheet flanked by two long α -helices. Whereas one of the best studied enzymes from the GMC family, glucose oxidase, has an open wide active site, the accessibility to the active site of *PeAAO* is restricted by an additional loop that forms a funnel-like hydrophobic channel. This channel connects the solvent with the FAD cofactor through a bottleneck formed by three aromatic residues, Tyr92, Phe397, and Phe501, which limit substrate diffusion to the active site. The active site is located in front of the re-side of the isoalloxazine ring of FAD and near two histidines, His502 and His546, postulated as catalytic residues (Ferreira et al. 2006) (Fig. 2).

Sequence comparison of the AAO-PDH clade members of the GMC family revealed that Tyr92 and Phe501 located in the hydrophobic substrate access channel of *PeAAO* are conserved among many AAOs, but absent in the PDH clade (Sützl et al. 2019). The loss of the narrow hydrophobic channel was suggested as the starting point for widening the substrate entrance site to accept other substrates like sugars by PDH. Another feature to discern AAO and PDH is the covalently bound FAD in PDH. The covalently bound FAD cofactor is usually related to higher redox potentials (Heuts et al. 2009) probably resulting in an altered substrate spectrum and thus contributing to the differentiation of PDHs and AAOs.

In 2020, the crystal structure of *MtAAOx* from *T. thermophilus* (formerly *Myceliophthora thermophila*) was solved and compared with the structure of *PeAAO* (Kadowaki et al. 2020). *MtAAOx* was expressed and secreted by *A. nidulans* and crystallized in glycosylated form. Comparative analysis of the two structures has revealed that whereas the FAD-binding domain is strongly conserved in *MtAAOx* and *PeAAO*, the substrate-binding domain and the substrate channel to the active site differ. *MtAAOx* possesses a wider substrate access channel connecting the active site with the solvent. The conserved catalytic histidines His579 and His622 are located in the easily accessible hydrophobic active site. Remarkably, three aromatic residues in the hydrophobic bottleneck present in *PeAAO*, Tyr92, Phe397, and Phe501, are not conserved in the *MtAAOx* structure. Furthermore, the *MtAAOx* structure has a unique Ca^{2+} binding pocket located near the catalytic site entrance. This element was not reported previously for any AA3_2 enzyme, and seems to stabilize the structure of the thermostable *MtAAOx* (Kadowaki et al. 2020).

Fig. 2 Catalytic site of *PeAAO*. View of the active site of *PeAAO* (PDB entry 3FIM) with FAD depicted in yellow, catalytic His502 and His546 depicted in gray, and Tyr92 (green) and Phe501 (blue) located in the hydrophobic substrate access channel



Like other GMC enzymes, the catalytic cycle of AAOs can be divided into two half-reactions, reductive and oxidative. In the reductive half-reaction, the alcohol substrate is oxidized via two-electron abstraction and the reduced form of the flavin cofactor is formed; in the oxidative half-reaction, reduced FAD is reoxidized whereas molecular oxygen is reduced to hydrogen peroxide. Crystal structures, kinetic and spectroscopic analyses, QM/MM studies, docking experiments, and site-directed mutagenesis performed by A. T. Martinez and colleagues have provided a detailed picture of substrate access in the active site and catalytic mechanism of *PeAAO* which are summarized in the recent review (Serrano et al. 2020). The alcohol substrate interacts with Phe397 in the funnel-like hydrophobic channel, which induces a rearrangement of the loop that limits the access to the active site (Carro et al. 2018a). This rearrangement promotes the substrate in a catalytically relevant position in the active site, where it is stabilized through an aromatic stacking interaction with Tyr92 (Ferreira et al. 2015b). Docking of *p*-methoxybenzyl alcohol and QM/MM calculations demonstrated that the pro-*R* hydrogen of the substrate is located 2.4 Å away from the N5 position of the isoalloxazine ring of the cofactor, enabling hydride transfer and yielding the reduced FAD during the reductive half-reaction. In the same time, the hydroxyl proton of the substrate is abstracted by the catalytic base His502, assisted by His546, which forms an H bond with the alcohol (Hernandez-Ortega et al. 2012c). Although proton abstraction and hydride transfer are highly coupled, they occur through non-synchronous concerted mechanism. Unlike other GMC oxidoreductases, no stable alkoxide intermediate was detected, which suggests that AAO follows a different mechanism for alcohol oxidation (Hernandez-Ortega et al. 2011a).


Molecular oxygen accesses the buried FAD by diffusion through the narrow hydrophobic channel, where Phe501 was proposed to assist the O₂ molecule to approach the flavin C4a position (Hernandez-Ortega et al. 2011b). The structure of *MtAAOx* from *T. thermophilus* reveals a wider channel, which might explain the 202-times lower O₂ reactivity of this oxidase than *PeAAO*. Interestingly, the replacement of Phe501 through alanine in *PeAAO* resulted in reduction of O₂ reactivity (Hernandez-Ortega et al. 2011b).

In the oxidative half-reaction in *PeAAO*, reduced FAD first reacts with molecular oxygen to form a superoxide anion radical and neutral flavin semiquinone. Subsequently, one hydride (originally abstracted from the alcohol substrate in the reductive half-reaction) from the flavin N5 and one proton originating either from the catalytic His502 or from the solvent are transferred to the superoxide anion to furnish H₂O₂ (Carro et al. 2018c). The question, whether the first electron transfer is coupled to the proton transfer remains open. Differently from flavin monooxygenases (Paul et al. 2021; Toplak et al. 2021) and some flavin oxidases (Sucharitakul et al. 2008; Wongnate and Chaiyen 2013), a C4a-hydroperoxyflavin is not stabilized in *PeAAO* (Serrano et al. 2020).

Substrate spectrum

Originally declared as aryl-alcohol oxidases, the enzymes belonging to this group accept a quite broad range of substrates (Fig. 3). However, substrate preferences differ for different AAOs. Activity of AAOs from *Pleurotus* strains was found to depend on the position and properties of aromatic ring substituting groups (Guillen et al. 1992; Jankowski et al. 2020). Whereas an electron-withdrawing substituent like nitro group at the *para*-position of the aromatic ring had a negative effect on enzyme activity (Guillen et al. 1992), an electron-donating substituent like methoxy-group at this position enhanced enzyme activity compared to unsubstituted benzyl alcohol (Galperin et al. 2016; Guillen et al. 1992; Jankowski et al. 2020; Varela et al. 2000). Other AAOs like *rCcAAO* from *C. cinerea*, BAO from *B. cinerea*, and AOx from *A. terreus* oxidized benzylic alcohols with a methoxy-substitution at the *para*- or *meta*-position of the aromatic ring equally well (Fig. 4A). Interestingly, the presence of *para*-hydroxy group differently affected the activity of AAOs: In comparison with benzyl alcohol, *p*-hydroxybenzyl alcohol was oxidized with a negligible activity by *PeAAO* (Guillen et al. 1992), but faster by *C. cinerea rCcAAO* (Tamaru et al. 2018). Whereas vanillyl alcohol was barely oxidized by *PeAAO2* from *P. eryngii* P34, isovanillyl alcohol was oxidized by the same enzyme with an almost 3-times higher activity than benzyl alcohol (Jankowski et al. 2020) (Fig. 4B). Differently, *C. cinerea rCcAAO* catalyzed oxidation of both vanillyl alcohol and isovanillyl alcohol with similar activities, which were only slightly lower than activity towards benzyl alcohol (Chakraborty et al. 2014; Tamaru et al. 2018). AAO of *Bjerkandera* BOS55 prefers mono- and dichlorinated anisyl alcohols as substrates (de Jong et al. 1994). Polyunsaturated aliphatic alcohols like *trans,trans*-2,4-hexadien-1-ol or 2,4-heptadien-1-ol were recognized as favorable substrates of many AAOs (Guillen et al. 1992; Jankowski et al. 2020; Tamaru et al. 2018). Non-conjugated aliphatic allylic alcohols like *trans*-2-hexen-1-ol, *trans*-2-hepten-1-ol, *trans*-2-octen-1-ol, and *trans*-2-cis-6-nonadien-1-ol were oxidized by *PeAAO2* from *P. eryngii* P34 to a much lesser extent (Jankowski et al. 2020). Of the aromatic alcohols, 2-naphthalenemethanol was identified as most easily oxidized by *Pleurotus* AAOs followed by 1-pyrenemethanol (Guillen et al. 1992; Jankowski et al. 2020). 9-Anthracenemethanol was not accepted as substrate at all, probably due to steric constraints. Recently, piperonyl alcohol, cumic alcohol, and 2-thiophenemethanol were identified as AAO substrates as well (Jankowski et al. 2020). These results indicate that the substrate scope of AAOs is dependent on the chemical properties and reactivity of the substrate and is controlled by the size and properties of the active sites. Obviously, a comparative structure-function analysis including more structural information from various AAOs will enable the identification of

Fig. 3 Substrates accepted by AAOs. Alcohols are depicted in blue, aldehydes in red



positions or regions in the active sites responsible for different substrate preferences and provide a basis for rational or semi-rational protein engineering of different AAOs.

In rare cases, AAOs catalyze the oxidation of aldehyde substrates to the corresponding acids. Aldehyde oxidation catalyzed by AAOs was proposed to mainly proceed via the *gem*-diols formed by aldehyde hydration, and thus following the mechanism analogous to alcohol oxidation (Ferreira et al. 2010). Cleavage of one of the *gem*-diol O–H bonds involves the catalytic His502 as a base, and the presence of electron-withdrawing or electron-donating ring substituting groups

seems to stabilize or destabilize the *gem*-diol intermediate. In contrast to the substituted aromatic alcohols, *PeAAO* demonstrated the highest activity towards the aromatic aldehydes with electron-withdrawing substituents like nitro groups or chlorine at the *para*- or the *ortho*-positions. Although benzaldehyde was accepted as substrate, the highest activity of *PeAAO* was observed with *p*-chlorobenzaldehyde and *p*-nitrobenzaldehyde. The presence of an electron-donating methoxy-group at the *para*-position resulted in low or no activity like with veratraldehyde and *p*-anisaldehyde (Ferreira et al. 2010).




Fig. 4 Substrate specificity of AAOs towards *m*- and *p*-anisyl alcohol (**A**) and vanillyl and isovanillyl alcohol (**B**). **A** Relative activity of AAOs from *B. cinerea* (Goetghebeur et al. 1992), *B. adusta* (Romero et al. 2009), *C. cinerea* (Tamaru et al. 2018), *A. terreus* (Chakraborty et al. 2014), *P. ostreatus* (Sannia et al. 1991), and *P. eryngii* (Guillen et al. 1992) towards *m*-anisyl alcohol. Activity towards *p*-anisyl alcohol was

set to 100%. **B** Relative activity of AAOs from *B. cinerea* (Goetghebeur et al. 1992), *B. adusta* (Romero et al. 2009), *C. cinerea* (Tamaru et al. 2018), *P. ostreatus* (Sannia et al. 1991), *P. eryngii* P34 (Jankowski et al. 2020), and *P. eryngii* (Guillen et al. 1992) towards vanillyl alcohol. Activity towards isovanillyl alcohol was set to 100%

Heterologous expression

Although identified almost 60 years ago and widespread in nature, only a limited number of AAOs have been characterized so far. Since production levels of AAOs in their original host organisms are usually low and their purification from protein mixtures is tedious, overexpression of AAOs in heterologous microorganisms appears attractive to obtain sufficient amounts of enzymes for purification and biochemical characterization. Significant efforts have been undertaken to achieve high-level expression of AAOs in several microbial hosts, including *Escherichia coli*, *Aspergillus nidulans*, *Pichia pastoris*, *Saccharomyces cerevisiae*, and *C. cinerea* (Table 1). However, their heterologous expression seems to be quite challenging and often suffers from low yields. The first heterologous expression of an AAO was reported in 2001 by Varela and colleagues (Varela et al. 2001). They produced *PeAAO* in *A. nidulans* using an alcohol dehydrogenase promoter of *A. nidulans* (*alcA*) with a maximal volumetric activity of 400–500 mU/mL culture. A two-step chromatographic purification procedure yielded 3 mg of active enzyme from 1 L of *A. nidulans* culture (Ferreira et al. 2005). Expression of the *PeAAO* gene without its signal peptide sequence in *E. coli* under the control of the *tac*-promoter resulted in formation of inclusion bodies. After in vitro refolding and purification, 45 mg of active AAO per liter of *E. coli* culture was obtained (Ruiz-Duenas et al. 2006). However, non-glycosylated *PeAAO* expressed in *E. coli* showed lower pH and temperature stability compared to the glycosylated enzyme isolated from *P. eryngii* or expressed in *A. nidulans* (Ruiz-Duenas

et al. 2006). To further improve AAO production level and to establish an appropriate expression host suitable for directed evolution, *S. cerevisiae* was investigated as host for the expression of *PeAAO* (Vina-Gonzalez et al. 2015). First, different secretion signal peptides of *S. cerevisiae* were investigated to promote secretion of *PeAAO*. The native secretion signal sequence of *PeAAO* was replaced by either the prepro-leader sequence from the mating α -factor (prepro α) or the prepro-leader sequence from the K₁ killer toxin (preproK) or chimeras of both, respectively. The prepro α -AAO variant was produced at the highest yield of 1.5 U/L followed by the construct with a chimeric leader sequence pre α proK-AAO secreted at 0.5 U/L. This difference might be explained by differences in efficiencies of cleavage of the signal peptide sequences. The prepro α sequence contains both the KEX2 and STE13 cleavage sites, while the preproK sequence carries only the KEX2 cleavage site. When high amounts of a heterologous protein bearing the α -factor prepro-leader sequence are expressed in *S. cerevisiae*, the amount of STE13 protease in the Golgi might be not enough for efficient cleavage (Ahmad et al. 2014). Since the proK peptide was efficiently cleaved off by KEX2 protease, the pre α proK-AAO variant was used to further improve the *PeAAO* secretion and activity by directed evolution. Several *PeAAO* mutant libraries were constructed by conventional directed evolution and using the method named Mutagenic Organized Recombination Process by Homologous In Vivo Grouping (MORPHING) in *S. cerevisiae* (Vina-Gonzalez et al. 2015). Out of 5000 clones screened, the FX7 variant with the mutation H91N demonstrated a 96-fold improved total activity compared to the

Table 1 Heterologous expression of AAOs in different hosts

Name	Origin	Expression host	Vol. Activity (U/L)	Yield (mg/L)	Reference
<i>PeAAO</i>	<i>Pleurotus eryngii</i>	<i>Aspergillus nidulans</i>	–	3 ^a	Ferreira et al. (2005)
<i>PeAAO</i>	<i>Pleurotus eryngii</i>	<i>Aspergillus nidulans</i>	400–500	–	Varela et al. (2001)
<i>PeAAO</i>	<i>Pleurotus eryngii</i>	<i>Escherichia coli</i>	–	45 ^b	Ruiz-Duenas et al. (2006)
<i>PeAAO_FX7</i>	<i>Pleurotus eryngii</i>	<i>Saccharomyces cerevisiae</i>	48	2	Vina-Gonzalez et al. (2015)
<i>PeAAO_FX9</i>	<i>Pleurotus eryngii</i>	<i>Saccharomyces cerevisiae</i>	–	4.5	Vina-Gonzalez et al. (2018)
<i>PeAAO_FX9</i>	<i>Pleurotus eryngii</i>	<i>Pichia pastoris</i>	1378	25.5	Vina-Gonzalez et al. (2018)
<i>PeAAO2</i>	<i>Pleurotus eryngii</i> P34	<i>Pichia pastoris</i>	7250	315	Jankowski et al. (2020)
AAO	<i>Pleurotus sapidus</i>	<i>Coprinopsis cinerea</i>	125	–	Galperin et al. (2016)
rAOx	<i>Aspergillus terreus</i> MTCC6324	<i>Escherichia coli</i>	–	–	Chakraborty et al. (2014)
rCcAAO	<i>Coprinopsis cinerea</i>	<i>Pichia pastoris</i>	–	–	Tamaru et al. (2018)
MtAAOx	<i>Thermothelomyces thermophilus</i> M77	<i>Aspergillus nidulans</i>	–	8.4	Kadowaki et al. (2020)
AAO	<i>Thermothelomyces thermophilus</i>	<i>Aspergillus nidulans</i>	1906 ^c	–	Liu and Wilkins (2020)
UmAAO	<i>Ustilago maydis</i>	<i>Pichia pastoris</i>	–	1000	Couturier et al. (2016)

^a Purified enzyme

^b Purified and refolded from inclusion bodies

^c Activity measured with DCPIP as final electron acceptor

parental pre α proK-AAO variant. FX7 was expressed at a concentration of 2 mg/L as a heavily glycosylated protein (~50% glycosylation) and possessed biochemical properties comparable to those of PeAAO expressed in *E. coli*, while pH and thermal stabilities were significantly enhanced. To further improve PeAAO secretion in *S. cerevisiae*, in vivo shuffling of the FX7 variant with six other improved variants was combined with subsequent MORPHING of the pre α proK leader sequence (Vina-Gonzalez et al. 2018). The most active clone FX9 with 4 mutations in the pre α proK peptide and 2 mutations, H91N and L170M, in the mature protein was secreted at concentration of 4.5 mg/L during cultivations in shake flasks. Expression of the FX9 variant in *P. pastoris* under control of the methanol-inducible *AOX1* promoter yielded 25.5 mg/L of enzyme (final activity of 1378 U/L) from a 5 L fed-batch culture (Vina-Gonzalez et al. 2018). The *P. pastoris*-expressed FX9 variant showed different kinetic parameters and lower thermal stability than its counterpart expressed in *S. cerevisiae*. These differences were attributed to the poor glycosylation of the FX9 variant expressed in *P. pastoris* (degree of glycosylation not measurable) compared to heavy glycosylation when expressed in *S. cerevisiae* (60% glycosylation). Another *Pleurotus* AAO from *P. sapidus* was heterologously expressed in *C. cinerea* under control of the *Agaricus bisporus gpdII* promoter up to a volumetric activity of 125 U/L (Galperin et al. 2016). Recently, heterologous expression of PeAAO2 from *P. eryngii* P34 in *P. pastoris* was reported with an exceptional high yield of 315 mg/L and a volumetric activity of 7250 U/L after fed-batch cultivation in 3 L (Jankowski et al. 2020). Although mature PeAAO2 and the FX9 variant of PeAAO differ only in 9 amino acid residues located on or near the protein's surface, PeAAO2 was much better expressed than FX9 and more stable at acidic and basic pH values. The higher pH stability might be due to glycosylation, since PeAAO2 was strongly glycosylated (30% N-glycosylation) in *P. pastoris*. Two further AAOs, UmAAO from *Ustilago maydis* and rCcAAO from *C. cinerea*, were also expressed in *P. pastoris* (Couturier et al. 2016; Tamaru et al. 2018). UmAAO was secreted with the help of the α -factor prepro-leader peptide and expressed with a C-terminal (His)₆-tag. Around 1 g of recombinant UmAAO per liter of culture medium was produced which is the highest yield of an AAO reported so far. rCcAAO was also fused to the α -factor prepro-leader peptide for secretion, but unfortunately no information on expression yields were provided. Kadowaki and coworkers reported on heterologous expression of MtAAOx from *T. thermophilus* M77 in *A. nidulans* (Kadowaki et al. 2020). The MtAAOx gene was expressed using a glucoamylase promoter and a heterologous signal peptide. After three chromatographic steps, 8.4 mg of MtAAOx was obtained from 1 L of culture. The group of Wilkins reported on heterologous expression of an AAO from *T. thermophilus* in *A. nidulans* as well (Liu et al. 2020; Pardo-Planas et al.

2017). Process optimization and up-scaling of protein production to 3-L stirred-tank bioreactor resulted in volumetric activity of 1906 U/L after 48 h (Liu and Wilkins 2020). However, no sequence information of the enzyme was provided and activity of this enzyme towards veratryl alcohol was solely measured with 2,6-dichlorophenol indophenol (DCPIP) as final electron acceptor. The question if this enzyme is identical or highly similar to MtAAOx remains open.

The second AAO heterologously expressed in *E. coli* is rAOx from *A. terreus* MTCC6324 (Chakraborty et al. 2014). rAOx expressed with both N- and C-terminal (His)₆-tag under control of the T7 promoter in *E. coli* BL21(DE3) predominantly formed inclusion bodies. After purification and in vitro refolding of aporAOx with its cofactor FAD, active rAOx was obtained, but no information on achieved yields were provided.

Biotechnological application

The ability of AAOs to reduce molecular oxygen to hydrogen peroxide coupled to alcohol oxidation makes them interesting candidates for biotechnological purposes (Fig. 5). On the one hand, AAOs releasing H₂O₂ are used to support reactions catalyzed by peroxidases and peroxygenases. On the other hand, AAO-catalyzed oxidation reactions provide a valuable tool for the synthesis of building blocks for chemical synthesis and flavors. Finally, there are processes in which AAOs are applied for both, biocatalytic oxidation and for in situ H₂O₂ formation to support other enzymes involved in multi-enzyme cascades for the production of bio-based precursors for polymers. Furthermore, using methods of protein engineering AAOs was optimized for application as effective biocatalysts. In the following, we will give a short overview of the processes and reactions involving AAOs.

Dye decolorization and pulp biobleaching

AAOs lead to H₂O₂ production and hence can support various enzymes during dye decolorization and lignin degradation. Synthetic dyes are widely used in the paper and textile industry causing large amounts of contaminated wastewater. Efficient wastewater treatment is thus of great importance to remove these harmful compounds being toxic to humans and aquatic organisms (Ledakowicz and Pazdzior 2021). Besides physical and chemical treatments, biological methods relying on fungi, bacteria, or isolated enzymes are of high importance. Among others, fungal ligninolytic enzymes, such as laccases, LiP, and MnP, have been broadly used for dye decolorization (Wesenberg et al. 2003). Only one report describes the direct participation of AAO in dye decolorization: AAO isolated from *Sphingobacterium* sp. ATM caused the degradation of the textile dyes Red 5B, Orange 3R, and Direct Blue GLL




Fig. 5 AAOs for biotechnological purposes

within 8 h (Tamboli et al. 2011). Other investigations revealed that AAOs contribute to the decolorization process by generating H₂O₂ for the dye degrading enzymes. For example, AAO from *Geotrichum candidum* Decl was not able to degrade textile dyes like Red 5B, but supported its decolorization by delivering H₂O₂ to the dye peroxidase DyP. Moreover, AAO was suggested to prevent the polymerization of dye products produced by DyP peroxidase, which facilitated their further degradation (Kim et al. 2001). A combination of dye decolorizing peroxidase DyP from *Mycetinis scorodonius* and AAO from *P. sapidus* resulted in a more efficient

decolorization of the dye Reactive Blue 5 in comparison to the DyP directly supplied with external H₂O₂. This effect was attributed to the low but constant production of H₂O₂ by AAO, which allows to overcome the possible DyP inhibition by high H₂O₂ concentrations (Galperin et al. 2016). This two-enzyme system does not require additional equipment and online quantification of H₂O₂ concentration and thus has a potential to replace the systems for in situ H₂O₂ supply like feeding or electrochemical peroxide production.

AAOs have been shown to contribute also to the pulp bleaching process to obtain white paper pulp. Original pulp

color is impacted by the chromophores of lignin. Thus, a set of hydrolases and oxidoreductases involved in lignin degradation have been tested for removal of these chromophores and pulp lightening. Whereas *P. cinnabarinus* laccase combined with the redox mediator 1-hydroxybenzotriazol (HBT) demonstrated the strongest delignification of pulp, the same laccase without mediator also caused pulp lightening that was increased in the presence of AAO (Sigoillot et al. 2005). Similar to the effect on dye decolorization, the positive effect of AAO was attributed to their ability to prevent repolymerization of the lignin phenoxy radicals produced by laccase (Sigoillot et al. 2005).

Synthesis of flavors and valuable building blocks

Flavors and fragrances are widely used in food and beverages, as well as in the cosmetic, detergent, and pharmaceutical industries. They are currently mainly produced by chemical synthesis. Due to the increasing consumer demand for “natural” flavors, more and more biotechnological processes relying on enzymes or microorganisms are established for the production of these compounds. Vanillin and benzaldehyde (cherry flavor), the leading compounds in the flavor and fragrance industry, can be formed, for example, via AAO-catalyzed oxidation of the corresponding alcohols, which makes AAOs interesting candidates for the synthesis of flavors and fragrances. For instance, recombinant *PeAAO* from *P. eryngii* catalyzes the oxidation of *trans*-2-hexen-1-ol to *trans*-2-hexenal, a major component responsible for the fresh flavor of fruits and vegetables. *trans*-2-Hexenal is widely applied in the flavor and fragrance industry as Green Note in food, beverages, and perfumes. In an attempt to enhance the biocatalytic performance of *PeAAO* that can be limited by the poor solubility and low diffusion rates of O₂ into the aqueous reaction media, the reaction was conducted in a continuous slug-flow microreactor (van Schie et al. 2018). Additionally, catalase was added to decompose the produced H₂O₂ to H₂O and O₂. With this setup, 10 mM *trans*-2-hexen-1-ol were completely converted to *trans*-2-hexenal within the residence time of 40 min, which corresponds to a turnover number (TN) of 32,400. By increasing residence time in a longer flow reactor, TNs of over 300,000 were reached (van Schie et al. 2018). Further increase of substrate concentration was however limited by low solubility of *trans*-2-hexen-1-ol in aqueous media. The use of two liquid-phase systems allowed not only to increase substrate solubility but also to avoid product inhibition (de Almeida et al. 2019). Dodecane and *trans*-2-hexen-1-ol itself as organic phase ensured the best results among the organic solvents tested. At a 1:1 ratio of aqueous to dodecane phase, full conversion of 500 mM *trans*-2-hexen-1-ol was achieved after 24 h. This corresponds to a TN of 650,000. When *trans*-2-hexen-1-ol was used as organic phase at a 4:1 ratio and both, *PeAAO* and catalase, were added six

times in course of the reaction, 2.6 M *trans*-2-hexenal were produced within 14 days, which corresponded to a TN of 2,200,000. These examples demonstrate high robustness of the recombinant *PeAAO* from *P. eryngii*. In another study, *PeAAO2* from *P. eryngii* P34 was shown to catalyze the oxidation of piperonyl alcohol to piperonal with the second highest catalytic efficiency among the substrates investigated with this enzyme (Fig. 5) (Jankowski et al. 2020). Piperonal, also termed as “heliotropin,” has a sweet, floral odor and is used in the perfumery and cosmetic industry, but also in the food industry as vanilla aroma. Piperonal is also used as an intermediate in the production of insecticides, pesticides, and pharmaceutical agents for, e.g., treatment of Alzheimer’s disease (Brum et al. 2019; Santos et al. 2004; Wang et al. 2019).

AAOs are also of potential interest for the synthesis of chiral secondary alcohols. Docking of *p*-methoxybenzyl alcohol in the active site of *PeAAO* has revealed a stereoselective hydride abstraction from the pro-*R* C α position (Hernandez-Ortega et al. 2012b). The stereoselective hydride abstraction from the *S*-enantiomer would lead to the corresponding ketone, while the *R*-enantiomer would remain unaffected. Chiral secondary alcohols are essential building blocks for the production of, e.g., pharmaceutical compounds. However, activity of *PeAAO* with secondary aromatic alcohols is much lower than with primary alcohols probably due to the narrow substrate access channel. To enhance activity of *PeAAO* towards secondary alcohols, rational protein design and directed evolution approaches have been applied (Serrano et al. 2019b; Vina-Gonzalez et al. 2019). The *PeAAO* variant F501A, where phenylalanine limiting substrate diffusion was replaced by the smaller alanine (Hernandez-Ortega et al. 2012b), was used as a starting enzyme for adaptive-protein energy landscape exploration simulations and combinatorial saturation mutagenesis (Serrano et al. 2019b). As a result, the double mutant I500M/F501W with increased activity and stereoselectivity was identified. Introducing the mutations I500M and F501W into the *PeAAO* variant FX9 with improved secretion (see above), followed by structure-guided evolution including in vivo site-directed recombination as the final step, resulted in the LanDo variant with a more than 800-fold enhanced activity towards 1-(*p*-methoxyphenyl)-ethanol compared to FX9 (Vina-Gonzalez et al. 2019). The catalytic efficiency (k_{cat}/K_m) of the LanDo variant with 1-(*p*-methoxyphenyl)-ethanol was 10-fold higher than of the I500M/F501W variant. The *S*-enantiomer of racemic 1-(*p*-methoxyphenyl)-ethanol was almost completely oxidized by the LanDo variant, which resulted after 2 h in *R*-enantiomer with *ee* >99%. This example demonstrates the potential of AAOs as biocatalyst for the synthesis of chiral secondary alcohols via deracemization.

Another potential field of AAO application lies in the production of compounds with beneficial effects on human health or drug metabolites. *PeAAO2* from *P. eryngii* P34 catalyzes

the oxidation of cumic alcohol to cuminaldehyde with the highest activity among the substrates tested with this enzyme. Cuminaldehyde is the major constituent of seed oil of *Cuminum cyminum* (Lee 2005; Li and Jiang 2004) with potential therapeutic effects (Morshedi et al. 2015). For example, cuminaldehyde demonstrated anticancer and antidiabetic properties (Lee 2005; Patil et al. 2013; Tsai et al. 2016).

Recently, a self-sufficient unspecific peroxygenase (UPO)/aryl-alcohol oxidase biocatalyst for the synthesis of dextrorphan, a human metabolite of the antitussive drug dextromethorphan, was reported (de Santos et al. 2020). The bifunctional fusion consisted of a UPO catalyzing the transformation of dextromethorphan to dextrorphan and an AAO supplying UPO with H₂O₂ through the oxidation of primary aromatic alcohols. The engineered SoLo variant of UPO from *Agrocybe aegerita* with reduced peroxidase activity towards aromatic alcohols and the PeAAO variant FX9 were chosen as fusion partners. At first, several enzyme fusion libraries were constructed consisting of fusions with different partner order and rigid or flexible linkers of different length and compositions between UPO and AAO. For secretion of the fusion construct in *S. cerevisiae*, two evolved leader sequences were tested, the pre α proK leader peptide for AAO and an evolved signal peptide for UPO. Five UPO-AAO fusions with the highest activities were purified and biochemically characterized. To exclude unwanted interactions of AAO substrates with the UPO partner, several primary aromatic alcohols were tested, and *p*-fluorobenzyl alcohol, 3-methoxybenzyl alcohol, and *p*-methoxybenzyl alcohol led to the best results. By using UPO-AAO fusion with *p*-fluorobenzyl alcohol as sacrificial substrate, 2 mM dextrorphan was obtained from 2 mM dextromethorphan after 24 h, and a TTN of 48,300 was achieved. Since oxidation of *p*-fluorobenzyl alcohol proceeded much faster than dextromethorphan hydroxylation, accumulated H₂O₂ most likely inactivated the UPO partner. Feeding of *p*-fluorobenzyl alcohol at 0.5 mM/h resulted in increased TTN for the synthesis of dextrorphan of over 62,000. The construction of UPO-AAO fusions for the synthesis of pharmaceutical or chemical compounds further expands the scope of application of AAOs.

Production of bio-based precursors for polymers

In times of an emerging bioeconomy, the sustainable production of bio-based polymer precursors or platform chemicals becomes more and more important. The use of bio-based 2,5-furandicarboxylic acid (FDCA) for the production of the biopolymer polyethylene furanoate (PEF), which could replace polyethylene terephthalate (PET), strongly contributes to establishing a sustainable bioeconomy. The chemical synthesis of FDCA takes place at high temperature and pressure, and involves organic solvents and metal catalysts (Sajid et al. 2018). The oxidation of 5-hydroxymethylfurfural (5-HMF) to FDCA

proceeds via 2,5-diformylfuran (DFF) and 5-formylfuran carboxylic acid (FFCA). Enzymatic routes from 5-HMF to FDCA, involving, for example, AAO (Fig. 6), can be conducted at ambient temperature without the need of polluting co-solvents. 5-HMF can be obtained by acid-catalyzed dehydration of fructose. Dijkman and Fraaije (2014) described a HMF oxidase (HMFO) from *Methylovorus* sp. strain MP688, a member of the GMC oxidoreductase family with AAO activity, which was expressed in *E. coli* and oxidized 5-HMF to FFCA as the main product (92%), but only minor amounts of FDCA (8%) were formed after 5 h of reaction. Recently, two other HMFOs from *P. nitroreducens* and *Pseudomonas* sp. strain 11/12A, both heterologously expressed in *E. coli*, were shown to catalyze oxidation of 5-HMF to up to 99% FDCA after 96 h (Vinambres et al. 2020). Carro and coworkers found that PeAAO expressed in *E. coli* also catalyzed the conversion of 5-HMF to FFCA, but no subsequent oxidation of FFCA to FDCA was detected. To complete the second reaction, an unspecific peroxygenase (UPO) from *A. aegerita* was added after oxidation of 5-HMF to FFCA yielding 91% FDCA after 120 h. Another approach was based on the combined action of three fungal enzymes, AAO, UPO, and galactose oxidase (GAO), for the conversion of 5-HMF to FDCA (Karich et al. 2018). AAO catalyzed the oxidation of 5-HMF to DFF along with the production of H₂O₂ to be used by UPO for the oxidation of 5-HMF to 5-hydroxymethyl-2-furancarboxylic acid (HMFCFA) and to FDCA via the intermediates DFF and FFCA. GAO provided additional H₂O₂ for UPO while oxidizing 5-HMF to DFF and particularly HMFCFA to FFCA. Starting with 10 mM 5-HMF, 7.9 mM FDCA was formed in this three-enzyme approach after 24 h. Serrano and colleagues recently reported that PeAAO expressed in *E. coli* catalyzes also the oxidation of FFCA to FDCA, with a 100-fold lower reaction rate compared to 5-HMF or DFF oxidation (Serrano et al. 2019a). They found that the oxidation of FFCA was negatively affected by the presence of H₂O₂ particularly at concentrations over 400 μ M, although the enzyme was quite stable in the presence of peroxide. By applying catalase together with PeAAO, 5-HMF was completely oxidized to FDCA after 12 days, which confirmed the inhibitory effect of H₂O₂ on FFCA oxidation. When using PeAAO variant I500M/F501W in catalase-assisted conversion of 15 mM 5-HMF to FDCA, a nearly 2-fold increase in total turnover number (TTN) was achieved.

Vina-Gonzalez and colleagues further engineered PeAAO for improved conversion of 5-HMF to FDCA (Vina-Gonzalez et al. 2020). Combinatorial saturation mutagenesis of the variant FX9 at positions involved in substrate positioning in the active site, Ile500 and Phe501 in the first round, and Tyr92 and His546 in the second round, leading to the Bantha variant with the additional mutation F501W as the most active enzyme. The catalytic efficiency of the Bantha variant was 3-fold higher compared to PeAAO wild-type. Formation of FDCA was 6-fold higher when using the Bantha variant and




Fig. 6 Enzymatic oxidation of HMF and MMF to FDCA involving AAO

reached 3.0% after 2 days of reaction starting from 2 mM 5-HMF as substrate.

An alternative route for the production of FDCA is based on 5-methoxymethylfurfural (5-MMF) as starting compound, which is more stable upon storage than 5-HMF. For this route, another three-enzyme system was chosen that consisted of AAO, UPO, and methanol oxidase (MOX) (Carro et al. 2018b). AAO catalyzed the oxidation of 5-MMF to MMFA, while UPO catalyzed the two-step oxidation of MMFA to FDCA via the intermediate FFCA. The activity of UPO was supported by H_2O_2 generated by AAO and additionally in the MOX-mediated oxidation of methanol to formaldehyde. After 120 h, 98% FDCA were formed starting with 1.5 mM 5-MMF. These examples illustrate the usefulness of AAOs for valorization of products from lignocellulose biorefinery.

Concluding remarks and future perspectives

More than 60 years after the discovery of AAO in *T. versicolor*, a large number of AAO encoding genes were identified in the genomes of many wood-decaying fungi. Their functions in nature and the general catalytic mechanism are considered deciphered, and the reactions catalyzed by AAOs and summarized here impressively demonstrate the high potential of these enzymes for biotechnological purposes. Advances in protein engineering have prompted the construction of AAO variants with altered substrate scope and increased stereoselectivity, which opens new perspectives for synthetic chemistry. Realization of AAO-catalyzed reactions in two-phase systems and in a flow reactor illustrates the high robustness of these biocatalysts and their potential applicability in technical processes at high scale.

However, structural and mechanistic aspects have been tackled mainly based on the crystal structure and molecular modelling of one representative - *PeAAO* from *P. eryngii*. In

most cases, this enzyme and mutants thereof were successfully used for the synthesis of valuable compounds. So far, only a few other AAOs were purified and characterized in respect to their biochemical properties and substrate spectra. Since AAO sequences demonstrate low identity, three-dimensional structures of other members would provide more insights into catalysis and substrate recognition, and allow to rationalize differences in substrate spectra and properties of these enzymes. In 2020, the second structure of an AAO originating from *T. thermophilus* has been solved, which revealed several significant differences compared to the structure of *PeAAO*. Structural information on other AAOs and their comparison could build a broader background for enzyme engineering to promote recognition and oxidation of industrially relevant compounds with high activity and selectivity. However, heterologous expression of AAOs in recombinant hosts remains a challenging task and seems to limit their broader application and protein engineering. Although significant efforts have been undertaken to enhance heterologous expression of AAOs, no straightforward strategy has been established yet. Engineering of the signal peptide facilitated protein secretion of *PeAAO* in *S. cerevisiae* in the culture medium, but also mutations in the protein sequence positively influenced gene expression (Vina-Gonzalez et al. 2015, 2018). Another intriguing question is why expression of *PeAAO* in *P. pastoris* failed, whereas the homologous *PeAAO2* with only seven different amino acids was expressed at a concentration of 315 mg/L. Obviously, more deep understanding of the effect of these amino acids on transcription, translation, protein folding, secretion, and stability is required and will remain in focus of future studies.

The scope of AAO applications can be expanded by their exploiting as supplier of H_2O_2 for various peroxide-dependent enzymes. A contribution to enzyme-mediated delignification of plant biomass in biorefinery concepts seems plausible since ligninolytic peroxidases relying on H_2O_2 play a major role in

lignin degradation in nature. Along with their use in biobleaching and biopulping in the pulp and paper industry, AAOs might also support peroxidases in polymer synthesis, bioremediation of phenolic compounds, peroxidase-based biosensors, and biocatalysis (Regalado et al. 2004). Reaction engineering will focus on developing substrate-feeding strategies to control the level of H₂O₂ released by AAOs in order to avoid its negative effect on heme-containing peroxidases and peroxygenases. Combinations of AAOs with other biocatalysts in multi-step cascade reactions offer alternative routes for the synthesis of various molecules and provide exciting future perspectives to access novel compounds that cannot be synthesized by classical chemical routes.

Thus, the concepts described in this review build a solid background for further development of AAOs towards applications with efficiencies approaching economic viability.

Author contribution VU and KK conceived and wrote the manuscript. VU and KK conceived and designed the figures. All authors read and approved the final manuscript.

Funding Open access funding enabled and organized by Projekt DEAL. The scientific activities of the Bioeconomy Science Center were financially supported by the Ministry of Innovation, Science and Research of North Rhine-Westphalia, Germany, within the framework of the NRW-Strategieprojekt BioSC (No. 313/323-400-002 13).

Availability of data and material All data on which the conclusions were drawn are presented in this study.

Code availability Not applicable.

Declarations

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

Consent to participate Not applicable.

Consent for publication Not applicable.

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ahmad M, Hirz M, Pichler H, Schwab H (2014) Protein expression in *Pichia pastoris*: recent achievements and perspectives for heterologous protein production. *Appl Microbiol Biotechnol* 98(12):5301–5317. <https://doi.org/10.1007/s00253-014-5732-5>
- Asada Y, Watanabe A, Ohtsu Y, Kuwahara M (1995) Purification and characterization of an aryl-alcohol oxidase from the lignin-degrading basidiomycete *Phanerochaete chrysosporium*. *Biosci Biotechnol Biochem* 59(7):1339–1341. <https://doi.org/10.1271/bbb.59.1339>
- Becker J, Wittmann C (2019) A field of dreams: lignin valorization into chemicals, materials, fuels, and health-care products. *Biotechnol Adv* 37(6):107360. <https://doi.org/10.1016/j.biotechadv.2019.02.016>
- Berka RM, Grigoriev IV, Otiillar R, Salamov A, Grimwood J, Reid I, Ishmael N, John T, Darmond C, Moisan MC, Henrissat B, Coutinho PM, Lombard V, Natvig DO, Lindquist E, Schmutz J, Lucas S, Harris P, Powlowski J, Bellemare A, Taylor D, Butler G, de Vries RP, Allijn IE, van den Brink J, Ushinsky S, Storms R, Powell AJ, Paulsen IT, Elbourne LD, Baker SE, Magnuson J, Laboissiere S, Clutterbuck AJ, Martinez D, Wogulis M, de Leon AL, Rey MW, Tsang A (2011) Comparative genomic analysis of the thermophilic biomass-degrading fungi *Myceliophthora thermophila* and *Thielavia terrestris*. *Nat Biotechnol* 29(10):922–927. <https://doi.org/10.1038/nbt.1976>
- Bhat SG, Vaidyanathan CS (1976) Purification and properties of L-4-hydroxymandelate oxidase from *Pseudomonas convexa*. *Eur J Biochem* 68(2):323–331. <https://doi.org/10.1111/j.1432-1033.1976.tb10818.x>
- Bourbonnais R, Paice MG (1988) Veratryl alcohol oxidases from the lignin-degrading basidiomycete *Pleurotus sajor-caju*. *Biochem J* 255(2):445–450. <https://doi.org/10.1042/bj2550445>
- Brückmann M, Termonia A, Pasteels JM, Hartmann T (2002) Characterization of an extracellular salicyl alcohol oxidase from larval defensive secretions of *Chrysomela populi* and *Phratora vitellinae* (Chrysomelina). *Insect Biochem Mol Biol* 32(11):1517–1523. [https://doi.org/10.1016/s0965-1748\(02\)00072-3](https://doi.org/10.1016/s0965-1748(02)00072-3)
- Brum JOC, Neto DCF, de Almeida JSFD, Lima JA, Kuca K, França TCC, Figueroa-Villar JD (2019) Synthesis of new quinoline-piperonal hybrids as potential drugs against Alzheimer's disease. *Int J Mol Sci* 20(16):3944. <https://doi.org/10.3390/ijms20163944>
- Bugg TDH, Ahmad M, Hardiman EM, Rahmanpour R (2011) Pathways for degradation of lignin in bacteria and fungi. *Nat Prod Rep* 28(12):1883–1896. <https://doi.org/10.1039/c1np00042j>
- Carro J, Martinez-Julvez M, Medina M, Martinez AT, Ferreira P (2017) Protein dynamics promote hydride tunnelling in substrate oxidation by aryl-alcohol oxidase. *Phys Chem Chem Phys* 19(42):28666–28675. <https://doi.org/10.1039/c7cp05904c>
- Carro J, Amengual-Rigo P, Sancho F, Medina M, Guallar V, Ferreira P, Martinez AT (2018a) Multiple implications of an active site phenylalanine in the catalysis of aryl-alcohol oxidase. *Sci Rep* 8(1):8121. <https://doi.org/10.1038/s41598-018-26445-x>
- Carro J, Fernandez-Fueyo E, Fernandez-Alonso C, Canada J, Ullrich R, Hofrichter M, Alcalde M, Ferreira P, Martinez AT (2018b) Self-sustained enzymatic cascade for the production of 2,5-furandicarboxylic acid from 5-methoxymethylfurfural. *Biotechnol Biofuels* 11:10. <https://doi.org/10.1186/s13068-018-1091-2>
- Carro J, Ferreira P, Martinez AT, Gadda G (2018c) Stepwise hydrogen atom and proton transfers in dioxygen reduction by aryl-alcohol oxidase. *Biochemistry* 57(11):1790–1797. <https://doi.org/10.1021/acs.biochem.8b00106>
- Casieri L, Varese GC, Anastasi A, Prigione V, Svobodova K, Marchisio VF, Novotny C (2008) Decolorization and detoxication of reactive

- industrial dyes by immobilized fungi *Trametes pubescens* and *Pleurotus ostreatus*. *Folia Microbiol* 53(1):44–52. <https://doi.org/10.1007/s12223-008-0006-1>
- Cavener DR (1992) GMC Oxidoreductases - a newly defined family of homologous proteins with diverse catalytic activities. *J Mol Biol* 223(3):811–814. [https://doi.org/10.1016/0022-2836\(92\)90992-S](https://doi.org/10.1016/0022-2836(92)90992-S)
- Chakraborty M, Goel M, Chinnadayala SR, Dahiya UR, Ghosh SS, Goswami P (2014) Molecular characterization and expression of a novel alcohol oxidase from *Aspergillus terreus* MTCC6324. *PLoS One* 9(4):19. <https://doi.org/10.1371/journal.pone.0095368>
- Couturier M, Mathieu Y, Li A, Navarro D, Drula E, Haon M, Grisel S, Ludwig R, Berrin JG (2016) Characterization of a new aryl-alcohol oxidase secreted by the phytopathogenic fungus *Ustilago maydis*. *Appl Microbiol Biotechnol* 100(2):697–706. <https://doi.org/10.1007/s00253-015-7021-3>
- Daniel G, Volc J, Filonova L, Plíhal O, Kubátová E, Halada P (2007) Characteristics of *Gloeophyllum trabeum* alcohol oxidase, an extracellular source of H₂O₂ in brown rot decay of wood. *Appl Environ Microbiol* 73(19):6241–6253. <https://doi.org/10.1128/aem.00977-07>
- de Almeida TP, van Schie MMCH, Ma A, Tieves F, Younes SHH, Fernandez-Fueyo E, Arends IWCE, Riul A, Hollmann F (2019) Efficient aerobic oxidation of *trans*-2-hexen-1-ol using the aryl alcohol oxidase from *Pleurotus eryngii*. *Adv Synth Catal* 361(11):2668–2672. <https://doi.org/10.1002/adsc.201801312>
- de Gonzalo G, Colpa DI, Habib MHM, Fraaije MW (2016) Bacterial enzymes involved in lignin degradation. *J Biotechnol* 236:110–119. <https://doi.org/10.1016/j.jbiotec.2016.08.011>
- de Jong E, de Vries FP, Field JA, van der Zwan RP, de Bont JAM (1992a) Isolation and screening of basidiomycetes with high peroxidative activity. *Mycol Res* 96:1098–1104. [https://doi.org/10.1016/S0953-7562\(09\)80121-4](https://doi.org/10.1016/S0953-7562(09)80121-4)
- de Jong E, van Berkel WJ, van der Zwan RP, de Bont JA (1992b) Purification and characterization of vanillyl-alcohol oxidase from *Penicillium simplicissimum*. A novel aromatic alcohol oxidase containing covalently bound FAD. *Eur J Biochem* 208(3):651–657. <https://doi.org/10.1111/j.1432-1033.1992.tb17231.x>
- de Jong E, Cazemier AE, Field JA, de Bont JA (1994) Physiological role of chlorinated aryl alcohols biosynthesized de novo by the white rot fungus *Bjerkandera* sp. strain BOS55. *Appl Environ Microbiol* 60(1):271–277. <https://doi.org/10.1128/AEM.60.1.271-277.1994>
- de Santos PG, Lazaro S, Vina-Gonzalez J, Hoang MD, Sanchez-Moreno I, Glieder A, Hollmann F, Alcalde M (2020) Evolved peroxygenase-aryl alcohol oxidase fusions for self-sufficient oxyfunctionalization reactions. *ACS Catal* 10(22):13524–13534. <https://doi.org/10.1021/acscatal.0c03029>
- Dijkman WP, Fraaije MW (2014) Discovery and characterization of a 5-hydroxymethylfurfural oxidase from *Methylovorus* sp. strain MP688. *Appl Environ Microbiol* 80(3):1082–1090. <https://doi.org/10.1128/aem.03740-13>
- Dijkman WP, de Gonzalo G, Mattevi A, Fraaije MW (2013) Flavoprotein oxidases: classification and applications. *Appl Microbiol Biotechnol* 97(12):5177–5188. <https://doi.org/10.1007/s00253-013-4925-7>
- Ewing TA, Dijkman WP, Vervoort JM, Fraaije MW, van Berkel WJ (2014) The oxidation of thiols by flavoprotein oxidases: a biocatalytic route to reactive thiocarbonyls. *Angew Chem Int Ed Engl* 53(48):13206–13209. <https://doi.org/10.1002/anie.201407520>
- Ewing TA, Gygli G, Fraaije MW, van Berkel WJH (2020) Vanillyl alcohol oxidase. *Enzymes* 47:87–116. <https://doi.org/10.1016/bs.enz.2020.05.003>
- Farmer VC, Henderson MEK, Russell JD (1960) Aromatic-alcohol-oxidase activity in the growth medium of *Polystictus versicolor*. *Biochem J* 74(2):257–262. <https://doi.org/10.1042/bj0740257>
- Fernandez IS, Ruiz-Duenas FJ, Santillana E, Ferreira P, Martinez MJ, Martinez AT, Romero A (2009) Novel structural features in the GMC family of oxidoreductases revealed by the crystal structure of fungal aryl-alcohol oxidase. *Acta Crystallogr D Biol Crystallogr* 65(Pt 11):1196–1205. <https://doi.org/10.1107/S0907444909035860>
- Ferreira P, Medina M, Guillen F, Martinez MJ, Van Berkel WJH, Martinez AT (2005) Spectral and catalytic properties of aryl-alcohol oxidase, a fungal flavoenzyme acting on polyunsaturated alcohols. *Biochem J* 389:731–738. <https://doi.org/10.1042/Bj20041903>
- Ferreira P, Ruiz-Duenas FJ, Martinez MJ, van Berkel WJ, Martinez AT (2006) Site-directed mutagenesis of selected residues at the active site of aryl-alcohol oxidase, an H₂O₂-producing ligninolytic enzyme. *FEBS J* 273(21):4878–4888. <https://doi.org/10.1111/j.1742-4658.2006.05488.x>
- Ferreira P, Hernandez-Ortega A, Herguedas B, Rencoret J, Gutierrez A, Martinez MJ, Jimenez-Barbero J, Medina M, Martinez AT (2010) Kinetic and chemical characterization of aldehyde oxidation by fungal aryl-alcohol oxidase. *Biochem J* 425:585–593. <https://doi.org/10.1042/Bj20091499>
- Ferreira P, Carro J, Serrano A, Martinez AT (2015a) A survey of genes encoding H₂O₂-producing GMC oxidoreductases in 10 *Polyporales* genomes. *Mycologia* 107(6):1105–1119. <https://doi.org/10.3852/15-027>
- Ferreira P, Hernandez-Ortega A, Lucas F, Carro J, Herguedas B, Borrelli KW, Guallar V, Martinez AT, Medina M (2015b) Aromatic stacking interactions govern catalysis in aryl-alcohol oxidase. *FEBS J* 282(16):3091–3106. <https://doi.org/10.1111/febs.13221>
- Galperin I, Javeed A, Luig H, Lochnit G, Ruhl M (2016) An aryl-alcohol oxidase of *Pleurotus sapidus*: heterologous expression, characterization, and application in a 2-enzyme system. *Appl Microbiol Biotechnol* 100(18):8021–8030. <https://doi.org/10.1007/s00253-016-7567-8>
- Goetghebeur M, Nicolas M, Brun S, Galzy P (1992) Purification and properties of benzyl alcohol oxidase from *Botrytis cinerea*. *Biosci Biotechnol Biochem* 56(2):298–303. <https://doi.org/10.1271/bbb.56.298>
- Guillen F, Evans CS (1994) Anisaldehyde and veratraldehyde acting as redox cycling agents for H₂O₂ production by *Pleurotus eryngii*. *Appl Environ Microbiol* 60(8):2811–2817
- Guillen F, Martinez AT, Martinez MJ (1990) Production of hydrogenperoxide by aryl-alcohol oxidase from the ligninolytic fungus *Pleurotus eryngii*. *Appl Microbiol Biotechnol* 32(4):465–469. <https://doi.org/10.1007/Bf00903784>
- Guillen F, Martinez AT, Martinez MJ (1992) Substrate-specificity and properties of the aryl-alcohol oxidase from the ligninolytic fungus *Pleurotus eryngii*. *Eur J Biochem* 209(2):603–611. <https://doi.org/10.1111/j.1432-1033.1992.tb17326.x>
- Hall M (2020) Flavoenzymes for biocatalysis. *Enzymes* 47:37–62. <https://doi.org/10.1016/bs.enz.2020.05.001>
- Hammel KE (1997) Fungal degradation of lignin. In: Cadisch G, Giller KE (eds) *Driven by nature: plant litter quality and decomposition*. CAB International, Oxon, pp 33–45
- Hernandez-Ortega A, Borrelli K, Ferreira P, Medina M, Martinez AT, Guallar V (2011a) Substrate diffusion and oxidation in GMC oxidoreductases: an experimental and computational study on fungal aryl-alcohol oxidase. *Biochem J* 436(2):341–350. <https://doi.org/10.1042/BJ20102090>
- Hernandez-Ortega A, Lucas F, Ferreira P, Medina M, Guallar V, Martinez AT (2011b) Modulating O₂ reactivity in a fungal flavoenzyme: involvement of aryl-alcohol oxidase Phe-501 contiguous to catalytic histidine. *J Biol Chem* 286(47):41105–41114. <https://doi.org/10.1074/jbc.M111.282467>
- Hernandez-Ortega A, Ferreira P, Martinez AT (2012a) Fungal aryl-alcohol oxidase: a peroxide-producing flavoenzyme involved in lignin degradation. *Appl Microbiol Biotechnol* 93(4):1395–1410. <https://doi.org/10.1007/s00253-011-3836-8>
- Hernandez-Ortega A, Ferreira P, Merino P, Medina M, Guallar V, Martinez AT (2012b) Stereoselective hydride transfer by aryl-

- alcohol oxidase, a member of the GMC superfamily. *ChemBioChem* 13(3):427–435. <https://doi.org/10.1002/cbic.201100709>
- Hernandez-Ortega A, Lucas F, Ferreira P, Medina M, Guallar V, Martinez AT (2012c) Role of active site histidines in the two half-reactions of the aryl-alcohol oxidase catalytic cycle. *Biochemistry* 51(33):6595–6608. <https://doi.org/10.1021/bi300505z>
- Iida K, Cox-Foster DL, Yang XL, Ko WY, Cavener DR (2007) Expansion and evolution of insect GMC oxidoreductases. *BMC Evol Biol* 7:75. <https://doi.org/10.1186/1471-2148-7-75>
- Jankowski N, Koschorreck K, Urlacher VB (2020) High-level expression of aryl-alcohol oxidase 2 from *Pleurotus eryngii* in *Pichia pastoris* for production of fragrances and bioactive precursors. *Appl Microbiol Biotechnol* 104(21):9205–9218. <https://doi.org/10.1007/s00253-020-10878-4>
- Kadowaki MAS, Higasi PMR, de Godoy MO, de Araujo EA, Godoy AS, Prade RA, Polikarpov I (2020) Enzymatic versatility and thermostability of a new aryl-alcohol oxidase from *Thermothelomyces thermophilus* M77. *Biochim Biophys Acta* 1864(10):129681. <https://doi.org/10.1016/j.bbagen.2020.129681>
- Karich A, Kleeborg SB, Ullrich R, Hofrichter M (2018) Enzymatic preparation of 2,5-furandicarboxylic acid (FDCA)-a substitute of terephthalic acid-by the joined action of three fungal enzymes. *Microorganisms* 6(1):12. <https://doi.org/10.3390/microorganisms6010005>
- Kim SJ, Suzuki N, Uematsu Y, Shoda M (2001) Characterization of aryl alcohol oxidase produced by dye-decolorizing fungus, *Geotrichum candidum* Dec1. *J Biosci Bioeng* 91(2):166–172. <https://doi.org/10.1263/jbb.91.166>
- Large AT, Connock MJ (1993) Aromatic alcohol oxidase and aldehyde oxidase activities in the digestive gland of 3 species of terrestrial gastropod (*Helix aspersa*, *Arion ater*, *Limax flavus*). *Comp Biochem Physiol B: Biochem Mol Biol* 104(3):489–491. [https://doi.org/10.1016/0305-0491\(93\)90271-6](https://doi.org/10.1016/0305-0491(93)90271-6)
- Ledakowicz S, Pazdzior K (2021) Recent achievements in dyes removal focused on advanced oxidation processes integrated with biological methods. *Molecules* 26(4):870. <https://doi.org/10.3390/molecules26040870>
- Lee HS (2005) Cuminaldehyde: aldose reductase and α -glucosidase inhibitor derived from *Cuminum cyminum* L. seeds. *J Agric Food Chem* 53:2446–2450. <https://doi.org/10.1021/jf048451g>
- Levasseur A, Drula E, Lombard V, Coutinho PM, Henrissat B (2013) Expansion of the enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes. *Biotechnol Biofuels* 6:41. <https://doi.org/10.1186/1754-6834-6-41>
- Levasseur A, Lomascolo A, Chabrol O, Ruiz-Duenas FJ, Boukhris-Uzan E, Piumi F, Kues U, Ram AF, Murat C, Haon M, Benoit I, Arfi Y, Chevret D, Drula E, Kwon MJ, Gouret P, Lesage-Meessen L, Lombard V, Mariette J, Noirot C, Park J, Patyshakuliyeva A, Sigoillot JC, Wiebenga A, Wosten HA, Martin F, Coutinho PM, de Vries RP, Martinez AT, Klopp C, Pontarotti P, Henrissat B, Record E (2014) The genome of the white-rot fungus *Pycnoporus cinnabarinus*: a basidiomycete model with a versatile arsenal for lignocellulosic biomass breakdown. *BMC Genomics* 15:486. <https://doi.org/10.1186/1471-2164-15-486>
- Li R, Jiang ZT (2004) Chemical composition of the essential oil of *Cuminum cyminum* L. from China. *Flavour Fragrance J* 19:311–313. <https://doi.org/10.1002/ffj.1302>
- Liu ES, Wilkins MR (2020) Process optimization and scale-up production of fungal aryl alcohol oxidase from genetically modified *Aspergillus nidulans* in stirred-tank bioreactor. *Bioresour Technol* 315:123792. <https://doi.org/10.1016/j.biortech.2020.123792>
- Liu ES, Li MX, Abdella A, Wilkins MR (2020) Development of a cost-effective medium for submerged production of fungal aryl alcohol oxidase using a genetically modified *Aspergillus nidulans* strain. *Bioresour Technol* 305:123038. <https://doi.org/10.1016/j.biortech.2020.123038>
- Mann V, Large A, Khan S, Malik Z, Connock MJ (1989) Aromatic alcohol oxidase - a new membrane-bound H₂O₂-generating enzyme in alimentary tissues of the slug *Arion ater*. *J Exp Zool* 251(3):265–274. <https://doi.org/10.1002/jez.1402510302>
- Martin C, Binda C, Fraaije MW, Mattevi A (2020) The multipurpose family of flavoprotein oxidases. *Enzymes* 47:63–86. <https://doi.org/10.1016/bs.enz.2020.05.002>
- Martinez AT, Rencoret J, Nieto L, Jimenez-Barbero J, Gutierrez A, del Rio JC (2011) Selective lignin and polysaccharide removal in natural fungal decay of wood as evidenced by in situ structural analyses. *Environ Microbiol* 13(1):96–107. <https://doi.org/10.1111/j.1462-2920.2010.02312.x>
- Mathieu Y, Piumi F, Valli R, Aramburu JC, Ferreira P, Faulds CB, Record E (2016) Activities of secreted aryl alcohol quinone oxidoreductases from *Pycnoporus cinnabarinus* provide insights into fungal degradation of plant biomass. *Appl Environ Microbiol* 82(8):2411–2423. <https://doi.org/10.1128/Aem.03761-15>
- Morshedi D, Aliakbari F, Tayaranian-Marvian A, Fassihi A, Pan-Montojo F, Pérez-Sánchez H (2015) Cuminaldehyde as the major component of *Cuminum cyminum*, a natural aldehyde with inhibitory effect on alpha-synuclein fibrillation and cytotoxicity. *J Food Sci* 80:H2336–H2345. <https://doi.org/10.1111/1750-3841.13016>
- Muheim A, Leisola MSA, Schoemaker HE (1990a) Aryl-alcohol oxidase and lignin peroxidase from the white-rot fungus *Bjerkandera adusta*. *J Biotechnol* 13(2-3):159–167. [https://doi.org/10.1016/0168-1656\(90\)90101-G](https://doi.org/10.1016/0168-1656(90)90101-G)
- Muheim A, Waldner R, Leisola MSA, Fiechter A (1990b) An extracellular aryl-alcohol oxidase from the white-rot fungus *Bjerkandera adusta*. *Enzym Microb Technol* 12(3):204–209. [https://doi.org/10.1016/0141-0229\(90\)90039-S](https://doi.org/10.1016/0141-0229(90)90039-S)
- Pardo-Planas O, Prade RA, Wilkins MR (2017) High-yield production of aryl alcohol oxidase under limited growth conditions in small-scale systems using a mutant *Aspergillus nidulans* strain. *J Ind Microbiol Biotechnol* 44(2):247–257. <https://doi.org/10.1007/s10295-016-1884-x>
- Patil SB, Takalikar SS, Joglekar MM, Haldavnekar VS, Arvindekar AU (2013) Insulinotropic and β -cell protective action of cuminaldehyde, cuminol and an inhibitor isolated from *Cuminum cyminum* in streptozotocin-induced diabetic rats. *Br J Nutr* 110:1434–1443. <https://doi.org/10.1017/S0007114513000627>
- Paul CE, Eggerichs D, Westphal AH, Tischler D, van Berkel WJH (2021) Flavoprotein monooxygenases: versatile biocatalysts. *Biotechnol Adv*:107712. <https://doi.org/10.1016/j.biotechadv.2021.107712>
- Regalado V, Perestelo F, Rodriguez A, Carnicero A, Sosa FJ, De la Fuente G, Falcon MA (1999) Activated oxygen species and two extracellular enzymes: laccase and aryl-alcohol oxidase, novel for the lignin-degrading fungus *Fusarium proliferatum*. *Appl Microbiol Biotechnol* 51(3):388–390. <https://doi.org/10.1007/s002530051407>
- Riyadi FA, Tahir AA, Yusof N, Sabri NSA, Noor M, Akhri F, Othman N, Zakaria Z, Hara H (2020) Enzymatic and genetic characterization of lignin depolymerization by *Streptomyces* sp. S6 isolated from a tropical environment. *Sci Rep* 10(1):9. <https://doi.org/10.1038/s41598-020-64817-4>
- Romero E, Ferreira P, Martinez AT, Martinez MJ (2009) New oxidase from *Bjerkandera* arthroconidial anamorph that oxidizes both phenolic and nonphenolic benzyl alcohols. *Biochim Biophys Acta* 1794(4):689–697. <https://doi.org/10.1016/j.bbapap.2008.11.013>
- Ruiz-Duenas FJ, Ferreira P, Martinez MJ, Martinez AT (2006) In vitro activation, purification, and characterization of *Escherichia coli* expressed aryl-alcohol oxidase, a unique H₂O₂-producing enzyme. *Protein Expr Purif* 45(1):191–199. <https://doi.org/10.1016/j.pep.2005.06.003>
- Sajid M, Zhao XB, Liu DH (2018) Production of 2,5-furandicarboxylic acid (FDCA) from 5-hydroxymethylfurfural (HMF): recent progress

- focusing on the chemical-catalytic routes. *Green Chem* 20(24): 5427–5453. <https://doi.org/10.1039/c8gc02680g>
- Sannia G, Limongi P, Cocca E, Buonocore F, Nitti G, Giardina P (1991) Purification and characterization of a veratryl alcohol oxidase enzyme from the lignin degrading basidiomycete *Pleurotus ostreatus*. *Biochim Biophys Acta* 1073(1):114–119. [https://doi.org/10.1016/0304-4165\(91\)90190-r](https://doi.org/10.1016/0304-4165(91)90190-r)
- Santos AS, Pereira N, Da Silva IM, Sarquis MIM, Antunes OAC (2004) Peroxidase catalyzed microbiological oxidation of isosafrol into piperonal. *Process Biochem* 39:2269–2275. <https://doi.org/10.1016/j.procbio.2003.11.019>
- Serrano A, Calvino E, Carro J, Sanchez-Ruiz MI, Canada FJ, Martinez AT (2019a) Complete oxidation of hydroxymethylfurfural to furandicarboxylic acid by aryl-alcohol oxidase. *Biotechnol Biofuels* 12(1):217. <https://doi.org/10.1186/s13068-019-1555-z>
- Serrano A, Sancho F, Vina-Gonzalez J, Carro J, Alcalde M, Guallar V, Martinez AT (2019b) Switching the substrate preference of fungal aryl-alcohol oxidase: towards stereoselective oxidation of secondary benzyl alcohols. *Catal Sci Technol* 9(3):833–841. <https://doi.org/10.1039/c8cy02447b>
- Serrano A, Carro J, Martinez AT (2020) Reaction mechanisms and applications of aryl-alcohol oxidase. *Enzymes* 47:167–192. <https://doi.org/10.1016/bs.enz.2020.05.005>
- Sigoillot C, Camarero S, Vidal T, Record E, Asther M, Perez-Boada M, Martinez MJ, Sigoillot JC, Asther M, Colom JF, Martinez AT (2005) Comparison of different fungal enzymes for bleaching high-quality paper pulps. *J Biotechnol* 115(4):333–343. <https://doi.org/10.1016/j.jbiotec.2004.09.006>
- Sucharitakul J, Prongjit M, Haltrich D, Chaiyen P (2008) Detection of a C4a-hydroperoxyflavin intermediate in the reaction of a flavoprotein oxidase. *Biochemistry* 47(33):8485–8490. <https://doi.org/10.1021/bi801039d>
- Sütlz L, Laurent CVFP, Abrera AT, Schütz G, Ludwig R, Haltrich D (2018) Multiplicity of enzymatic functions in the CAZy AA3 family. *Appl Microbiol Biotechnol* 102:2477–2492. <https://doi.org/10.1007/s00253-018-8784-0>
- Sütlz L, Foley G, Gillam EMJ, Boden M, Haltrich D (2019) The GMC superfamily of oxidoreductases revisited: analysis and evolution of fungal GMC oxidoreductases. *Biotechnol Biofuels* 12:118. <https://doi.org/10.1186/s13068-019-1457-0>
- Tamaru Y, Umezawa K, Yoshida M (2018) Characterization of an aryl-alcohol oxidase from the plant saprophytic basidiomycete *Coprinosia cinerea* with broad substrate specificity against aromatic alcohols. *Biotechnol Lett* 40(7):1077–1086. <https://doi.org/10.1007/s10529-018-2534-3>
- Tamboli DP, Telke AA, Dawkar VV, Jadhav SB, Govindwar SP (2011) Purification and characterization of bacterial aryl alcohol oxidase from *Sphingobacterium* sp. ATM and its uses in textile dye decolorization. *Biotechnol Bioprocess Eng* 16(4):661–668. <https://doi.org/10.1007/s12257-011-0031-9>
- Toplak M, Matthews A, Teufel R (2021) The devil is in the details: the chemical basis and mechanistic versatility of flavoprotein monooxygenases. *Arch Biochem Biophys* 698:108732. <https://doi.org/10.1016/j.abb.2020.108732>
- Tsai KD, Liu YH, Chen TW, Yang SM, Wong HY, Cherng J, Chou KS, Cherng JM (2016) Cinnamaldehyde from *Cinnamomum verum* induces cell death through targeting topoisomerase 1 and 2 in human colorectal adenocarcinoma COLO 205 cells. *Nutrients* 8:1–17. <https://doi.org/10.3390/nu8060318>
- van Schie MMCH, de Almeida TP, Laudadio G, Tieves F, Fernandez-Fueyo E, Noel T, Arends IWCE, Hollmann F (2018) Biocatalytic synthesis of the Green Note *trans*-2-hexenal in a continuous-flow microreactor. *Beilstein J Org Chem* 14:697–703. <https://doi.org/10.3762/bjoc.14.58>
- Varela E, Bockle B, Romero A, Martinez AT, Martinez MJ (2000) Biochemical characterization, cDNA cloning and protein crystallization of aryl-alcohol oxidase from *Pleurotus pulmonarius*. *Biochim Biophys Acta* 1476(1):129–138. [https://doi.org/10.1016/S0167-4838\(99\)00227-7](https://doi.org/10.1016/S0167-4838(99)00227-7)
- Varela E, Guillen F, Martinet AT, Martinez MJ (2001) Expression of *Pleurotus eryngii* aryl-alcohol oxidase in *Aspergillus nidulans*: purification and characterization of the recombinant enzyme. *Biochim Biophys Acta* 1546(1):107–113. [https://doi.org/10.1016/S0167-4838\(00\)00301-0](https://doi.org/10.1016/S0167-4838(00)00301-0)
- Vina-Gonzalez J, Alcalde M (2020) Directed evolution of the aryl-alcohol oxidase: beyond the lab bench. *Comput Struct Biotechnol J* 18:1800–1810. <https://doi.org/10.1016/j.csbj.2020.06.037>
- Vina-Gonzalez J, Gonzalez-Perez D, Ferreira P, Martinez AT, Alcalde M (2015) Focused directed evolution of aryl-alcohol oxidase in *Saccharomyces cerevisiae* by using chimeric signal peptides. *Appl Environ Microbiol* 81(18):6451–6462. <https://doi.org/10.1128/Aem.01966-15>
- Vina-Gonzalez J, Elbl K, Ponte X, Valero F, Alcalde M (2018) Functional expression of aryl-alcohol oxidase in *Saccharomyces cerevisiae* and *Pichia pastoris* by directed evolution. *Biotechnol Bioeng* 115(7):1666–1674. <https://doi.org/10.1002/bit.26585>
- Vina-Gonzalez J, Jimenez-Lalana D, Sancho F, Serrano A, Martinez AT, Guallar V, Alcalde M (2019) Structure-guided evolution of aryl alcohol oxidase from *Pleurotus eryngii* for the selective oxidation of secondary benzyl alcohols. *Adv Synth Catal* 361(11):2514–2525. <https://doi.org/10.1002/adsc.201900134>
- Vina-Gonzalez J, Martinez AT, Guallar V, Alcalde M (2020) Sequential oxidation of 5-hydroxymethylfurfural to furan-2,5-dicarboxylic acid by an evolved aryl-alcohol oxidase. *Biochim Biophys Acta* 1868(1): 140293. <https://doi.org/10.1016/j.bbapap.2019.140293>
- Vinambres M, Espada M, Martinez AT, Serrano A (2020) Screening and evaluation of new hydroxymethylfurfural oxidases for furandicarboxylic acid production. *Appl Environ Microbiol* 86(16). <https://doi.org/10.1128/AEM.00842-20>
- Wang S, Bao L, Song D, Wang J, Cao X (2019) Heterocyclic lactam derivatives containing piperonyl moiety as potential antifungal agents. *Bioorg Med Chem Lett* 29:126661. <https://doi.org/10.1016/j.bmcl.2019.126661>
- Wesenberg D, Kyriakides I, Agathos SN (2003) White-rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnol Adv* 22(1-2):161–187. <https://doi.org/10.1016/j.biotechadv.2003.08.011>
- Wongnate T, Chaiyen P (2013) The substrate oxidation mechanism of pyranose 2-oxidase and other related enzymes in the glucose-methanol-choline superfamily. *FEBS J* 280(13):3009–3027. <https://doi.org/10.1111/febs.12280>
- Heuts DPHM, Scrutton NS, McIntire WS, Fraaije MW (2009) What's in a covalent bond? On the role and formation of covalently bound flavin cofactors. *FEBS J* 276(13):3405–3427. <https://doi.org/10.1111/j.1742-4658.2009.07053.x>
- Regalado C, García-Almendárez BE, Duarte-Vázquez MA (2004) Biotechnological applications of peroxidases. *Phytochem Rev* 3(1):243–256. <https://doi.org/10.1023/B:PHYT.0000047797.81958.69>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.