

Peculiar Type II supernovae from blue supergiants

Io K. W. Kleiser,^{1*} Dovi Poznanski,^{1,2*†} Daniel Kasen,^{2,3} Timothy R. Young,⁴
Ryan Chornock,⁵ Alexei V. Filippenko,¹ Peter Challis,⁵ Mohan Ganeshalingam,¹
Robert P. Kirshner,⁵ Weidong Li,¹ Thomas Matheson,⁶ Peter E. Nugent²
and Jeffrey M. Silverman¹

¹Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

²Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

³Department of Physics, University of California, Berkeley, CA 94720, USA

⁴Department of Physics and Astrophysics, University of North Dakota, 213 Witmer Hall, Grand Forks, ND 58202-7129, USA

⁵Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁶National Optical Astronomy Observatory, 950 North Cherry Avenue, Tucson, AZ 85719, USA

Accepted 2011 March 14. Received 2011 March 12; in original form 2010 December 17

ABSTRACT

The vast majority of Type II supernovae (SNeII) are produced by red supergiants, but SN 1987A revealed that blue supergiants (BSGs) can produce members of this class as well, albeit with some peculiar properties. This best-studied event revolutionized our understanding of SNe and linking it to the bulk of Type II events is essential. We present here the optical photometry and spectroscopy gathered for SN 2000cb, which is clearly not a standard SNeII and yet is not a SN 1987A analogue. The light curve of SN 2000cb is reminiscent of that of SN 1987A in shape, with a slow rise to a late optical peak, but on substantially different time-scales. Spectroscopically, SN 2000cb resembles a normal SNeII, but with ejecta velocities that far exceed those measured for SN 1987A or normal SNeII, above $18\,000\text{ km s}^{-1}$ for $H\alpha$ at early times. The red colours, high velocities, late photometric peak and our modelling of this object all point towards a scenario involving the high-energy explosion of a small-radius star, most likely a BSG, producing $0.1 M_{\odot}$ of ^{56}Ni . Adding a similar object to the sample, SN 2005ci, we derive a rate of ~ 2 per cent of the core-collapse rate for this loosely defined class of BSG explosions.

Key words: supernovae: general – supernovae: individual: SN 2000cb – supernovae: individual: SN 1987A – supernovae: individual: SN 2005ci – supernovae: individual: SN 1998A – supernovae: individual: SN 1999em.

1 INTRODUCTION

Massive stars that retain their hydrogen envelope during their evolution end their lives as Type II supernovae (SNeII). The vast majority of these are characterized by a fast (few days) rise to a flat light curve, most pronounced in the reddest optical bands, with a duration of 80–100 d. This ‘plateau’ phase, for which they have been named SNeII-P, is interpreted as the recession of the photosphere as the ejecta expand and cool (e.g. Kirshner et al. 1973; Barbon, Ciatti & Rosino 1979). The spectra of SNeII-P are typically dominated by strong P Cygni profiles of hydrogen lines, as well as iron absorption features (e.g. see the review by Filippenko 1997).

Due to the relatively simple physics driving their optical evolution, SNeII-P were the first SNe to be suggested as distance in-

dicators, via the so-called expanding photosphere method (EPM; Kirshner & Kwan 1974). While this method and its descendants depend on modelling, Hamuy & Pinto (2002) suggested an empirical relation between luminosity and ejecta velocity, as measured from iron absorption lines, that proved compelling when combined with the dust-extinction correction (Nugent et al. 2006; Poznanski et al. 2009; Olivares 2010; Poznanski, Nugent & Filippenko 2010).

Over the past decade, ~ 20 pre-explosion locations of SNeII-P have been directly imaged with the *Hubble Space Telescope* or deep ground-based images, yielding five detections of progenitor stars, all of which were red supergiants (RSGs), and many limits on stars with masses in the range $7.5\text{--}15 M_{\odot}$ (Smartt et al. 2009, and references therein). These masses are somewhat at odds with those derived from explosion modelling, which tend to be higher, closer to $15\text{--}25 M_{\odot}$ (e.g. Nadyozhin 2003; Utrobin, Chugai & Pastorello 2007; Utrobin & Chugai 2009), though Dessart, Livne & Waldman (2010) push towards lower masses of less than $\sim 20 M_{\odot}$.

*E-mail: io.kleiser@berkeley.edu (IKWK); dovi@berkeley.edu (DP)

†Einstein Fellow.

SNeII-P show some diversity, but overall they tend to be a fairly homogeneous group. However, there are intriguing relatives to this class of objects, most notably SN 1987A, which exploded in the nearby Large Magellanic Cloud and whose progenitor was a compact blue supergiant (BSG) star (Arnett et al. 1989, and references therein). Given the unique contribution of SN 1987A to the study of core-collapse SNe, particular attention to objects that resemble it is warranted. The most similar published SN was SN 1998A (Pastorello et al. 2005). While its light-curve shape was nearly identical to that of SN 1987A, SN 1998A was more luminous and bluer, and its spectra showed higher expansion velocities. Pastorello et al. (2005) attribute these differences to a higher energy BSG explosion.

SN 2000cb was discovered on 2000 April 27.4 (UT dates are used throughout this paper) by Papenkova & Li (2000) in the spiral galaxy IC 1158 at $\alpha = 16^{\text{h}}01^{\text{m}}32^{\text{s}}.15$, $\delta = +1^{\circ}42'23''.0$ (J2000) as part of the Lick Observatory Supernova Search (Li et al. 2000; Filippenko et al. 2001). An unfiltered image obtained with the Katzman Automatic Imaging Telescope (KAIT) on April 24.4 shows the supernova, while there is no detection to a limiting magnitude of 18.7 in an image taken on April 9.5. The SN is at a projected distance of about 4.3 kpc from the host-galaxy centre, off one of the outer arms of this spiral galaxy. Jha et al. (2000) classified it spectroscopically as a SNIa on April 28.4 and noted high expansion velocities measured from the hydrogen absorption features (up to $18\,300\text{ km s}^{-1}$ for $\text{H}\alpha$), as confirmed on April 29.3 by Aldering & Conley (2000). Some peculiarities of SN 2000cb and its similarity to SN 1987A were noted by Hamuy (2001) and as a result, both SNe were excluded from a sample of SNeII-P used as standardizable candles (Hamuy 2004).

In this paper, we present photometric and spectroscopic data on SN 2000cb, all taken within 160 d after the explosion. We investigate its properties and compare it to the prototypical SNeII SN 1999em, SN 1987A and SN 1998A. As we show, while SN 2000cb shares some characteristics with SN 1987A and SN 1998A, it does not appear to closely match any of our comparison objects, further expanding the range of known possible outcomes of massive stellar death. Various arguments, in addition to our model fit to the bolometric evolution, point to a BSG progenitor that produced a strong explosion with a small envelope and a significant amount of ^{56}Ni .

2 DATA ACQUISITION AND REDUCTION

Photometric observations are given in Table 1. Data were obtained using the KAIT, the robotic 0.76-m reflecting telescope at the Lick Observatory, between 2000 April 29.41 and September 15.14, both in *BVI* filters and as unfiltered observations that were made as part of the routine SN search. Following automatic on-site flat-fielding and bias subtraction, images were reduced using the KAIT pipeline (see Ganeshalingam et al. 2010). Galaxy subtraction was performed using template images from several months after the SN had faded beyond detection and the DAOPHOT package in IRAF¹ was used to perform point spread function fitting photometry on the SN as well as on comparison stars in the field. Fig. 1 shows the SN, its host and the comparison stars whose magnitudes are listed in Table 2. All magnitudes are on the Vega-based system.

¹ The IRAF (Image Reduction and Analysis Facility): distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under cooperative agreement with the National Science Foundation.

Table 1. Unfiltered and *BVI* photometry of SN 2000cb.

JD	Phase ^a	Clear (mag)	B (mag)	V (mag)	I (mag)
245 1658.96	3	18.72 (06)	–	–	–
245 1661.97	6	17.74 (06)	–	–	–
245 1663.92	8	17.83 (06)	18.78 (06)	18.04 (03)	17.66 (06)
245 1665.96	10	17.42 (06)	–	–	–
245 1667.93	12	–	18.78 (07)	17.80 (04)	17.30 (06)
245 1670.96	15	17.06 (06)	–	–	–
245 1691.93	36	16.33 (06)	–	–	–
245 1692.88	37	–	17.73 (03)	16.62 (02)	15.87 (02)
245 1693.93	38	16.27 (06)	–	–	–
245 1696.87	41	16.11 (06)	17.83 (05)	16.61 (02)	15.80 (03)
245 1699.88	44	16.11 (06)	–	–	–
245 1700.85	45	–	17.87 (02)	16.56 (02)	15.75 (02)
245 1702.87	47	16.24 (06)	–	–	–
245 1706.85	51	16.14 (06)	17.98 (05)	16.59 (02)	15.67 (01)
245 1713.82	58	–	18.05 (04)	16.61 (02)	15.63 (02)
245 1716.85	61	16.04 (06)	–	–	–
245 1717.84	62	–	18.08 (04)	16.56 (02)	15.59 (02)
245 1719.81	64	16.19 (06)	–	–	–
245 1721.82	66	–	18.14 (04)	16.56 (01)	15.57 (02)
245 1722.84	67	16.18 (06)	–	–	–
245 1726.79	71	16.11 (06)	–	–	–
245 1728.77	73	–	18.19 (05)	16.57 (02)	15.58 (01)
245 1731.78	76	16.10 (06)	–	–	–
245 1735.72	80	–	18.20 (04)	16.63 (02)	15.61 (02)
245 1740.75	85	16.20 (06)	–	–	–
245 1742.73	87	–	18.38 (06)	16.66 (02)	15.66 (02)
245 1743.78	88	16.23 (13)	–	–	–
245 1747.72	92	16.21 (06)	–	–	–
245 1749.69	94	–	18.55 (06)	16.85 (02)	15.74 (02)
245 1752.72	97	16.34 (06)	–	–	–
245 1756.69	101	16.61 (06)	18.78 (09)	17.18 (02)	15.93 (02)
245 1760.68	105	16.90 (06)	–	–	–
245 1770.69	115	16.98 (06)	–	–	–
245 1774.66	119	16.89 (07)	–	–	–
245 1781.66	126	–	19.37 (16)	17.90 (05)	16.52 (03)
245 1788.66	133	–	19.37 (18)	18.14 (19)	–
245 1795.64	140	–	19.32 (20)	18.20 (10)	16.68 (03)
245 1802.64	147	–	–	18.21 (15)	16.80 (07)

Note. The values in parentheses are 1σ measurement uncertainties in hundredths of mag.

^aRelative to JD = 245 1656.

We correct our data for Galactic extinction of $E(B - V)_{\text{Gal}} = 0.114$ mag from Schlegel, Finkbeiner & Davis (1998), using the extinction law of Cardelli, Clayton & Mathis (1989), with a slope of $R_V = 3.1$. The unfiltered photometry of SNe with the KAIT is usually dominated by *R*-band light (e.g. Li et al. 2003). This appears to be a good approximation for SN 2000cb; the unfiltered data lie between the *V* and *I* light curves throughout most of the lifetime of the SN. We therefore correct the unfiltered data for Galactic extinction as though they were acquired with an *R*-band filter. The distance to SN 2000cb is 30 ± 7 Mpc (Springob et al. 2009) and its measured redshift is $z = 0.0064$. We find the explosion date to be 2000 April 21.5 (JD 245 1656) \pm 4.1 d by extrapolating the unfiltered light curve using a cubic fit to the first five points.

Our light curve of SN 1999em was first published by Leonard et al. (2002). We adopt a total extinction value $E(B - V)_{\text{total}} = 0.1$ mag from spectral fitting (Baron et al. 2000) and a Cepheid distance of 11.7 ± 1.0 Mpc (Leonard et al. 2003), with a redshift of $z = 0.0024$. We assume the explosion date of 1999 October 23.5 found by Jones et al. (2009) using the EPM.




Figure 1. Colour composite of g -, r - and i -band SDSS images of SN 2000cb overlain on a KAIT V -band image. The SDSS observation serendipitously caught the SN on 2000 May 4, shortly after the discovery. The SN is located at a projected distance of 4.3 kpc from the nucleus of its host galaxy, IC 1158.

The light curve of SN 1987A (Hamuy et al. 1988) is corrected for extinction using $E(B - V)_{\text{Gal}} = 0.075$ mag (Schlegel et al. 1998) and $E(B - V)_{\text{host}} = 0.1$ mag (Woosley et al. 1987). We adopt a distance of 50 ± 5.2 kpc to the Large Magellanic Cloud (Storm et al. 2004) and an explosion date of 1987 February 23.316 (Arnett et al. 1989).

Two early-time spectra of SN 2000cb, around days 7 and 9 after the explosion, were obtained using the FAST spectrograph (Fabricant et al. 1998) on the 1.5-m Tillinghast telescope at the F. L. Whipple Observatory (slit width 3 arcsec, resolution ~ 7 Å). Beginning about 40 d after the explosion, seven additional spectra were taken with the Kast spectrograph on the Shane 3-m reflecting telescope at the Lick Observatory (Miller & Stone 1993). The first

five were obtained using only the red side of the dual-arm Kast spectrograph and the last two were taken with both sides. With a slit width of 2 arcsec, the typical resolution (full width at half-maximum intensity) was ~ 6 Å on the blue side ($\lambda \lesssim 5500$ Å) and ~ 8 Å on the red side. The slit was in all cases aligned close to the parallactic angle (Filippenko 1982) in order to minimize the effects of atmospheric dispersion. Table 3 provides a log of the spectroscopic observations.

All spectra were reduced using techniques detailed by Foley et al. (2003) and Matheson et al. (2008), among others. IRAF CCD processing was performed and the data were sky-subtracted and extracted optimally (Horne 1986). The observations were wavelength-calibrated by fitting low-order polynomials to arc-lamp spectra. Night-sky lines were cross-correlated with a template sky spectrum and small wavelength shifts were applied when appropriate. Using the spectra of standard stars, the data were then flux-calibrated and corrected for telluric absorption lines (Wade & Horne 1988; Bessell 1999; Matheson et al. 2000). Information for the Kast data comes from our SN data base (Silverman et al., in preparation).

3 PHOTOMETRY

In the top panel of Fig. 2, we compare the light curves of SN 2000cb and SN 1999em. SN 2000cb is fainter by 2–3 mag in each band at early times and then ascends during the first 30 d into the plateau. During the photospheric phase, SN 2000cb follows this prototypical SNIIP reasonably well until day 90, when it begins to drop off the plateau. This happens about 20 d earlier than for SN 1999em, but the decline is not as rapid. As a result, SN 2000cb is overluminous by up to ~ 1 mag, depending on the band, from day 110 until the latest observation at day 147. Our photometric data are listed in Table 1.

The assumption that SNIIP have essentially the same colour evolution during the photospheric phase is one method often used to determine reddening (e.g. Leonard et al. 2002). Such a fit requires a value of $E(B - V)_{\text{host}} = 0.2$ mag with $R_V = 2.6$ to be applied to SN 2000cb, as shown in the bottom panel of Fig. 2. This allows us to match the light curves and colour curves of the two objects reasonably well during the plateau phase, roughly between days 30

Table 2. Comparison stars for the field of SN 2000cb.

Star	α_{J2000}	δ_{J2000}	B (mag)	V (mag)	I (mag)	N_{calib}^a
1	16:01:24.72	+1:45:12.95	18.454 (005)	16.948 (007)	15.039 (004)	3 ^b
2	16:01:37.57	+1:45:02.35	16.034 (010)	15.355 (011)	14.562 (011)	4
3	16:01:29.31	+1:44:54.15	17.965 (009)	17.269 (014)	16.468 (013)	4 ^b
4	16:01:37.68	+1:44:01.86	16.058 (011)	15.245 (010)	14.370 (005)	4
5	16:01:42.60	+1:43:54.82	15.331 (012)	14.637 (009)	13.873 (011)	4
6	16:01:37.97	+1:43:15.13	16.444 (015)	15.609 (013)	14.721 (004)	4
7	16:01:40.84	+1:43:09.37	16.944 (009)	16.305 (008)	15.547 (006)	4
8	16:01:41.42	+1:42:49.59	17.800 (012)	17.156 (003)	16.482 (007)	4 ^b
9	16:01:25.05	+1:42:36.87	16.987 (013)	16.394 (012)	15.638 (004)	3 ^c
10	16:01:26.98	+1:42:14.91	17.544 (014)	16.849 (008)	16.039 (011)	4
11	16:01:25.51	+1:42:29.21	18.994 (006)	17.610 (015)	15.882 (012)	3 ^b
12	16:01:35.25	+1:41:02.21	16.905 (009)	16.229 (007)	15.420 (003)	4
13	16:01:22.75	+1:40:58.56	18.695 (030)	17.391 (000)	15.837 (001)	2 ^b
14	16:01:34.33	+1:39:49.91	18.143 (016)	17.348 (017)	16.321 (013)	2
15	16:01:34.61	+1:45:25.55	15.842 (012)	14.952 (003)	14.006 (012)	4

Note. 1σ uncertainties are in units of 10^{-3} mag.

^aNumber of calibration epochs.

^bOne calibration fewer in the V band.

^cOne calibration fewer in the I band.

Table 3. Journal of spectroscopic observations of SN 2000cb.

UT date	Phase (d)	Range (Å)	Exposure (s)	Airmass ^a	Observer(s) ^b	Instrument
2000 April 28.37	6.9	3720–7540	2 × 1200	1.2	PB	FAST
2000 April 30.38	8.9	3720–7540	1200	1.2	ZB	FAST
2000 May 31.372	39.9	4250–6950	900	1.3	AF, AC	Kast
2000 June 27.265	66.8	4240–6950	1200	1.3	AF, AC	Kast
2000 July 6.377	75.9	4250–6960	1200	2.5	AF, RC, MM	Kast
2000 July 28.308	97.8	4250–6940	900	2.2	AF, RC	Kast
2000 August 26.198	126.7	4240–6950	1500	1.8	DL, WL	Kast
2000 September 6.216	137.7	3300–7760	1800	2.8	AF, WL, RC	Kast
2000 September 26.136	157.6	3300–7800	1800	2.2	AF, WL, RC, MP	Kast

Note. Spectra were obtained using the Kast spectrograph on the 3-m Shane telescope at the Lick Observatory and the FAST spectrograph on the 1.5-m Tillinghast telescope at the Whipple Observatory.

^aAirmass corresponds to the middle of the exposure.

^bAC = Alison Coil, AF = Alex Filippenko, DL = Douglas Leonard, MM = Maryam Modjaz, MP = Mark Phillips, PB = Perry Berlind, RC = Ryan Chornock, WL = Weidong Li, and ZB = Zoltan Balog.




Figure 2. Top panel: light curves of SN 2000cb and SN 1999em in the B , V and I bands, assuming no host-galaxy extinction for SN 2000cb. Bottom panel: the same light curves, but with SN 2000cb matched to SN 1999em using $E(B - V)_{\text{host}} = 0.2$ mag and $R_V = 2.6$. While assuming that significant dust extinction can cause the light curves to match for a fraction of the SN lifetimes, it is inconsistent with our spectroscopic limits and the location of the SN within its host galaxy. Note that there is a systematic uncertainty in the scaling, mostly due to the distance to SN 2000cb, that is not known to better than ~ 0.5 mag.

and 90 after the explosion. However, such a significant extinction value is inconsistent with the absence of detectable narrow Na I D absorption (see Section 4) and it seems unlikely considering the position of the SN near the outskirts of its host galaxy. We therefore assume throughout the rest of this paper that SN 2000cb suffered no host-galaxy extinction, though the presence of some extinction would not appreciably alter any of our conclusions.

In Fig. 3, we compare the photometric evolution of SN 2000cb and SN 1987A. While the SNe show similar qualitative trends such as a slow rise time, a plateau-like phase and a shallow drop-off (as also noted by Hamuy 2001), they clearly behave differently.




Figure 3. Similar to Fig. 2, but comparing the light curves to those of SN 1987A. While there is some qualitative resemblance in the photometric evolution, particularly at later times, the SNe clearly develop on different time-scales. The uncertainty for SN 1987A is dominated by 0.1 mag associated with the distance-modulus correction.

Fig. 4 shows $B - V$ and $V - I$ colour curves of SN 2000cb compared with those of SN 1999em, SN 1987A and SN 1987A. At early times, SN 2000cb appears to follow SN 1987A and both are redder in $B - V$ than SN 1999em beyond anything that could be ascribed to extinction, which indicates a lower effective temperature. In $B - V$, SN 2000cb joins the bluer objects from about day 30 until day 60, when the colours of all four are comparable. Until day 100 or so, SN 2000cb is slightly redder than SN 1999em and SN 1987A, but perhaps not significantly. As it falls off the plateau, SN 2000cb begins to evolve quickly towards the blue. In $V - I$, SN 2000cb again tends to be redder than the three comparison SNe between days 50 and 100, but overall these objects all show broadly similar colour evolution.

4 SPECTROSCOPY

Spectra obtained between days 7 and 157 after the explosion are shown in Fig. 5. All exhibit features characteristic of SNeII during the photospheric phase, including strong $\text{H}\alpha$ P Cygni profiles as well as $\text{H}\beta$ absorption, Na I D $\lambda\lambda 5890, 5896$, Fe II $\lambda\lambda 4924, 5018, 5169$ (hereinafter, Fe triplet), and perhaps Ba and Sc lines. A number




Figure 4. Colour curves of SN 2000cb compared with those of SN 1999em, SN 1998A and SN 1987A. The three objects exhibit not only similar basic trends, but also some significant differences.

of these absorption features are labelled in the figure based on line identifications for SN 1999em (Leonard et al. 2002) and SN 1998A (Pastorello et al. 2005).

There are indications that the equivalent width of narrow Na I D $\lambda\lambda 5890, 5896$ absorption correlates with the amount of dust extinction suffered by a SN, despite significant scatter (e.g. Barbon et al. 1990; Richmond et al. 1994; Turatto, Benetti & Cappellaro 2003; see also Poznanski et al., in preparation). Regardless of the scatter, there seems to be a correlation between high extinction and Na I D absorption. We detect no absorption in any of the individual spectra of SN 2000cb nor in a stack of all nine spectra.

For the stack, we normalize the spectra, crop a region of 60 Å around 5893 Å, subtract a fitted line to remove any continuum con-

tribution, and combine the spectra. The resulting average spectrum has a standard deviation of 5 per cent in flux and yet we do not detect any narrow Na I D absorption. For a typical resolution element of about 5 Å, we thus get a 3σ limit of 0.47 Å on the equivalent width, following equation (4) of Leonard et al. (2001); this corresponds to $E(B - V) < 0.12$ mag (using the relation of Barbon et al. 1990) or $A_V < 0.31$ mag for the best-fitting value to the photometry of $R_V = 2.6$. Such a low $E(B - V)$ is inconsistent with the value 0.2 derived by matching the photometry to SN 1999em and it indicates that SN 2000cb most likely suffered no significant host-galaxy extinction.

The spectra of SN 2000cb are overall very similar to those of typical SNeII, but direct comparisons to SN 1999em and SN 1987A reveal several significant differences. Fig. 6 shows individual spectra of SN 2000cb plotted against spectra at similar phases for each of these objects.

At early times, SN 2000cb closely resembles SN 1987A: their continua match well and the strengths of the prominent hydrogen features are comparable. Both are redder than SN 1999em, which also has much weaker features. The features of SN 2000cb tend to be significantly blueshifted compared to either object throughout their evolution in our sample. Around day 67, all three objects are very similar spectroscopically, although SN 2000cb has much deeper absorption troughs for H α and even more strikingly for H β , which is barely discernible in the other two objects. At about day 127, SN 2000cb again matches both comparisons well; however, most of the Fe (and other metal) absorption lines are weaker. SN 2000cb still exhibits strong hydrogen compared to SN 1999em, but at this phase SN 1987A shows features of comparable depth. The equivalent widths of the hydrogen absorption lines for SN 2000cb and SN 1987A exceed those of SN 1999em by a factor of 8–10 at early times and at later times measurements of SN 2000cb are about twice as great as those of SN 1987A, while features of SN 1999em are even weaker. We also note that the feature at about 5600 Å in SN 2000cb, identified as Na I D $\lambda\lambda 5890, 5896$, has a profile that differs from that of both SN 1999em and SN 1987A in the day 67 comparison, but matches reasonably well at later times. This feature may be a blend with He I $\lambda 5876$.

SN 1998A, identified by Pastorello et al. (2005) as a higher energy analogue of SN 1987A, is a very close spectroscopic match to




Figure 5. Spectroscopic evolution of SN 2000cb, with prominent features identified.




Figure 6. Optical spectra of SN 2000cb at days 9, 66 and 126 compared with spectra of SN 1999em and SN 1987A at similar phases. Each is shown in the supernova’s rest frame. Note the high expansion velocities of SN 2000cb, the early-time similarity of SN 2000cb to SN 1987A and the prominence of the H α absorption feature in the day 67 spectrum.

SN 2000cb, as shown in Fig. 7; its spectra are almost identical at all other available phases as well. The depths of the hydrogen features are comparable, while SN 1999em and SN 1987A differ greatly. It is interesting to note the spectroscopic similarity between SN 2000cb and SN 1998A in spite of the fact that they are not nearly as close photometrically, the light curve of SN 1998A being nearly identical to that of SN 1987A but brighter by about 0.7 mag.

Fig. 8 illustrates the evolution of the H α profile beginning at \sim 40 d after the explosion. In addition to the higher velocity and velocity dispersion of the absorption minimum, the emission peak of SN 2000cb is blueshifted significantly more than those of the other




Figure 7. Comparison of the SN 1998A spectrum at \sim 62 d after the explosion with that of SN 2000cb at a similar phase.





Figure 8. H α feature in spectra of SN 2000cb, SN 1999em and SN 1987A, showing the velocity evolution and line profiles. All spectra are scaled by their H α emission strength and corrected for redshift.

two SNe. The H α emission profile also exhibits a slight excess on the blue side. This structure in the P Cygni profile is reminiscent of the ‘Bochum event’ observed in SN 1987A (Hauschik & Dachs 1987; Utrobin, Chugai & Andronova 1995). A similar feature, tentatively attributed to the same phenomenon by Leonard et al. (2002), also appears briefly in the profile of SN 1999em, but it is not present over as much time in that object’s evolution (only at \sim 101 d in our spectra).

We have measured the velocities of the most prominent absorption features of all four SNe, as shown in Fig. 9. The Fe triplet velocities (shown in the plot as the average of the velocities measured for Fe II $\lambda\lambda$ 4924, 5018, 5169) of SN 2000cb and SN 1998A are similar and typically exceed those of SN 1999em by \sim 1000–1500 km s $^{-1}$ and those of SN 1987A by \sim 2000–2500 km s $^{-1}$. However, they appear to follow roughly the empirical relation (indicated for each object by the dashed lines in the same figure) found by Nugent et al. (2006) for SNeII-P: $V(50) = V(t)(t/50)^{0.464 \pm 0.017}$, where $V(50)$ is the velocity 50 d after the explosion and $V(t)$ is the velocity at some time t in days. The H α and H β absorption velocities of SN 2000cb are more remarkable. H α absorption velocities exceed 18 000 km s $^{-1}$ within 10 d after the explosion and remain \sim 3000–5000 km s $^{-1}$ greater than for both SN 1999em and SN 1987A. The hydrogen velocities measured for SN 1998A are almost as high, lower than those of SN 2000cb by about 500–1000 km s $^{-1}$. Dessart & Hillier (2008) suggest that in SN 1987A-like objects, H α velocities overestimate those of the photosphere by a




Figure 9. Velocity evolution of SN 2000cb compared to SN 1999em, SN 1998A and SN 1987A as measured from $H\alpha$ (top panel), $H\beta$ (middle panel) and the averaged Fe triplet (bottom panel). The dashed lines follow the relation found by Nugent et al. (2006).

factor of 2 at most phases due to non-local thermodynamic equilibrium effects. SN 1987K (Filippenko 1988) and SN 2003bg (Hamuy et al. 2009) also had very high $H\alpha$ absorption velocities at early times. However, the spectra of these objects soon developed He I lines, leading to the ‘SNIib’ designation. Our spectral sequence in Fig. 5 shows that SN 2000cb is not a member of this class.

Poznanski et al. (2010) present a correlation between $H\beta$ and $Fe\ II$ velocities for H-dominated spectra of SNeII-P: $V_{Fe\ II} = (0.84 \pm 0.05)V_{H\beta}$. Fig. 10 shows this relation with our velocity measurements; it was derived for spectra taken between 5 to 40 d after the explosion because of a tendency for $H\beta$ absorption to become weak and blend with metal lines at later times. Our sample of SN 2000cb contains only one usable spectrum within this time frame (from day 40), but it is informative to examine the ratio of Fe and H velocities at a wider range of epochs in the context of this correlation. SN 1999em unsurprisingly follows this relation, as it was one of the SNe used to constrain it, especially at earlier times (and therefore at higher velocities). SN 2000cb and SN 1987A are clear outliers, while SN 1998A deviates in a similar way but to a lesser extent.

5 BOLOMETRIC PROPERTIES AND MODEL FIT

The first question posed by SN 2000cb is whether its light curve was powered primarily by the energy deposited in the initial explosion (as in typical SNeII-P) or by the subsequent radioactive decay of ^{56}Ni (as in SN 1987A and stripped-envelope SNeIb/SNeIc). The




Figure 10. $H\beta$ versus Fe velocities for SN 2000cb, SN 1998A, SN 1999em and SN 1987A. The shaded region marks the 1σ relation found by Poznanski et al. (2010) for early-time spectra. Velocities measured earlier than day 40 (where the relation was defined) are marked with the open symbols. Both SN 1987A and SN 2000cb consistently lie well outside this region, while SN 1999em remains near it even at later times. SN 1998A is consistent with the relation at early times but subsequently diverges from it.

slow rise time strongly disfavours the former possibility. The explosion shock wave deposits thermal energy throughout the stellar envelope, including the outermost layers, and thus radiation begins escaping from the ejecta immediately after shock breakout. The peak in the B -band light curve occurs not long thereafter, when the outer layers have cooled to temperatures $\lesssim 10\,000$ K. Model light-curve calculations of SNeII-P find rise times of ~ 10 d in the B band (Eastman et al. 1994; Baklanov, Blinnikov & Pavlyuk 2005; Kasen & Woosley 2009; Dessart & Hillier 2010), significantly shorter than the ~ 35 d B -band rise time of SN 2000cb. It is possible that varying the structure of the hydrogen envelope could lengthen the rise time, but detailed modelling would be needed to explore such effects. It is far more likely that the explosion energy was mostly spent on the expansion from a small progenitor radius, analogously to SN 1987A.

In order to derive bolometric properties, we fit a blackbody spectrum to the optical photometry of SN 2000cb, SN 1987A and SN 1999em, with the luminosity, radius and temperature being the free parameters. The resulting fits are shown in Fig. 11. We assume that the three objects differ from perfect blackbodies to a similar extent. This may introduce a systematic uncertainty in our absolute values, but it allows for comparisons to be made between the SNe. We find for SN 2000cb a bolometric rise time of about 70 d and peak luminosity of 10^{42} erg s^{-1} . Our derived values for SN 1987A and SN 1999em are generally consistent with previous estimates. While all three SNe have similar radii at most times, SN 1999em is clearly hotter and brighter at early times (as expected from its colours), similar to other well-studied SNeII-P (e.g. Dessart et al. 2008). SN 2000cb and SN 1987A reach a similar peak luminosity, but SN 2000cb reaches it about 20 d earlier.

These long rise times are readily explained by a radioactive energy source, as ^{56}Ni is typically concentrated towards the centre of the ejecta and radiation needs time to diffuse to the surface. Using the scaling relation for the effective diffusion time $t \propto (\kappa M_{\text{env}}/v)^{1/2}$ (Arnett 1979), we can estimate the ejected mass, M_{env} , and the




Figure 11. Bolometric properties for SN 1987A, SN 1999em and SN 2000cb, based on blackbody fits to the optical photometry. The thickness of the stripes marks 1σ uncertainties. Lines are (approximate) model fits for SN 2000cb and SN 1987A generated using the code presented by Young (2004) using $E = 1\text{ B}$, $R = 3 \times 10^{12}\text{ cm}$ and $M_{56\text{Ni}} = 0.07 M_{\odot}$ for SN 1987A, and $E = 2.0\text{ B}$, $R = 3 \times 10^{12}\text{ cm}$ and $M_{56\text{Ni}} = 0.1 M_{\odot}$ for SN 2000cb.

explosion energy of SN 2000cb from those inferred for SN 1987A: $M_{87A} \approx 14 M_{\odot}$, $E_{87A} \approx 1.1\text{ B}$ (Blinnikov et al. 2000; see also Woosley 1988; Shigeyama & Nomoto 1990; Utrobin et al. 1995), where $1\text{ B} = 10^{51}\text{ erg}$. The rise time of SN 2000cb is only 20 per cent faster than that of SN 1987A and the expansion velocities are a factor of ~ 1.7 higher. Adopting a similar opacity κ for both SNe would give an ejected mass of $M_{\text{env}} \approx 16.5 M_{\odot}$ and an explosion energy $E \approx M_{\text{env}} v^2 \approx 4\text{ B}$. Preliminary results from PHOENIX modelling indicate that the spectra of SN 2000cb are compatible with an ejecta mass of $M_{\text{env}} \approx 10 M_{\odot}$ and the same explosion energy of 4 B (Baron et al., in preparation). These values are significantly higher than the hydrogen envelope masses and energies typically inferred for SNeII-P ($M_{\text{env}} \approx 8 M_{\odot}$, $E \approx 1\text{ B}$), though Utrobin et al. (2007) do find higher envelope masses ($\sim 18 M_{\odot}$) for both SN 1987A and SN 1999em.

Assuming a radioactive energy source, we can also estimate the mass of ^{56}Ni ejected in the explosion using Arnett’s law (Arnett 1979), which states that the luminosity at peak equals the instantaneous rate of radioactive energy deposition. The luminosity we measure implies an ejected ^{56}Ni mass for SN 2000cb of $\sim 0.14 M_{\odot}$; however, because of radiative diffusion this is actually an upper bound, as some of this energy was deposited earlier in the light curve. If we interpret the slope of the light curve at $t > 120\text{ d}$ as being due to ^{56}Co decay, then we find a ^{56}Ni mass of $0.1 \pm 0.02 M_{\odot}$. This matches the highest values measured from the decay rates of SNeII-P at nebular phases (Hamuy 2003), while the inferred value for SN 1987A is $0.07 M_{\odot}$.

The lack of a thermal (i.e. non-radioactive) contribution to the light curve of SN 2000cb – or at the very least its dimness relative to SN 1999em at early times – provides an upper limit on the progenitor star radius. Both analytic and numerical models find that the thermal luminosity of SNeII-P on the plateau scales roughly as $L \propto E^{5/6} M^{-1/2} R^{2/3}$, where R is the initial stellar radius (Popov 1993; Kasen & Woosley 2009). Comparing our above estimates of the properties of SN 2000cb to those inferred for SN 1999em ($E_{99\text{em}} \approx 0.7\text{ B}$, $M_{99\text{em}} \approx 15 M_{\odot}$, $R_{99\text{em}} \approx 1000 R_{\odot}$; Baklanov et al. 2005), we find that the lower bolometric luminosity of SN 2000cb and its significantly higher energy imply an initial stellar radius 10–15 times smaller, $R \approx 5 \times 10^{12}\text{ cm}$, similar to that of SN 1987A, $R_{87A} \approx 3 \times 10^{12}\text{ cm}$ (Woosley 1988). Using the values for SN 1999em as found by Utrobin et al. (2007) gives similar results. These scalings have been determined under the assumption that the photosphere is near the recombination temperature of hydrogen, $T_{\text{rec}} \approx 6000\text{ K}$, which is consistent with our best-fitting blackbody temperature during the plateau.

We use the one-dimensional Lagrangian radiation-hydrodynamics code presented by Young (2004) in order to model the light curves of SN 2000cb and SN 1987A. The pre-SN model for both objects is taken from Woosley & Weaver (1995), with an initial mass of $20 M_{\odot}$. The code assumes spherical symmetry and that the neutron star has a mass set as an inner gravitational boundary. The simulation is initiated by artificially placing thermal and kinetic energy in the inner zones, extracted from the formation of the neutron star. The energy injection produces a density discontinuity that propagates as an outward-moving shock wave that takes several hours to exit the star. The simulation follows the progression of the SN material as it expands and releases the energy deposited by the shock.

Fig. 11 shows our fits to SN 1987A and SN 2000cb. For both SNe, we reproduce the time-scales and approximate maximum in the light curves. Included in the simulation is a calculation of the deposition of the energy from the decay of radioactive ^{56}Ni . One of the model parameters used to fit the light curve is the mixing of the radioactive material within the ejecta. The more extensive the Ni mixing is, the sooner the effects of the heating will be observed. For SN 2000cb, we find it necessary to mix ^{56}Ni uniformly out to $15 M_{\odot}$ of the ejecta. This allows an early heating of the material and subsequent earlier peak in the light curve. In addition, since the code does not take into account non-thermal opacity sources, an opacity floor is used to simulate continued opacity due to non-thermal ionization. The effect on the opacity of the large mass of ^{56}Ni and its substantial mixing is emulated by a higher opacity floor. In our model for SN 1987A, more substantial mixing would help smooth the sharp peak in the luminosity.

Fitting the velocity of the Fe lines in SN 2000cb requires an explosion energy of almost twice that of SN 1987A. As a consequence of the fast expansion, the debris becomes thinner faster, so a larger ejected mass of $17.5 M_{\odot}$ is necessary to harness the radioactive heating. This is 10 per cent higher than our fit to SN 1987A. The radii and temperatures we obtain are qualitatively consistent with the blackbody fits to the photometry except at early times, where the temperature from the model is significantly too high – similar to SN 1999em but unlike what we measure for these very red SNe. The basic input parameters for SN 1987A are $E_{87A} = 1\text{ B}$, $R_{87A} = 3 \times 10^{12}\text{ cm}$ and $M_{56\text{Ni}} = 0.07 M_{\odot}$. For SN 2000cb, we find $E_{00\text{cb}} = 2.0\text{ B}$, $R_{00\text{cb}} = 3 \times 10^{12}\text{ cm}$ and $M_{56\text{Ni}} = 0.1 M_{\odot}$. Here again we find that SN 2000cb had a relatively small initial radius consistent with a BSG, a significant explosion energy (although not as high as derived from the scaling above) and a substantial amount of ^{56}Ni . It

Table 4. Unfiltered photometry of SN 2005ci.

JD	Phase ^a	Clear (mag)
245 3523.86	3	20.16 (15)
245 3531.81	11	18.78 (13)
245 3532.85	12	18.75 (13)
245 3533.83	13	18.76 (13)
245 3534.76	14	18.86 (13)
245 3536.80	16	18.51 (16)
245 3540.82	20	18.31 (13)
245 3545.79	25	18.18 (13)
245 3549.78	29	17.94 (13)
245 3553.79	33	17.75 (13)
245 3557.78	37	17.63 (13)
245 3561.76	41	17.72 (15)
245 3565.75	45	17.53 (13)
245 3567.68	47	17.68 (15)

Note. The values in parentheses are 1σ measurement uncertainties in hundredths of mag.

^aRelative to JD = 245 3521.

seems plausible that the relatively high ^{56}Ni yield of SN 2000cb is related to its higher explosion energy, consistent with the correlation proposed by Nadyozhin (2003).

While there is typically some degeneracy between the energy and envelope mass derived for SNe (as we can often constrain only their ratio), we find for SN 2000cb that both the mass and the energy were substantial. The massive ejecta are required in order to trap the gamma rays emitted by the decaying ^{56}Ni . A smaller envelope would entail more escape, pushing our ^{56}Ni mass estimates even higher.

A possibility which we have not taken into consideration is that the asymmetry in the explosion and ejecta may have played a significant role for this object. This would require extensive modelling and is beyond the scope of this paper; moreover, we would also need to explain why most SNeII do not have such bizarre properties or any sign of significant asphericity (e.g. Leonard et al. 2001).

6 SN 2005ci AND THE RATE OF BSG EXPLOSIONS

While a few possible identifications have been made (e.g. Schmitz & Gaskell 1988; Woodings et al. 1998; see a list in Pastorello et al. 2005), the only published secure analogue to SN 1987A is SN 1998A. Here, in addition to SN 2000cb, we report that SN 2005ci (Madison & Li 2005; Modjaz et al. 2005) may be of the same ilk. The peculiarity of this object was already noted by the Caltech Core-Collapse Project (Arcavi et al. 2009). Unfortunately, we have only partial coverage of the unfiltered light curve from the KAIT (see Table 4). Nevertheless, the resemblance to SN 2000cb is striking. While SN 2005ci is fainter by ~ 1.2 mag (to some extent this could be due to extinction, which we cannot constrain), the rise times are nearly identical, as can be seen in the top panel of Fig. 12. The bottom panel shows a spectrum of SN 2005ci at ~ 12 d after the explosion compared to that of SN 2000cb at a similar phase. The two spectra are qualitatively similar, although the features of SN 05ci are much weaker. The velocities of the hydrogen absorption minima are measured to be $\sim 15\,000$ and $\sim 13\,000$ km s $^{-1}$ for H α and H β , respectively. These match well the velocities of SN 2000cb at a comparable phase (Fig. 9) and they are significantly higher than those measured for SN 1999em and SN 1987A.




Figure 12. Comparison between SN 2000cb and SN 2005ci. Unfiltered light curves (top panel) and spectra (bottom panel).

This additional identification raises the number of such SNe discovered by the Lick Observatory SN search to two out of 52 SNeII-P discovered over the period studied by Li et al. (2011), in a volume-limited sample. The corresponding rate is $\sim 4_{-2.6}^{+5.2}$ per cent that of normal SNeII-P, where the dominant error is derived from Poisson statistics. Since SNeII-P are the most abundant SNe in the Universe, this is not an insignificant fraction, of the order of 2 per cent of all core-collapse SNe. The rate we derive is consistent with the one determined by Smartt et al. (2009) – less than ~ 3 per cent – by counting all SNe within about 28 Mpc and assuming the sample is reasonably complete and representative.

It would be of interest to study the metallicities of the locations of these SNe, given the possibility that massive stars are more likely to explode as BSGs than as RSGs at low metallicities (e.g. Langer 1991; Chieffi et al. 2003). The location of SN 2000cb in the outskirts of its host spiral galaxy suggests a relatively low metallicity, in fact. Recently, Anderson et al. (2010) have measured the metallicity of H II regions near the positions of core-collapse SNe, including one in the general vicinity of SN 2000cb, for which they find an oxygen abundance (8.70 ± 0.13 from the O3N2 method) that is close to typical for SNeII and SNeII-P. However, that H II region was more than 3 kpc away from the SN, so it may not give an accurate estimate of the metallicity in the progenitor’s immediate environment.

7 CONCLUSION

We have presented the data and analysis of SN 2000cb, a peculiar SN, with an atypical light curve and extreme photospheric velocities. SN 2000cb strongly resists our attempts to cast it into the

standard SNII-P category while also avoiding identity with SN 1987A. Photometrically, it is intrinsically redder than SN 1999em and fainter by about 0.5 mag on the plateau. It rises to the maximum brightness much more slowly than a normal SNII-P, yet this rise is faster than that of SN 1987A and proceeds with a distinctively different colour and luminosity evolution. The main spectroscopic peculiarity lies in the exceptionally high velocities of its features at all times, most notably for the hydrogen absorption.

Our modelling, as well as extensive comparisons to other SNe, favours a high-energy explosion of a relatively small-radius star, most probably a BSG. We derive a rate for BSG explosions that is of the order of 2 per cent of the rate of all core-collapse SNe.

SNeII-P are the most abundant SNe in the Universe and originate from the explosion of RSGs, the most abundant evolved massive stars. SN 1987A is the keystone of the modern SNII theory, despite being the explosion of a BSG. Linking peculiar SNe, such as SN 1987A and SN 2000cb, to regular SNeII is critical for elucidating the processes that bring a star from formation to obliteration.

ACKNOWLEDGMENTS

We thank Eddie Baron, Luc Dessart, Dan Maoz and the referee (Stephen Smartt) for useful discussions and comments on this manuscript; Mario Hamuy and Andrea Pastorello for sharing their data; and the following for assistance with observations: Zoltan Balog, Perry Berlind, Alison Coil, Douglas Leonard, Maryam Modjaz and Mark Phillips. DP acknowledges support from an Einstein Fellowship and from the US Department of Energy Scientific Discovery through Advanced Computing (SciDAC) programme under contract DE-FG02-06ER06-04. AVF's SN group at UC Berkeley is supported by the NSF grant AST-0908886 and by the TABASGO Foundation. The SN research at the Harvard College Observatory is supported by NSF grant AST-0907903. The construction and ongoing operation of the KAIT were made possible by donations from the Sun Microsystems, Inc., Hewlett-Packard Company, AutoScope Corporation, Lick Observatory, NSF, University of California, Sylvia & Jim Katzman Foundation and TABASGO Foundation. The Kast spectrograph at the Lick Observatory resulted from a generous donation made by Bill and Marina Kast. We are grateful to the dedicated staff at the Lick and F. L. Whipple Observatories.

REFERENCES

Aldering G., Conley A., 2000, *IAU Circ.*, 7410, 3
 Anderson J. P., Covarrubias R. A., James P. A., Hamuy M., Haberman M., 2010, *MNRAS*, 407, 2660
 Arcavi I. et al., 2009, *BAAS*, 214, #604.01
 Arnett W. D., 1979, *ApJ*, 230, L37
 Arnett W. D., Bahcall J. N., Kirshner R. P., Woosley S. E., 1989, *ARA&A*, 27, 629
 Baklanov P. V., Blinnikov S. I., Pavlyuk N. N., 2005, *Astron. Lett.*, 31, 429
 Barbon R., Ciatti F., Rosino L., 1979, *A&A*, 72, 287
 Barbon R., Benetti S., Rosino L., Cappellaro E., Turatto M., 1990, *A&A*, 237, 79
 Baron E. et al., 2000, *ApJ*, 545, 444
 Bessell M. S., 1999, *PASP*, 111, 1426
 Blinnikov S., Lundqvist P., Bartunov O., Nomoto K., Iwamoto K., 2000, *ApJ*, 532, 1132
 Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
 Chieffi A., Domínguez I., Höflich P., Limongi M., Straniero O., 2003, *MNRAS*, 345, 111
 Dessart L., Hillier D. J., 2008, *MNRAS*, 383, 57
 Dessart L., Hillier D. J., 2010, *MNRAS*, 405, 2141
 Dessart L. et al., 2008, *ApJ*, 675, 644

Dessart L., Livne E., Waldman R., 2010, *MNRAS*, 408, 827
 Eastman R. G., Woosley S. E., Weaver T. A., Pinto P. A., 1994, *ApJ*, 430, 300
 Fabricant D., Cheimets P., Caldwell N., Geary J., 1998, *PASP*, 110, 79
 Filippenko A. V., 1982, *PASP*, 94, 715
 Filippenko A. V., 1988, *AJ*, 96, 1941
 Filippenko A. V., 1997, *ARA&A*, 35, 309
 Filippenko A. V., Li W. D., Treffers R. R., Modjaz M., 2001, in Paczyński B., Chen W.-P., Lemme C., eds, *ASP Conf. Ser. Vol. 246, Small-Telescope Astronomy on Global Scales*. Astron. Soc. Pac., San Francisco, p. 121
 Foley R. J. et al., 2003, *PASP*, 115, 1220
 Ganeshalingam M. et al., 2010, *ApJS*, 190, 418
 Hamuy M. A., 2001, PhD thesis, Univ. Arizona
 Hamuy M., 2003, *ApJ*, 582, 905
 Hamuy M., 2004, in Freedman W. L., ed., *Measuring and Modeling the Universe*. Carnegie Observatories, Pasadena
 Hamuy M., Pinto P. A., 2002, *ApJ*, 566, L63
 Hamuy M., Suntzeff N. B., Gonzalez R., Martin G., 1988, *AJ*, 95, 63
 Hamuy M. et al., 2009, *ApJ*, 703, 1612
 Hanuschik R. W., Dachs J., 1987, *A&A*, 182, L29
 Horne K., 1986, *PASP*, 98, 609
 Jha S., Challis P., Kirshner R., Berlind P., 2000, *IAU Circ.*, 7410, 2
 Jones M. I. et al., 2009, *ApJ*, 696, 1176
 Kasen D., Woosley S. E., 2009, *ApJ*, 703, 2205
 Kirshner R. P., Kwan J., 1974, *ApJ*, 193, 27
 Kirshner R. P., Oke J. B., Penston M. V., Searle L., 1973, *ApJ*, 185, 303
 Langer N., 1991, *A&A*, 243, 155
 Leonard D. C., Filippenko A. V., Ardila D. R., Brotherton M. S., 2001, *ApJ*, 553, 861
 Leonard D. C. et al., 2002a, *PASP*, 114, 35 [Erratum: 114, 1291]
 Leonard D. C. et al., 2002b, *AJ*, 124, 2490
 Leonard D. C., Kanbur S. M., Ngeow C. C., Tanvir N. R., 2003, *ApJ*, 594, 247
 Li W. D. et al., 2000, in Holt S. S., Zhang W. W., eds, *Cosmic Explosions*. Am. Inst. Phys., New York, p. 103
 Li W., Filippenko A. V., Chornock R., Jha S., 2003, *ApJ*, 586, L9
 Li W. et al., 2011, *MNRAS*, 412, 1473
 Madison D. R., Li W., 2005, *IAU Circ.*, 8541, 1
 Matheson T., Filippenko A. V., Ho L. C., Barth A. J., Leonard D. C., 2000, *AJ*, 120, 1499
 Matheson T. et al., 2008, *AJ*, 135, 1598
 Miller J. S., Stone R. P. S., 1993, *Lick Obs. Tech. Rep. 66*. Lick Observatories, Santa Cruz
 Modjaz M., Kirshner R., Challis P., Calkins M., 2005, *IAU Circ.*, 8542, 2
 Nadyozhin D. K., 2003, *MNRAS*, 346, 97
 Nugent P. et al., 2006, *ApJ*, 645, 841
 Olivares F. et al., 2010, *ApJ*, 715, 833
 Papenkova M., Li W. D., 2000, *IAU Circ.*, 7410, 1
 Pastorello A. et al., 2005, *MNRAS*, 360, 950
 Popov D. V., 1993, *ApJ*, 414, 712
 Poznanski D. et al., 2009, *ApJ*, 694, 1067
 Poznanski D., Nugent P. E., Filippenko A. V., 2010, *ApJ*, 721, 956
 Richmond M. W., Treffers R. R., Filippenko A. V., Paik Y., Leibundgut B., Schulman E., Cox C. V., 1994, *AJ*, 107, 1022
 Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
 Schmitz M. F., Gaskell C. M., 1988, in Kafatos M., Michalitsianos A. G., eds, *Supernova 1987A in the Large Magellanic Cloud*. Cambridge Univ. Press, Cambridge, p. 112
 Shigeyama T., Nomoto K., 1990, *ApJ*, 360, 242
 Smartt S. J., Eldridge J. J., Crockett R. M., Maund J. R., 2009, *MNRAS*, 395, 1409
 Springob C. M., Masters K. L., Haynes M. P., Giovanelli R., Marinoni C., 2009, *ApJS*, 182, 474
 Storm J., Carney B. W., Gieren W. P., Fouqué P., Latham D. W., Fry A. M., 2004, *A&A*, 415, 531
 Turatto M., Benetti S., Cappellaro E., 2003, in Hillebrandt W., Leibundgut B., eds, *From Twilight to Highlight: The Physics of Supernovae*. Springer-Verlag, Berlin, p. 200

Utrobin V. P., Chugai N. N., 2009, *A&A*, 506, 829
Utrobin V. P., Chugai N. N., Andronova A. A., 1995, *A&A*, 295, 129
Utrobin V. P., Chugai N. N., Pastorello A., 2007, *A&A*, 475, 973
Wade R. A., Horne K., 1988, *ApJ*, 324, 411
Woodings S. J., Williams A. J., Martin R., Burman R. R., Blair D. G., 1998, *MNRAS*, 301, L5

Woosley S. E., 1988, *ApJ*, 330, 218
Woosley S. E., Weaver T. A., 1995, *ApJS*, 101, 181
Woosley S. E., Pinto P. A., Martin P. G., Weaver T. A., 1987, *ApJ*, 318, 664
Young T. R., 2004, *ApJ*, 617, 1233

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.