

Perfect Matching Preservers

Richard A. Brualdi

Department of Mathematics
University of Wisconsin - Madison
Madison, WI 53706, USA
brualdi@math.wisc.edu

Martin Loeb1*

Department of Applied Mathematics
Institute for Theoretical Computer Science
Charles University
Malostranské náměstí 25, 118 00
Prague, Czech Republic
loeb1@kam.mff.cuni.cz

Ondřej Pangrác*

Department of Applied Mathematics
Institute for Theoretical Computer Science
Charles University
Malostranské náměstí 25, 118 00
Prague, Czech Republic
pangrac@kam.mff.cuni.cz

Submitted: Nov 15, 2004; Accepted: Oct 19, 2006; Published: Oct 31, 2006

AMS Subject Classification: 05C20, 05C50, 05C70

Abstract

For two bipartite graphs G and G' , a bijection $\psi : E(G) \rightarrow E(G')$ is called a (perfect) matching preserver provided that M is a perfect matching in G if and only if $\psi(M)$ is a perfect matching in G' . We characterize bipartite graphs G and G' which are related by a matching preserver and the matching preservers between them.

*Supported by Ministry of Education of Czech Republic as project LN00A056.

1 Introduction

A subset $M \subseteq E(G)$ of the edge set $E(G)$ of a graph G is called a *matching* provided that no two edges in M have a vertex in common. A *perfect matching* M is a matching with the property that each vertex of G is incident with an edge in M . For k a positive integer, a graph G is *k-extendable* provided that G has a matching of size k and every matching in G of size at most k can be extended to a perfect matching in G .

In this paper we characterize the bipartite graphs G and G' that are related by a matching preserver and so, with appropriate labeling of edges, have the same perfect matchings. We will achieve this by a full description of matching preservers defined as follows: A bijection $\psi : E(G) \rightarrow E(G')$ is *matching preserving*, or is a *matching preserver*, provided that M is a perfect matching in G if and only if $\psi(M)$ is a perfect matching in G' . Matching preservers for bipartite graphs G were investigated in [2] (see also [1]) in the context of the diagonals of a matrix and the associated diagonal hypergraph. Let A be the bi-adjacency matrix of G . Then A is a $(0, 1)$ -matrix, and the matchings of G are in one-to-one correspondence with the permutation matrices P satisfying $P \leq A$ (entrywise order). The property that G is 1-extendable is equivalent to the property that the bi-adjacency matrix A has *total support*. The property that G is connected and 1-extendable is equivalent to the property that A is *fully indecomposable*. See [3] for a discussion of these matrix properties. The vertices of the hypergraph mentioned above correspond to the edges of G (the positions of the 1's in A) and the hyperedges are the perfect matchings of G (the permutations matrices $P \leq A$, more properly, the set of the n positions of P that are occupied by 1's).

Let G be a connected, 1-extendable, bipartite graph with parts X and Y of size n . The edges of G are pairs xy of vertices with $x \in X$ and $y \in Y$. Let u and v be vertices belonging to different parts of G such that $\{u, v\}$ forms a vertex cut of G . Thus there are bipartite graphs G_1 with parts $X_1 \subseteq X, Y_1 \subseteq Y$, and G_2 with parts $X_2 \subseteq X, Y_2 \subseteq Y$, such that $X_1 \cap X_2 = \{u\}$ and $Y_1 \cap Y_2 = \{v\}$ and each edge of G belongs to either G_1 or G_2 (if uv is an edge of G , then uv is the only common edge of G_1 and G_2). Let G' be the bipartite graph G' obtained from G by replacing each occurrence of u in an edge of G_1 with v and each occurrence of v in an edge of G_1 with u (the neighbors of u and v in G_1 are interchanged). Then G' is a bipartite graph with parts $Y_1 \cup (X_2 \setminus \{u\})$ and $X_1 \cup (Y_2 \setminus \{v\})$. We say that the graph G' is obtained from G by a *bi-twist* with respect to the vertices u and v , and G' is a bi-twist of G (again with respect to vertices u and v). It is easy to verify that *bi-twists preserve both cycles and perfect matchings* [2].

In the language of matrices, a bi-twist is described as follows. Let A be the *labeled bi-adjacency* of order n of G . By this we mean that the 1's of the ordinary bi-adjacency matrix (the 1's correspond to the edges of G) are replaced by distinct elements of some set. Since $\{u, v\}$ is a vertex cut of G , we may choose an ordering for the rows and columns of A , with u corresponding to the first row and v corresponding to the first column, so

that A has the form

$$\left[\begin{array}{c|cc} * & \alpha & \beta \\ \hline \gamma & A_1 & O \\ \hline \delta & O & A_2 \end{array} \right], \quad (1)$$

where

$$(i) \left[\begin{array}{c|c} * & \alpha \\ \hline \gamma & A_1 \end{array} \right] \text{ and } (ii) \left[\begin{array}{c|c} * & \beta \\ \hline \delta & A_2 \end{array} \right] \quad (2)$$

are labeled bi-adjacency matrices of G_1 and G_2 , respectively. Since G is 1-extendable and so has a perfect matching, the matrices A_1 and A_2 are square. A labeled bi-adjacency matrix of G' is the matrix obtained from (1) by replacing (2)(i) with its transpose

$$\left[\begin{array}{c|c} * & \gamma^T \\ \hline \alpha^T & A_1^T \end{array} \right] \quad (3)$$

resulting in the matrix

$$\left[\begin{array}{c|cc} * & \gamma^T & \beta \\ \hline \alpha^T & A_1^T & O \\ \hline \delta & O & A_2 \end{array} \right]. \quad (4)$$

This matrix operation is called *partial transposition* in [1] and [2]. It follows from (1) that in order that a bipartite graph with parts of size n have a bi-twist, its (labeled) adjacency matrix must have a p by q zero submatrix and a complementary q by p zero submatrix for some positive integers p and q with $p + q = n - 1$.

In the language of bipartite graphs, the conjecture in [1] and [2] can be stated as follows.

Conjecture 1.1 *Let G and G' be two 1-extendable, bipartite graphs and let $\psi : E(G) \rightarrow E(G')$ be a matching preserver. Then there is a sequence of bi-twists of G resulting in a graph isomorphic to G' and ψ is induced by this isomorphism.*

As bi-twists do not suffice to describe all matching preservers between bipartite graphs, this conjecture is not true.

Example 1.2 Let G be the bipartite graph with labeled bi-adjacency matrix

$$A = \begin{bmatrix} a & b & c & 0 & 0 & 0 \\ d & e & f & 0 & 0 & 0 \\ 0 & 0 & k & l & m & 0 \\ 0 & 0 & r & s & t & 0 \\ w & 0 & 0 & 0 & u & v \\ z & 0 & 0 & 0 & x & y \end{bmatrix}$$

with parts of size $n = 6$. No bi-twist of G (partial transposition of A) is possible. Yet the bipartite graph G' with labeled bi-adjacency matrix

$$B = \begin{bmatrix} a & b & c & 0 & 0 & 0 \\ d & e & f & 0 & 0 & 0 \\ 0 & 0 & u & v & w & 0 \\ 0 & 0 & x & y & z & 0 \\ m & 0 & 0 & 0 & k & l \\ t & 0 & 0 & 0 & r & s \end{bmatrix}$$

has the same collection of matchings as G . In fact, in both cases, the set of matchings is the union of the sets of matchings corresponding to the two labeled adjacency matrices

$$\begin{bmatrix} a & b & 0 & 0 & 0 & 0 \\ d & e & 0 & 0 & 0 & 0 \\ 0 & 0 & k & l & 0 & 0 \\ 0 & 0 & r & s & 0 & 0 \\ 0 & 0 & 0 & 0 & u & v \\ 0 & 0 & 0 & 0 & x & y \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 & b & c & 0 & 0 & 0 \\ 0 & e & f & 0 & 0 & 0 \\ 0 & 0 & 0 & l & m & 0 \\ 0 & 0 & 0 & s & t & 0 \\ w & 0 & 0 & 0 & 0 & v \\ z & 0 & 0 & 0 & 0 & y \end{bmatrix}$$

□

The operation in Example 1.2 in going from G to G' is an instance of what we call bi-transposition and which we now define. It is the only other operation in addition to bi-twists that is needed in order to describe matching preservers.

Let G_1, G_2, G_3 be bipartite graphs with bipartitions (V_1^i, V_2^i) , and having pairwise disjoint vertex sets. We further assume that $|V_1^i| = |V_2^i| + 1$. Let a_i, b_i be vertices from the part V_1^i of G_i , $i = 1, 2, 3$. Let G be the bipartite graph obtained from G_1, G_2, G_3 by identifying the vertices in each of the three pairs $\{b_1, a_2\}$, $\{b_2, a_3\}$, and $\{b_3, a_1\}$. Let G' be the bipartite graph obtained from G_1, G_2, G_3 by identifying the vertices in each of the three pairs $\{b_1, a_3\}$, $\{b_2, a_1\}$, and $\{b_3, a_2\}$. Then graph G' is said to be obtained from G by a *bi-transposition* of G_1, G_2 and G_3 (see Figure 1).

It is straightforward to verify that the operation of bi-transposition also preserves both cycles and perfect matchings, but cannot be replaced by the bi-twists. The following is the main result of this paper. It is proved in the last section.

Theorem *Let G and G' be two 1-extendable, bipartite graphs and let $\psi : E(G) \rightarrow E(G')$ be a matching preserver. Then there is a sequence of bi-twists and bi-transpositions of G resulting in a graph isomorphic to G' and ψ is induced by this isomorphism.*




Figure 1: Bi-transposition.

2 Preliminaries

In this section we review some facts that will be used in our proof of Theorem 4.1. Let G and G' be 1-extendable bipartite graphs, and suppose that $\psi : E(G) \rightarrow E(G')$ is a matching preserver. By Theorem 2.4 of [2], and it is not difficult to prove, there is a bijection between the components of G and G' such that ψ induces a matching preserver between corresponding components. Hence we may restrict our attention to connected, 1-extendable bipartite graphs—in matrix terms, to fully indecomposable matrices.

The next lemma follows from the inductive structure of a nearly decomposable matrix (see [3]), equivalently from the ear structure of elementary bipartite graphs (see [4]). For convenience, we give a short self-contained proof.

Lemma 2.1 *Let $G = (V, E)$ be a 1-extendable, connected bipartite graph. Then G has a perfect matching M such that for each edge e of M , the vertices of e do not form a cut in G .*

Proof: It suffices to show that if $\{u, v\}$ is a vertex cut such that $e = uv$ is an edge, then the graph $G \setminus e$ obtained from G by deleting edge e is 1-extendable and connected. By recursively deleting such edges we arrive at a 1-extendable, connected bipartite graph G' , where G' has a perfect matching M and for all edges $e' = u'v'$ of G' , in particular for those in M , $\{u', v'\}$ is not a cut of G' and hence not a cut of G .

Since 1-extendable, connected graphs are always 2-connected, it suffices to show that $G \setminus e$ is 1-extendable. There are subgraphs G_1 and G_2 such that $V(G_1) \cup V(G_2) = V(G)$, $V(G_1) \cap V(G_2) = \{u, v\}$, $E(G_1) \cup E(G_2) = E(G)$ and $E(G_1) \cap E(G_2) = \{e\}$. Each perfect matching not containing e has both of the edges incident with $\{u, v\}$ contained in the same G_i .

Let e' be an edge of $G \setminus e$. Let M' be a perfect matching of G containing e' . If $e \notin M'$, then M' is also a perfect matching of G' and contains e' . Assume that $e \in M'$, and that e.g. e' is an edge of G_1 . Let M'' be the restriction of M' to a perfect matching of G_1 . Let $f \neq e$ be an edge of G_2 incident with $\{u, v\}$ and let N be a perfect matching of G containing f . Then N contains a perfect matching N_2 of G_2 . Thus $(M'' \setminus \{e\}) \cup N_2$ is a perfect matching of $G \setminus e$ containing e' , and this completes the proof. □

We now review a classical theorem of Whitney [8]. Let G be a 2-connected graph with vertex cut $\{u, v\}$. There are subgraphs G_1 and G_2 such that $V(G_1) \cup V(G_2) = V(G)$, $V(G_1) \cap V(G_2) = \{u, v\}$, $E(G_1) \cup E(G_2) = E(G)$ and $E(G_1) \cap E(G_2) = \{uv\}$ or \emptyset depending on whether or not uv is an edge of G . Define a graph G' as follows: Let G' be the graph obtained from G by replacing each occurrence of u in an edge of G_1 with v and each occurrence of v in an edge of G_1 with u (the neighbors of u and v in G_1 are interchanged). Then G' is obtained from G by a *twist*, and G' is a twist of G (again with respect to vertices u and v). (If uv is an edge of G , then it is also an edge of G' .) It was proved by Whitney [8] that each graph with the same cycles as the 2-connected graph G —that is, a graph that is 2-*isomorphic* to G —can be obtained from G by a sequence of twists. Truemper [6] simplified the proof and obtained a bound on the number of twists needed.

Theorem 2.2 *Let G be a 2-connected graph with $n \geq 2$ vertices, and let H be a graph 2-isomorphic to G . Then G can be transformed into a graph G^* which is isomorphic to H by a sequence of at most $n - 2$ twists.*

The technique of Truemper uses the concept of generalized cycles. A graph G is a *generalized cycle* with constituents G_1, G_2, \dots, G_k ($k \geq 2$) provided that the following hold:

- (i) each G_i is a connected subgraph of G having nonempty edge set E_i ; additionally, if $k = 2$ then both G_1 and G_2 contain at least three vertices;
- (ii) the edge sets E_i , $1 \leq i \leq k$, partition the edge set $E(G)$, and each G_i has exactly two vertices in common with $\cup_{j \neq i} G_j$ (these vertices are called the *contact vertices* of G_i);
- (iii) replacing each G_i by an edge joining the contact vertices of G_i produces an ordinary cycle.

The generalized cycle G is a connected graph. If $k \geq 3$ and each G_i has only two vertices (since G_i is connected, these two vertices are joined by an edge), then G is an ordinary cycle.

The first assertion in the next lemma is due to Tutte [7]; the second assertion is due to Truemper [6]. In the lemma, a G_i consisting of a single edge is regarded as 2-connected.

Lemma 2.3 *If a graph G is 2-connected but not 3-connected, then there exists a representation of G as a generalized cycle where each constituent is 2-connected.*

Moreover, let G be a 2-connected, generalized cycle as above. If $\psi : E(G) \rightarrow E(H)$ is a 2-isomorphism of G to H , then H is a generalized cycle with constituents H_1, H_2, \dots, H_k , where H_i is the subgraph of H induced by $\psi(E_i)$ for $1 \leq i \leq k$.

3 Directed graphs

There is a well-known correspondence between matchings in a bipartite graph G and circuits in a directed graph (*digraph*) D constructed from G and a specified perfect matching of G . This correspondence can be easily understood by using adjacency matrices. Let $M = \{u_1v_1, u_2v_2, \dots, u_nv_n\}$ be a perfect matching of G and let $A = [a_{ij}]$ be the biadjacency matrix of G where $a_{ij} = 1$ if and only if u_iv_j is an edge of G , $1 \leq i, j \leq n$. Thus A has all 1's on its main diagonal and these 1's correspond to the edges of M . The matrix $A - I_n$ is the adjacency matrix of a digraph $D(G, M)$. The digraph can also be understood as obtained from G by orienting each edge from one part of G to its other part, and then contracting all of the edges of M .

A *circuit* of a digraph is a circular sequence of distinct edges such that the terminal vertex of each edge is the initial vertex of the edge that follows. As such, a circuit may be identified with its collection of edges, since its circular arrangement is unique. Similarly, we may identify a *path* in a digraph with its collection of edges.

Let M' be another perfect matching in G . Then $(M \setminus M') \cup (M' \setminus M)$ is a collection of pairwise vertex disjoint cycles of G of even length whose edges alternate between M and M' . In $D(G, M)$ these cycles correspond to pairwise vertex-disjoint circuits (not necessarily a spanning set since M and M' may have edges in common). Using the matching M , we may reverse this construction to obtain, given a collection of pairwise vertex-disjoint circuits of $D(G, M)$, a perfect matching M' of G . Thus, *there is a one-to-one correspondence between perfect matchings in G and collections of pairwise-vertex disjoint circuits in $D(G, M)$* . This well-known observation allows us to reformulate our problem in terms of digraphs and pairwise vertex-disjoint circuits.

A digraph is *strongly connected* provided that for each ordered pair of vertices u, v , there is a path from u to v . The 1-extendability of the connected bipartite graph is equivalent to the strong connectivity of $D(G, M)$. We formalize this well-known property in the next lemma (see e.g. [3]). (In matrix terms this property is usually stated as: A $(0, 1)$ -matrix A of order n with all 1's on its main diagonal is fully indecomposable if and only if the matrix $A - I_n$ is irreducible.

Lemma 3.1 *Let G be a connected bipartite graph and let M be a perfect matching of G . Then G is 1-extendable if and only if the digraph $D(G, M)$ is strongly connected.*

The analogue of Whitney's theorem for digraphs was proved by Thomassen [5]. First, recall that an *isomorphism*, respectively, an *anti-isomorphism*, of a digraph D onto a digraph D' is a bijection $f : V(D) \rightarrow V(D')$ such that, for all $u, v \in V(D)$, there is an arc in D from vertex u to vertex v if and only if there is an arc in D' from vertex $f(u)$ to vertex $f(v)$, respectively, from $f(v)$ to $f(u)$.

A directed twist of a digraph D is defined in a similar way to a twist in a graph. Let D_1, D_2 be subgraphs of D of order at least 3, such that $V(D_1) \cup V(D_2) = V(D)$, $V(D_1) \cap V(D_2) = \{u, v\}$, $E(D_1) \cup E(D_2) = E(D)$. Let D' be obtained from D by replacing arcs of the form uw, wu, vw , and wv by, respectively, wv, vw, wu and uw for each $w \in V(D_2)$ and then reversing the direction of all the remaining arcs of D_2 . Then

D' is obtained from D by a *directed twist* (or *di-twist*), with respect to the vertices u and v , and D' is a di-twist of D (again with respect to the vertices u and v). Clearly, D and D' have the same circuits and D is strongly connected if and only if D' is.

In the language of matrices, a directed twist is described as follows. Let A be the adjacency matrix of the digraph D where the vertices have been ordered so that u and v come first followed by the remaining vertices of D_1 and then the remaining vertices of D_2 . Thus A has the form

$$\left[\begin{array}{c|c|c|c} a & b & \alpha_1 & \alpha_2 \\ \hline c & d & \beta_1 & \beta_2 \\ \hline \gamma_1 & \delta_1 & A_1 & O \\ \hline \gamma_2 & \delta_2 & O & A_2 \end{array} \right], \quad (5)$$

where

$$(i) \left[\begin{array}{c|c|c} a & b & \alpha_1 \\ \hline c & d & \beta_1 \\ \hline \gamma_1 & \delta_1 & A_1 \end{array} \right] \quad \text{and} \quad (ii) \left[\begin{array}{c|c|c} a & b & \alpha_2 \\ \hline c & d & \beta_2 \\ \hline \gamma_2 & \delta_2 & A_2 \end{array} \right] \quad (6)$$

are the adjacency matrices of D_1 and D_2 , respectively. An adjacency matrix of the digraph D' is obtained from (5) by replacing (6)(ii) in (5) with

$$\left[\begin{array}{c|c|c} a & c & \delta_2^T \\ \hline b & d & \gamma_2^T \\ \hline \beta_2^T & \alpha_2^T & A_2^T \end{array} \right] \quad (7)$$

Thomassen [5] proved an analogue of Whitney's theorem for digraphs, applying Whitney's theorem to the underlying graph. If D is a digraph, then G_D denotes the *underlying graph* of G .

Theorem 3.2 *Let D and D' be two strongly connected digraphs with 2-connected underlying graphs G_D and $G_{D'}$. Let $\varphi : E(D) \rightarrow E(D')$ be a bijection such that φ and φ^{-1} preserve circuits. Then there exist a sequence of di-twists of D resulting in a digraph D^* such that φ is induced by an isomorphism or anti-isomorphism of D^* onto D' .*

Note that the requirement of the 2-connectivity of the underlying graphs is necessary only for G_D . That $G_{D'}$ is 2-connected then follows from Whitney's theorem.

Let D be a digraph. Then D is a *generalized circuit* provided D is strongly connected and the underlying graph G_D is a generalized cycle. Let D be a generalized circuit such that the constituents of its underlying graph G are G_1, G_2, \dots, G_k , and the contact vertices

in G_i are u_i and v_i , $i = 1, 2, \dots, k$. Then the corresponding digraphs D_1, D_2, \dots, D_k are called the *constituents of the generalized circuit* D , and the vertices u_i, v_i in D_i are called its *contact vertices*, $i = 1, 2, \dots, k$. Note that $v_i = u_{i+1}$ where the subscripts are interpreted modulo k . Moreover, in the rest of the paper, we work only with generalized circuits with the underlying graphs of all the constituents 2-connected.

Since D is assumed to be strongly connected, it follows that, for each constituent D_i , either D_i is strongly connected or the digraph obtained from D_i by contracting each strong component to a vertex contains a path with initial vertex corresponding to the strong component containing u_i and final vertex corresponding to the strong component containing v_i , or the other way around. The following lemma is now easily verified.

Lemma 3.3 *If D is a generalized circuit, then D has a circuit C containing all of the contact vertices, and passing through all of its constituents.*

In a generalized circuit D , we always assume that its constituents have been labeled D_1, D_2, \dots, D_k in such a way that the circuit C in Lemma 3.3 comes into D_i at u_i ($=v_{i-1}$) and leaves D_i at v_i ($=u_{i+1}$).

Now we state a directed analogue of the second assertion in Lemma 2.3.

Lemma 3.4 *Let D be a generalized circuit with constituents D_1, D_2, \dots, D_k and let the underlying graphs of all D_i be 2-connected, $i = 1, 2, \dots, k$. Let E_i denote the edge-set of D_i . Let $\varphi : E(D) \rightarrow E(D')$ be a bijection such that φ and φ^{-1} preserve circuits. Then D' is a generalized circuit with constituents D'_1, D'_2, \dots, D'_k (not necessarily ordered in this way), where D'_i is the subgraph of D' induced by $\varphi(E_i)$ for $1 \leq i \leq k$.*

Proof: The corresponding underlying graphs G_D and $G_{D'}$ are 2-isomorphic by Theorem 3.2. The lemma now follows by applying Lemma 2.3. □

The contact vertices of the generalized circuit D are partitioned into three types. If there is a circuit in D_i meeting one of its contact vertices u_i, v_i but not the other, then we say that that contact vertex is *heavy in D_i* . If there is no circuit in D_i containing a particular contact vertex, then we call that contact vertex *light in D_i* ; in this case, the strong component of D_i containing the contact vertex contains no other vertex. If a contact vertex in D_i is neither light nor heavy, then it is called *cyclic in D_i* ; if e.g. u_i is cyclic in D_i , then there is a circuit containing u_i , and each circuit in D_i containing u_i also contains v_i . Note that *if one of the contact vertices is cyclic in D_i , then the second one clearly cannot be light*.

Let D be a generalized circuit with constituents D_i (each of them has 2-connected underlying graph) and contact vertices u_i, v_i , $i = 1, 2, \dots, k$. Let σ be a permutation of $\{1, 2, \dots, k\}$ and let $\varepsilon \in \{-1, +1\}^k$. Let the digraph $D^{\sigma, \varepsilon}$ be obtained from D by rearranging the constituents in the following way: First, assume a directed graph consisting of disjoint components D_1, D_2, \dots, D_k . Then, for $i = 1, 2, \dots, k$, if $\varepsilon_i = -1$ reverse the orientation of the edges of D_i and set $x_i = v_i$, $y_i = u_i$; if $\varepsilon_i = +1$ set $x_i = u_i$ and $y_i = v_i$.

Finally, we identify vertices $y_{\sigma(i)}$ and $x_{\sigma(i+1)}$ (modulo k), for $i = 1, 2, \dots, k$. Moreover, we call the rearrangement $D^{\sigma, \varepsilon}$ an *admissible rearrangement* of D if the following property holds:

(*) If v_i is heavy or cyclic in D_i and u_{i+1} is heavy or cyclic in D_{i+1} , then D_{i+1} follows D_i in $D^{\sigma, \varepsilon}$ as it does in D or D_i follows D_{i+1} , $i = 1, 2, \dots, k$. If, in addition, v_i (u_{i+1} , respectively) is actually heavy in D_i (D_{i+1} , respectively) then v_i (u_{i+1} , respectively) is one of the contact vertices of D_i and D_{i+1} in $D^{\sigma, \varepsilon}$.

It follows, in particular, that an admissible rearrangement of a circuit produces either the circuit itself or its reversal (as, by our convention, a set of edges).

The definition of an admissible rearrangement defines implicitly a partition of the constituents of D into *superconstituents*, where each superconstituent is a maximal sequence of consecutive constituents of D with the property that no inner contact vertex is light. It follows that a common contact vertex of two different superconstituents is light in at least one of them.

It is straightforward to verify the following lemma.

Lemma 3.5 *Let D be a generalized circuit. A rearrangement of the constituents of D does not change the set of circuits incident to any vertex of D if and only if it is an admissible rearrangement.*

The following theorem is a first step towards characterizing bijections between the edges of two digraphs that preserve the union of vertex-disjoint circuits.

Theorem 3.6 *Let D and D' be two strongly connected digraphs and let $\varphi : E(D) \rightarrow E(D')$ be a bijection. Then φ induces a bijection between unions of vertex-disjoint circuits of D and unions of vertex-disjoint circuits of D' (that is, φ and φ^{-1} preserve unions of vertex-disjoint circuits) if and only if, starting with D , there is a sequence of admissible rearrangements of generalized circuits that results in a digraph D^\dagger such that φ is induced by an isomorphism or anti-isomorphism of D^\dagger onto D' .*

Proof: It follows from Lemma 3.5 that admissible rearrangements do not change the set of vertex-disjoint circuits. We now consider the converse. First we note that the converse holds when D (and hence D') is a circuit. Let D be a minimal counterexample, and let Γ be a generalized circuit of D with the underlying graphs of all its constituents 2-connected; Γ exists by Lemma 2.3. Lemma 3.4 implies that the image Γ' of Γ under φ is a generalized circuit of D' whose constituents Γ'_i are images under φ of the constituents Γ_i of Γ . Let Γ_i^* be the strongly connected digraph obtained from Γ_i by adding a new edge from its contact vertex v_i to its contact vertex u_i , and let $\Gamma_i'^*$ be defined in a similar way using contact vertices u'_i and v'_i . Let φ_i^* be the extension of $\varphi|_{\Gamma_i}$ to Γ_i^* which maps the edge (v_i, u_i) to the edge (v'_i, u'_i) . In particular, φ_i^* is a bijection from the edges of Γ_i^* to those of $\Gamma_i'^*$ preserving the set of unions of vertex-disjoint circuits of Γ_i^* . Hence, by minimality of D , each φ_i^* is realized by admissible rearrangements of constituents of generalized circuits of Γ_i^* . Each admissible rearrangement of Γ_i^* corresponds to an admissible rearrangement of Γ_i . It follows that the generalized circuit Γ' is obtained by a rearrangement of the

constituents of the generalized circuit whose constituents are admissible rearrangements of the constituents of Γ . By Lemma 3.5, this rearrangement must be admissible, since a non-admissible rearrangement would change the set of unions of vertex-disjoint circuits. \square

By Theorem 3.6, to fully characterize mappings that preserve unions of vertex-disjoint circuits, it suffices to describe how admissible rearrangements can be carried out. To do this, we introduce two operations, that of a cyclic di-twist and of a supertransposition.

A di-twist of a strongly connected digraph D with parts D_1 and D_2 is *cyclic* if both twist vertices are light in one of the parts, say in D_1 (here we regard D as a generalized circuit with two constituents each with at least three vertices). What this means is the following: Let $V(D_1) \cap V(D_2) = \{u, v\}$ where u and v are light in D_1 . Then $\{u\}$ and $\{v\}$ are strong components of D_1 with, say, each edge of D_1 incident with u directed from u and each edge of D_1 incident with v directed to v . Then the di-twist with respect to the parts D_1 and D_2 is a *cyclic di-twist*. It follows directly that a *cyclic di-twist of a digraph preserves unions of vertex-disjoint circuits*.

Now let D be a generalized circuit. Let S and S' be consecutive superconstituents and let u and v (respectively, u' and v') be the contact vertices of S (respectively, of S'), where $u' = v$. Further, let T be the union of all remaining superconstituents with contact vertices u_T and v_T where $u_T = v'$ and $v_T = u$. Assume that u , u' , and u_T are light in, respectively, S , S' , and T . Then the operation of *supertransposition* of superconstituents S and S' results in the digraph D' constructed by transposing S and S' in the following way: D' is obtained from the disjoint union of S , S' , and T by identifying u with v' , u' with v_T and u_T with v . It is straightforward to verify that supertransposition does not change vertex-disjoint unions of circuits. Our goal is now to show that cyclic di-twists and supertransposition can be used to completely describe edge bijections between strongly connected digraphs which preserve vertex-disjoint unions of circuits.

Consider a generalized circuit D with constituents D_1, \dots, D_k and superconstituents S_1, S_2, \dots, S_l . We associate with D an object called an *auxiliary circuit* that captures the relationship between the superconstituents. The auxiliary circuit has length l and an edge e_i , joining the contact vertices u_i, v_i of S_i , $i = 1, 2, \dots, l$; thus, each superconstituent is contracted to an edge joining its two contact vertices, and these edges are of three types. Edge e_i is a *fat-edge* provided that neither u_i nor v_i is a light vertex of S_i (fat-edges have no direction); e_i is a *thin-edge* and is directed from u_i to v_i provided that v_i is a light vertex of S_i while u_i is not; e_i is a *two-way edge* directed to both u_i and v_i provided both u_i and v_i are light vertices of S_i . Important properties of the auxiliary circuit, following from the definition of superconstituents, are:

- AC1. There is at least one edge directed to each of the vertices of an auxiliary circuit.
Hence:
- AC2. If an auxiliary circuit has at least one fat-edge, then it has a two-way edge.
- AC3. If an auxiliary circuit contains no two-way edges, then it is just an ordinary directed circuit.

We now define admissible rearrangements for auxiliary circuits. A *two-way path* is a path of the auxiliary circuit which starts and ends with a two-way edge. An *admissible rearrangement* of the edges of an auxiliary circuit C is an auxiliary circuit C' obtained by arbitrarily rearranging the fat-edges amongst themselves, arbitrarily rearranging the two-way paths amongst themselves, and removing thin-edges from their places and reinserting them in other places with orientations reversed (with respect to the clockwise orientation). Note that fat-edge rearrangements and two-way path rearrangements applied to an auxiliary circuit always result in an auxiliary circuit; a thin-edge shift requires that the property that at least one edge is directed to each of the vertices be maintained. The next lemma shows that two auxiliary circuits, with the same number of edges of each of the three types and with at least one two-way edge, are admissible rearrangements of one another.

Lemma 3.7 *Let C and C' be auxiliary circuits containing the same number of edges of each type, with at least one two-way edge. For C and C' , let there be assigned labels so that for each of the three types of edges, the edges have the same set of labels in C as in C' . Then C' can be obtained from C by fat-edge rearrangements, two-way path rearrangements, and thin-edge shifts.*

Proof: First we rearrange the labeled fat-edges and rearrange the labeled two-way paths of C so that the fat edges and the two-way edges agree with their cyclic positioning in C' . As remarked above, the result is an auxiliary circuit. Then by shifting the labeled thin-edges of C , we get them to be in the right position in C' : if the orientation of a thin-edge needs to be changed, then one shift is sufficient; otherwise, we shift the thin edge next to a two-way edge, thereby changing its orientation and then shift it to its right place, thereby changing its orientation back to what it was. □

We next consider how a cyclic di-twist of a generalized circuit D relates to the corresponding auxiliary circuit C . Let T_1 and T_2 be a partition of the superconstituents of D into two consecutive parts, where $V(T_1) \cap V(T_2) = \{u, v\}$. The di-twist with respect to T_1 and T_2 is cyclic provided that u and v are light in one of the parts, say T_1 . Denote by P_i the set of edges of the auxiliary circuit corresponding to the superconstituents in T_i , $i = 1, 2$. Then P_1 and P_2 partition the edges of the auxiliary circuit into two paths (consisting of the three types of edges). Since u and v are light in T_1 , the edge of P_1 at u , respectively, v is either a two-way edge or a thin edge directed to u , respectively, to v . Conversely, when we have such a partition of the auxiliary circuit with these properties, the di-twist with respect to corresponding parts of D is cyclic.

Lemma 3.8 *Let C and C' be auxiliary circuits containing the same number of edges of each type, with at least one two-way edge. For C and C' , let there be assigned labels so that for each of the three types of edges, the edges have the same set of labels in C as in C' . Then C' can be obtained from C by a sequence of cyclic di-twists.*

Proof: First consider an exchange of two fat-edges edges e_i and e_j . Let S_i and S_j be corresponding superconstituents. Since no fat-edges are consecutive in C , $C \setminus \{e_i, e_j\}$ consists of two paths P_1 and P_2 . Denote by T_m the union of superconstituents corresponding to edges of P_m , $m = 1, 2$. Then the exchange of e_i and e_j can be constructed by two cyclic di-twists: first twist with parts T_1 and $S_i \cup T_2 \cup S_j$ and then twist with parts T_2 and $S_i \cup T_1 \cup S_j$.

Now consider an exchange of two two-way paths e_i and e_j . Their exchange by cyclic di-twists is similar to that for fat-edges. Let S_i and S_j be the corresponding superconstituents. First suppose that the two-way paths e_i, e_j are consecutive so that $P = C \setminus \{e_i, e_j\}$ is a path. Denote by T the union of the corresponding superconstituents. Then the exchange of e_i and e_j can be carried out by the cyclic di-twist with parts T and $S_i \cup S_j$. Now suppose that e_i, e_j are not consecutive. Let P_1 and P_2 be the two paths of $C \setminus \{e_i, e_j\}$, and let T_1 and T_2 be the corresponding unions of superconstituents. Then the exchange of e_i and e_j can be carried out by two cyclic di-twists: first twist with parts T_1 and $S_i \cup T_2 \cup S_j$ and then twist with parts T_2 and $S_i \cup T_1 \cup S_j$.

Finally, we show how a thin-edge shift can be carried out by cyclic di-twists. Let e be a thin-edge, and let v be the vertex of C (and so a contact vertex of the corresponding generalized circuit) to which e is to be shifted. Let S_e be the superconstituent corresponding to e . Finally, let P_1 and P_2 be the paths from v to the closest vertex of e in both directions of C , and let T_1 and T_2 be the corresponding unions of superconstituents. Then the shift of e can be constructed by two cyclic di-twists: first twist with parts T_1 and $T_2 \cup S_e$ and then twist with parts T_2 and $T_1 \cup S_e$. □

We now apply the previous two lemmas to admissible rearrangements of generalized circuits. An admissible rearrangement of a generalized circuit is an *admissible superconstituent rearrangement* provided that it only rearranges the superconstituents.

Lemma 3.9 *Let D be a generalized circuit. Then each admissible superconstituent rearrangement D' of D can be accomplished by a sequence of cyclic di-twists and supertranspositions.*

Proof: Let C and C' be the auxiliary circuits corresponding to D and D' , respectively. Then the superconstituent rearrangement induces a bijection between the edges of C and those of C' that preserves edge types. If the auxiliary circuit C of D contains at least one two-way edge, then by Lemma 3.7, C' may be obtained from C by a sequence of fat-edge exchanges, two-way edge exchanges, and thin-edge shifts. By Lemma 3.8, this can be accomplished by a sequence of cyclic di-twists. Suppose that C does not contain any two-way edges. Then C is an ordinary circuit, and C' is also an ordinary circuit obtained by a rearrangement of the edges of C . But then C' can be obtained from C by a sequence of transpositions of consecutive edges. Each such transposition of edges can be accomplished by a supertransposition. □

We now characterize bijections between the edges of digraphs that preserve unions of vertex-disjoint circuits.

Theorem 3.10 *Let D and D' be two strongly connected digraphs, and let $\varphi : E(D) \rightarrow E(D')$ be a bijection. Then φ induces a bijection between unions of vertex-disjoint circuits of D and unions of vertex-disjoint circuits of D' if and only if, starting with D , there is a sequence of cyclic di-twists and supertranspositions that results in a digraph D^\dagger such that φ is induced by an isomorphism or anti-isomorphism of D^\dagger onto D' .*

Proof: Each admissible rearrangement consists of an admissible rearrangement of superconstituents and admissible rearrangements within a superconstituent. Moreover, admissible rearrangements within a superconstituent can be accomplished by cyclic di-twists. Hence the theorem follows by Theorem 3.6 and Lemma 3.9. □

4 Perfect Matching Preservers

We can now prove the main result of this paper.

Theorem 4.1 *Let G and G' be two 1-extendable, bipartite graphs and let $\psi : E(G) \rightarrow E(G')$ be a matching preserver. Then there is a sequence of bi-twists and bi-transpositions of G resulting in a graph isomorphic to G' and ψ is induced by this isomorphism.*

Proof: By Lemma 2.1, G has a perfect matching M with the property that the two vertices of each edge do not form a cut of G . A bi-twist of the bipartite graph G corresponds to a cyclic twist of the digraph $D(G, M)$, and it follows from the definition of the supertransposition, in particular from the assumption of the lightness of the connecting vertices, that bi-transposition of G corresponds to a supertransposition of $D(G, M)$. As observed previously, bi-twists and bi-transpositions preserve perfect matchings.

For the converse, assume that $\psi : E(G) \rightarrow E(G')$ is a matching preserver. Let $M' = \psi(M)$. Let $\varphi : E(D(G, M)) \rightarrow E(D(G', M'))$ be the bijection naturally determined by ψ . From the correspondence between matchings in G , respectively, G' , and unions of vertex-disjoint circuits of $D(G, M)$, respectively $D(G', M')$, φ induces a bijection between pairwise vertex-disjoint circuits of $D(G, M)$ and those of $D(G', M')$. Hence by Theorem 3.10, $D(G', M')$ may be obtained from $D(G, M)$ by cyclic twists and supertranspositions, and so G' may be obtained from G by bi-twists and bi-transpositions. □

References

- [1] Richard A. Brualdi, Linear Preservers and Diagonal Hypergraphs. *Linear Alg. Applics.*, 373 (2003), 51–65.

- [2] Richard A. Brualdi and Jeffrey A. Ross, Matrices with isomorphic diagonal hypergraphs. *Discrete Math.*, 33 (1981), 123-138.
- [3] Richard A. Brualdi and Herbert J. Ryser, *Combinatorial Matrix Theory*. Cambridge University Press, 1991.
- [4] L. Lovász and M.D. Plummer, *Matching Theory*. Annals of Discrete Math., 29, North-Holland, 1986.
- [5] Carsten Thomassen, Whitney's 2-Switching Theorem, Cycle Space, and Arc Mappings of Directed Graphs. *J. Comb. Theory Ser. B* 46(3) (1989) 257–291.
- [6] K. Truemper, On Whitney's 2-isomorphism theorem for graphs. *J. Graph Theory* (1980) 43–49.
- [7] W.T. Tutte, *Connectivity in Graphs*. University of Toronto Press, Chapter 11, 1996.
- [8] H. Whitney, 2-isomorphic graphs. *Amer. J. Math.* 55 (1933), 245–254.