



# Performance Index and PSII Connectivity Under Drought and Contrasting Light Regimes in the CAM Orchid *Phalaenopsis*

Nathalie Ceusters<sup>1</sup>, Roland Valcke<sup>2</sup>, Mario Frans<sup>1</sup>, Johan E. Claes<sup>3</sup>, Wim Van den Ende<sup>4</sup> and Johan Ceusters<sup>1,5\*</sup>

<sup>1</sup> Department of Biosystems, Division of Crop Biotechnics, Research Group for Sustainable Crop Production & Protection, KU Leuven, Geel, Belgium, <sup>2</sup> Molecular and Physical Plant Physiology, UHasselt, Diepenbeek, Belgium, <sup>3</sup> Department of Microbial and Molecular Systems, Bioengineering Technology TC, KU Leuven, Geel, Belgium, <sup>4</sup> Laboratory of Molecular Plant Biology, Department of Biology, KU Leuven, Leuven, Belgium, <sup>5</sup> Centre for Environmental Sciences, Environmental Biology, UHasselt, Diepenbeek, Belgium

## OPEN ACCESS

### Edited by:

Marouane Baslam,  
Niigata University, Japan

### Reviewed by:

Jim Leebens-Mack,  
University of Georgia, United States  
Jie He,  
Nanyang Technological University,  
Singapore

### \*Correspondence:

Johan Ceusters  
johan.ceusters@kuleuven.be

### Specialty section:

This article was submitted to  
Plant Abiotic Stress,  
a section of the journal  
Frontiers in Plant Science

**Received:** 20 March 2019

**Accepted:** 18 July 2019

**Published:** 06 August 2019

### Citation:

Ceusters N, Valcke R, Frans M, Claes JE, Van den Ende W and Ceusters J (2019) Performance Index and PSII Connectivity Under Drought and Contrasting Light Regimes in the CAM Orchid *Phalaenopsis*. *Front. Plant Sci.* 10:1012. doi: 10.3389/fpls.2019.01012

Crassulacean acid metabolism (CAM) is a specialized mode of photosynthesis characterized by improved water use efficiency mediated by major nocturnal CO<sub>2</sub> fixation. Due to its inherent metabolic plasticity CAM represents a successful physiological strategy for plant adaptation to abiotic stress. The present study reports on the impact of drought stress and different light intensities (PPFD 50 and 200 μmol m<sup>-2</sup> s<sup>-1</sup>) on the photosynthetic performance of the obligate CAM orchid *Phalaenopsis* “Edessa” by integrating diel gas exchange patterns with assessments of the light reactions by analyzing fast chlorophyll a fluorescence induction. Parameters such as PI<sub>abs</sub> (performance index), different energy fluxes per active reaction centre (RC) reflecting the electron flow from photosystem II to photosystem I and the energetic communication between PSII complexes defined as connectivity were considered for the first time in a CAM plant. A higher PS II connectivity for plants grown under low light ( $\rho \sim 0.51$ ) compared to plants grown under high light ( $\rho \sim 0.31$ ) brought about similar specific energy fluxes of light absorbance, dissipation and processing through the electron transport chain, irrespective of the light treatment. With a 25% higher maximum quantum yield and comparable biomass formation, low light grown plants indeed proved to process light energy more efficiently compared to high light grown plants. The performance index was identified as a very reliable and sensitive parameter to indicate the onset and progress of drought stress. Under restricted CO<sub>2</sub> availability (due to closed stomata) leaves showed higher energy dissipation and partial inactivation of PSII reaction centres to reduce the energy input to the electron transport chain and as such aid in avoiding overexcitation and photodamage. Especially during CAM idling there is a discrepancy between continuous input of light energy but severely reduced

availability of both water and CO<sub>2</sub>, which represents the ultimate electron acceptor. Taken together, our results show a unique flexibility of CAM plants to optimize the light reactions under different environmental conditions in a dual way by either attenuating or increasing energy flux.

**Keywords:** chlorophyll fluorescence, crassulacean acid metabolism, *Phalaenopsis*, performance index, PSII connectivity, specific energy fluxes


## INTRODUCTION

Plants thrive in a variety of environments, each associated with certain characteristics and limitations (e.g., rainforest, desert, and arctic conditions). Due to their sessile nature they need to be extremely adaptable to their continuously changing environment. Crassulacean acid metabolism (CAM), which is characterized by an optimized water use efficiency by taking up CO<sub>2</sub> predominantly at night is an important physiological strategy for plant adaptation (Cushman and Borland, 2002; Borland et al., 2011; Ceusters et al., 2017). Traditionally, diel CAM has been defined within a four-phase framework to describe the photosynthetic performance (Osmond, 1978): (i) phase I – open stomata in the dark and external CO<sub>2</sub> fixation via phosphoenolpyruvate carboxylase (PEPC) into C<sub>4</sub> acids (mostly malic acid); (ii) phase 2 – open stomata at the start of the light period and external CO<sub>2</sub> fixation by combination of PEPC and ribulose-1,5-bisphosphate carboxylase-oxygenase (RubisCO); (iii) phase 3 – closed stomata in the middle of the day whilst malic acid is decarboxylated [catalyzed by NAD(P)-malic enzyme (ME) or phosphoenolpyruvate carboxykinase] and refixation of CO<sub>2</sub> by RubisCO; and (iv) phase 4 – stomata open toward the end of the day and external CO<sub>2</sub> is fixed via RubisCO and/or PEPC. An ubiquitous feature of CAM is the integration of circadian and metabolite control over nocturnal C<sub>4</sub> and daytime C<sub>3</sub> carboxylation processes, hereby providing plasticity for optimizing carbon gain and water use by extending or curtailing the period of net CO<sub>2</sub> uptake over any 24-h period (Borland et al., 1999; Ceusters et al., 2009a, 2010; Haydon et al., 2011; Ceusters et al., 2014; Yang et al., 2015). When experiencing severe drought stress CAM plants enter the stage of CAM-idling, showing no net CO<sub>2</sub> uptake and recycling respiratory CO<sub>2</sub> behind closed stomates during the complete diel cycle to minimize any further water loss (Ting, 1985; Ceusters et al., 2009a,b).

Plant photosynthesis is generally characterized by two linked key processes, i.e., (1) light energy capture and use in the electron transport chains to provide energy and reducing power and (2) CO<sub>2</sub> sequestration into triose phosphates (Figure 1). Non-destructive methods like chlorophyll fluorescence and leaf gas exchange analyses can easily be applied to investigate photosynthetic performance. Chlorophyll fluorescence analyses allow to obtain detailed information about the primary light mediated reactions of photosynthesis. Basically, light energy absorbed by the sun can be dissipated by either photochemical (i.e., photosynthesis) or non-photochemical processes (i.e., heat or fluorescence) (Maxwell and Johnson, 2000). As these processes occur in close competition, abiotic stress factors such as

deviations of the optimal temperature, light and water availability can easily cause imbalances between the absorption of light energy by chlorophyll and the subsequent use of energy in CO<sub>2</sub> fixation (Figure 1; Cornic and Massacci, 1996; Falk et al., 1996). Evaluation of the response of the photosynthetic apparatus to different conditions has generally been performed by analyzing parameters describing the maximum quantum efficiency of photosystem II (PSII) photochemistry ( $F_v/F_m$ ), the effective quantum yield of PSII photochemistry ( $\Phi_{PSII}$ ), electron transport rate (ETR), photochemical quenching (qP) and non-photochemical quenching (qN). In addition different authors suggested that (1) the performance index ( $PI_{abs}$ ) parameter, which quantifies the overall functionality of the electron flow through PSII, might be a sensitive parameter of plant homeostasis; and (2) that it is also interesting to evaluate specific energy fluxes per reaction centre (RC) i.e., absorption (ABS/RC), trapping ( $TR_0/RC$ ), electron transport ( $ET_0/RC$ ), reduction of end electron acceptors ( $RE_0/RC$ ) and dissipation ( $DI_0/RC$ ) to obtain information about the response of the photosynthetic apparatus under different environmental conditions (Krüger et al., 1997; Rapacz, 2007; Zivcak et al., 2008, 2014a,b; Campos et al., 2014; Chondrogiannis and Grammatikopoulos, 2016). The specific energy fluxes measured per active PSII reaction centre are all interconnected and can be considered as expressing the behavior of the PSII system when all RCs are open (i.e., at the onset of excitation). In healthy plants (Figure 1A), part of the flux of absorbed photons (ABS) will be channeled as trapping flux to the RC (TR) and another part of this excitation energy is dissipated, mainly as heat (DI). An electron transport will be created within the RCs (ET) and further to PSI (RE), ultimately leading to CO<sub>2</sub> fixation (Strasser et al., 2000). Another intriguingly topic is the possibility of energetic communication (also known as “grouping” or “connectivity” and “overall grouping probability”) between different PSII complexes (Stirbet, 2013), allowing the transfer of excitation energy from a closed PSII RC to an open (active) PSII RC (Figures 1B,C). Zivcak et al. (2014a) demonstrated that the omission of connectivity can lead to misinterpretation of the JIP-test results in mature barley plants. The JIP test, which stands for the major inflection points in the fluorescence induction curve, is based on the theory of energy fluxes in the photosynthetic apparatus (Strasser et al., 2000; Stirbet and Govindjee, 2011). The concept of PSII connectivity is still under discussion in scientific literature and has not yet been taken into account in studies with CAM plants which are exceptionally well adapted to environmental stresses.

In CAM ecophysiology, chlorophyll fluorescence has already been explored to some extent regarding adaptation and survival strategy in response to variations in the natural



environment. Measurements of chlorophyll fluorescence quenching confirmed the existence of diverse photoprotective mechanisms in CAM involving non-radiative excess energy dissipation and xanthophyll cycling (Winter et al., 1990; Adams and Demmig-Adams, 1996; Barker and Adams, 1997; Masrahi et al., 2015). Changes in the electron transport rates in CAM, related to CO<sub>2</sub> reduction rates in chloroplasts, were shown to result primarily in alterations in the rates of non-radiative energy dissipation so that consequently the reduction state of the primary electron acceptor of PSII was maintained at relatively low and constant levels (Winter and Demmig, 1987). The extra requirements of ATP for CAM plants are reflected by high values of total qN, which can be twice as much as compared to C3 (Keiller et al., 1994; Heber et al., 1996; Skillman and Winter, 1997). Chlorophyll fluorescence measurements also revealed the negative impact of abiotic stress, related to water and light availability, on the photosynthetic

capacity of different CAM plants by significantly reducing photosynthetic electron transport rate and the overall effective quantum yield of PSII photochemistry, despite an increased capacity in dissipating excess energy (Osmond, 1982; de Mattos et al., 1999; Pollet et al., 2009; Arias-Moreno et al., 2017; Zheng et al., 2019).

The purpose of the present work was to gain more insight in the functioning of the photosynthetic apparatus in the CAM orchid *Phalaenopsis* “Edessa” under different light conditions and under drought stress, by evaluating performance indices (PI<sub>abs</sub>) and specific energy fluxes as described above. Therefore a detailed study was performed integrating diel gas exchange patterns with analyses of fast chlorophyll *a* fluorescence induction. Under both environmental conditions, specific fluxes were calculated without and with connectivity to investigate the importance of energetic communication between PSII complexes in the CAM orchid *Phalaenopsis* “Edessa.”

## MATERIALS AND METHODS

### Plant Material and Experimental Setup

*Phalaenopsis* “Edessa” is an obligate CAM plant and belongs to the family of the Orchidaceae. Vegetative plants of 16 weeks old were cultivated in a growth room with a day/night temperature of 28°C, a relative humidity of 75% and a 12-h photoperiod with photosynthetic photon flux density (PPFD) of 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$  at leaf level. Watering was performed twice a week; one time with a nutrient solution Peters 20N-8.7P-16.6K of 1  $\text{mS cm}^{-1}$  and once with water. Following a 2 months acclimation to these conditions, plants were divided into two groups to investigate the possible effect of light intensity on the photosynthetic performance: (1) PPFD of 50  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (low light: LL) and (2) PPFD of 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (high light: HL). As total diel  $\text{CO}_2$  uptake for *Phalaenopsis* has earlier been noticed to saturate around 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , this light intensity is further considered as high light treatment for *Phalaenopsis* “Edessa” (Guo et al., 2012). To determine chlorophyll content and titratable acidity samples were taken from the upper one-third of young fully expanded leaves ( $n = 5$  plants) at week 0, 4, 6, and 10. In addition a drought experiment was performed with other plants kept at 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , which had also been divided into two groups: (1) control group which continued to receive watering twice a week and (2) a treatment group which was attributed to drought stress by withholding water. Samples were taken from the upper one-third of young fully expanded leaves ( $n = 5$  plants) to determine chlorophyll content and titratable acidity at week 0, 4, and 6. Leaf discs ( $n = 5$  plants) were punched out just beneath the sample position to determine relative water content (RWC). At week six, the oldest and most wilted leaves of the treatment group were also included to analyze chlorophyll *a* fluorescence.

### Gas Exchange Measurements and Titratable Acidity

For both light and drought experiments, gas exchange was measured on the youngest fully expanded leaves after 10 and 6 weeks, respectively, using a LCi Portable Photosynthesis System (ADC BioScientific Ltd., United Kingdom). The top part of the leaf was enclosed in a broad leaf chamber (6.25  $\text{cm}^2$ ) and the incoming air was passed through a 20-l bottle to buffer short-term fluctuations in the  $\text{CO}_2$  concentration. Gas exchange data were collected over a 24-h period with measurements obtained at 15-min intervals ( $n = 3$  plants). By integrating specific areas under the  $\text{CO}_2$  exchange curves, net gas exchange was calculated per phase as well as total net gas exchange during the 24-h time course. Nocturnal  $\text{CO}_2$  uptake was also assessed by analyzing the nocturnal increase in titratable acidity ( $\Delta\text{H}^+$ ) by titration of methanol extracts against 0.005 M NaOH with phenolphthalein as indicator.

### Relative Water Content Determination

Relative water content was determined by sampling the youngest fully expanded leaves ( $n = 5$  plants) and calculated with the formula: (fresh mass – dry mass)/(turgid mass – dry mass) according to Ceusters et al. (2008). Leaf parts (95  $\text{mm}^2$ ) were

floated in plastic tubes filled with demineralized water for 6 h in darkness at 4°C to determine turgid mass.

### Determination of Chlorophyll Content and Plant Growth

For the calculation of leaf chlorophyll contents, plant pigments were extracted by immersing leaf material in *N,N*-dimethylformamide (DMFA) at room temperature for 72 h in darkness ( $n = 5$  plants). The supernatant was used to determine absorbance at 647 nm ( $A_{647}$ ) and 664 nm ( $A_{664}$ ) (Genesys 10S UV-VIS, Thermo Fisher Scientific, United States). These data were used to calculate the content of chlorophyll *a*, chlorophyll *b* and total chlorophyll by means of the empirical formulas:  $C_a = 11.65 A_{664} - 2.69 A_{647}$ ;  $C_b = 20.81 A_{647} - 4.53 A_{664}$ ;  $C = C_a + C_b$  (Moran and Porath, 1980; Porra et al., 1989; Wellburn, 1994).

Differences in plant growth, after growing for 10 weeks under different light conditions, were evaluated on basis of differences in above ground fresh and dry mass (g). Above ground plant material was separated from the roots, weighed and dried at 70°C to a constant mass after 8 days ( $n = 5$  plants).

### Chlorophyll *a* Fluorescence

Chlorophyll *a* fluorescence measurements were carried out by means of a Handy PEA fluorometer (Plant Efficiency Analyser, Hansatech, King’s Lynn, United Kingdom) and were taken on the adaxial side, always on the left side of the main vein, of young fully developed leaves ( $n = 15$  plants). Measurements were performed using a saturating pulse of 3 000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , pulse duration of 1 s, and fixed gain (1.2 $\times$ ). Induction curve analysis (Handy PEA software V1.10) allowed to evaluate the effectiveness of fluorescence saturation during measurements. Before carrying out measurements, leaves were always allowed to dark adapt for 20 min using light-withholding clips (Hansatech). The light level of the saturating pulse and the minimal dark period had experimentally been determined for *Phalaenopsis* “Edessa” beforehand in order to obtain true values for the chlorophyll *a* fluorescence parameters. During the light experiment chlorophyll measurements were taken at 08.00 h (Phase II) on week 0, week 4, week 6, and week 10. All chlorophyll fluorescence data was assembled at 08.00 h because additional measurements at 12.00 h (Phase III) revealed no significant differences for these parameters in the different CAM phases (**Supplementary Table 1**). To study the progression of drought stress, chlorophyll fluorescence measurements were also taken at 08.00 h on week 0, week 4, and week 6.

The measured fast chlorophyll fluorescence induction curves ( $F_0$  to  $F_m$ ) were analyzed by the JIP test, which is based on the theory of energy fluxes in the photosynthetic apparatus (Strasser et al., 2000; Stirbet and Govindjee, 2011). When receiving light, part of the flux of photons absorbed by PSII antenna pigments (ABS) is dissipated (DI), mainly as heat, and another part is converted to redox energy by reducing the electron acceptor  $Q_A$  (TR). This electron acceptor is then reoxidized creating an electron transport (ET) until the reduction of the end electron acceptors at the PSI electron

acceptor side (RE) (**Figure 1A**). This stepwise flow of energy through PSII can also be expressed per RC, defined as following specific energy fluxes: ABS/RC,  $Tr_0$ /RC,  $Di_0$ /RC,  $Et_0$ /RC, and  $Re_0$ /RC. All these specific fluxes refer to time zero, i.e., the onset of fluorescence induction. In a logarithmic time scale, fast chlorophyll fluorescence induction curves have a typical shape which shows the steps O, J, I, P (Strasser and Strasser, 1995; Srivastava et al., 1999; Strasser et al., 2000, 2004, 2010), making it possible to collect following cardinal points: maximal fluorescence intensity ( $F_m$ , when all RCs are closed), minimum fluorescence intensity ( $F_0$ , when all RCs are open), fluorescence intensity at 2 and 30 ms ( $F_J$  and  $F_I$ , respectively) and at 50 and 300  $\mu$ s ( $F_{50 \mu s}$  and  $F_{300 \mu s}$ ). These primary data were used to calculate chlorophyll fluorescence parameters describing maximum quantum efficiency of PSII ( $F_v/F_m$ ), performance index ( $PI_{abs}$ ) and the afore mentioned specific energy fluxes (Chondrogiannis and Grammatikopoulos, 2016). Definitions and equations of the measured and calculated JIP parameters are described in **Table 1**.

Since fluorescence induction data may be affected by the existence of PSII excitonic connectivity (**Figures 1B,C**), i.e., transfer of excitation energy from a closed PSII RC to an open (active) PSII RC (Stirbet, 2013), it was interesting to take this process into account. Connectivity parameters were calculated to allow comparison between results without and with taking connectivity into account (**Table 1**). Based on Zivcak et al. (2014a) the curvature constant (C) of the initial phase of the O-J curve (from 0.05 to 2 ms) was used to correct the values of the specific

fluxes for connectivity, i.e., multiplying the specific flux values by  $1 + C$  (**Table 1**).

## Statistical Analyses

Where appropriate, data were analyzed using the statistical software package IBM SPSS Statistics V23. Before carrying out statistical tests, normality of the data was checked by means of the Kolmogorov-Smirnoff statistic ( $p > 0.01$ ). Since our goal was to interpret the chlorophyll fluorescence data in function of the specific treatments (control/drought and different light intensities) and exclude non-treatment effects (plant age, time effects, ...), results were always compared between treatments from the same age by independent sample *t*-test or Tukey's Studentized range test, both at the 1% probability level. All replicates considered in our study were independent biological replicates originating from different plants.

## RESULTS

### Light


#### Gas Exchange Measurements and Titratable Acidity

For both investigated PPFD conditions (i.e., LL:  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$  and HL:  $200 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) a traditional diel CAM pattern was observed (**Figure 2**). A complete 24 h CAM cycle occurred with nocturnal  $\text{CO}_2$  fixation (Phase I) and net  $\text{CO}_2$  loss during the day (Phase III), accompanied by two intermediate phases (Phases II and IV). During Phase I, extending from 20.00 h

**TABLE 1** | Definitions and equations of measured and calculated parameters derived from fast fluorescence kinetics.

Parameter	Name and basic physiological interpretation
<b>Basic JIP-test parameters</b>	
$V_t = (F_t - F_{50 \mu s}) / (F_m - F_{50 \mu s})$	Relative variable fluorescence at time t ( $V_J$ , $V_I$ at 2, 30 ms)
$M_0 = dV/dt_0$	Approximate value of the initial slope of relative variable fluorescence curve $V_t$ (for $F_0 = F_{50 \mu s}$ )
<b>Specific energy fluxes (per active PSII reaction centre)</b>	
$ABS/RC = (M_0/V_J) \times (1/\phi P_0)$	Absorbed photon flux per RC
$Tr_0/RC = M_0/V_J$	Trapped excitation flux (leading to $Q_A$ reduction) of absorbed photons per RC
$Et_0/RC = (M_0/V_J) \times (1 - V_J)$	Electron transport flux (from reduced $Q_A$ to $Q_B$ ) per RC
$Re_0/RC = (M_0/V_J) \times (1 - V_I)$	Electron flux reducing end electron acceptors at the PSI acceptor side, per RC
$Di_0/RC = (ABS/RC - Tr_0/RC)$	Dissipated energy flux per RC
<b>Performance index</b>	
$PI_{ABS} = RC/ABS \times [\phi P_0 / (1 - \phi P_0)] \times [\Psi E_0 / (1 - \Psi E_0)]$	Performance index for energy conservation from photons absorbed by PSII antenna, to the reduction of $Q_B$
<b>Quantum yields and probabilities</b>	
$\phi P_0 = F_v/F_m$	Maximum quantum efficiency of primary PSII photochemistry; maximum efficiency at which light absorbed by PSII is used for reduction of $Q_A$
$\Psi E_0 = 1 - V_J$	Probability with which a PSII trapped electron is transferred from reduced $Q_A$ to $Q_B$
<b>Connectivity among PSII units</b>	
$W_E = 1 - [(F_J - F_{300 \mu s}) / (F_J - F_{50 \mu s})]^{1/5}$	Model-derived value of relative variable fluorescence at 100 $\mu$ s calculated for unconnected PSII units
$W = (F_{100 \mu s} - F_{50 \mu s}) / (F_J - F_{50 \mu s})$	Relative variable fluorescence at 100 $\mu$ s
$C = (W_E - W) / [V_J \times W \times (1 - W_E)]$	Curvature constant of initial phase of the O-J curve
$\rho_{2G} = C \times [F_{50 \mu s} / (F_J - F_{50 \mu s})]$	Overall grouping probability between PSII units when they are all open
$\rho = [\rho_{2G} \times (F_m / F_{50 \mu s} - 1)] / [1 + \rho_{2G} \times (F_m / F_{50 \mu s} - 1)]$	Connectivity parameter
$\omega = \rho \times [(F_m - F_{50 \mu s}) / F_m]$	Probability of the connectivity among PSII units when they are all closed

Based on information presented by Strasser et al. (1995, 2000, 2004, 2010), Strasser and Stirbet (2001), Stirbet and Govindjee (2011), Stirbet (2013).



**FIGURE 2** | Net 24 h CO<sub>2</sub> uptake ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) for young fully expanded leaves of *Phalaenopsis* "Edessa" 10 weeks after growing under different light conditions i.e., PPFD of  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$  (blue) and PPFD of  $200 \mu\text{mol m}^{-2} \text{s}^{-1}$  (orange). The dark period is indicated in gray. Gas exchange curves are representative of three replicate runs with SE < 15%.

to 8.00 h, nocturnal CO<sub>2</sub> uptake of LL grown plants was only approximately one third that of plants grown under HL. This difference was also reflected in the nocturnal increase of titratable acidity which was already significantly higher for HL plants after 4 weeks (Figure 3). In addition, during Phase II, CO<sub>2</sub> uptake was lowered by 80% when grown under LL after 10 weeks. Phase IV was also delayed with 2 h under LL, resulting in only 44% CO<sub>2</sub> uptake in comparison to plants grown under HL. As a consequence, total daily CO<sub>2</sub> gain of HL plants was three times higher compared to LL plants ( $129 \pm 20 \text{ mmol m}^{-2}$  and  $41 \pm 19 \text{ mmol m}^{-2}$  respectively; Figure 2).

### Chlorophyll a Fluorescence Parameters


Comparing the two light conditions, no significant differences were observed for the maximum quantum efficiency of PSII photochemistry ( $F_v/F_m$ ) and for the performance index for

absorption ( $PI_{\text{abs}}$ ) (Figure 4 and Table 1). However, regarding the specific energy fluxes (per active RC) (Figure 1A and Table 1), significantly higher values were obtained for plants grown under HL (Figure 4A). After 6 weeks, a significant increase was noticed for photon absorption (ABS/RC) for plants grown under HL, concomitantly with a slight increase in the activity of energy dissipation ( $Di_0/RC$ ). A significant rise was also calculated for the electron trapping efficiency ( $Tr_0/RC$ ) after 10 weeks grown under HL. For the activity of electron transport within the reaction centre ( $Et_0/RC$ ) and the flow of electrons further than PSII ( $Re_0/RC$ ) significant differences were observed from week 4. Also for these parameters, the highest values were calculated for plants grown under HL.


To estimate the connectivity among PSII units, the first part of fast chlorophyll fluorescence induction curve was used (from 0.05 to 2 ms) as described above (Strasser and Stirbet, 2001; Joliot and Joliot, 2003; Stirbet, 2013). From week 6, calculated values of parameters associated with connectivity (Table 1), i.e., overall grouping probability of PSII units –  $p_{2G}$ , connectivity parameter –  $p$ , probability of connectivity among PSII units –  $\omega$  (as defined by Strasser and Stirbet, 2001), were ~1.5 times higher in plants grown under LL compared to those grown under HL (Figure 5). After correction for connectivity, i.e., multiplying the specific flux values by  $1 + C$  (Zivcak et al., 2014a), all values obtained under LL became higher (Figure 4B) and eliminated the significant differences in the activity of energy dissipation ( $Di_0/RC$ ) and electron transport ( $Et_0/RC$ ) originating from not considering connectivity. At week 6 photon absorption (ABS/RC) was equally for both light conditions and trapping efficiency was significantly increased in LL plants when taking connectivity into account.

### Chlorophyll Content and Plant Growth


After 10 weeks of growing under different light conditions, significantly lower values in content of chlorophyll *a*, chlorophyll *b* and total chlorophyll were observed for plants grown under the highest light condition (Figure 6). The Chl *a/b* ratio remained



**FIGURE 3** | Nocturnal increase (between 20.00 h and 08.00 h) in titratable acidity ( $\text{mmol H}^+ \text{m}^{-2}$ ) in young fully expanded leaves of *Phalaenopsis* "Edessa" at 0, 4, 6, and 10 weeks after growing under different light conditions, i.e., PPFD of  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$  (blue) and PPFD of  $200 \mu\text{mol m}^{-2} \text{s}^{-1}$  (orange). Data are means  $\pm$  SD ( $n = 5$  plants). Asterisks indicate significant differences per week between different light conditions at  $P < 0.01$  according to the independent sample *t*-test.



**FIGURE 4** | “Spider plots” of selected chlorophyll a fluorescence transients parameters measured at 08.00 h in young fully expanded leaves of *Phalaenopsis* “Edessa” at 0, 4, 6, and 10 weeks after growing under different light conditions, i.e., PPFD of 50  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (blue) and PPFD of 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (orange). Parameters are calculated without connectivity (A) and corrected for connectivity between PSII units (B). All values are shown as percent of PPFD of 50  $\mu\text{mol m}^{-2} \text{s}^{-1}$  ( $n = 15$  plants). Asterisks indicate significant differences between different light conditions at  $P < 0.01$  according to the independent sample  $t$ -test.



unaffected by light condition. Both fresh and dry mass of the shoots showed an overall similar plant growth for plants grown under LL and HL after 10 weeks (Table 2).

## Drought


### Gas Exchange Measurements and Titratable Acidity

The  $\text{CO}_2$  uptake pattern of *Phalaenopsis* “Edessa” grown under standard conditions (PPFD 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ; 12-h photoperiod), watered twice a week, fitted the four-phase CAM framework as earlier described (Figure 7). After 6 weeks of



withholding water no net  $\text{CO}_2$  uptake occurred, indicating that the plants were in the CAM-idling mode with closed stomata during the complete diel cycle. In accordance with the gas exchange data, the nocturnal increase in titratable acidity was reduced by >80% when grown under drought stress (Figure 8A).

### Chlorophyll a Fluorescence Parameters and Relative Water Content

To indicate the onset of water stress, RWC was measured and showed a significant decrease from 4 weeks of withholding water supply (Figure 8B). During progression of drought the







**TABLE 2** | Biomass on fresh and dry mass (g) basis for above ground material of *Phalaenopsis* "Edessa" grown for 10 weeks under different light conditions, i.e., PPFD 50  $\mu\text{mol m}^{-2} \text{s}^{-1}$  or 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ .


	PPFD ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ )	
	50	200
Fresh	94 $\pm$ 13	74 $\pm$ 5
Dry	4.4 $\pm$ 0.6	4.0 $\pm$ 0.4

Data are means  $\pm$  SD ( $n = 5$  plants) and no significant differences between the different light conditions were detected at  $P < 0.01$  according to the independent sample  $t$ -test.

maximal quantum efficiency of PSII photochemistry ( $F_v/F_m$ ) remained unaffected (Figure 9 and Table 1). In contrast, the  $PI_{\text{abs}}$  parameter was already significantly lower after 4 weeks of drought stress in comparison to control plants (Figure 9). Drought stress also affected the specific energy fluxes within

and related to PSII (Figure 1A and Table 1). After 4 weeks of drought stress, a significant increase was observed for photon absorption ( $ABS/RC$ ) compared to control plants (Figure 9A). Also for the trapping efficiency ( $Tr_0/RC$ ) and the flow of electrons further than PSII ( $Re_0/RC$ ), significantly higher values were obtained at week 4 in the leaves of plants exposed to drought stress (Figure 9A). At week ten  $Tr_0/RC$  was significantly higher in the most wilted leaves whilst  $Re_0/RC$  and  $Et_0/RC$  remained unaffected. Activity of energy dissipation ( $Di_0/RC$ ) showed a significant increase in the most wilted leaves compared to the measurements in plants grown under control conditions.

The calculated connectivity parameters (Table 1) of measurements performed at week 6 in leaves of control plants were  $\sim 1.3$  times higher compared to measurements performed on the youngest fully expanded leaves of plants exposed to drought stress, and  $\sim 7.6$  times higher compared to measurements on the most wilted leaves of plants exposed to



drought stress (Figure 10). As a consequence, the increase of the calculated fluxes after correction for connectivity, was the highest in the control leaves and the lowest in the most wilted leaves (Figure 9B). Taking connectivity into account showed an increased energy dissipation ( $Di_0/RC$ ) for drought stressed plants and indicated an equal trapping efficiency ( $Tr_0/RC$ ) for drought and wilted leaves at week six (Figure 9B).


### Chlorophyll Content

Overall, the content of photosynthetic pigments remained unaffected during the 6 weeks of drought stress. No significant

differences were observed for Chl *a*, Chl *b*, and total Chl between control and drought stressed plants (Figure 11).

## DISCUSSION

Chlorophyll fluorescence measurements in CAM plants mainly report photochemical and non-photochemical quenching ( $qP$  and  $qN$ ), the maximum quantum efficiency ( $F_v/F_m$ ) and the effective quantum yield ( $\Phi_{PSII}$ ) of PSII photochemistry (Table 1; Winter and Demmig, 1987; Winter et al., 1990; Keiller et al., 1994;



Adams and Demmig-Adams, 1996; Heber et al., 1996; Barker and Adams, 1997; Skillman and Winter, 1997; de Mattos et al., 1999; Lin and Hsu, 2004; Masrahi et al., 2015; Liu et al., 2016; Arias-Moreno et al., 2017). A detailed investigation of the performance index, specific energy fluxes related to PSII processing and the concept of connectivity between PSII units (Figure 1 and Table 1) revealed important PSII related adaptations concerning the initial light reactions in CAM plants under different environmental conditions.


## Increased PSII Connectivity Under Light Limitation

Chlorophyll fluorescence measurements were performed to investigate the effects of low light (LL: PPFD  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) and high light (HL: PPFD  $200 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) on the photosynthetic performance of the CAM plant *Phalaenopsis* “Edessa.” Our detailed energy flux analyses focused on the efficiency of energy capture and the electron transport in PSII system when all RCs are open (i.e., at the onset of excitation). Comparing these parameters between LL and HL (Figure 4), stresses the importance of the energetic communication between PSII complexes defined as connectivity when evaluating specific energy fluxes in CAM plants (Figure 1). Without taking connectivity into account (Figure 4A), all values of specific energy fluxes were significantly lower after 10 weeks of growing under LL compared to the HL treatment, indicating an affected energy processing through PSII upon light limitation. However, taking into account higher connectivity of LL grown plants (Figure 5), higher values were recorded for all energy fluxes (Figure 4B) and as such these plants seemed to process light energy more efficiently compared to plants grown under the high light conditions. This was also reflected in a 25% higher maximum quantum yield for plants under LL as indicated by relating the daily  $\text{CO}_2$  gain of a leaf to the amount of photons absorbed per day (Skillman, 2008). Plants grown under LL realized an overall similar biomass on both fresh and dry mass basis (Table 2) despite a triple decrease in  $\text{CO}_2$  fixation in the measured source leaves for plants under LL (Figure 2). However,

HL grown plants were typically characterized by several layers of leaf pairs largely shading each other whilst LL grown plants showed an improved architecture with less underlying sink leaves and larger source leaves. Under LL, total chlorophyll content in source leaves after 10 weeks was also 140% in comparison to contents of plants grown under HL (Figure 6C), which was also reflected by the higher photon absorption flux (ABS/RC) for plants grown under LL (Figure 4B). The increase in total chlorophyll content was mainly due to a significant increase in Chl *a* (Figure 6A), which can reflect an increase in active RC’s. As a consequence, the higher connectivity in plants grown under LL is consistent with the absence of a difference in the performance index  $\text{PI}_{\text{abs}}$  (Figure 4 and Table 1), due to their more efficient energy processing. Without considering connectivity an increase in  $\text{PI}_{\text{abs}}$  would be expected for HL, since both trapping efficiency and electron transport increased significantly by HL and only the parameter RC/ABS decreased. Taking connectivity into account changed the concentration of reaction centres per chlorophyll (RC/ABS) and trapping efficiency ( $\text{Tr}_0/\text{RC}$ ) in the opposite direction, whilst the other parameters remained unaffected and no difference was observed for  $\text{PI}_{\text{abs}}$ . As such, the question can be raised whether a higher connectivity for plants grown under low light conditions might underpin the adaptive radiation of CAM plants from full sun to deep shade conditions. Several species from the Bromeliaceae are excellent examples of CAM representatives thriving in light limited environments such as wet cloud forests and shaded understoreys of tropical rainforests (Pierce et al., 2002; Ceusters et al., 2011).

## Increased Energy Dissipation and Partial Inactivation of PSII Reaction Centres to Reconcile Energy and Carbon Requirements During Drought Stress

To evaluate the effects of drought stress on the photosynthetic performance in the CAM plant *Phalaenopsis* “Edessa,” chlorophyll fluorescence parameters (Table 1) were compared between control plants (watered twice a week) and drought stressed plants. In contrast to the maximum quantum efficiency



of PSII ( $F_v/F_m$ ) which remained unaffected, the performance index  $PI_{\text{abs}}$  reflected the reduction of RWC for drought stressed plants (Figure 9). These results corroborate investigations in C3 plants where the performance index  $PI_{\text{abs}}$  has already been proposed a reliable and sensitive parameter to evaluate plant homeostasis with regard to moderate abiotic stress (Zivcak et al., 2008; Campos et al., 2014; Jedmowski et al., 2015). As indicated in Table 1, the performance index takes into account three dependent characteristics: the concentration of reaction centres per chlorophyll (RC/ABS), a parameter related to primary photochemistry ( $\varphi P_0$ ) and a parameter related to electron transport ( $\psi E_0$ ) (Strasser et al., 2004). As a consequence, changes in environmental conditions which influence either antenna properties (ABS/RC), trapping efficiency ( $Tr_0/RC$ ) or electron transport beyond  $Q_A$  in PSII ( $Et_0/RC$  and  $Re_0/RC$ ) (Figure 1) will have an impact on the current state of plant performance and  $PI_{\text{abs}}$  will decrease under stress. Mathematically, the observed decrease in performance index could not be explained by merely looking at the three separate parameters on which  $PI_{\text{abs}}$  is based (Table 1). Only the trapping efficiency ( $Tr_0/RC$ ) significantly increased under drought whilst RC/ABS was only significantly different between control and drought, and the parameter  $Et_0/RC$  remained unaffected (Figure 9A). Taking connectivity into account eliminated the observed significant difference for  $Tr_0/RC$  between drought and wilted but could not fully

explain the observed significant decrease in  $PI_{\text{abs}}$  (Figure 9B). From these data it can be concluded that other components or processes, such as oxygen-evolving complex (OEC); transition from active to non- $Q_A$ -reducing centres; carotenoids and PSI may also influence the performance index (Figure 1). Different studies revealed that drought can indeed result in damage to the OEC and block further electron transport. This phenomenon is mostly characterized by an increase in the specific fluxes for trapping, absorption and dissipation (Guissé et al., 1995; Wang et al., 2012; Guha et al., 2013).

Our data also demonstrate that enhanced protection of photochemistry in the CAM plant *Phalaenopsis* “Edessa” is also achieved by adjusting the energy distribution between photosystems. Two possible explanations can be presented for the observed increase in ABS/RC under drought stress (Figure 9B): (1) inactivation of some PSII RC’s, considering that the ABS/RC ratio is calculated as the total number of photons absorbed by Chl molecules in all RC’s divided by total number of active RC’s, or (2) an increase in antenna size (van Heerden et al., 2007; Gomes et al., 2012). The latter seems to be less plausible since no significant differences were obtained in the content of photosynthetic pigments between the two treatments (Figure 11). The inactivation of some PSII RC’s, becoming non- $Q_A$ -reducing centres, can explain the observed increase in  $Tr_0/RC$  under drought. Down-regulation of photochemical

activity together with a significant increase in thermal dissipation may play a critical role in protecting plants exposed to drought stress from over-excitation and photodamage (van Heerden et al., 2007; Zivcak et al., 2014b; Wang et al., 2017; Kalaji et al., 2018). As such, the performance index depicts a downregulation of PSII related energy fluxes as a physiological response to drought stress. Especially for CAM plants during CAM idling the Calvin-Benson cycle, for which the end products of the electron transport chain (ATP and NADH) are indispensable (Figure 1A), is seriously compromised by carbon and water limitation (Figure 7). In this physiological mode of survival, inactivation of several PSII RC's adds in overcoming severe drought periods by attenuating energy fluxes in consistence with the very low CO<sub>2</sub> availability, due to stomatal closure during day and night, in the leaf mesophyll cells.

## CONCLUSION

The results presented in this paper suggest a clear physiological role for PSII connectivity (energetic communication between PSII complexes) in the CAM plant *Phalaenopsis* "Edessa." A higher connectivity for plants grown under low light ( $p \sim 0.51$ ) compared to plants grown under high light ( $p \sim 0.31$ ) resulted in similar specific energy fluxes irrespective of the light treatment. With a 25% higher maximum quantum yield and comparable biomass formation, low light grown plants proved to process light energy more efficiently compared to high light grown plants. As such, taking into account PSII connectivity might help to better understand the adaptation of CAM plants to changing environmental conditions. The obtained results also indicate that PI<sub>abs</sub> is a sensitive parameter to detect drought stress in the CAM plant *Phalaenopsis* "Edessa." Besides an enhanced thermal dissipation drought stressed plants also seemed to attenuate the electron flow from PSII toward PSI by partial inactivation of PSII reaction centres. This strategy might help to reconcile the light reactions and the carbon fixation reactions which are seriously compromised by carbon and water limitation during CAM-idling and as such minimize chances of photodamage to occur. Further

## REFERENCES

- Adams, W. W., and Demmig-Adams, B. (1996). "Energy dissipation and the xanthophyll cycle in CAM plants," in *Crassulacean acid Metabolism. Ecological studies (Analysis and Synthesis)*, Vol. 114, eds K. Winter, and J. A. C. Smith, (Berlin: Springer).
- Arias-Moreno, D. M., Jiménez-Bremont, J. F., Maruri-López, I., and Delgado-Sánchez, P. (2017). Effects of catalase on chloroplast arrangement in *Opuntia streptacantha* chlorenchyma cells under salt stress. *Sci. Rep.* 7:8656. doi: 10.1038/s41598-017-08744-x
- Barker, D. H., and Adams, W. W. (1997). The xanthophyll cycle and energy dissipation in differently oriented faces of the cactus *Opuntia macrorhiza*. *Oecologia* 109, 353–361. doi: 10.1007/s004420050093
- Borland, A. M., Hartwell, J., Jenkins, G. I., Wilkins, M. B., and Nimmo, H. G. (1999). Metabolite control overrides circadian regulation of phosphoenolpyruvate carboxylase kinase and CO<sub>2</sub> fixation in crassulacean acid metabolism. *Plant Physiol.* 121, 889–896. doi: 10.1104/pp.121.3.889
- Borland, A. M., Zambrano, V. A. B., Ceusters, J., and Shorrocks, K. (2011). The photosynthetic plasticity of crassulacean acid metabolism: an evolutionary innovation for sustainable productivity in a changing world. *New Phytol.* 191, 619–633. doi: 10.1111/j.1469-8137.2011.03781.x

research into the specific energy fluxes and the connectivity between different PSII units in CAM plants under combinations of abiotic stress factors is highly encouraged. This will enable to expand our physiological knowledge about the interplay of the light and carbon fixation reactions in CAM plants under changing environmental conditions.

## DATA AVAILABILITY

The raw data supporting the conclusions of this manuscript will be made available by the authors, without undue reservation, to any qualified researcher.

## AUTHOR CONTRIBUTIONS

NC, RV, and JC proposed the conceptual framework for the study and performed the data collection and analysis. NC and MF performed the experimental analyses. NC, RV, JEC, WVdE, and JC interpreted the data and wrote the manuscript.

## FUNDING

This research was supported by the Research Fund KU Leuven.

## ACKNOWLEDGMENTS

Microflor NV is acknowledged for supplying the plant materials.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2019.01012/full#supplementary-material>

- Campos, H., Trejo, C., Peña-Valdivia, C. B., García-Nava, R., Conde-Martínez, F. V., and Cruz-Ortega, M. D. R. (2014). Photosynthetic acclimation to drought stress in *Agave salmiana* otto ex Salm-Dyck seedlings is largely dependent on thermal dissipation and enhanced electron flux to photosystem I. *Photosynth. Res.* 122, 23–39. doi: 10.1007/s11120-014-0008-6
- Ceusters, J., Borland, A. M., Ceusters, N., Verdoodt, V., Godts, C., and De Proft, M. P. (2010). Seasonal influences on carbohydrate metabolism in the CAM bromeliad *Aechmea* 'Maya': consequences for carbohydrate partitioning and growth. *Ann. Bot.* 105, 301–309. doi: 10.1093/aob/mcp275
- Ceusters, J., Borland, A. M., and De Proft, M. P. (2009a). Drought adaptation in plants with crassulacean acid metabolism involved the flexible use of different storage carbohydrate pools. *Plant Signal. Behav.* 4, 212–214. doi: 10.4161/psb.4.3.7813
- Ceusters, J., Borland, A. M., Londers, E., Verdoodt, V., Godts, C., and De Proft, M. P. (2009b). Differential usage of storage carbohydrates in the CAM bromeliad *Aechmea* 'Maya' during acclimation to drought and recovery from dehydration. *Physiol. Plant.* 135, 174–184. doi: 10.1111/j.1399-3054.2008.01186.x
- Ceusters, J., Borland, A. M., Godts, C., Londers, E., Croonenborghs, S., Van Goethem, D., et al. (2011). Crassulacean acid metabolism under severe light

- limitation: a matter of plasticity in the shadows? *J. Exp. Bot.* 62, 283–291. doi: 10.1093/jxb/erq264
- Ceusters, J., Borland, A. M., Taybi, T., Frans, M., Godts, C., and De Proft, M. P. (2014). Light quality modulates metabolic synchronization over the diel phases of crassulacean acid metabolism. *J. Exp. Bot.* 65, 3705–3714. doi: 10.1093/jxb/eru185
- Ceusters, J., Londers, E., Verdoodt, V., Ceusters, N., and De Proft, M. P. (2008). Seasonal impact on physiological leaf damage risk of *Aechmea* hybrid under greenhouse conditions. *Sci. Hortic.* 118, 242–245. doi: 10.1016/j.scienta.2008.06.006
- Ceusters, N., Van den Ende, W., and Ceusters, J. (2017). “Exploration of sweet immunity to enhance abiotic stress tolerance in plants: lessons from CAM,” in *Progress in Botany*. eds M. C. Francisco, U. Lüttge, and R. Matyssek, (Berlin: Springer), 78.
- Chondrogiannis, C., and Grammatikopoulos, G. (2016). Photosynthesis in developing leaf of juveniles and adults of three Mediterranean species with different growth forms. *Photosynth. Res.* 130, 427–444. doi: 10.1007/s11120-016-0276-4
- Cornic, G., and Massacci, A. (1996). “Leaf photosynthesis under drought stress,” in *Photosynthesis and the Environment*, Edn 1, ed. N. R. Baker, (Netherlands: Springer).
- Cushman, J. C., and Borland, A. M. (2002). Induction of crassulacean acid metabolism by water limitation. *Plant Cell Environ.* 25, 295–310. doi: 10.1046/j.0016-8025.2001.00760.x
- de Mattos, E. A., Herzog, B., and Lüttge, U. (1999). Chlorophyll fluorescence during CAM –phases in *Clusia minor* L. under drought stress. *J. Exp. Bot.* 50, 253–261. doi: 10.1093/jxb/50.331.253
- Falk, S., Maxwell, D. P., Laudenschlager, D. E., and Huner, N. P. A. (1996). “Photosynthetic adjustment to temperature,” in *Photosynthesis and the Environment*, Edn 1, ed. N. R. Baker, (Netherlands: Springer).
- Gomes, M. T. G., da Luz, A. C., dos Santos, M. R., Batitucci, M. D. C. P., Silva, D. M., and Falqueto, A. R. (2012). Drought tolerance of passion fruit plants assessed by the OJIP chlorophyll a transient. *Sci. Hortic.* 142, 49–56. doi: 10.1016/j.scienta.2012.04.026
- Guha, A., Sengupta, D., and Reddy, A. R. (2013). Polyphasic chlorophyll a fluorescence kinetics and leaf protein analyses to track dynamics of photosynthetic performance in mulberry during progressive drought. *J. Photochem. Photobiol. B Biol.* 119, 71–83. doi: 10.1016/j.jphotobiol.2012.12.006
- Guissé, B., Srivastava, A., and Strasser, R. J. (1995). “Effects of high temperature and water stress on the polyphasic chlorophyll a fluorescence transient of potato leaves,” in *Photosynthesis: from light to biosphere*, ed. P. Mathis, (Dordrecht: Kluwer Academic Publishers), 913–916.
- Guo, W. J., Lin, Y. Z., and Lee, N. (2012). Photosynthetic light requirements and effects of low irradiance and daylength on *Phalaenopsis amabilis*. *J. Am. Soc. Hortic. Sci.* 137, 465–472. doi: 10.21273/jashs.137.6.465
- Haydon, M. J., Bell, L. J., and Webb, A. A. R. (2011). Interactions between plant circadian clocks and solute transport. *J. Exp. Bot.* 62, 2333–2348. doi: 10.1093/jxb/err040
- Heber, U., Kaiser, W. M., and Neimanis, S. (1996). “Regulation of crassulacean acid metabolism in *Kalanchoë pinnata* as studied by gas exchange and measurements of chlorophyll fluorescence,” in *Crassulacean acid metabolism. Ecological studies (Analysis and Synthesis)*, Vol. 114, eds K. Winter, and J. A. C. Smith, (Berlin: Springer).
- Jedrowski, C., Ashoub, A., Momtaz, O., and Brüggemann, W. (2015). Impact of drought, heat and their combination on chlorophyll fluorescence and yield of wild barley (*Hordeum spontaneum*). *J. Bot.* 2015, 120868. doi: 10.1155/2015/120868
- Joliet, P., and Joliet, A. (2003). Excitation transfer between photosynthetic units: the 1964 experiment. *Photosynth. Res.* 76, 241–245.
- Kalaji, H. M., Rackova, L., Paganova, V., Swoczyna, T., Rusinowski, S., and Sitko, K. (2018). Can chlorophyll-a fluorescence parameters be used as bio-indicators to distinguish between drought and salinity stress in *Tillia cordata* Mill? *Environ. Exp. Bot.* 152, 149–157. doi: 10.1016/j.envexpbot.2017.11.001
- Keiller, D. R., Slocombe, S. P., and Cockburn, W. (1994). Analysis of chlorophyll a fluorescence in C3 and CAM forms of *Mesembryanthemum crystallinum*. *J. Exp. Bot.* 45, 325–334. doi: 10.1093/jxb/45.3.325
- Krüger, G. H. J., Tsimilli-Michael, M., and Strasser, R. J. (1997). Light stress provokes plastic and elastic modifications in structure and function of photosystem II in camellia leaves. *Physiol. Plant.* 101, 265–277. doi: 10.1034/j.1399-3054.1997.1010203.x
- Lin, M. J., and Hsu, B. D. (2004). Photosynthetic plasticity of *Phalaenopsis* in response to different light environments. *J. Plant Physiol.* 161, 1259–1268. doi: 10.1016/j.jplph.2004.05.009
- Liu, Y. C., Liu, C. H., Lin, Y. C., Lu, C. H., Chen, W. H., and Wang, H. L. (2016). Effect of low irradiance on the photosynthetic performance and spiking of *Phalaenopsis*. *Photosynthetica* 54, 259–266. doi: 10.1007/s11099-016-0079-z
- Masrahi, Y. S., Al-Turki, T. A., and Sayed, O. H. (2015). Photosynthetic adaptation and survival strategy of *Duvalia velutina* in an extremely arid environment. *Photosynthetica* 53, 555–561. doi: 10.1007/s11099-015-0143-0
- Maxwell, K., and Johnson, G. N. (2000). Chlorophyll fluorescence – a practical guide. *J. Exp. Bot.* 51, 659–668. doi: 10.1093/jxb/51.345.659
- Moran, R., and Porath, D. (1980). Chlorophyll determination in intact tissues extracted with N,N-dimethylformamide. *Plant Physiol.* 65, 478–479. doi: 10.1104/pp.65.3.478
- Osmond, C. B. (1978). Crassulacean acid metabolism. a curiosity in context. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 29, 379–414. doi: 10.1093/jxb/ern052
- Osmond, C. B. (1982). “Carbon cycling and stability of the photosynthetic apparatus in CAM,” in *Crassulacean acid metabolism*, eds I. P. Ting, and M. Gibbs, (Rockville: American Society of Plant Physiologists), 112–127.
- Pierce, S., Winter, K., and Griffiths, H. (2002). The role of CAM in high rainfall cloud forests: an in situ comparison of photosynthetic pathways in Bromeliaceae. *New Phytol.* 25, 1181–1189. doi: 10.1046/j.1365-3040.2002.00900.x
- Pollet, B., Stepe, K., Van Labeke, M. C., and Lemeur, R. (2009). Diurnal cycle of chlorophyll fluorescence in *Phalaenopsis*. *Photosynthetica* 47, 309–312. doi: 10.1007/s11099-009-0048-x
- Porra, R. J., Thompson, W. A., and Kriedemann, P. E. (1989). Determination of accurate extinction coefficients and simultaneous equation for assaying chlorophylls a and b extracted with four different solvents: Verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochim. Biophys. Acta* 975, 384–394. doi: 10.1016/s0005-2728(89)80347-0
- Rapacz, M. (2007). Chlorophyll a fluorescence transient during freezing and recovery in winter wheat. *Photosynthetica* 45, 409–418. doi: 10.1007/s11099-007-0069-2
- Skillman, J. B. (2008). Quantum yield variation across the three pathways of photosynthesis: not yet out of the dark. *J. Exp. Bot.* 59, 1647–1661. doi: 10.1093/jxb/ern029
- Skillman, J. B., and Winter, K. (1997). High photosynthetic capacity in a shade-tolerant crassulacean acid metabolism plant. *Plant Physiol.* 113, 441–450. doi: 10.1104/pp.113.2.441
- Srivastava, A., Strasser, R. J., and Govindjee. (1999). Greening of peas: parallel measurements of 77K emission spectra, OJIP chlorophyll a fluorescence transient, period four oscillation of the initial fluorescence level, delayed light emission, and P700. *Photosynthetica* 37, 365–392.
- Stirbet, A. (2013). Excitonic connectivity between photosystem II units: what is it, and how to measure it? *Photosynth. Res.* 116, 189–214. doi: 10.1007/s11120-013-9863-9
- Stirbet, A., and Govindjee. (2011). On the relation between the Kautsky effect (chlorophyll a fluorescence induction) and Photosystem II: Basics and applications of the OJIP fluorescence transient. *J. Photochem. Photobiol. B Biol.* 104, 236–257. doi: 10.1016/j.jphotobiol.2010.12.010
- Strasser, B. J., and Strasser, R. J. (1995). “Measuring fast fluorescence transients to address environmental questions: The JIP-test,” in *Photosynthesis: From Light to Biosphere*, ed. P. Mathis, (Dordrecht: Kluwer Academic Publishers), 977–980.
- Strasser, R. J., Srivastava, A., and Govindjee, G. (1995). Polyphasic chlorophyll a fluorescence transients in plants and cyanobacteria. *Photochem. Photobiol.* 61, 32–42. doi: 10.1111/j.1751-1097.1995.tb09240.x
- Strasser, R. J., Srivastava, A., and Tsimilli-Michael, M. (2000). “The fluorescence transient as a tool to characterize and screen photosynthetic samples,” in *Probing photosynthesis: Mechanisms Regulation and Adaptation*, eds M. Yunus, U. Pathre, and P. Mohanty, (London: Taylor and Francis), 445–483.

- Strasser, R. J., and Stirbet, A. D. (2001). Estimation of the energetic connectivity of PSII centres in plants using the fluorescence rise O-J-I-P. Fitting of experimental data to three different PSII models. *Math. Comput. Simulation* 56, 451–461.
- Strasser, R. J., Tsimilli-Michael, M., Qiang, S., and Goltsev, V. (2010). Simultaneous in vivo recording of prompt and delayed fluorescence and 820 nm reflection changes during drying and after rehydration of the resurrection plant *Haberlea rhodopensis*. *Biochim. Biophys. Acta Bioenerg.* 1797, 1313–1326. doi: 10.1016/j.bbabi.2010.03.008
- Strasser, R. J., Tsimilli-Michael, M., and Srivastava, A. (2004). “Analysis of the chlorophyll fluorescence transient,” in *Chlorophyll a fluorescence: a signature of photosynthesis*, eds G. C. Papageorgiou, and Govindjee, (Netherlands: Springer Press), 321–362. doi: 10.1007/978-1-4020-3218-9\_12
- Ting, I. P. (1985). Crassulacean acid metabolism. *Annu. Rev. Plant Physiol.* 36, 595–622.
- van Heerden, P. D. R., Swanepoel, J. W., and Krüger, G. H. J. (2007). Modulation of photosynthesis by drought in two desert scrub species exhibiting C3-mode CO2 assimilation. *Environ. Exp. Bot.* 61, 124–136. doi: 10.1016/j.envexpbot.2007.05.005
- Wang, Y., Xu, C., Wu, M., and Chen, G. (2017). Characterization of photosynthetic performance during reproductive stage in high-yield hybrid rice LYJP exposed to drought stress probed by chlorophyll a fluorescence transient. *Plant Growth Regul.* 81, 489–499. doi: 10.1007/s10725-016-0226-3
- Wang, Z. X., Chen, L., Ai, J., Qin, H. Y., Liu, Y. X., Xu, P. L., et al. (2012). Photosynthesis and activity of photosystem II in response to drought stress in Amur Grape (*Vitis amurensis* Rupr.). *Photosynthetica* 50, 189–196. doi: 10.1007/s11099-012-0023-9
- Wellburn, A. R. (1994). The spectral determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. *J. Plant Physiol.* 144, 307–313. doi: 10.1016/s0176-1617(11)81192-2
- Winter, K., and Demmig, B. (1987). Reduction state of Q and nonradiative energy dissipation during photosynthesis in leaves of a crassulacean acid metabolism plant, *kalanchoë daigremontiana* hamet et perr. *Plant Physiol.* 85, 1000–1007. doi: 10.1104/pp.85.4.1000
- Winter, K., Lesch, M., and Diaz, M. (1990). Changes in xanthophyll-cycle components and in fluorescence yield in leaves of a crassulacean-acid-metabolism plant, *clusia rosea* Jacq., throughout a 12-hour photoperiod of constant irradiance. *Planta* 182, 181–185. doi: 10.1007/BF00197108
- Yang, X., Cushman, J. C., Borland, A. M., Edwards, E. J., Wullschleger, S. D., Tuskan, G. A., et al. (2015). A roadmap for research on crassulacean acid metabolism (CAM) to enhance sustainable food and bioenergy production in a hotter, drier world. *New Phytologist.* 207, 491–504. doi: 10.1111/nph.13393
- Zheng, L., Ceusters, J., and Van Labeke, M. C. (2019). Light quality affects light harvesting and carbon sequestration during the diel cycle of crassulacean acid metabolism in *Phalaenopsis*. *Photosynth. Res.* doi: 10.1007/s11120-019-00620-1 [Epub ahead of print].
- Zivcak, M., Brestic, M., Kalaji, H. M., and Govindjee. (2014a). Photosynthetic responses of sun- and shade-grown barley leaves to high light: is the lower PSII connectivity in shade leaves associated with protection against excess of light? *Photosynth. Res.* 119, 339–354. doi: 10.1007/s11120-014-9969-8
- Zivcak, M., Kalaji, H. M., Shao, H. B., Olsovska, K., and Brestic, M. (2014b). Photosynthetic proton and electron transport in wheat leaves under prolonged moderate drought stress. *J. Photochem. Photobiol. B Biol.* 137, 107–115. doi: 10.1016/j.jphotobiol.2014.01.007
- Zivcak, M., Brestic, M., Olsovska, K., and Slamka, P. (2008). Performance index as a sensitive indicator of water stress in *Triticum aestivum* L. *Plant Soil Environ.* 54, 133–139. doi: 10.17221/392-pse

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Ceusters, Valcke, Frans, Claes, Van den Ende and Ceusters. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.