

Performance of Wheat (*Triticum aestivum* L.) as Influenced by Application of Single Super Phosphate Alone and In Combination with Different Organic Fertilizers in Calcareous Soil of Swabi, Pakistan

Aftab JAMAL¹, Waqar YOUNAS¹, Muhammad FAWAD²

¹Department of Soil and Environmental Sciences, The University of Agriculture Peshawar, Pakistan, ²Department of Weed Sciences, The University of Agriculture Peshawar, Pakistan.

✉ : aftabses98@gmail.com

ABSTRACT

The influence of Single Super Phosphate (SSP) alone and in combination with different organic fertilizers were evaluated in a field study on wheat (*Triticum aestivum* L.) production conducted at farmer field located in Swabi, Pakistan during 2017-18. Results exposed that among all the treatments, combined application of SSP at 100kg ha⁻¹ along with 6 t ha⁻¹ Poultry manure significantly improved the growth as well as yield parameters of wheat crop. On the other hand, spike length remained unaffected by all the treatments in the prevailing soil and environmental condition. Combined use of SSP and PM were strongly suggested for obtaining optimum yield of wheat crop.

Article History

Received : 29.08.2018

Accepted : 08.10.2018

Keywords

Single super phosphate,
Poultry manure,
Swabi,
Pakistan

Research Article

Pakistan'ın Swabi Bölgesindeki Kireçli Topraklara Uygulanan Süper Fosfat ve Super Fosfat ile Organik Gübre Kombinasyonlarının Buğdayın (*Triticum aestivum* L.) Performansına Etkisi

ÖZET

2017-2018 yıllarında Swabi (Pakistan)'da yetiştirici şartlarında yürütülen bu çalışmada, Sadece Süper Fosfat (SSP), Organik ve Süper Fosfat ile farklı organik gübre (SSP+Organic Manure) karışımlarının buğday verim ve performansına etkisi araştırılmıştır. Uygulanan gübre kombinasyonlarından hektara 100 kg SSP ve 6 t kümes hayvanı gübresesi (SSP+PM) kombinasyonunun buğday verimi ve büyüme parametrelerini önemli ölçüde iyileştirdiği sonucuna varılmıştır. Diğer yandan, başak uzunluğunun mevcut toprak ve çevre koşullarında herhangi bir uygulamadan etkilenmediği görülmüştür. Çalışma sonucuna göre; optimum düzeyde buğday verimi elde etmek için SSP ve PM'nin birlikte kullanımı önerilmiştir.

Makale Tarihi

Geliş Tarihi : 29.08.2018

Kabul Tarihi : 08.10.2018

Anahtar Kelimeler

Tek süper fosfat,
Kanath gübresesi,
Swabi,
Pakistan

Araştırma Makalesi

To cite: Aftab JA, Younas W, Fawad M 2019. Performance of Wheat (*Triticum aestivum* L.) as Influenced By Application of Single Super Phosphate Alone and In Combination with Different Organic Fertilizers In Calcareous Soil of Swabi, Pakistan. KSÜ Tar Doğa Derg 22(1): 14-18. DOI: 10.18016/ksutarimdog.v22i39650.455876.

INTRODUCTION

Wheat technically known as *Triticum aestivum* L. belongs to the family Poaceae, one of the most important cereal crops grown all over the world. Pakistan ranked 6th among top wheat producing countries in the world and 3rd largest producer in Asia. In cereal crop production wheat ranked 1st in Pakistan, contributes greatly to growth and survival of peoples of the country (GOP, 2016). It is a predominant staple food and a major daily diet providing carbohydrates, dietary fibers, proteins and vitamins (Economic survey 2001; Hussain *et al.* 2004; FAO, 2009). About 80 % of Pakistani farmers cultivated wheat on an area of 9.23 million hectares with total production of 25.5 million tons during the winter season 2016. Wheat contributed

10 % to the total agriculture and 2.1 % to the total country gross domestic product (GDP) (MINFA, 2016). Wheat yield in Pakistan is 2 tons ha⁻¹ which is comparatively very low as compare to United Kingdom and Germany yield 8 and 7 tons ha⁻¹ respectively. The major constraints of low wheat productivity in Pakistan are inappropriate nutrients application, intensive cropping system, unpredictable weather condition and erosion that considerably deplete the soil fertility and ultimately affect the wheat productivity. Continues use of chemical fertilizers and improper soil and nutrients management create catastrophic effects on our soil fertility (Arif *et al.* 2014). Moreover, Continuous application of chemical fertilizers causes soil health problems even if applied in balanced

proportion (Zia *et al.*, 2000). For this purpose, agriculture scientists are now trying to develop an agriculture system which not only lower the production cost but also conserve the natural resources (Abbas *et al.*, 2012). Organic fertilizers like FYM, compost and PM either in combination with each other or with mineral fertilizers render greater beneficial effects on growth as well as yield parameters of crop (Gowda, *et al.* 2010).

Pakistan soils are generally low in organic matter (less than 1 %) firstly because of the arid climate of the country and secondly because of very less amount of OM added during crop cultivation (Abbas *et al.*, 2012) and it can be increased through the utilization of both organic and mineral fertilizers in combination (Bodruzzaman *et al.* 2010). Organic fertilizers may be the best substitute for chemical fertilizers because the importance of organic fertilizers cannot be ignored. Organic fertilizers have many characteristics. They supply a variety of nutrients to the crop as well as add organic matter content to the soil which in turn improves the soil physical characteristics. The positive interaction between the combination of organic and mineral fertilizers was reported by (Pong and Laty, 2000). Now-a-day artificially synthesized fertilizers are the trend and commonly in use but it is also fact that increased the expenses of farmers to such a level that an average farmer can't avail those sources. Apart from these uses of artificially synthesized fertilizer badly affect the soil fertility from various angles and contaminates the environment at various levels. The combination of organic manures with chemical fertilizers can be helpful to wash out the worse effects of the long-lasting use of synthetically prepared fertilizers (Jama *et al.* 1997).

Keeping in mind the importance of organic fertilizers the present study was launched to evaluate the effect of SSP alone and in combination with different organic fertilizers on wheat crop yield.

MATERIALS and METHODS

Site Description

A field study was launched at farmer field in village Dagi, District Swabi, lies between 34° 7', 48" N and 72° 28', 11" E of Khyber Pakhtunkhwa, Pakistan.

Experimental design and treatments

The experiment was laid out in randomized complete block design (RCBD) having three replications. Each plot size was kept to 20 m². A total of eight treatments (given below) were utilized in the experiment. Phosphorous (P) as Single Super phosphate (SSP) was applied at rate of 100 kg ha⁻¹ alone and in combination with different organic fertilizers. Farm yard manure (FYM) and compost was applied at the rate of 10 t ha⁻¹ and poultry manure (PM) was applied at the rate of 6 t ha⁻¹ (Recommended doses for under study soil).

T1	Control (No. fertilizers)
T2	Single super phosphate (SSP) (at rate of 100kg ha ⁻¹)
T3	Compost (at rate of 10 t ha ⁻¹)
T4	Farm yard manure (FYM) (at rate of 10 t ha ⁻¹)
T5	Poultry manure (PM) (at rate of 6 t ha ⁻¹)
T6	SSP + Compost
T7	SSP + FYM
T8	SSP + PM

Methodology

The wheat variety Manthar 03 was used as test crop during the experiment and was sown with seed rate of 120 kg seed ha⁻¹ and maintained row to row distance of 30 cm. P as SSP was applied at the rate of 100 kg ha⁻¹. A week before sowing of seeds, SSP and organic fertilizers were applied. At sowing time Recommended doses of nitrogen at the rate of 120 kg ha⁻¹ and potassium as sulphate of potassium (SOP) was applied at the rate of 60 kg ha⁻¹. Nitrogen was applied in two splits, first dose at sowing time and second at the time of first irrigation in the form of urea. The crop was irrigated at continuous intervals according to the need of crop and the weeds which emerged during crop growth were manually pulled from whole of the experimental area. The growth parameters like plant height and spike length (cm) were recorded at maturity stage of crop. Wheat yield parameters like 1000- grain weight, grain yield and biological yield (t ha⁻¹) were recorded after harvesting of crop.

Analysis of soil and organic fertilizers

Before the experiment a composite soil sample was taken from the field at 0-15 cm depth and physiochemical properties like pH, EC (Richards, 1954), soil texture (Gee and Bauder 1986), lime content (U.S. Salinity Laboratory Staff. 1954), and OM (Nelson and Sommer, 1982) of the experimental soil were determined. The soil was also analyzed for total N (Bremner, 1996), AB-DTPA- extractable P and K, (Soltanpour, 1985). The under-study soil was found silty clay loam in texture, alkaline in reaction with pH value (8.12), EC (1.4 dsm⁻¹), low in organic matter (0.69) and highly calcareous in nature (CaCO₃ 16.8%). The available total N 0.05 %, available P 3.5 mg kg⁻¹ and exchangeable K 120 mg kg⁻¹ were recorded respectively (Table1). The same physiochemical characteristics of the same soil series was also reported by Jamal and Jamal (2018).

Statistical Analysis

The collected data were statistically analyzed and recorded. Least significant difference (LSD) at 5 percent probability level was computed to compare treatment means (Steel *et al.* 1997). Organic fertilizers were also subjected to nutrient analysis (Ryan *et al.* 2001) and the results are presented in (Table 2).

Table 1. Some soil physico-chemical properties (0-15 cm depth)

Property	Value
Clay (%)	19
Silt (%)	62
Sand (%)	19
Textural class	Silty Clay Loam (SCL)
pH (1:5; Soil:Water)	8.1
Organic matter (%)	0.69
EC (dsm ⁻¹)	1.4
Lime (CaCO ₃) (%)	16.8
AB-DTPA P (mg kg ⁻¹)	3.5
Total N (%)	0.05
AB-DTPA K (mg kg ⁻¹)	120

Table 2. Chemical analysis of Organic fertilizers used in the study

Organic fertilizers	Nitrogen (%)	Phosphorous (%)	Potash (%)
Farm yard manure	0.55	0.39	0.62
Compost	0.62	0.4	0.56
Poultry manure	1.33	0.79	0.36

Table 3. Wheat plant height, spike length, 1000-grain weight, grain yield and biological yield as influenced by Single Super Phosphate alone and in combination with different organic fertilizers.

Treatments	Plant Height (cm)	Spike length (cm)	1000-grain weight (g)	Grain Yield(t ha ⁻¹)	Biological yield (t ha ⁻¹)
T1= Control	52.33 c	8.15	30.00 d	2.11 d	5.5 e
T2 =SSP	71.66 ab	8.19	41.67 bc	4.08 abc	10.66 b
T3=Compost	65.33 abc	8.17	39.67 bc	3.66 bc	8.14 cd
T4=FYM	62.00 bc	8.08	38.00 c	3.41 c	7.37 d
T5=PM	67.33 ab	8.11	40.33 bc	3.98 abc	9.06 c
T6=SSP + Compost	74.33 ab	8.03	44.33 ab	4.48 abc	12.81 a
T7 =SSP + FYM	72.33 ab	8.19	42.67 abc	4.04 abc	11.88 ab
T8=SSP + PM	78.33 a	8.08	47.00 a	4.63 a	13.11 a
LSD at 0.05	13.64	N.S	5.19	0.88	1.50

*Means with different letter (s) in columns are significantly different at P<0.05

1000-Grain weight (g)

Single super phosphate either alone or in combination with organic fertilizers significantly improved 1000-grain weight of wheat as compared with control; however, the application SSP in combination with PM (T8) produced the maximum 1000-grain weight of 47 g followed by 44.33g produced by the T6 received the combination of SSP and compost (Table3). Published literature showed that PM along with chemical fertilizers increased wheat 1000-grain yield significantly. Abbas *et al.* (2012) reported that PM at 6 t ha⁻¹ along with recommended doses of N, P, K significantly increased wheat 1000-grain weight, similarly in another study Akhtar *et al.* (2000) investigated that organic fertilizers and chemical fertilizers in combination significantly improved yield contributing factor in wheat than the chemical fertilizer applied alone.

RESULTS and DISCUSSION

Plant height and Spike length (cm)

Application of single super phosphate alone and in combination with different organic fertilizers significantly increased wheat plant height as compared with control (Table 3). Maximum plant height of 78.33 cm was observed with the combined application of SSP +PM closely followed by SSP + compost application treatment. However, the spike length was not significantly affected by the applied treatments and the maximum spike length of 8.19 cm was observed in T2 and minimum of 8.03 cm in T6 (Table 3). Published literature revealed that crop growth parameters may be improved with organic and inorganic fertilizers application (Dixit & Gupta (2000); Selvakumari *et al.* (2000) and Khoshgoftarmanesh and Kalbasi (2002). Delden, (2001) stated that wheat plant height can be increased with application of organic and inorganic fertilizers. However, the nonsignificant effect of organic fertilizers on wheat spike length was also reported by Jamal and Fawad (2018) in the prevailing soil and environmental condition.

Grain yield (t ha⁻¹)

Wheat grain yield was significantly increased with application of SSP and organic fertilizers either alone or in combination as compared with the control treatment. The highest grain yield of 4.63 t ha⁻¹ was observed in T8 (SSP + PM) which was followed by T6 (SSP + Compost) having grain yield 4.48 t ha⁻¹. The minimum grain yield 2.11 t ha⁻¹ was observed in T1 (control) (Table 3). Maximum grain yield in SSP + PM plot might be due to higher nutrients content in PM than FYM and Compost (Jamal and Fawad 2018). The increase in grain yield was related to improvement in some of yield contributing factors (Table 3). Our result was in lined with (Khanam *et al.*, 2001; Rees and Castle, 2002; and Reddy *et al.* 2005).

Biological yield (t ha⁻¹)

Similar to other parameters maximum biological yield 13.11 t ha⁻¹ was observed in T8 (SSP + PM) which was followed by T6 (SSP + Compost) having biological yield 12.81 t ha⁻¹ (Table 3), however no statistical difference was found for biological yield between T8 and T6. Among all the organic fertilizers, PM performed better in improvement wheat growth parameters as well as yield parameters. The highest biological yield in T8 may be due to the fact that PM directly supplies the essential nutrient like NPK to the plants more than other fertilizer source. Abbas *et al.*, (2012) suggested that combine application of PM at 6 t ha⁻¹ and recommended doses of NPK produced optimum wheat yield. Our result was also in lined with the findings of (Ghosh *et al.* 2004; Sarwer *et al.*, 2008). The significant effect of PM in increasing biological yield of wheat was also reported by Jamal and Fawad (2018) for the same soil and environmental condition.

CONCLUSION

It was clearly concluded from the study that application of Single Super Phosphate in combination with Poultry manure significantly increased wheat yield in the prevailing soil and environmental conditions. Both growth and yield parameters of wheat may be better improved with combination of SSP at 100 kg ha⁻¹ and poultry manure at 6 t ha⁻¹ respectively. Hence it is recommended that SSP + PM may be applied in combination for obtaining the optimum yield of wheat crop.

REFERENCES

- Abbas G, Khattak JZKA, Mir M, Ishaque M, Hussain H, Wahedi M, Ahmed MS, Ullah A 2012. "Effect of organic manures with recommended dose of NPK on the performance of wheat (*Triticumaestivum* L.)." *J. Anim. Plant. Sci.* 22(3): 683-687.
- Akhtar Z, Ali A, Akhtar, T, Yousaf M 2000. Comparative effect of phosphate fertilizer and farmyard manure on Kohistan-97. Integrated Plant Nutrition Management Symposium, 15-20 April, Islamabad.
- Arif M, Tasneem M, Bashir F, Yaseen G, Iqbal RM 2014. Effect of integrated use of organic manures and inorganic fertilizers on yield and yield components of rice. *J. Agric. Res.* 52(2): 197-206.
- Bodruzzaman M, Meisner CA, Sadat MA, Hossain MI 2010. Long-term effects of applied organic manures and inorganic fertilizers on yield and soil fertility in a wheat-rice cropping pattern. In *Proceedings of the 19th World Congress of Soil Science, 10-15 Aug, Brisbane, Australia.*
- Bremner JM 1996. Nitrogen total in D.L. (*Methods of Soil Analysis* Part 3. Sparks Ed. *Amer.Soc. Agron* : 1085-1022.
- Delden AV 2001. Yield and growth components of potato and wheat under organic nitrogen management. *Agron. J.*, 93(6): 1370–1385.
- Dixit KG, Gupta BR 2000. Effect of farmyard manure, chemical and biofertilizers on yield and quality of rice (*Oryza sativa* L.) and soil properties. *Journal of the Indian Society of Soil Science*, 48(4): 773-780.
- Economic Survey. 2001. Government of Pakistan, Economic Advisor's Wing, Finance Division, Islamabad.
- FAO 2009. Food balance sheets. Food and agriculture organization, Rome.
- Ghosh PK, Ramesh KK, Bandyopadhyay AK, Tripathi KM, Hati AK, Misra CL, Acharya 2004. Comparative effectiveness of cattle manure, poultry manure, phospho compost and fertilizer-NPK on three cropping systems in vertisols of semi-arid tropics. I. Crop yields and system performance. *Bioresource. Technology.* 95(1) :77-83.
- Gee, Bauder 1986. Particle-size analysis. (Methods of soil analysis. Part 1. Sparks Ed. *Amer.Soc. Agron*). 383-411.
- Gowda C, BiradarPatil NK, Patil BN, Awaknavar JS, Ninganur BT, Hunje R 2010. Effect of organic manures on growth, seed yield and quality of wheat. *Karnataka Journal of Agricultural Sciences*, 21(3): 21-26.
- GOP 2016. Pakistan becomes world's No. 6 wheat producing country. Dawn Newspaper 18 Oct, 2016.
- Hussain, T, Abbas S, Khan MA, Scrimshaw NS 2004. Lysine of wheat flour improves selected indices of the nutritional status of predominantly cereal eating families in Pakistan. *Food Nutr. Bull.* 25(2): 114-122.
- Jama B, Swinkels RA, Buresh RJ 1997. Agronomic and economic evaluation of organic and inorganic sources of phosphorus in western Kenya. *Agronomy Journal*, 89(4): 597-604.
- Jamal A, Jamal H 2018. Assessment and Distribution of Macro and Micro Nutrients in Different Soil Series of District Swabi, Khyber Pakhtunkhwa, Pakistan. *Journal of Horticulture and Plant Research.* 2:23-32
- Jamal A, Fawad M 2018. Application of different organic manures in optimizing optimum yield for wheat in calcareous soil. *World News of Natural Sciences.* 20: 23-30.
- Khanam M, Rahman MM, Islam MR, Islam MR 2001. Effect of manures and fertilizers on the growth and yield of BRRI Dhan 30. *Pak. J.Bio. Sci.* 4(2): 172-174.
- Khoshgoftarmanesh AH, Kalbasi M 2002. Effect of municipal waste leachate on soil properties and growth and yield of rice. *Communications in Soil Science and Plant Analysis*, 33(13-14): 2011-2020.
- MINFA 2016. Govt. of Pakistan, Ministry of Food and Agric. Econ. Wing, Islamabad.

- Nelson DW, Sommers L 1982. Total carbon, organic carbon, and organic matter1, (*Methods of Soil Analysis Part 2*. Sparks Ed. *Amer.Soc. Agron*). 539-579.
- Pong XP, laty J 2000.Organic farming challenge of timing N availability to crop N requirements. *Soil Sci. Soc* 64(1): 247-253.
- Richard 1954. Diagnosis and improvement of Saline and Alkali soils, USDA Handbook.No:60. US Govt.Press, Washington, DC 160.
- Ryan J, Estefan G, Rashid A 2001. Soil and Plant Analysis Laboratory Manual. International Centre for Agricultural Research in Dry Areas (ICARDA) and the National Agricultural Research Center (NARC), Aleppo, Syria and Islamabad, Pakistan Aleppo. 172p.
- Rees R, Castle K 2002. Nitrogen recovery in soils amended wit organic manures combined with inorganic fertilizers. *Agron. J.*, 22(7) :739–746.
- Reddy SS, Shivaraj B, Reddy VC, Ananda MG 2005.Direct effect of fertilizer and residual effect of organic manure on yield and nutrient uptake of maize (*Zea mays* L.) in groundnut-maize cropping system. *Crop Res. Hsiao*, 29(3) : 390-395.
- Soltanpour PN 1985. Use of AB-DTPA soil test to evaluate elemental variability and toxicity.1985 *Commun. Soil Sci. Plant Anal.* 16(3): 323-338.
- Sarwar G, Hussain N, Schmeisky H, Muhammad S, Ibrahim M, Safdar E 2008. Use of compost an environment friendly technology for enhancing rice-wheat production in Pakistan. *Pak. J. Bot.* 40(1):1553-1558.
- Selvakumari G, Baskar M, Jayanthi, D, Mathan KK. 2000. Effect of integration of fly ash with fertilizers and organic manures on nutrient availability, yield and nutrient uptake of rice in Alfisols. *Journal of the Indian Society of Soil Science*, 48(2): 268-278.
- Steel RGD, Torrie JH, Dickey DA 1997. Principles and Procedures of Statistics (A Biometrical Approach, 3rd Ed. *McGraw Hill Book Company*, New York) 172-177.
- US Salinity Laboratory Staff 1954. Carbonate and bicarbonate by titration with acid. Diagnosis and improvement of saline and alkali soils, 98.
- Zia MS, Mann RA, Aslam M, Khan MA, Hussain F 2000. The role of green manuring in sustaining rice-wheat production. *Integrated Plant Nutrition Management Symposium*, 15-20 April, Islamabad.