

3

Pesticide Contamination inside Farm and Non- Farm Homes

Brian Curwin, Misty Hein, Wayne Sanderson, Marcia Nishioka, Stephen Reynolds,
Elizabeth Ward, Michael Alavanja

Journal of Occupational and Environmental Hygiene 2005, 2(7):357-367

Abstract

Twenty-five farm (F) households and 25 non-farm (NF) households in Iowa were enrolled in a study investigating pesticide contamination inside homes. Air, surface wipe and dust samples were collected. Samples from 39 homes (20 F and 19 NF) were analyzed for atrazine, metolachlor, acetochlor, alachlor, and chlorpyrifos. Samples from 11 homes (5 F and 6 NF) were analyzed for glyphosate and 2,4-D. Greater than 88% of the air and 74% of the wipe samples were below the limit of detection (LOD). Among the air and wipe samples, chlorpyrifos was detected most frequently in homes. In the dust samples, all the pesticides were detected in greater than 50% of the samples except acetochlor and alachlor, which were detected in less than 30% of the samples. Pesticides in dust samples were detected more often in farm homes except 2,4-D, which was detected in 100 percent of the farm and non-farm home samples. The average concentration in dust was higher in farm homes versus non-farm homes for each pesticide. Further analysis of the data was limited to those pesticides with at least 50% of the dust samples above the LOD. All farms that sprayed a pesticide had higher levels of that pesticide in dust than both farms that did not spray that pesticide and non-farms, however, only atrazine and metolachlor were significantly higher. The adjusted geometric mean pesticide concentration in dust for farms that sprayed a particular pesticide ranged from 94 to 1300 ng/g compared to 12 to 1000 ng/g for farms that did not spray a particular pesticide and 2.4 to 320 ng/g for non-farms. The distributions of the pesticides throughout the various rooms suggest that the agricultural herbicides atrazine and metolachlor are potentially being brought into the home on the farmer's shoes and clothing. These herbicides are not applied in or around the home but they appear to be getting into the home para-occupationally. For agricultural pesticides take-home exposure may be an important source of home contamination.

Introduction

Farmers are the biggest users of pesticides applying approximately 1.2 billion pounds in 1999; herbicides accounted for the largest proportion of this amount with approximately 534 million pounds applied (1). A wide variety of agricultural pesticides are used on farms including herbicides, crop insecticides, livestock insecticides, fungicides, and fumigants. Crop herbicides are used the most with approximately 50 to 93 % of farmers reporting their use (2-4).

In farm homes, families may be exposed to pesticides through home contamination even though they may not participate in farming activities involving pesticide use.

Residential environments in proximity to farm operations where pesticides are used may be contaminated through a variety of routes including airborne spread, tracking of contaminated soil into the home, and through deposition on the clothing of applicators.

Indirect inhalation and dermal exposure of families to pesticides may occur through redistribution of pesticides via indoor air to surfaces (due to volatilization/condensation and resuspension/settling). A study by Lewis et al (5) which collected air, dust and surface wipe samples, documents rapid translocation of diazinon and chlorpyrifos within the home following indoor and outdoor home applications. Families are also exposed to pesticides through food and in homes that have been sprayed with pesticides.

The potential for exposure of children living on farms to pesticides is a serious concern. Several studies have found an association between *in utero* and postnatal household pesticide exposure and childhood leukemia (6-8). Differences in children's physiology, behavior patterns and hygiene may result in significantly greater exposures to

environmental contaminants than adults (9). Small children spend much of their time on the floor or ground and are very likely to come into contact with pesticide residues on carpets or uncovered floors when playing inside, and yard dirt when playing outside. Children may also be more susceptible than adults to the toxic effects of pesticides, due to the sensitivity of developing organ systems (10). Older children, through their increased mobility and ability to assist with farm work, may have opportunities for direct pesticide exposure. Although the public health importance of preventing injury to farm children has been well-recognized, the hazards of exposure to pesticides and other chemicals to children in the farm environment have received relatively little attention.

Studies have found that farm homes have a greater frequency of detectable residues of pesticides and higher concentrations of pesticides in dust than in reference homes, potentially leading to greater exposure to pesticides among family members (11-13). Pesticide urine concentrations among the children of farmers and farm workers have been shown to be elevated when compared to children of non-farm families (11, 14). These studies generally investigated organophosphate and other insecticides. To date, no studies investigating herbicide contamination in farm homes have been conducted.

Recent EPA-funded studies have shown that transport of lawn-applied pesticides in the residential environment can lead to elevated levels of those pesticides in the home within a short period of time after application. For example, Nishioka et al (15, 16) measured the distribution of the herbicide 2,4-D in homes within a week of a lawn application, and showed that transport mechanisms were dominated by track-in from active dogs, the home-owner's contaminated shoes, and the children's shoes when worn

indoors. Lewis et al (5) found that chlorpyrifos residues in indoor air and in carpet dust were higher within a few days of an exterior residential application than before the application, and suggested that track-in was the principal source of these residues.

It should be noted that to date studies investigating agricultural pesticide exposures among farm families have focused on insecticides, particularly organophosphates, while agricultural herbicide studies have not been conducted. Studies investigating track-in of herbicides have been so far confined to non agricultural, residential applications in non-farm homes. The primary purpose of this paper is to investigate farm home pesticide contamination to seven pesticides, six of which are herbicides, and to describe the sources of pesticide contamination in farm homes. A comparison of pesticide contamination will be made among farm homes and reference homes. This paper offers unique information on pesticide exposure among farm families not previously studied by investigating herbicide exposure, four of which are not used residentially, which offers insight into para-occupational exposure pathways in the home.

Methods

Study Population

In the spring and summer of 2001, 25 farm (F) households and 25 non-farm (NF) households in Iowa were enrolled in a study investigating agricultural pesticide contamination inside homes. Participant recruitment has been described previously (17). Briefly, participants were recruited from participants of the Agricultural Health Study in Keokuk and Mahaska counties, the Keokuk Country Rural Health Study in Iowa, and through word of mouth. To be eligible for the study, each home had to have

at least one child 8 years old or less. In addition, non-farm homes had to be located on land that was not used for farming and have no person in the home working in agriculture or commercial pesticide application, and the farm homes had to be using during the spring of 2001 at least one of the 7 target pesticides: atrazine, acetochlor, alachlor, chlorpyrifos, metolachlor, glyphosate, and 2,4-D. These pesticides were selected because of their extensive use in Iowa agriculture. Six of the pesticides are corn or soybean herbicides, while chlorpyrifos is an insecticide used on corn. The study protocol was reviewed and approved by the Institutional Review Boards of the National Institute for Occupational Safety and Health, University of Iowa and the National Cancer Institute.

Sample Collection

Between May and August of 2001, each home was visited on two occasions. The first visit was shortly after a spraying event, and the second visit was approximately 4 weeks later (mean 4 weeks, range 3 to 5 weeks). A three-part questionnaire was administered to either parent at each home on the first visit. The information was updated on the second visit. Part 1 dealt with parental information. Part 2 dealt with child information and included questions about whether children handled pesticides, performed other farm chores, or had access to treated fields. Part 3 dealt with household information, including residential pesticide use in and around the home, and proximity of the house to treated fields. In addition to the three-part questionnaire, a fourth part was administered to the principal farmer in the farm homes only, about those factors that may influence home contamination, including farm activities, agricultural pesticide use, crops, agricultural practice, and use of personal protective equipment (PPE) since the start of the growing season and throughout the study period. The questionnaire gathered

information from the start of the 2001 growing season until the last home visit, and generally reflected the early 2001 growing season. With respect to home, yard and garden use of residential pesticides, home owners were asked about their use in the month and year prior to the first visit, and the month between visits. Data on the pesticide applied, farm practices, farm demographics and household pesticide use have been reported previously (17).

Environmental samples, including surface wipe, dust, and air, were collected at each visit. Wipe samples were collected from the steering wheel and driver's seat of the primary family vehicle, and from the kitchen counter, top of the washing machine, and from various rooms with hard surface floors inside the home. Dust samples were collected from carpet where available, including wall-to-wall carpet, area rugs, or floor mats from the entrance way, father's change area, laundry room, child's bedroom and child's playroom. When floors from these rooms did not have a carpet or rug, a wipe sample was collected. A single 24-hour air sample was collected from the living room of each home and an additional 24-hour air sample was collected outside, near the home. Dust and wipe samples from 39 homes (20 F and 19 NF) were analyzed for atrazine, metolachlor, acetochlor, alachlor, and chlorpyrifos. Dust and wipe samples from 11 homes (5 F and 6 NF) were analyzed for glyphosate and 2,4-D. All air samples were analyzed for atrazine, metolachlor, acetochlor, alachlor, chlorpyrifos and 2,4-D.

Dust samples were collected from carpets using a high-volume surface sampler (HVS-3, Cascade Stamp Sampling Systems (CS3) Inc., Sandpoint, ID) using the American Society for Testing Material (ASTM) Standard Practice for Collection of Dust from Carpeted Floors for Chemical Analysis (18).

The wiping procedure consisted of sampling a 1 ft x 1 ft (900 cm²) area using two 4 inch x 4 inch (103.2 cm²) Johnson and Johnson SOF-WICK[®] sponges (Physician Sales and Services, Cincinnati, OH) sequentially. The first sponge was moistened with 10 ml of 100% isopropanol and four adjacent but slightly overlapping wipes of approximately 8 cm width were taken in one direction. The sponge was folded after each pass so that a clean surface was available for each wipe. The sponge was placed in an amber glass jar covered with a PTFE lined cap. The second sponge was then moistened with 10 ml of 100% isopropanol and four more adjacent 8 cm wide wipes were taken in a similar manner but in a direction perpendicular to the first wipe. The second sponge was added to the jar containing the first sponge. To sample the steering wheel, one sponge was wrapped around the steering wheel and half the wheel was wiped. The sponge was folded in half, and the second half of the steering wheel was wiped in the same manner. The procedure was repeated with the second sponge, starting with the second half of the steering wheel. The sponges were treated the same way as above. Polyurethane foam (PUF) moistened with 6 ml of isopropanol was used in the same manner to sample for glyphosate and 2,4-D.

Air samples were collected for 24 hours on OSHA Versatile Sampler (OVS-2) sorbent tubes (SKC, Eighty Four, PA) containing XAD-2 resin with an 11 mm quartz fiber pre-filter and polyurethane foam (PUF). The nominal flow rate for the sampling pump was 1 Lpm. Pumps were pre- and post-calibrated each sampling day with the OVS-2 media in line using a Bios DriCal DC-Lite[®].

Approximately one field blank sample for every 20 air and wipe samples was submitted for analysis along with the field samples. The blank samples were handled in the same manner as the field samples. All blank samples were below the analytical LOD for all pesticides tested.

All samples were transported from the field in a cooler and transferred to a refrigerator where they were stored for a few days until shipment to the laboratory. At the laboratory, samples were stored in a freezer from three to six months until analysis.

Sample Analysis

Air and wipe samples were analyzed by DataChem Laboratories (Salt Lake, UT) and the dust samples were analyzed by Battelle Memorial Institute (Columbus, OH). The limits of detection and recovery efficiencies are reported in Table 1.

Air

The OVS-2 tubes were separated into two sections with the front ring, filter and front resin section placed in one 4 ml vial, and the middle PUF separator and back resin section placed in another 4 ml vial. Each vial was desorbed with 2 ml of diazomethane desorption solution.

Wipe

The Sof-Wick sponges were desorbed in their shipping containers with 40 ml of isopropanol, after which an aliquot of each sample was poured into a GC vial for analysis. Liquid standards were used for quantitation. The PUF sponges were desorbed in their shipping containers with 75 ml of methanol. All air and the Sof-Wick sponge wipe samples were analyzed using a gas chromatograph equipped with an electron capture detector using a 30m DB-1701 column programmed from 130-270 °C. The

PUF samples were analyzed using a gas chromatograph equipped with an electron capture detector using a 30m DB-608 column programmed from 90-270 °C.

Dust

Dust analyses for acetochlor, atrazine, metolachlor, alachlor, chlorpyrifos (non-acidic or neutral pesticides) included extraction of 0.5 g aliquots of the dust with 12 mL of 1:1 (v:v) hexane:acetone using sonication. The extract was cleaned up using sequential elution on a silica SPE cartridge. Samples were analyzed using GC/MS in the multiple ion detection mode with a 30 m DB-1701 column programmed from 160-280 °C.

Analyses for 2,4-D (acidic pesticide) included sonication extraction of 1 g of dust with 25 mL of a 70:30 (v:v) mix of acetonitrile and 0.1 M sodium acid phosphate buffer at pH=3 (19). The extract was further cleaned prior to analysis using a C18 SPE cartridge. The extract was derivatized using diazomethane. The extracts were analyzed as described above, using a 30 m RTx-5 ms column programmed from 180-280 °C.

Table 1. Pesticide detection limits by sample type

Pesticide	Air (ng/sample)		Surface wipe (ng/sample)		Dust (ng/g) ^A	
	LOD	% Recovery	LOD	% Recovery	LOD	% Recovery
Atrazine	200	90 – 100	4000	100	1.5	98
Metolachlor	10	90 – 100	100	99	0.7	86
Chlorpyrifos	10	90 – 100	80	103	1.5	83
Acetochlor	10	90 – 100	100	100	1.5	80
Alachlor	10	90 – 100	40	100	1.5	82
Glyphosate	n/a	n/a	400	90 - 100	0.7	91
2,4-D	200	90 – 100	700	90 - 100	0.7	104

Abbreviations: n/a = not applicable

^A Based on an extracted sample of 0.5 g of dust for atrazine, metolachlor, chlorpyrifos, acetochlor, alachlor, and glyphosate and 1.0 g of dust for 2,4-D.

Analyses for glyphosate (acidic pesticide) included sonication extraction of 0.5 g of dust with 12 mL of deionized water following addition of isotopically-labelled glyphosate and acidification of the dust with concentrated H₃PO₄. The extract was partitioned with neutral solvents for cleanup, then evaporated to a small volume under vacuum and then lyophilized overnight. The residue was derivatized with a 2:1 mixture of trifluoroacetic acid and trifluoroethanol, and then extracted into dichloromethane. Samples with chromatographic interferences to glyphosate were further cleaned up using sequential elution on a silica SPE cartridge. Samples were analyzed using GC/MS in the multiple ion detection mode with a 30 m RTx-5 ms column programmed from 120-180 °C.

Data Analysis

Pesticide levels were reported in ng/sample for air and surface wipe samples and in ng/g for dust samples by the analyzing laboratories and were not corrected for recovery efficiency. The analytical limit of detection (LOD) varied by pesticide and sample type (Table 1). The LODs for the dust samples were based on an extracted sample of 0.5 g of dust for the neutral pesticide and glyphosate analyses and 1 g of dust for the 2,4-D analysis. Eighty percent of the dust samples provided at least these amounts. The rest of the samples contained less than 0.5 g of dust. These low mass samples were analyzed in all instances with the exception of one dust sample which only produced sufficient dust for the 2,4-D analysis. However, the LODs would be higher for these samples due to the smaller amounts of dust. Therefore, LODs for samples less than 0.25 g were adjusted proportionally: the LOD for a sample with mass < 0.0625 g was adjusted by a factor of 10, a sample with mass < 0.125 g by a factor of 4, and a sample with mass < 0.25 g by a factor of 2, unless all other full mass samples from the household were also non-detects.

The number of air samples with detectable levels of pesticide was small, therefore additional analyses were not performed on these samples. The percent of wipe and dust samples above the LOD were computed separately for farm and non-farm samples.

Rates of detection were compared between farm and non-farm samples using generalized estimating equations (GEE) methods to account for the correlated nature of samples taken within the same household. The GENMOD procedure in SAS, which fits generalized linear models, was used to compute the odds ratio for detecting a positive sample for farm homes versus non-farm homes. Models specified a logit link function, an exchangeable correlation matrix, and household as a repeated effect.

For surface wipe samples, pesticide levels reported in ng/sample were standardized to ng/cm² using the area associated with each sample. Since less than half of the wipe samples had analytes present above the LOD, only the range of the detectable samples was reported. For dust samples, pesticide levels reported as 'below the LOD' were replaced with one-half of the LOD (20) prior to analysis. Pesticide levels in dust reported as ng/g were standardized to ng/cm² using the total mass (in grams) and area associated with each sample. Both the distributions of the pesticide concentration in dust (ng/g) and the pesticide concentration in carpet (ng/cm²) were highly skewed to the right, therefore a natural log transformation was applied to these concentrations prior to statistical analysis. The geometric mean (GM) and geometric standard deviation (GSD) were only reported when at least 50 percent of the samples overall were above the LOD.

Since each household was sampled on two visits and more than one dust sample was obtained at each visit, resulting in correlated dust samples, mixed-effects models were used to determine statistical significance. In these models, household was treated as a

random effect and group (farm, non-farm), visit (visit 1, visit 2), and room (child's bedroom, child's play room, laundry room, father's change room, and entrance way) were treated as fixed effects. For farm households, crop spray records were used to determine whether the pesticide was sprayed in the 7 days preceding the visit (yes, no). The seven day cutoff was intended to focus on more recent pesticide applications, rather than applications that occurred more than one week prior to the visit. Household covariates (Table 2) included the age of the home (< 60 , ≥ 60 years), home/lawn/garden sprayed with pesticide in the last month/year (yes, no), the age of the carpets (< 8 , ≥ 8 years), frequency of carpet vacuuming ($< \text{once per week}$, $\geq \text{once per week}$), own a dog (yes, no), own a cat (yes, no), presence of doormats (yes, no), and proximity to farm fields (< 0.5 , ≥ 0.5 mile). Cutpoints for the age of the home and carpets and frequency of vacuuming were selected to approximately divide the households equally. Crop demographics, pesticide use and application practices, use of PPE, and children's farm activities were presented previously (17). Additional covariates were tested one at a time, after adjusting for group, spray status, visit and room. All mixed models were fit using the MIXED procedure in SAS assuming a compound symmetric covariance structure. Model residuals were assessed for departures from normality. For dust samples obtained from farm homes, estimates of within- and between-household variability, after adjusting for spray status, visit, and room were computed using the MIXED procedure in SAS assuming a compound symmetric covariance structure.

All significance testing was performed at the 0.05 level of significance. When comparing geometric means for more than two categories, p-values were adjusted using the Tukey-Kramer adjustment for multiple comparisons. All statistical analyses were performed using SAS system software, version 8.2 (SAS Institute, Inc., Cary, NC).

Table 2: Household covariates

Variable	Household type	
	Non-farm (n = 25)	Farm (n = 25)
Age of home (years)		
Median (range)	30 (1 – 111)	84 (<1 – 139)
Percent of homes sprayed with insecticides in the		
last month	12%	40%
last year	28%	52%
Percent of lawns treated with pesticides in the		
last month	28%	12%
last year	38%	32%
Percent of gardens sprayed with pesticides in the ^A		
last month	17%	14%
last year	33%	43%
Age of carpet (years)		
Median (range)	6 (1 – 40)	10 (<1 – 40)
Vacuum < 1 time per week, %	24%	21%
Own a dog, %	40%	76%
Own a cat, %	48%	68%
Have doormats, %	68%	80%
Percent of homes < 0.5 miles from farm fields	44%	100%

^A Limited to 12 non-farm and 21 farm homes that reported having a garden.

Results

Air samples

A total of 99 indoor and 98 outdoor air samples were obtained and analyzed for atrazine, metolachlor, chlorpyrifos, acetochlor, alachlor, and 2,4-D. Eighty-nine percent of the indoor air and 99% of the outdoor air samples were below the LOD for the six pesticides tested. Chlorpyrifos was detected in indoor air samples taken from six farm (range: 0.04 - 0.23 $\mu\text{g}/\text{m}^3$) and two non-farm homes (range: 0.01 – 0.05 $\mu\text{g}/\text{m}^3$) and acetochlor was detected in one indoor air sample taken from a farm home (0.04 $\mu\text{g}/\text{m}^3$). All indoor air samples were below the LOD for atrazine, metolachlor, alachlor and 2,4-D. None of the homes had detectable levels in any of the air samples taken outside the home, except for one farm home which had a single sample positive for metolachlor (0.1 $\mu\text{g}/\text{m}^3$).

Wipe samples

A total of 203 house and 153 vehicle wipe samples were obtained for the neutral pesticide analysis and 82 house and 48 vehicle wipe samples were obtained for the glyphosate/2,4-D analysis (Table 3). A majority of these samples were below the LOD for the pesticides tested. For house wipe samples, atrazine was detected in only a single non-farm sample, metolachlor in only 4 farm samples, and acetochlor in only 7 farm samples. All house wipe samples were below the LOD for alachlor, glyphosate, and 2,4-D. Chlorpyrifos was the most commonly detected pesticide in both house (F 23% versus NF 22%) and vehicle wipe samples (F 21% versus NF 7.9%, odds ratio (OR) = 3.1, 95% confidence interval (CI) = 1.04 – 9.1). Acetochlor was detected more often in farm vehicle wipe samples (F 13% versus NF 5.3%) and metolachlor was detected significantly more often in farm vehicle wipe samples (F 12% versus NF 1.3%, OR = 9.8, 95% CI = 1.1-87). Atrazine and alachlor were rarely detected and glyphosate and 2,4-D were never detected in vehicle wipe samples.

Dust samples

A total of 295 dust samples (sample mass: range = 0.01 – 204 g, median = 2.8 g) were obtained from carpet (sample area: range = 0.3 – 7.4 m², median = 1.1 m²) inside the homes. After adjusting for visit and room, farm homes had a significantly higher geometric mean carpet dust loading than non-farm homes (F 2.7 versus NF 1.5 g/m², p-value = 0.026). The unadjusted geometric mean concentration (ng/g) of each pesticide sampled in dust was higher in farm homes compared to non-farm homes, although only significantly for atrazine and metolachlor. The difference, however, becomes even more apparent when standardizing for area sampled (ng/cm²), due to the fact that farm homes had more dust than non-farm homes.

Table 3. Percent and range of wipe and dust samples greater than or equal to the limit of detection.^A

Sample type Pesticide	Non-farm			Farm			Odds ratio ^C	
	n	n > LOD (%)	Range ^B	n	n > LOD (%)	Range ^B	e ^β	95% CI
<i>House wipe</i>								
Atrazine	95	1 (1.1%)	160	108	0 (0%)	---	---	---
Metolachlor	95	0 (0%)	---	108	4 (3.7%)	0.85-8.5	---	---
Chlorpyrifos	95	21 (22%)	0.22-3.8	108	25 (23%)	0.32-25	1.1	0.44-2.8
Acetochlor	95	0 (0%)	---	108	7 (6.5%)	0.32-2.5	---	---
Alachlor	95	0 (0%)	---	108	0 (0%)	---	---	---
Glyphosate	39	0 (0%)	---	43	0 (0%)	---	---	---
2,4-D	39	0 (0%)	---	43	0 (0%)	---	---	---
<i>Vehicle wipe</i>								
Atrazine	76	0 (0%)	---	77	3 (3.9%)	38-410	---	---
Metolachlor	76	1 (1.3%)	5.2	77	9 (12%)	9.8-680	9.8	1.1-87
Chlorpyrifos	76	6 (7.9%)	0.43-11	77	16 (21%)	0.23-9.3	3.1	1.04-9.1
Acetochlor	76	4 (5.3%)	1.3-6.2	77	10 (13%)	0.79-39	2.7	0.68-11
Alachlor	76	1 (1.3%)	3.3	77	2 (2.6%)	1.2-1.3	2.0	0.19-21
Glyphosate	26	0 (0%)	---	22	0 (0%)	---	---	---
2,4-D	26	0 (0%)	---	22	0 (0%)	---	---	---
<i>Dust</i>								
Atrazine	114	30 (26%)	0.0017-0.077	116	91 (78%)	0.00039-17	9.4	3.5-25
Metolachlor	114	59 (52%)	0.00073-1.3	116	80 (69%)	0.0011-9.8	2.1	1.1-3.9
Chlorpyrifos	114	92 (81%)	0.00021-3.6	116	97 (84%)	0.00049-10	1.2	0.37-3.6
Acetochlor	114	17 (15%)	0.00054-1.4	116	34 (29%)	0.00086-2.6	2.1	0.84-5.5
Alachlor	114	5 (4.4%)	0.00027-0.012	116	12 (10%)	0.00085-0.046	2.3	0.51-11
Glyphosate	33	28 (85%)	0.0012-13	31	31 (100%)	0.0081-2.7	---	---
2,4-D	33	33 (100%)	0.0041-1.9	32	32 (100%)	0.00099-5.3	---	---

Abbreviations: n = number of samples; LOD = limit of detection; CI = confidence interval

^A Samples from 20 farm and 19 non-farm homes were analyzed for atrazine, metolachlor, chlorpyrifos, acetochlor, and alachlor. Samples from 5 farm and 6 non-farm homes were analyzed for glyphosate and 2,4-D.

^B Range of samples greater than or equal to the LOD (ng/cm²), reported to two significant figures. The ng/cm² value for dust was calculated by multiplying the ng/g value reported by the laboratory by the amount (g) of dust collected per cm² of carpet sampled.

^C e^β = the odds ratio, defined as the odds of a farm sample being above the LOD divided by the odds of a non-farm sample being above the LOD, obtained from the GENMOD procedure in SAS assuming an exchangeable correlation matrix.

A total of 230 dust samples from 20 farm and 19 non-farm homes were analyzed for atrazine, metolachlor, chlorpyrifos, acetochlor, and alachlor (Table 3). Compared to the wipe samples, dust samples were more likely to detect pesticide residues with a majority

of the dust samples above the LOD for atrazine, metolachlor, chlorpyrifos, glyphosate and 2,4-D. A pesticide residue was detected significantly more often in dust samples from farm homes compared to non-farm homes for atrazine (OR = 9.4, 95% CI = 3.5 – 25) and metolachlor (OR = 2.1, 95% CI = 1.1 – 3.9). Detection rates were similar between farm homes and non-farm homes for chlorpyrifos in dust samples. A total of 65 dust samples from 5 farm and 6 non-farm homes were analyzed for glyphosate and 2,4-D (Table 3). Glyphosate was detected marginally more often in dust samples from farm homes while 2,4-D was detected in every dust sample.

Dust samples were categorized as belonging to a non-farm home, a farm home that did not apply the pesticide in the 7 days preceding the visit, and a farm home that applied the pesticide in the 7 days preceding the visit. Acetochlor and alachlor were excluded from this analysis since greater than 50% of the dust samples for these pesticides were below the LOD. Geometric means, after adjusting for visit and room, for each of these groups are presented in Table 4. Atrazine and metolachlor were significantly higher in dust from farm homes that reported applying these pesticides in the 7 days preceding the visit compared to farm homes that did not apply these pesticides and non-farm homes. In addition, the concentration of atrazine in dust was significantly higher in farm homes that did not apply atrazine compared to non-farm homes. Chlorpyrifos and glyphosate were higher, but not significantly, in dust from farm homes that applied these pesticides in the 7 days preceding the visit compared to farm homes that did not apply them and non-farm homes. However, there were only two farms that reported having sprayed chlorpyrifos prior to a visit, and one non-farm, located within a ¼ mile of a farm and in close proximity to a field, had unusually high levels of glyphosate in dust (n=7, GM = 2100 ng/g). If this non-farm is excluded, then farm homes that sprayed glyphosate

within 7 days preceding the visit had significantly greater concentrations of glyphosate in dust than non-farm homes. The spray effect analysis for 2,4-D included dust samples when 2,4-D was applied to crops in the 30 days preceding the farm-visit since there were only 2 farm-visits where 2,4-D was applied to crops in the 7 days preceding the visit. 2,4-D levels were similar in dust from farm homes that sprayed 2,4-D in the 30 days preceding the visit compared to farm homes that did not spray 2,4-D. 2,4-D was higher, but not significantly, in farm homes compared to non-farm homes. Acetochlor was only applied to crops at 5 farms prior to visits; alachlor was not applied to crops at any of the farms, and since less than 50% of the dust samples were above the analytical LOD for both acetochlor and alachlor, additional analyses were not performed for these pesticides.

The distributions of five of the pesticides in the homes are shown in Table 5. In general, for atrazine and metolachlor, the entrance way, father's change area and laundry room had the highest levels of pesticide in dust for farm homes that sprayed these pesticides within the 7 days preceding sampling, whereas in non-farms, the entrance way and child's bedroom had the highest pesticide levels in dust. Chlorpyrifos levels in dust were similar in all rooms but highest in the child's bedroom for both farm and non-farm households. A room effect could not be assessed in farm homes that sprayed chlorpyrifos in the 7 days preceding the visit due to sample size limitations. Glyphosate levels in dust were highest in the child's bedroom for both farm and non-farm homes, while 2,4-D concentrations in dust were highest in the entrance way for non-farms and highest in the change area for farms.

Table 4. Dust sample results from the spray effect analysis. ^A

Pesticide Spray category	n	% > LOD	Pesticide residue in dust (ng/g)				Pesticide residue in carpet (ng/cm ²)				
			GM	GSD	Adjusted GM ^B	95% CI	GM	GSD	Adjusted	GM	95% CI
Atrazine											
Non-farm						1.1 –					0.00018 –
Farm – no spray	114	26	2.3	6.0	2.4 ^{C,D}	5.1	0.00035	12	0.00035 ^{F,G}		0.00068
Farm – spray	58	64	16	11	26 ^{C,E}	11 –	0.0042	16	0.0055 ^{F,H}		0.0025 –
within 7 days	58	93	170	11	94 ^{D,E}	59	0.048	12	0.039 ^{G,H}		0.012
						41 –					0.017 –
						220					0.087
Metolachlor											
Non-farm						3.2 –					0.00043 –
Farm – no spray	114	52	5.7	14	5.9 ^I	11	0.00088	24	0.00089 ^K		0.0018
Farm – spray	95	64	9.9	13	12 ^J	6.0 –	0.0032	23	0.0037 ^L		0.0017 –
within 7 days	21	90	310	20	240 ^{I,J}	22	0.042	32	0.043 ^M		0.0078
						69 –					0.0099 –
						840					0.19
Chlorpyrifos											
Non-farm						14 –					0.0021 –
Farm – no spray	114	81	30	8.5	33	80	0.0046	14	0.0050		0.012
Farm – spray	111	83	39	11	41	17 –	0.011	13	0.012		0.0051 –
within 7 days	5	100	73	2.1	106	96	0.011	4.7	0.021		0.029
						17 –					0.0018 –
						680					0.23
Glyphosate											
Non-farm						21 –					0.0022 –
Farm – no spray	33	85	140	2.1	110	610	0.023	34	0.018		0.15
Farm – spray	18	100	920	2.1	1000	140 –	0.22	2.9	0.28		0.024 –
within 7 days	13	100	1100	2.4	1300	7400	0.33	5.0	0.45		3.2
						180 –					0.039 –
						9700					5.3
2,4-D											
Non-farm						100 –					0.013 –
Farm – no spray	33	100	330	3.3	320	1000	0.056	4.9	0.053		0.22
Farm – spray	20	100	340	2.7	850	240 –	0.082	3.7	0.20		0.038 –
within 30 days	12	100	1700	4.0	730	3100	0.41	14	0.25		1.0
						190 –					0.045 –
						2800					1.4

Abbreviations: n = number of samples; LOD = limit of detection; GM = geometric mean; GSD = geometric standard deviation; CI = confidence interval

^A All results reported to two significant figures. Samples reported as below the LOD were assigned ½ LOD prior to statistical analysis.

^B Adjusted for visit (visit 1, visit 2) and room (child's bedroom, child's play room, laundry room, father's change room, and entrance way).

^{C, D, F–J, L} Tukey-Kramer adjusted p-value < 0.001; ^{E, M} Tukey-Kramer adjusted p-value < 0.01; ^K Tukey-Kramer adjusted p-value < 0.05. Means with the same letter are significantly different.

After adjusting for visit, room, and spray status, none of the pesticides were related to any of the household covariates except for atrazine. The concentration of atrazine in dust was significantly higher in farm homes only that reported using an insecticide inside the home within the year prior to sampling after adjusting for visit, room, and spray status. For dust samples from farm households, the effects of agricultural practices on pesticide concentration were also evaluated. However, due to small sample sizes for each pesticide and a lack of variability among some practices, only a limited analysis of atrazine was performed. Atrazine, applied to crops at 16 out of 20 farms in the neutral pesticide analysis, was applied by the farmer at 10 farms and by a custom applicator at 6 farms. Higher atrazine levels in household dust were observed at farm-visits where atrazine was applied by the farmer compared to farm-visits where atrazine was applied by a custom applicator (adjusted GM 370 versus 27 ng/g, p-value = 0.0013). Farmers self-applying atrazine reported similar spray practices, so it was not possible to perform tests of significance for many of these variables. Higher atrazine levels in household dust were marginally associated with the use of a closed cab, however, after adjusting for the number of acres sprayed, the difference was not significant (adjusted GM closed cab 610 versus open cab 290 ng/g, p-value = 0.46).

For each dust sample, the pesticide concentration in dust (ng/g) was converted to a pesticide loading on the carpet (ng/cm²) using the mass and area associated with each sample. The effect of spraying in this analysis was similar to the spray effect in the pesticide concentration in dust analysis (Table 4). The rooms were not equally dusty, however, with the entrance way having the highest amount of dust per unit area compared to the other rooms. Consequently, pesticide loadings on the carpet in the entrance way tend to be higher than loadings from the other rooms.

Table 5. Dust sample results from the room effect analysis.

Pesticide Group	Number of samples	Room in house (Adjusted GM ^A , ng/g)				
		Child's bedroom	Child's playroom	Laundry room	Father's change area	Entrance way
Atrazine						
Non-farm	114	2.7	2.0	1.3 ^B	0.85 ^C	4.5 ^{B, C}
Farm – no spray	58	8.9 ^D	15	24	76 ^D	35
Farm – spray within 7 days	58	100 ^E	140	530	740 ^E	340
Metolachlor						
Non-farm	114	41 ^{F, G, H}	1.8 ^{F, I}	0.50 ^{G, J}	0.40 ^{H, K}	15 ^{I, J, K}
Farm – no spray	95	30 ^{L, M}	6.5	1.4 ^{L, N}	3.0 ^M	23 ^N
Farm – spray within 7 days	21	55	9.2	1200	1400	350
Chlorpyrifos						
Non-farm	114	52 ^O	32	33	12 ^O	31
Farm	116	77 ^{P, Q}	22 ^P	39	56	22 ^Q
Glyphosate						
Non-farm	33	510	8.6	260	60	260
Farm	31	1500 ^R	1400 ^S	n/a	1400	550 ^{R, S}
2,4-D						
Non-farm	33	450	120 ^T	83	270 ^U	740 ^{T, U}
Farm	32	660	610	1300	1600	850

Abbreviations: GM = geometric mean; n/a = not available

^A All results reported to two significant figures. Samples reported as below the LOD were assigned ½ LOD prior to statistical analysis. Geometric mean is adjusted for visit (visit 1, visit 2). Significance testing was performed within each group.

^{B-E, M, N, P, S-U} Tukey-Kramer adjusted p-value < 0.05; ^{F-H, J, K} Tukey-Kramer adjusted p-value < 0.0001; ^{I, L, O, Q, R} Tukey-Kramer adjusted p-value < 0.01. Means with the same letter are significantly different.

For example, in farm homes that sprayed atrazine in the 7 days preceding the visit, atrazine levels in carpet dust (ng/g) were higher, although not significantly, in the father's change room compared to the entrance way (least squares geometric mean (LSGM) = 740 versus 340 ng/g, respectively). However, after standardizing to unit area

(ng/cm²), atrazine loadings on the carpet were higher at the entrance way compared to the father's change area (LSGM = 0.59 versus 0.16 ng/cm², respectively).

The within-household (GSD_w) and between-household (GSD_b) variance components expressed as geometric standard deviations for five of the pesticides are provided in Table 6 for both pesticide concentration in dust (ng/g) and pesticide concentration in carpet (ng/cm²) for dust samples from farm households. Variance components, computed after adjusting for visit, room, and spray status, were not markedly changed by the addition of other exposure determinants to the model.

Discussion

While there are a few studies that have investigated the take-home pesticide issue and pesticide home contamination in rural and agricultural environments, most of these studies have examined organophosphate pesticides, while this study looked at several pesticides not generally measured in these previous studies. Chlorpyrifos has been studied frequently and can serve as a benchmark.

In a study conducted by the Centers for Disease Control and Prevention (CDC) for the Arizona Department of Health Services, dust was collected from 152 homes and 25 schools and tested for the presence of 43 pesticides (21). Chlorpyrifos had a geometric mean (GM) of 113 ng/g. Curl et al (22) collected 156 house dust samples from farm worker households and found a GM level of 50 ng/g for chlorpyrifos. This compares

with chlorpyrifos concentrations of 40 and 30 ng/g for farm and non-farm homes respectively in our study.

Table 6. Within- and between-household variance components^A for pesticide levels in dust samples obtained from farm households.

Sample type Pesticide	Number of farms	Number of samples	Within-household		Between-household	
			GSD _w	%	GSD _b	%
Dust (ng/g)						
Atrazine	20	116	4.6	45	5.4	55
Metolachlor	20	116	10.4	81	3.1	19
Chlorpyrifos	20	116	3.8	33	6.5	67
Glyphosate	5	31	1.8	77	1.4	23
2,4-D	5	32	2.1	22	4.2	78
Dust (ng/cm ²)						
Atrazine	20	116	7.2	84	2.4	16
Metolachlor	20	116	16.2	86	3.1	14
Chlorpyrifos	20	116	4.7	37	7.5	63
Glyphosate	5	31	3.2	67	2.2	33
2,4-D	5	32	3.7	45	4.2	55

Abbreviations: GSD_w = estimated geometric standard deviation of the within-household distribution; GSD_b = estimated geometric standard deviation of the between-household distribution; % is the percent of the random effect variance attributable to that source.

^A Variance components were computed using the MIXED procedure in SAS after adjusting for visit, room, and spray status.

Farm homes in this study are clearly more contaminated than non-farm homes. Other studies have found similar results. Simcox et al (13) measured pesticide levels in house dust of both farm homes and reference homes and found that farm homes had significantly higher levels of pesticide in dust. Bradman et al (12) found a higher pesticide levels in dust between farm worker homes and non-farm worker homes.

Differences in pesticide levels in dust seen between non-farm homes, farm homes that did not spray the pesticide, and farm homes that did spray the pesticide were greater for the strictly agricultural pesticides (e.g. atrazine and metolachlor) versus pesticides that have both residential and agricultural uses (chlorpyrifos, glyphosate and 2,4-D). This would be expected since these latter pesticides are commonly used in residential settings. Chlorpyrifos, glyphosate, and 2,4-D appear to be ubiquitous in both the non-farm and farm homes. Better than 80% of the dust samples in both farm and non-farm homes had detectable levels of these pesticides. This finding is comparable to other literature reports. Chlorpyrifos, for example, was detected in 81% of dust samples in Yuma County, Arizona (21) and in 98% and 82% of dust samples from agricultural and non-agricultural families respectively (13). It is interesting to note that chlorpyrifos was one of the most frequently detected pesticides in the current study but was applied at only 2 farms (17).

One potential source of pesticides in farm homes results from farmer take-home mechanism. When atrazine or metolachlor was applied to crops on the farm, concentrations of these pesticides tended to be higher in dust in the entrance way, laundry room, and change room – rooms where dirt would be tracked in, or the farmer's clothes would be deposited. These pesticides have agricultural uses only, and therefore would not be used in or around the house. Chlorpyrifos, glyphosate and 2,4-D all have residential uses so that contamination may have multiple sources. This is supported by our finding that both farm homes and non farm homes have a high percent of detectable samples for these pesticides and their more even distribution throughout the homes. The higher levels of atrazine and metolachlor in the entrance way, laundry room and

change room would suggest that the farmer is bringing them home on clothing and shoes, supporting the notion of take-home pesticide exposure. Other studies have suggested the take-home pathway as the primary mechanism for contamination of the indoor environment (12, 13, 22-24). Acetochlor, which was sprayed by only a few farmers, and alachlor which was not sprayed at all, were not detected frequently enough to allow this analysis.

Spray drift as a source of pesticide residue cannot be ruled out. Even though both the indoor and outdoor air samples were largely non-detectable, they were taken a few days after an application, by which time pesticide in the air may have settled out. Koch et al (25) found that OP metabolite levels in children's urine samples were higher during the spray season in an agricultural region in the absence of parental work contact or residential proximity to treated fields. The authors suggest that spray drift may account for some of the observed increases.

Several factors that were anticipated to be associated with pesticide levels in dust were investigated for their affect on the pesticide levels. Only the use of an insecticide inside a farm home was found to be associated with atrazine in dust. Since atrazine is an herbicide, this association does not make sense, and may be spurious since farm homes that sprayed with an insecticide in the last year were also more likely to have sprayed atrazine prior to both visits. Age of carpet, frequency of carpet vacuuming, the presence of door mats, the age of the homes, and the presence of pets were not associated with pesticide levels in dust. It is unclear why none of these variables were associated, however, testing for an association between the pesticide level and some of the

household covariates was complicated due to confounding with farm/non-farm status or confounding with spray status (for example, all farms that sprayed the pesticide may have had 'old' carpet). In a simulated pesticide track-in study by Nishioka et al (19), 2,4-D levels in dust were lower when a door mat was present. In another study, the presence of a high activity dog was shown to be significantly correlated with 2,4-D levels in indoor house dust (16). The authors warn though that the sample sizes were small and caution should be exercised in interpreting the results. One reason why the door mats may not have reduced pesticide levels in dust in our study is that they may have acquired a high pesticide and dust loading after only a short time of use, becoming a reservoir for contamination as opposed to an element for reducing contamination. This speculation should be investigated further. In the case of dog activity, dogs on the farms were outdoor dogs only. In only one case in the farm homes with dogs, did the dog spend time both indoors and outdoors.

Distance to a treated field did not correlate with pesticide levels in dust in non-farm homes. Distance to a treated field was not analyzed for the farm homes, since all farm homes were reported to be within $\frac{1}{4}$ mile of a treated field. It may be that the distance categories ($< \frac{1}{4}$ mile, $\frac{1}{4}$ mile to $\frac{1}{2}$ mile, $\frac{1}{2}$ mile to $\frac{3}{4}$ mile, $\frac{3}{4}$ mile to 1 mile, > 1 mile) may not have permitted detection of differences. Simcox et al (13) found that pesticide levels in dust decreased with increasing distance, but considered much smaller distances (< 50 ft, 50-200 ft, > 200 ft). In Yuma County Arizona, however, although an association between pesticide levels in dust and proximity to treated fields was not investigated, the authors did investigate urine metabolite levels and proximity to a treated field and did not find an association (21).

Most of the analyses in this paper focused on the pesticide concentrations in dust. The wipe and air samples were not a particularly useful sample media for evaluation of low-level pesticides in homes in this study. This may largely be the result of higher LODs for the wipe and air samples in this study. Indeed, the dust sample values are orders of magnitude lower than the wipe values which are likely due to the better limits of detection for the dust analysis. Another factor may be the sampling method, particularly the use of polyurethane foam (PUF) for the wipe samples for 2,4-D and glyphosate. Of the dust samples analyzed for 2,4-D and glyphosate, 100% and 94% had detectable levels of 2,4-D and glyphosate, respectively. These pesticides were not detected in any wipe sample. One apparent problem with using PUF for wipe sampling is that PUF does not hold liquid very well. The amount of isopropanol added to the PUF had to be reduced to six ml from the 10 ml added to the Sof-Wick sponges. Even so, the isopropanol would run off the PUF, leaving the PUF fairly dry when wiping. Because of this, it is likely that not as much pesticide residue would be picked up from the surface. Further investigation is needed to confirm this assumption.

There are a few limitations to the analyses. Chlorpyrifos was not sprayed very often, so it is difficult to draw conclusions about the spray effect for chlorpyrifos. In the glyphosate/2,4-D analysis, there were only 5 farms and 6 non-farms available for the analysis. As a result of the small number of homes, the differences seen were not statistically significant. Only the acid form of 2,4-D was analyzed in the samples. In some farm-homes the ester form of 2,4-D may have been applied resulting in an underestimate of 2,4-D contamination. Testing some of the household covariates for a

relationship with pesticides was difficult due to confounding with farm/non-farm status or confounding with spray status. The LODs for the wipe and air samples are substantially higher than the LODs for the dust samples, making it difficult to compare the sample media. Lastly, for the dust data analysis, we considered the effect of spraying in the seven day period preceding the visit. The choice of seven days, although somewhat arbitrary, was intended to focus on more recent pesticide applications.

Conclusion

Farm homes have more pesticide residue inside than non-farm homes and farms that spray a particular pesticide are more likely to have higher levels of that pesticide inside the home than other homes. This is particularly apparent for the strictly agricultural herbicides atrazine and metolachlor. While these herbicides are not applied in or around the home, they appear to be getting into the home para-occupationally. It appears for agricultural pesticides that take-home exposure may be an important source of home contamination.

Acknowledgements

The authors would like to thank the many people who helped with this study, in particular Donald Booher for his technical assistance, Fran Guerra for her administrative assistance and Beth Whelan for her guidance. We would also like to thank Ann Stromquist, Jill Moore, and Matt Nonnenman, of the Great Plains Center for Agricultural Health, University of Iowa, and Charles Lynch and Patricia Gillette of the Agricultural Health Study, University of Iowa for their help in recruiting participants.

We would like to thank Marty Jones and Craig Taylor, of the Department of Occupational and Environmental Health, School of Public Health, University of Iowa, for their technical assistance. Finally, we would like to thank DataChem Laboratories for providing the air and wipe sampling analysis.

References

1. EPA. Pesticide industry sales and usage: 1998 and 1999 market estimates. Biological and Economic Analysis Division, Office of Pesticide Programs, Office of Prevention, Pesticides and Toxic Substances, U.S. Environmental Protection Agency, Washington, DC. 2002.
2. Reynolds, SJ, Merchant JA, Stromquist AM, Burmeister LF, Taylor C, Lewis MQ, and Kelly KM. (1998) Keokuk County Iowa rural health study: self reported use of pesticides and protective equipment. *J. Agricul. Safety Health* 1998; 1:67-77
3. Alavanja MC, Sandler DP, McMaster SB, Hoar-Zahm S, McDonnell CJ, Lynch CF, Pennybacker M, Rothman N, Dosemeci M, Bond AE, and Blair A. The agricultural health study. *Environ. Health Persp.* 1996; 104(4):362-369
4. Mandel JH, Carr WP, Hilmer T, Leonard PR, Halberg JU, Sanderson WT, and Mandel JS. Factors associated with safe use of agricultural pesticides in Minnesota. *J. Rural Health* 1996; 12(4):301-310
5. Lewis RG, Fortune CR, Blanchard FT, and Camann DE. Movement and deposition of two organophosphorus pesticides within a residence after interior and exterior applications. *J Air Waste Manage Assoc* 2001; 51:339-351
6. Ma X, Buffler PA, Gunier RB, Dahl G, Smith MT, Reinier K, et al. Critical windows of exposure to household pesticides and risk of childhood leukemia. *Environ. Health Persp* 2002; 110(9):955-960
7. Zahm SH, Ward MH. Pesticides and childhood cancer. *Environ. Health Persp.* 1998; 106(suppl 3):893-908
8. Daniels JL, Olshan AF, Savitz DA. Pesticides and childhood cancers. *Environ. Health Persp* 1997; 105:1068-1077
9. Garry VF. Pesticides and children. *Toxicol Applied Pharmacol* 2004; 198:152-163
10. Guyton AC. Textbook of Medical Physiology, 7th ed. W.B. Saunders Company, Philadelphia, PA. 1986

11. Fenske RA, Kissel JC, Lu C, Kalman DA, Simcox NJ, Allen EH, et al. Biologically based pesticide dose estimates for children in an agricultural community. *Environ Health Persp* 2000b; 108(6):515-520
12. Bradman MA, Harnly ME, Draper W, Seidel S, Teran S, Wakeham D, Neutra R. Pesticide exposure to children from California's Central Valley: Results of a pilot study. *J. Exp. Anal. Environ. Epidem* 1997; 7(2):217-234
13. Simcox NJ, Fenske RA, Wolz SA, Lee I-C, Kalman DA. Pesticides in household dust and soil: Exposure pathways for children of agricultural families. *Environ. Health. Persp.* 1995; 103(12):1126-1134
14. Loewenherz C, Fenske RA, Simcox NJ, Bellamy G, Kalman D. Biological monitoring of organophosphorus pesticide exposure among children of agricultural workers in central Washington State. *Environ. Health. Persp.* 1997; 105(12):1344-1353
15. Nishioka MG, Lewis RG, Brinkman MC, Burkholder HM, Hines CE, Menkedick JR . Distribution of 2,4-D in air and on surfaces inside residences after lawn applications: comparing exposure estimates from various media for young children. *Environ Health Persp* 2001; 109: 1185-1191.
16. Nishioka MG, Burkholder HM, Brinkman MC, Lewis RG. Distribution of 2,4-dichlorophenoxyacetic acid in floor dust throughout homes following homeowner and commercial lawn applications: Quantitative effects of children, pets and shoes. *Environ. Sci. Technol.* 1999; 33(9):1359-1365
17. Curwin BD, Sanderson WT, Reynolds SJ, Hein MJ, Alavanja MC. Pesticide use and practices in an Iowa farm family pesticide exposure study. *J. Agr. Safety Health* 2002; 8(4):423-433
18. ASTM. Standard practice for the collection of floor dust for chemical analysis. Standard Practice D5438-00. American Society for Testing and Materials, Philadelphia, PA. 2000
19. Nishioka MG, Burkholder HM, Brinkman MC, Gordon SM, Lewis RG. Measuring transport of lawn applied herbicide acids from turf to home: Correlation of dislodgeable 2,4-D turf residues with carpet dust and carpet surface residues. *Environ. Sci. Technol.* 1996; 30(11):3313-3320
20. Hornung RW, and Reed LD. 1990 Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg* 1990; 5: 46 – 51.
21. CDC. Pesticide exposure in children living in agricultural areas along the United States-Mexico boarder, Yuma County, Arizona, US Centers for Disease Control, National Center for Environmental Health, Health Studies Branch, Atlanta, GA 2002
22. Curl CL, Fenske RA, Kissel JC, Shirai JH, Moate TF, Griffith W, et al. Evaluation

of take-home organophosphorus pesticide exposure among agricultural workers and their children. *Environ. Health Perspect* 2002; 110(12):A787-A792

23. McCauley LA, Michaels S, Rothlein J, Muniz J, Lasarev M, Ebbert C. Pesticide exposure and self reported home hygiene. *Amer. Assoc. Occup. Health Nurses J.* 2003; 51(3):113-119

24. Thompson B, Coronado GD, Grossman JE, Puschel K, Solomon CC, Islas I, et al. Pesticide take-home pathway among children of agricultural workers: Study design, methods, and baseline findings. *J. Occup. Environ. Med.* 2003; 45:42-53

25. Koch D, Lu C, Fisker-Anderson J, Jolley L, Fenske RA. Temporal association of children's pesticide exposure and agricultural spraying: Report of a longitudinal biological monitoring study. *Environ Health Persp* 2002; 110(8):829-833