



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 9 Issue: VIII Month of publication: August 2021

DOI: <https://doi.org/10.22214/ijraset.2021.37402>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Pharmacognostic and Nootropic Aspects of *Withania Somnifera*: A Potential Herbal Drug as Memory Enhancer


Minakshee G. Nimbawar¹, Bhushan R. Gudalwar², Wrushali A. Panchale³, Ashish B. Wadekar⁴, Jagdish V. Manwar*⁵, Shubham K. Wadkute⁶, Ravindra L. Bakal⁷

^{1, 3, 4, 6, 7}IBSS's Dr. Rajendra Gode Institute of Pharmacy, Mardi Road, Amravati-444 602, MS, India

^{2, 5}IBSS's Dr. Rajendra Gode College of Pharmacy, Mardi Road, Amravati-444 602, MS, India

Abstract: *Withania somnifera* is a well-known herbal drug commonly used in India and Indian subcontinent. It contains a spectrum of diverse phytoconstituents that are responsible for number of significant biological effects. Various parts of plants especially roots are rich in source of active phytoconstituents. In the current COVID-19 situation, various national and international agencies are suggesting the use of plant for increasing immunity as well as memory booster. The drug is mainly used in the management of brain related disorders such as anxiety, depression, stress, etc. The mainly used as memory enhancer as it increases the neurogenesis process in the brain. In present paper, we have covered the pharmacognosy, phytochemistry, and therapeutic applications of the plant related to the brain.

Keywords: *Withania somnifera*; Memory enhancer; Roots; Stress.



Graphical abstract

I. INTRODUCTION

Despite of the availability of the *anglo-Indian* medicines, popularity of herbal drugs remained unaffected. This might be due to their wider acceptance and long history. Common public in India still using herbal medicines. People across the world were depend upon the products and plants obtained from the nature. In India, even today, people mostly from villages and tribal areas are depends on medicinal plants [1]. Plants or products thereof has become an integral part of our life. In the light of COVID-19 situation, WHO encouraging the people across the world to use medicinal plants. Some of the medicinal plants that are commonly used are *Withania somnifera*, *Hypericum perforatum*, *Aloe barbadensis*, *Oroxylum indicum*, *Capsicum annum*, etc [2-23].

Withania somnifera, also known as Indian Ginseng and Ashwagandha, is one of the most widely used medicinal plant across the world. Indian name Ashwagandha is derived from the sankrit language. In Sanskrit language, ashwagandha means horse's smell probably originated from the odor of its root, which resembles that of sweaty horse. The reason behind its popularity is the plant possesses enormous activities and is enrich source of energy. Almost every therapeutic activity is shown by this plant. In USA and most of the European countries, physicians prescribes tablet formulations or powder of *W. Somnifera* as an adjuvant medicine. The drug is capable of imparting long life, youthful vigor, and good intellectual powers [24].

The drug is clinically used for the treatment of loss of memory, nervous exhaustion, insomnia, consumption, etc. These traditional uses imply that Ashwagandha may possibly be useful at improving neurodegenerative diseases. Indeed, this herbal drug has been reported to exert various pharmacological effects such as anti-neuropsychiatric, Parkinson's disease, anxiety, cognitive and neurological disorders, immunomodulatory, anti-inflammatory, anti-tumor and anti-oxidant effects [25].

The drug also used to inhibit the development of tolerance and dependence on chronic use of various psychotropic drugs. Ashwagandha is commonly available as a churna, a fine powder that can be mixed with water, ghee or honey. In this review, we describe effects of Ashwagandha extracts, phytoconstituents of Ashwagandha (mainly withanolides: steroidal lactones with ergostane skeleton), its derivatives in the context of neurodegenerative diseases and potential therapeutic applications of the plant [26].

II. TAXONOMICAL CLASSIFICATION [27]

Kingdom: Plantae
 Subkingdom: Tracheobionta, Vascular plants;
 Super division: Spermatophyta, Seeds plants;
 Division: Angiosperma
 Class: Dicotyledons
 Order: Tubiflorae
 Family: Solanaceae
 Genus: Withania
 Species : *Withania somnifera*

III. SYNONYMS [28]

Language Name	Language Name
Sanskrit: <i>Ashwagandha</i>	Tamil: <i>Amukkura, amkulang, amukkuram-kilangu</i>
English: <i>Winter Cherry</i>	Karnataka: <i>Viremaddlinagadde, Pannaeru,</i>
Hindi: <i>Punir, asgandh</i>	Konkani: <i>Fatarfoda</i>
Bengali: <i>Ashvagandha</i>	Punjabi: <i>Asgand, isgand</i>
Gujrati: <i>Ghodakun, Ghoda, Asoda</i>	Marathi: <i>Asgund, asvagandha</i>
Telgu: <i>Pulivendram</i>	Rajasthani: <i>Chirpotan</i>

IV. BOTANICAL DESCRIPTION

Plant is a small, woody shrub that grows about two feet in height. An erect, evergreen, tomatoes shrub, 30-150 cm high, found throughout the drier parts of India in waste places and on bunds. Roots are stout fleshy, whitish brown; leaves simple ovate, glabrous, those in the floral region smaller and opposite; flowers inconspicuous, greenish or lubrid-yellow, in axillary, umbellate cymes; berries small, globose, orange-red when mature, enclosed in the persistent calyx; seeds yellow, reniform. The roots are the main portions of the plant used therapeutically. The bright red fruit is harvested in the late fall and seeds are dried for planting in the following spring [29-32].

V. ORIGIN AND DISTRIBUTION

Plant grows abundantly throughout India, especially in Madhya Pradesh, Uttar Pradesh, Punjab and North Western parts of India like Gujarat and Rajasthan. It can be found growing in Africa, the Mediterranean, Congo, South Africa, Egypt, Morocco, Jordan and Pakistan [33-34].

VI. PARTS USED

Whole plant, roots, leaves, stem, green berries, fruits, seeds, bark are used as a crude drug for various purpose (Fig. 1) [35].



Fig. 1: Plant (A) with fruits (B), roots (C) and marketed formulation (D).

VII. PHYTOCHEMISTRY

Various studies reported that most of the chemical constituents are present in the roots of the plant. The total alkaloidal content of the Indian roots has been revealed to vary between 0.13 and 0.31%. The leaves primarily contain withaferin A, a steroidal lactone is the most important withanolide. The fruits contain amino acids, a proteolytic enzyme, condensed tannins, and flavonoids. They contain a high ratio of free amino acids which are proline, valine, tyrosine, alanine, glycine, hydroxyproline, aspartic acid, glutamic acid, cystine and cysteine. Shoots primarily contain scopoletin and also they contain protein, calcium and phosphorous. Stem and bark consist of number of condensed tannins, flavonoids and free amino acids. The major biochemical constituents present in the root are shown in **Table 1** [36-45].

Table 1: Phytoconstituents of roots.

Class	Phytoconstituents
Alkaloids	Withanine, withananine, withasomnine, somniferine, tropeltigloate, somniferinine, somninine and nicotine somnine, somniferinine, withananine, pseudo-withanine, tropine, pseudo-tropine, 3-a-gloyloxytropene, choline, cuscohygrine, isopelletierine, anaferine
Steroidal lactones	Withaferin-A, withanone, withanolide-E, withanolide-F, withanolide-A, withanolideG, withanolide-H, withanolide-I, withanolide-J, withanolide-K, withanolides L, withanolide-M
Steroids	Cholesterol, b-sitosterol, stigmaterol, diosgenin, stigmastadien, sitoinsides VII, sitoinsides VIII, sitoinsides IX, sitoinsides X
Salts	Cuscohygrine, anahygrine, tropine, pseudotropine, anaferine
Flavonoids	Kaempferol, quercetin
N- containing compounds	Withanol, somnisol, and somnitol

VIII. BRAIN RELATED EFFECTS

A. Anxiety and Depression

Extracts of Ashwagandha produce GABA-like activity which may account for the herbs anxiolytic actions. The bioactive constituents produce calming effect. Excessive neuronal activity can lead to restlessness and insomnia, but GABA inhibits the number of nerve cells that the fire in brain and helps to induce sleep, uplift mood and reduce anxiety. It also exhibited an antidepressant effect and as a mood stabilizer [46-48].

B. Chronic Stress

Drug is a very popular herb for reduce stress. Chronic stress (CS) can result in a number of adverse physiologic conditions including cognitive deficit, immune suppression, sexual dysfunction, gastric ulceration, irregularities in glucose homeostasis, and changes in plasma corticosterone levels. Drug inhibited stress-induced gastric ulcer more effectively as compared to the standard drug ranitidine. Administration of ashwagandha with other drugs was found to offer protection against f biological, physical and chemical stressors [49-52].

C. Nootropic Effect

Ashwagandha may improve brain function, memory, reaction times and the ability to perform tasks. The bioactive components like Sitoindosides VII-X and withaferin isolated from aqueous methanol extract of roots slightly enhanced acetylcholinesterase (AChE) activity in the lateral septum and globus pallidus, and decreased AchE activity in the vertical diagonal band. These changes were accompanied by enhanced M1-muscarinic-cholinergic receptor binding in lateral and medial septum as well as in frontal cortices, whereas the M2- muscarinic receptor-binding sites were increased in a number of cortical regions including cingulated, frontal, parietal, and retrosplinal cortex.

Daily administration of drug for 6 days significantly improved memory consolidation On the basis of some findings, it is suggested that drug exhibits a nootropic-like effect [53-57].

D. Anti-parkinsonian Properties

Parkinson's disease is a neurodegenerative disease characterized by the selective loss of dopamine neurons of the substantia nigra pars compacta. Drug inhibited catalepsy and provide hope for treatment of Parkinson's disease. An Anti-parkinsonian effect of extract has been reported due to potent antioxidant, antiperoxidative and free radical quenching properties in various diseased conditions. Chronic treatment with root extract for a period of 4 weeks to reserpine treated animals significantly and dose dependently reduced the vacuous chewing movements and tongue protrusions. Administration of drug also showed an enhancement in phagocytic activity of peritoneal macrophages result in confirms the immunomodulatory activity [58-63].

IX. ANALYSIS OF *WITHANIA SOMNIFERA*

There are many analytical tools that are used for the analysis of various pharmaceutical formulations, herbal formulations, crude drugs and their extracts [64-82]. These methods includes Uv-spectrophotometry, gas chromatography, HPLC, HPTLC, etc [83-103].

X. CONCLUSION

Thus the drug *Withania somnifera* could be used as memory enhancer. Clinical studies of drug proved it as wonder drug for almost all types of memory related problems.

XI. ACKNOWLEDGMENTS

We express our sincere thanks to Shri. Yogendraji Gode and Dr. Yogeshji Gode, IBSS's Dr. Rajendra Gode Institute of Pharmacy, Amravati and Dr. Rajendra Gode College of Pharmacy, Amravati (India).

A. Disclosure of Conflict of Interest

The author declares no conflict of interest.

B. Disclosure of Conflict of Interest

The author declares no conflict of interest.


REFERENCES

- [1] Manwar J, et al. Comparative antioxidant potential of *Withania somnifera* based herbal formulation prepared by traditional and non-traditional fermentation processes. *Integrative Medicine Research*.2013; 2: 56–61.
- [2] Manmode R, et al. Effect of preparation method on antioxidant activity of ayurvedic formulation kumaryasava. *J Homeop Ayurv Med*. 2012; 1: 114.
- [3] More MP, Dongare PN, Patinge PA, Bakal RL, Motule AS. An overview on phytoconstitute and utilisation of *Lepidium sativum* linn (Garden Cress). *World Journal of Pharmacy and Pharmaceutical Science*. 2021; 10(1): 710-719.
- [4] Dongare PN, et al. An Overview on herbal cosmetics and cosmoceuticals. *Int J Pharm Sci Rev Res*. 2021; 68(1): 75-78.
- [5] Dongare PN, Motule AS, et al. An Overview on Anticancer drugs from Marine source. *World Journal of Pharmaceutical Research*. 2021; 10(1) : 950-956
- [6] Badukale NA, et al. Phytochemistry, pharmacology and botanical aspects of *Madhuca indica*: A review. *Journal of Pharmacognosy and Phytochemistry*. 2021; 10(2): 1280-1286.
- [7] Gudalwar BR, et al. *Allium sativum*, a potential phytopharmacological source of natural medicine for better health. *GSC Advanced Research and Reviews*. 2021; 06(03): 220–232.
- [8] Manwar J, Mahadik K, Paradkar A. Plackett–Burman design: A statistical method for the optimization of fermentation process for the yeast *Saccharomyces cerevisiae* isolated from the flowers of *Woodfordia fruticosa*. *Fermentation Technology*. 2013; 2: 109.
- [9] Wadekar AB, et al. Morphology, phytochemistry and pharmacological aspects of *Carica papaya*, an review. *GSC Biological and Pharmaceutical Sciences*. 2020; 14(03): 234-248.
- [10] Manwar J, et al. Isolation, biochemical and genetic characterizations of alcohol-producing yeasts from the flowers of *Woodfordia fruticosa*. *J Young Pharm*. 2013; 5(4): 191-194.
- [11] Manmode R, et al. Effect of preparation method on antioxidant activity of ayurvedic formulation kumaryasava. *J Homeop Ayurv Med*. 2012; 1: 114.
- [12] Khadatkar SN, et al. In-vitro anthelmintic activity of root of *Clitoria ternatea* linn. 2008; 4(13): 148-150.
- [13] Khadatkar SN, et al. Preparations and evaluation of microcapsules of capsaicin. *International Journal of Chemical Sciences*. 2007; 5(5): 2333-2341.
- [14] Malode GP, et al. Phytochemistry, pharmacology and botanical aspects of *Murraya Koenigii* in the search for molecules with bioactive potential - A review. *GSC Advanced Research and Reviews*. 2021; 06(03): 143–155.
- [15] Padgilwar S, et al. Traditional uses, phytochemistry and pharmacology of *Oroxylum Indicum*: A Review. *International Journal of Pharmaceutical and Phytopharmacological Research*. 2014; 3(6): 483-486.
- [16] Nikhare AM, et al. Morphological, Phytochemical and pharmacological aspects of *Syzygium Cumini*. *International Journal of Medical, Pharmaceutical and Biological Sciences*. 2021; 1(1): 1-11.
- [17] Padgilwar SS, Manwar JV. Relative Influence of adrenergic β -agonist and antagonist on the inflammation and their interaction with aspirin. *European Journal of Experimental Biology*. 2013; 3(1):467-472.

- [18] Parbat AY, et al. Ethnopharmacological review of traditional medicinal plants as immunomodulator. *World Journal of Biology Pharmacy and Health Sciences*. 2021; 06(02): 043–055.
- [19] Malode LL, et al. Potential of medicinal plants in management of diabetes: An updates. *GSC Advanced Research and Reviews*. 2021; 08(01): 149-159.
- [20] Sahare AY, et al. Antimicrobial activity of *Pseudarthria viscida* roots. *Asian Journal of Microbiology Biotechnology & Environmental Sciences*. 2008; 10(1): 135-136.
- [21] Dongare PN, et al. Recent development in novel drug delivery systems for delivery of herbal drugs: An updates. *GSC Advanced Research and Reviews*. 2021;
- [22] Sahare AY, et al. *Hypericum perforatum*: A Medicinal plant. *Plant Archives*. 2007; 7(2): 463-468.
- [23] Gudalwar BR, Panchale WA, Manwar JV, Nimbalwar MG, Badukale NA, Bakal RL. Pharmacognosy, phytochemistry and clinical applications of traditional medicinal plants as memory booster. *GSC Advanced Research and Reviews*. 2021.
- [24] Jana SN, Charan SM, Health Benefits and Medicinal Potency of *Withania somnifera*: A Review. *Int. J. Pharm. Sci. Rev. Res.*, 2018; 48(1): 22-29.
- [25] Gupta GL, Rana RM, Varalakshmi P. Immunomodulatory role of *Withania somnifera* root powder on experimental induced inflammation: An in vivo and in vitro study. *Vascul Pharmacol*, 2006; 44 (6): 406-410.
- [26] Umadevi M, Rajeshwari R, Rahale Sharmila C, Selvavenkadesh S, Pushpa R, Kumar Sampath K P, Bhowmik D. Traditional and medicinal uses of *Withania somnifera*, *The Pharma Innovation*, 2012; 9: 102-110.
- [27] Rizvi Faizi T, Razauddin Md, URRahman Saif Md, Jahan T, Naz Z, Kumar R, Kumar A, Ali M, Immunomodulatory effect of Ashwagandha against doxorubicin toxicity, *European Journal of Pharmaceutical and Medical Research*, 2016; 3(7): 463-467.
- [28] Kumar R, Singh J.K, Nath A, Ali Md, Kumar A, Sinha P, Effect of *Withania somnifera* on Estrogen, Cholesterol and Subcellular structure of ovary of chlorpyrifos exposed mice, *Caribbean Journal of Science and Technology*, 2013; 1: 061- 069.
- [29] Narinderpal K, Junaid N, Raman B, A review on pharmacological profile of *Withania somnifera* (Ashwagandha), Research and reviews: *Journal of Botanical Sciences*, 2013; 2: 6-14.
- [30] Mirjalili Hossein M, Moyano E, Bonfill M, Cusido M.R, Palazon J, Steroidal lactones from *Withania somnifera*, an ancient plant for novel medicine, *Molecules*, 2009; 14: 2373- 2393.
- [31] Mir Ahmad B, Khazir J, Mir A.N, Hasan ul T, Koul S. Botanical, chemical and pharmacological review of *Withania somnifera* (Indian ginseng): an ayurvedic medicinal plant, *Indian Journal of Drugs and Diseases*, 2012; 1: 2278-2958.
- [32] Gupta p, Goel R, Agarwal AV, Asif MH, Sangwan NS, Sangwan RS & Trivedi P. Comparative transcriptome analysis of different chemotypes elucidates withanolide biosynthesis pathway from medicinal plant *Withania somnifera*. *Scientific Reports*, 2015; 5: 1-13.
- [33] Khare CP. *Indian Medicinal Plants—An Illustrated Dictionary*. First Indian Reprint, Springer (India) Pvt. Ltd., New Delhi, 2007; 717- 718.
- [34] Anonymous. *Standardisation of Single Drugs of Unani Medicine*. Part III, 1st ed. Central Council for Research in Unani Medicine (CCRUM), New Delhi, 2007; 9-14.
- [35] Gaurav N, Kumar A, Tyagi M, Kumar D, Chauhan U.K, Singh A.P, Morphology of *Withania somnifera* (Distribution, Morphology, Phytosociology of *Withania somnifera* L. Dunal), *International Journal of Current Science Research*, 2015;1: 2454-5422.
- [36] Rao R, Rajput D.K, Nagaraju G, Adinarayana G, Opportunities and challenges in the cultivation of Ashwagandha (*Withania somnifera* (L.) Dunal), *Journal of Pharmacognosy*, 2012; 3: 88-91.
- [37] Jain R, Kachhwaha S, Kothari S.L, Phytochemistry, pharmacology, and biotechnology of *Withania somnifera* and *Withania coagulans*: A review, *Journal of Medicinal Plants Research*, 2012; 6:5388-5399.
- [38] Bara Kiran J, Soni R, Jaiswal S, Saksena P, Phytochemical study of the plant *Withania somnifera* against various diseases, *Journal of Agriculture and Veterinary Science*, 2016; 9: 109-112.
- [39] Srivastav Kumar A, Das P, Phytochemical extraction and characterization of roots of *Withania somnifera* for its antibacterial, antioxidant, antiinflammation and analgesic activity, *International Journal of Innovative Research and Development*, 2014; 3: 22-33.
- [40] Uddin R, Samiulla L, Singh V.K, Jamil S.S, Phytochemical and pharmacological profile of *Withania somnifera* Dunal: A review. *Journal of Applied Pharmaceutical Science*, 2012; 02: 170-175.
- [41] Kumar V, Dey A, Chatterjee Sunder S, *Phytopharmacology of Ashwagandha as an antidiabetic herb*, Springer International Publishing, 2017; 3: 37-68.
- [42] Ahmad M, Saleem S, Ahmad AS, Ansari MA, Yousuf S, Hoda MN, Neuroprotective effects of *Withania somnifera* on 6- hydroxydopamine induced Parkinsonism in rats, *Hum Exp Toxicol*, 2005; 24(3): 137-147.
- [43] Kumar A, Kulkarni SK, Effect of BR-16A (Mentat) a polyherbal formulation on drug-induced catalepsy in mice, *Indian J Exp Biol*, 2016; 44(1): 45-48.
- [44] Gupta Lal G, Rana A.C, Plant review *Withania somnifera* (Ashwagandha): A review, *Pharmacognosy Reviews*, 2017; 1: 129-136.
- [45] Majumder B, Role of protein vibration in antialzheimer's effect of Ashwagandha (*Withania somnifera*)-An analytical approach, *International Journal of Biophysics*, 2017; 7(3): 41- 47.
- [46] Khan MA, Subramanyaan M, Arora VK, Banerjee BD, Ahmed RS, Effect of *Withania somnifera* (Ashwagandha) root extract on amelioration of oxidative stress and autoantibodies production in collagen-induced arthritic rats, *J Complement Integr Med*, 2015; 12(2): 117-125.
- [47] Giri RK, Comparative study of anti-inflammatory activity of *Withania somnifera* (Ashwagandha) with hydrocortisone in experimental animals (Albino rats), *Journal of Medicinal Plants Studies*, 2016; 4(1): 78-83.
- [48] Singh A, Malhotra S, Subban R, Anti-inflammatory and analgesic agents from Indian medicinal plants, *International Journal of Integrative Biology*, 2008; 3: 57-72.
- [49] Verma S, Pharmacological activity of *Withania somnifera*, *World Journal of Pharmacy and Pharmaceutical Sciences*, 2016; 5: 602-605.
- [50] Ashok Amit G, Shende M.B, Chothe D.S, Antistress activity of Ashwagandha (*Withania somnifera* Dunal)-A review, *International Ayurvedic Medical Journal*, 2014; 2(3): 386- 393.
- [51] Trivedi Bihari A, Mahajan D, Ashwagandha-The powerful antistress herb, *World Journal of Pharmaceutical and Life Sciences*, 2017; 3: 177-180.
- [52] Singh N, Verma P, Pandey R.B, Gilca M, Role of *Withania somnifera* and treatment of cancer: An overview, *International Journal of Pharmaceutical Sciences and Drug Research*, 2011; 3(4): 274-279.
- [53] Chandrasekhar K, Kapoor J, Anishetty S, A prospective, randomized double blind, placebo controlled study of safety efficacy of a high concentration full-spectrum extract of Ashwagandha root in reducing stress and anxiety in adults, *Indian J Psychol Med*, 2012; 34(3): 274-279.

- [54] Singh G, Sharma PK, Dudhe R, Singh S, Biological activities of *Withania somnifera*, *Annals of Biological Research*, 2010; 1(3): 56-63.
- [55] Ziauddin M, Phansalkar N, Patki P, Diwanay S, Patwardhan B, Studies on the immunomodulatory effects of *Ashwagandha*, *Journal of Ethnopharmacology*, 1996; 50(2): 69-76.
- [56] Prakash J, Yadav SK, Chouhan S, Singh SP. Neuroprotective role of *Withania somnifera* root extract in maneb-paraquat induced mouse model of Parkinsonism. *Neurochem. Res.* 2013; 38: 972–980.
- [57] Singh Pal S, Tanwer SB, Khan M, Antifungal potential of *Ashwagandha* against some pathogenic fungi, *International Journal of Biopharmaceutics*, 2010; 1(2): 72-74.
- [58] Rizwana N, Antibacterial potential of *Withania somnifera* L. against human pathogenic bacteria, *African Journal of Microbiology Research*, 2012; 6(22): 4810-4815.
- [59] Patil SP, Maki S, Khedkar SA, Rigby AC, Chan C. Withanolide A and asiatic acid modulate multiple targets associated with amyloid-beta precursor protein processing and amyloid-beta protein clearance. *J. Nat. Prod.*, 2010; 73: 1196–1202.
- [60] Kurapati KR, Atluri VS, Samikkannu T, Nair MP. *Ashwagandha* (*Withania somnifera*) reverses beta-amyloid1–42 induced toxicity in human neuronal cells: Implications in HIV-associated neurocognitive disorders (HAND). *PLoS ONE*, 2013; 8: 77624.
- [61] Nakayama N, Tohda C. Withanoside IV improves hindlimb function by facilitating axonal growth and increase in peripheral nervous system myelin level after spinal cord injury. *Neurosci. Res.*, 2007; 58: 176–182.
- [62] Ahmad M, Saleem S, Ahmad AS, Ansari MA, Yousuf S, Hoda MN, Islam F. Neuroprotective effects of *Withania somnifera* on 6-hydroxydopamine induced Parkinsonism in rats. *Hum. Exp. Toxicol.*, 2005; 24: 137–147.
- [63] Kumar P, Kumar A. Possible neuroprotective effect of *Withania somnifera* root extract against 3-nitropropionic acid-induced behavioral, biochemical, and mitochondrial dysfunction in an animal model of Huntington's disease. *J. Med. Food*, 2009; 12: 591–600.
- [64] Motule AS, et al. Development and physicochemical evaluation of bilayered transdermal patches of ondansetron hydrochloride *Journal of Innovations in Pharmaceutical and Biological Sciences*. 2021; 8(3): 17-23.
- [65] Chaudhari KD, et al. Floating drug delivery system: An update of preparation and classification of formulation. *Ijppr.Human*. 2021; 21(1):207-220.
- [66] Chaudhari KD, et al. Comprehensive review on characterizations and application of gastro-retentive floating drug delivery system. *GSC Advanced Research and Reviews*. 2021; 07(01):035-044.
- [67] Dhamankar AK, et al. The novel formulation design of O/ of ketoprofen for improving transdermal absorption. *Int J of Pharm Tech Res.* 2009; 4(1Suppl): 1449-1457.
- [68] Jain CM, et al. Review on approaches for development and evaluation of extended-release tablets. Review on approaches for development and evaluation of extended-release tablets. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2021;10(4): 542-554.
- [69] Kadam CY, et al. Design and In vitro characterization of phase transition system using rivastigmine tartrate for nasal drug delivery system. *World Journal of Pharmaceutical Research*. 2018; 8(1): 815-829.
- [70] Malode GP, et al. Formulation and evaluation of a novel floating in situ gel system for the treatment of peptic ulcer. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2021; 10(4):416-1433.
- [71] Nimbawar MG, et al. A brief review on principle, preparation and properties of proniosomes: A provesicular drug delivery system. *World J Pharm Sci*. 2021; 9(5): 149-162.
- [72] Nimbawar MG, et al. An overview of characterizations and applications of proniosomal drug delivery system. *GSC Advanced Research and Reviews*, 2021, 07(02), 025–034.
- [73] Nimbawar MG, Upadhye K, Dixit G. Fabrication and evaluation of ritonavir proniosomal transdermal gel as a vesicular drug delivery system. *Pharmacophore*. 2016; 7(2):82-95.
- [74] Manwar J, Kumbhar DD, Bakal RL, Baviskar SR, Manmode RS. Response surface based co-optimization of release kinetics and mucoadhesive strength for an oral mucoadhesive tablet of cefixime trihydrate. *Bulletin of Faculty of Pharmacy. Cairo University*. 2016; 54: 227–235. <http://dx.doi.org/10.1016/j.bfopcu.2016.06.004>
- [75] Manwar JV, et al. Diclofenac Sodium Loaded Nanosized Ethosomes: An Investigation on Z-Average, Polydispersity and Stability. *J Pharm Res*. 2017; 1(3): 000115.
- [76] Nimbawar MG, et al. Fabrication and evaluation of ritonavir proniosomal transdermal gel as a vesicular drug delivery system. *Pharmacophore*. 2016; 7(2): 82-95.
- [77] Patil SS, et al. Ultrasound-Assisted Facile Synthesis of Nanostructured Hybrid Vesicle for the Nasal Delivery of Indomethacin: Response Surface Optimization, Microstructure, and Stability. *AAPS PharmSciTech*. 2019;20(3):97.
- [78] Pophalkar PB, et al. Development and evaluation of ondansetron medicated jelly. *World Journal of Pharmaceutical Research*. 2018; 7(19): 1252-1263.
- [79] Shubham Garibe, et al. Bioequivalence study of test formulations T1 and T2 Nadolol tablets USP with reference formulation in healthy adult, human subjects under fed conditions. *Ijppr.Human*. 2021; 20(2):20-28.
- [80] Suroshe RS, et al. Development and characterization of osmotic drug delivery system of model drug. *World Journal of Pharmaceutical Research*. 2018; 7(18): 1158-1171.
- [81] Vaidya VM, et al. Design and in vitro evaluation of mucoadhesive buccal tablets of terbutaline sulphate. *Int J PharmTech Res*. 2009; 1(3): 588-597.
- [82] Vohra M, et al. Bioethanol production: Feedstock and current technologies. *Journal of Environmental Chemical Engineering*. 2014; 2 (1):573-584.
- [83] Bakal RL, et al. Spectrophotometric estimation of amitriptyline HCl and chlordiazepoxide in pharmaceutical dosage form. *Indian Journal of Pharmaceutical Education and Research*. 2008; 42: 23–26.
- [84] Bakal RL, et al. Spectrophotometric estimation of amitriptyline HCL and chlordiazepoxide in tablet dosage form. *International Journal of Chemical Sciences*. 2007; 5(1):360–364.
- [85] Gulhane CA, et al. UV- Visible Spectrophotometric estimation of azithromycin and cefixime from tablet formulation by area under curve method. *World Journal of Pharmaceutical Sciences*.2021; 9(6): 163-168.
- [86] Manwar JV, et al. Development of newer RP-HPLC method for simultaneous estimation of cefixime and linezolid in bulk drugs and combined dosage form. *International Journal of Pharmacy and Life Sciences*. 2021;12(1):26-31.

- [87] Gulhane CA, et al. Liquid chromatographic method for simultaneous estimation of thiocolchicoside and etoricoxib from tablet formulation. *Asian Journal of Pharmaceutical Analysis*. 2021;11(2): 118-122.
- [88] Manwar JV, et al. Application of simultaneous equation method for the determination of azithromycin and cefixime trihydrate in tablet formulation. *Research Journal of Pharmacy and Technology*. 2017;10(1):108-112.
- [89] Bagade SB, et al. Simultaneous high performance thin layer chromatographic estimation of methocarbamol and nimesulide in combined dose tablet. *Journal of Pharmaceutical Research*. 2006;5(4):137-140.
- [90] Manmode RS, et al. Stability indicating HPLC method for simultaneous determination of methocarbamol and nimesulide from tablet matrix. *Der Chemica Sinica*. 2011;2(4):81-85.
- [91] Manwar J, Mahadik K, Paradkar A, et al. Gas chromatography method for the determination of non-ethanol volatile compounds in herbal formulation. *International Journal of Analytical and Bioanalytical Chemistry*. 2013; 3(1):12-17.
- [92] Manwar JV, et al. Experimental design approach for chromatographic determination of ketorolac tromethamine from bulk drug and tablet formulation. *Global Journal of Pharmacy & Pharmaceutical Sciences*. 2017;3(2):38-47.
- [93] Panchale WA, et al. Chromatographic analysis of famotidine, paracetamol and ibuprofen from tablet formulation. *Research Journal of Pharmacy and Technology*. 2019; 12:231-263.
- [94] Manwar JV, et al. Rapid RP-HPLC method for estimation of zidovudine from tablet dosage form. *Der Chemica Sinica*. 2011; 2(5): 152-156.
- [95] Manwar JV, Mahadik KR, Paradkar AR, et al. Determination of withanolides from the roots and herbal formulation of *Withania somnifera* by HPLC using DAD and ELSD detector. *Der Pharmacia Sinica*. 2012; 3: 41–46.
- [96] Panchale WA, et al. RP-HPLC method for simultaneous determination of escitalopram oxalate and flupentixol HCl in tablet dosage form. *GSC Biological and Pharmaceutical Sciences*. 2021; 14(01):169-174.
- [97] Panchale WA, et al. RP-HPLC method for simultaneous determination of metformin hydrochloride and linagliptine in pharmaceutical dosage form. *World Journal of Pharmaceutical and Medical Research*. 2021;7(5):234- 238.
- [98] Nimbokar SW, et al. Development and validation of RP-HPLC method for determination of zonisamide from tablet formulation. *World Journal of Pharmaceutical and Medical Research*. 2021;7(2):196-200.
- [99] Panchale WA, Bakal RL. First-order derivative spectrophotometric estimation of gemifloxacin mesylate and ambroxol HCl in tablet dosage form. *GSC Biological and Pharmaceutical Sciences*. 2021; 14(2):029-036.
- [100] Manwar JV, et al. Response surface based optimization of system variables for liquid chromatographic analysis of candesartan cilexetil. *Journal of Taibah University for Science*. 2017; 11:159–172.
- [101] Panchale WA, et al. Concurrent analysis of ambroxol HCl and salbutamol sulphate from tablet formulation by RP-HPLC. *GSC Biological and Pharmaceutical Sciences*. 2020; 13(03):197-202.
- [102] Panchale WA, et al. Simultaneous estimation of salbutamol sulphate and ambroxol HCl from their combined dosage form by UV-Vis spectroscopy using simultaneous equation method. *GSC Biological and Pharmaceutical Sciences*. 2020;13(03):127-134.
- [103] Sabhadinde AF, et al. Novel RP-HPLC method for simultaneous analysis of chlorthalidone and telmisartan from combined dosage form. *Ijppr.Human*. 2020; 20(1):491-502.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)