

Review Article

Pharmacological and other Bioactivities of the Genus *Polygonum* - A Review

Xiaoxv Dong¹, Jing Fu¹, Xingbin Yin¹, Xuechun Li¹, Bo Wang², Sali Cao¹, Jin Zhang¹, Hui Zhang¹, Yang Zhao¹ and Jian Ni*

¹School of Chinese Pharmacy, Beijing University of Chinese Medicine, Beijing, 100102, ²Institute of Materia Medica, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China

*For correspondence: **Email:** njtcm@263.net; **Tel:** 010-84738607

Received: 6 June 2014

Revised accepted: 14 September 2014

Abstract

The genus *Polygonum* encompasses approximately 300 species widely distributed around the world, among which 120 *Polygonum* species grow in China. Among the rich species, 81 of them are widely used in traditional Chinese and folk medicines. In recent years, several studies have found that the genus *Polygonum* possesses a variety of biological activities including antioxidation, antimicrobial, anti-tumor, anti-obesity, etc. However, there are still not enough systemic data on the chemical constituents and their pharmacological effects; hence, it would be useful to review current literature for available pharmacological activities of the genus and as well as its active ingredients. Both in-vitro and in-vivo studies have provided strong evidence for the therapeutic potential of the genus *Polygonum*. This review collates and examines information on the pharmacological effects of the genus *Polygonum* as well as its chemical constituents.

Keywords: *Polygonum*, Chemical constituents, Pharmacological properties, Flavonoids, Anthraquinones

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Polygonum is a medicinal large genus of *Polygonaceae*, it falls into about 300 species widely distributed around the world. The genus *Polygonum* contains many medicinal plants, such as *Polygonum multiflorum*, *Polygonum cuspidatum*, *Polygonum bistorta*, *Polygonum aviculare*, *Polygonum tinctorium*, etc [1]. Among the rich species, parts of the plant are recorded in traditional Chinese medicines and folk medicines. Additionally, many chemical constituents have been identified such as flavonoids, anthraquinones, stilbenes, glycolipids and terpenes [2]. For its enormous social and economic implications, it is urgent to understand

the mechanisms between the chemical constituents and the pharmacological effects. The primary objectives of this review are to summarize: the pharmacological effects and the main chemical constituents, and their structures.

Antioxidant effect

Most of the genus *Polygonum* have antioxidant and clears the body of excess free radicals [1]. 5,6-dihydropyranobenzopyronean 5,6-dihydropyranobenzopyrone (Fig. 1) isolated from *Polygonum amplexicaule* had a strong ability to scavenge oxygen free radicals [3]. Similarly, hydropiperoside B, vanicode A and vanicode E (Fig. 1) isolated from *Polygonum hydropiper* L also exhibited antioxidant activity [4].

Additionally, flavonoids and flavonoid glucoside also exhibited high antioxidant activity. Among these, 2''-O-(3,4,5-trihydroxybenzoyl) quercitrin (galloyl quercitrin) (Fig.1) showed the strongest antioxidant activity [5-7]. *Polygonum aviculare* L. extracts strongly exhibited antioxidant effects by free radical scavenging assays, superoxide radical scavenging assays, lipid peroxidation assays and hydroxyl radical induced DNA strand scission assays [8]. *Polygonum minus* extracts exhibited gastro protective activities. The mechanisms were attributed to the synthesis of antioxidant and PGE2 [9]. Three compounds including gallic acid, catechin, and 2,3,5,4'-tetrahydroxystilbene-2-O- β -D-glucoside (Fig. 1) isolated from *Polygonum multiflorum* showed strong antioxidant activity [10]. In addition, the stilbene glycoside from *Polygonum multiflorum* also exhibited strong antioxidant activity by increasing the activities of superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) [11]. What is more, the stilbene glycoside (Fig.1) also exerted its antioxidant activity by inhibiting the A/R-mediated elevation of MDA content [12]. Resveratrol and its liposomal isolated from *Polygonum cuspidatum* could protect the dopaminergic neurons, attributing their radical scavenging ability and antioxidant effects which could be attributed their radical scavenging ability and antioxidant effects [13,14]. The methanol and ethyl acetate extracts of *Polygonum tinctorium* also exhibited higher antioxidant activity [15]. *Polygonum maritimum* L. extracts presented a remarkable antioxidant scavenging effects on DPPH radical [16]. Similarly, the MEOH extract of *Polygonum sachalinensis* had antioxidant effects by free radical-scavenging activities [17].

Anti-inflammatory effect

The inhibitory effect of ethanol extract of *Polygonum cuspidatum* (PCE) on mouse ear inflammation induced by 12-O-tetradecanoylphorbol-13-acetate (TPA) was tested. The results showed PCE significantly reduced the ear edema in a dose-dependent manner [18]. Additionally, the mRNA expressions of TNF- α , IL-6, and C-reactive protein significantly decreased after treatment with the extract of *Polygonum cuspidatum* (containing 20 % resveratrol) [19]. Similarly, well-organized bands of collagen, fibroblasts and hair follicle increased, while the number of inflammatory cells reduced after wound healing rats were treated with the extract from *Polygonum cuspidatum* [20]. Furthermore, *Polygonum cuspidatum* extracts significantly inhibited FCA-induced joint swelling within 3 days in FCA-induced adjuvant arthritis model [21]. *Polygonum*

viviparum (PV) extracts could inhibit inflammation induced by lipopolysaccharide in RAW 264.7

RAW264.7 macrophages, and the mechanisms were related to might its haem oxygenase-1 induction and activation of the Nrf2 pathway [22]. *Polygonum viscosum* extracts possessed moderate anti-inflammatory activity on raw paw edema induced by carrageenan [23]. Quercetin-3-O- β -D-glucuronide (Fig 1) separated from *Polygonum perfoliatum* significantly suppressed ear edema induced by dimethyl benzene in mice [24]. Two compounds (5-glutenin-3-one, friedelanol) from *Polygonum bistorta* significantly largely suppressed the inflammatory response [25].

Antibacterial and antifungal effect

A fraction isolated from *Polygonum cuspidatum* possessed antibacterial effect against *Streptococcus mutans*, indicating the fraction might be useful for controlling dental biofilms and improving the cariostatic properties of fluoride without increasing its exposure [26]. In addition, the ethyl acetate fraction (polydatin, resveratrol, anthraglycoside B, emodin) the ethyl acetate fraction composed of polydatin, resveratrol, anthraglycoside B, and emodin (Fig. 1) also exhibited a significantly antibacterial effect against three of the five common foodborne bacteria which were *Bacillus cereus*, *Listeria monocytogenes*, and *Staphylococcus aureus* [27,28]. Through By bioassay-guided separation and analysis of antibacterial activity, the essential oils isolated from *Polygonum bistorta* inhibit *Paenibacillus larvae*, *Melissococcus plutonius* and *Bacillus subtilis* [29]. *Polygonum punctatum* extracts from the Brazilian medicine plant possessed antifungal effect. The sesquiterpene dialdehyde polygodial (Fig. 1) was the main active constituent [30]. The chloroform extract of *Polygonum aviculare* isolated from *Polygonum aviculare* had a significant antibacterial effect against *Escherichia coli*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Salmonella paratyphi*, *Shigella flexner*, *Staphylococcus aureus*, *Bacillus subtilis*, *Streptococcus pyogenes*, *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger* [31]. In addition, the gingival index significantly decreased after treatment with *Polygonum aviculare* extracts, indicating they could be useful for the therapy of gingivitis induced by bacteria. The results showed total flavonoids were the effective part of its antibacterial activity [32,33]. *Polygonum orientale* extracts also exhibited antibacterial effect, suggesting the extracts might be useful for controlling bacterial ring rot of potato disease [34]. *Polygonum capitatum*

extracts possessed bacteriostatic and bactericidal properties by evaluated *in vitro* [35]. Tryptanthrin and Kaempferol isolated from *Polygonum tinctorium* L significantly decreased the numbers of *Helicobacter pylori* colonies *in vivo* and *in vitro*, indicating they could be indicated for anti-*H. pylori* therapy [36]. Polygoumic acid (Fig. 1) from *Polygonum viscosum* could inhibit the growth of penicillin-resistant *Escherichia coli* (MIC = 0.05 mg/ml) and methicillin-resistant *Staphylococcus aureus* (MIC = 0.10 mg/ml) [37].

Anticancer effect

The extracts of *Polygonum hypoleucum* Ohwi possessed inhibitory effects on various tumor cells proliferation. Emodin was thought to be the main effective substances [38]. *Polygonum cuspidatum* extracts exhibited an antiproliferative effect on human lung cancer cells by inducing apoptosis and inhibiting cell growth in A549 and H1650 cell lines [39]. Similarly, resveratrol isolated from *Polygonum cuspidatum* significantly reduced Lewis lung tumor and tumor weight. Furthermore, it also prevented tumor growth and metastasis in lungs by inhibiting DNA synthesis of tumor cells and tumor-induced neovascularization in mice bearing highly metastatic Lewis lung carcinoma tumors [40]. What is more, *Polygonum cuspidatum* extracts significantly inhibited suspension growth, activated caspases, and induced anoikis in hepatocarcinoma cells in a dose- and time-dependent manner [41]. Additionally, *Polygonum cuspidatum* extracts also inhibited the proliferation of oral cancer cells by inducing caspase-dependent apoptosis, suggesting the extracts might be as a promising compound for the effective treatment of oral cancer [42].

Two new flavonoid glucuronides, quercetin-3-O- β -glucuronide and quercetin-3-O- α -rhamnosyl-(1 \rightarrow 2)- β -glucuronide (Fig. 1) isolated from *Polygonum amphibium* could induce apoptosis in Jurkat and HL60 cell lines by evaluated *in vitro* [43]. *Polygonum perfoliatum* extracts possessed cytotoxicity effects against human mammary carcinoma, human colon carcinoma, human hepatocellular carcinoma, human prostate carcinoma, and human erythroleukaemia cells [44]. *Polygonum aviculare* extracts (300, 350 and 400 ng/ μ l) also exhibited strong inhibitory effects on cell proliferation and induced apoptotic cell death in the MCF-7 breast cancer cells [45]. Phenylpropanoid esters of sucrose, vanicoside B and lapathoside A (Fig.1) isolated from *Polygonum lapathifolium* exhibited the inhibitory effects on the EBV-EA induction and anti-tumor-promoting effects on mouse two-stage skin

carcinogenesis [46]. The ethyl acetate extract and tryptanthrin (50 mg/kg) from *Polygonum tinctorium* Lour. exhibited anticancer activity on intestinal tumors induced by azoxymethane (AOM) [47].

Antiviral effect

Both ethanol extract and water extract of *Polygonum cuspidatum* significantly increase-d the expression of HBsAg and viral antigens., while inhibited the expression of HBeAg, suggesting *Polygonum cuspidatum* extracts could inhibit HBV in a stable HBV-producing cell line [48].

In addition, emodin isolated from *Polygonum cuspidatum* exhibited the potent antiviral activity by inhibiting CVB4 entry and replication, indicating it could be used as potential antiviral in the post-exposure prophylaxis for CVB4 infection [49]. *Polygonum tinctorium* extracts exhibited fairly strong antiviral effect against HIV-1 (EC₅₀ was 0.5 μ g/ml) [50]. Similarly, *Polygonum viscosum* extracts also possessed anti-HIV-1 activity. The main effective substances were quercetin 3-O-(6"-feruloyl)- β -D-galactopyranoside (Fig.1) and viscoazulone [51].

Lipid-regulating effect

Polygonum aviculare extracts (PAE) exhibited anti-obesity effects by suppressing lipogenesis in white adipose tissue and increasing antioxidant activity. Besides, its low toxicity in mice and its historical use suggested PAE might be used as a safe anti-obesity pharmaceutical [52]. *Polygonum hypoleucum* Ohwi extracts were found to exhibit favorable effects in alleviating metabolic syndrome, including obesity, hypertriglyceridemia, and the results showed emodin and emodin-8-O- β -D-glucopyranoside (Fig.1) were the main effective substances [53].

Plasma cholesterol, plasma triglyceride and low-density lipoprotein cholesterol increased, while very low-density lipoprotein cholesterol attenuated after treatment with a water-soluble fraction of *Polygonum multiflorum* (PMS), suggesting PMS might be applicable for the treatment of hyperlipidemia disease [54]. In a high fat/cholesterol rabbit model, polydatin from *Polygonum cuspidatum* obviously decreased the serum levels of total cholesterol (TC), triglyceride (TG), and low-density lipoprotein cholesterol. Meanwhile, the ratio of TC and the liver coefficient were also reduced [55]. In addition, resveratrol isolated from *Polygonum cuspidatum* could reduce the cholesteryl ester synthesis in

human hepatocytes by inhibiting ACAT activity in a dose-dependent manner [56].

Neuroprotective effect

The neuroprotective effects of *Polygonum multiflorum* extract (PME) and its two fractions (PME-I, PME-II) were studied in male C57BL/6 mice, the results showed PME-I had neuroprotective effect on damage of the substantia nigral dopaminergic system induced by PQMB in mice, indicating PCE could be beneficial in preventing Parkinson's disease [57]. Pretreatment with hexane extract from *Polygonum multiflorum* significantly decreased glutamate-induced neurotoxicity in a concentration-dependent manner and drastically inhibited glutamate-induced apoptosis [58].

In addition, the ethyl acetate extract from *Polygonum multiflorum* also had neuroprotective effects through both alleviation of extracellular regulated kinase (CRK) and p38 activation with increased activation of cAMP responsive element binding protein (CREB) under oxidative stress [59]. 85 % methanol extracts of *Polygonum cuspidatum* exhibited strong neuroprotective activity by a lipid peroxidation assay *in vitro* and an assay *in vivo* using a transient focal cerebra ischemia model. HPLC analysis for the key compound groups might be stilbene sand anthraquinones [60].

Estrogenic effect

Emodin and emodin 8-O- β -D-glucopyranoside (Fig.1) isolated from *Polygonum cuspidatum* could enhance proliferation of MCF-7 cell, the results demonstrated emodin might be useful for replacement therapy for human menoxenia and postmenopausal diseases [61]. *Polygonum hydropiper* extracts strongly altered the histological structures of both ovary-intact and OVX rats by inducing hyperplasia in places of luminal epithelium and degeneration of endometrial glands [62].


Nine other pharmacological effects

The hexane, ethylacetate and methanol extract of *Polygonum hydropiper* possessed significantly antinociceptive activity on acetic acid-induced writhing in mice [63]. What is more, a compound from the sprout of *Polygonum hydropiper* L. inhibited 70 % of the tyrosinase activity, suggesting it might be a new tyrosinase inhibitor alternative to cosmetic agents [64]. Emotion-8-O- β -D-glucoside (Fig 1) isolated from *Polygonum amplexicaule* Forb. (PAF) significantly promoted cell proliferation and differentiation of osteoblasts *in vitro* [65]. In addition, four anthraquinones and two stilbenes isolated from *Polygonum cuspidatum* also possessed strong inhibition of tyrosinase, indicating it might be useful as skin-whitening agents [66].


Table 1: Pharmacological effects of the Genus *Polygonum*

Pharmacological effect	Plant	Extract/Compound	<i>In vitro/in vivo</i>
Antioxidant Activity	<i>Polygonum amplexicaule</i> <i>Polygonum hydropiper</i> L.	5,6-dihydropyrano benzopyronean	<i>In vitro</i>
		Hydropiperoside B	<i>In vitro</i>
	<i>Polygonum aviculare</i> L. <i>Polygonum minus</i> <i>Polygonum multiflorum</i> <i>Polygonum cuspidatum</i> <i>Polygonum tinctorium</i> <i>Polygonum maritimum</i> <i>Polygonum sachalinensis</i>	vanicode A	<i>In vitro</i>
		vanicode E	<i>In vitro</i>
		Ethanol extracts	<i>In vitro</i>
		Ethyl acetate: methanol=1:1 (V/V)	<i>In vivo</i>
		Gallic acid	<i>In vitro</i>
		Catechin	<i>In vitro</i>
		2,3,5,4'-tetrahydroxystilbene2-O- β -D-glucoside	<i>In vitro</i>
		Stilbene glycoside	<i>In vivo</i>
		Resveratrol	<i>In vivo</i>
		methanol , ethyl acetate extracts	<i>In vitro</i>
		methanol extract	<i>In vitro</i>
MEOH extract	<i>In vitro</i>		
Anti-inflammatory activity	<i>Polygonum cuspidatum</i>	Ethanol extracts	<i>In vivo</i>
		Extracts (containing 20% resveratrol)	<i>In vivo</i>
		Ethyl acetate extracts	<i>In vivo</i>
	<i>Polygonum viviparum</i> <i>Polygonum viscosum</i> <i>Polygonum perfoliatum</i>	2-propanol extract	<i>In vivo</i>
		Sesquiterpenes	<i>In vivo</i>
		Flavonoid glycoside	<i>In vivo</i>
	<i>Polygonum perfoliatum</i>	Quercetin-3-O- β -D-glucuronide	<i>In vivo</i>


	<i>Polygonum bistorta</i>	5-glutinen-3-one Friedelanol	<i>In vitro</i> <i>In vitro</i>	
Antibacterial and antifungal activities	<i>Polygonum cuspidatum</i>	Resveratrol Emodin Physcion	<i>In vivo</i> <i>In vivo</i> <i>In vivo</i>	
	<i>Polygonum bistorta</i> <i>Polygonum punctatum</i> <i>Polygonum aviculare</i> .	Essential oils Sesquiterpene dialdehyde polygodial Chloroform extract	<i>In vitro</i> <i>In vitro</i> <i>In vitro</i>	
	<i>Polygonum orientale</i> <i>Polygonum capitatum</i> <i>Polygonum tinctorium</i> <i>Polygonum viscosum</i>	Water extracts Ethanol extracts Tryptanthrin , Kaempferol Polygosomic acid	<i>In vitro / in vivo</i> <i>In vitro</i> <i>In vitro / in vivo</i> <i>In vitro</i>	
	Plant	Extract/Compound	<i>In vitro/in vivo</i>	
Pharmacological effect Anticancer activity	<i>Polygonum hypoleucum</i> <i>Polygonum cuspidatum</i>	Emodin Ethanol extracts Resveratrol Methanol extracts	<i>In vitro</i> <i>In vivo</i> <i>In vivo</i> <i>In vitro</i>	
	<i>Polygonum amphibium</i> <i>Polygonum perfoliatum</i> <i>Polygonum aviculare</i> <i>Polygonum lapathifolium</i>	Glucuronides Methanol extracts Methanol extracts Phenylpropanoid esters of sucrose vanicoside B Lapathoside A	<i>In vitro</i> <i>In vitro</i> <i>In vitro</i> <i>In vivo</i> <i>In vivo</i> <i>In vivo</i>	
	<i>Polygonum tinctorium</i>	Ethyl acetate extracts Tryptanthrin	<i>In vitro</i> <i>In vitro</i>	
	Antiviral activity	<i>Polygonum cuspidatum</i>	Ethanol extracts Water extracts	<i>In vitro</i> <i>In vitro</i>
		<i>Polygonum tinctorium</i> <i>Polygonum viscosum</i>	Water extracts Quercetin 3-O-(6"-feruloyl)- β -D-galactopyranoside Viscoazulone	<i>In vitro</i> <i>In vitro</i> <i>In vivo</i> <i>In vitro</i>
		<i>Polygonum aviculare</i> <i>Polygonum hypoleucum</i>	Ethanol extracts Emodin Emodin-8-O- β -D-glucopyranoside	<i>In vivo</i> <i>In vivo</i> <i>In vivo</i>
	Lipid-regulating activity	<i>Polygonum multiflorum</i> <i>Polygonum cuspidatum</i>	Water extracts Polydatin Resveratrol	<i>In vivo</i> <i>In vivo</i> <i>In vitro</i>
Neuroprotective activity		<i>Polygonum multiflorum</i>	PME-I, PME-II Hexane extracts Ethyl acetate extracts	<i>In vivo</i> <i>In vitro</i> <i>In vitro</i>
	<i>Polygonum cuspidatum</i>	85% methanol extracts	<i>In vitro / in vivo</i>	
Estrogenic activity	<i>Polygonum cuspidatum</i>	Emodin Emodin 8-O- β -D-glucopyranoside	<i>In vitro</i> <i>In vitro</i>	
	<i>Polygonum hydropiper</i> <i>Polygonum hydropiper</i>	Crude root extracts Hexane extracts Ethyl acetate extracts Methanol extracts	<i>In vivo</i> <i>In vivo</i> <i>In vivo</i> <i>In vitro</i>	
	<i>Polygonum amplexicaule</i> <i>Polygonum cuspidatum</i>	Emotion-8-O- β -D-glucoside Anthraquinones	<i>In vitro</i> <i>In vitro</i>	




5,6-dihydropyrano benzopyrone




Catechin




hydroperoside B $R_1 = \text{OCH}_3$, $R_2 = \text{trans-}p\text{-courmaroyl}$, $R_3 = \text{Ac}$, $R_4 = \text{H}$
 vanicode A. $R_1 = R_4 = \text{H}$, $R_2 = \text{trans-}p\text{-courmaroyl}$, $R_3 = \text{Ac}$
 vanicode E. $R_1 = \text{H}$, $R_2 = \text{trans-}p\text{-courmaroyl}$, $R_3 = R_4 = \text{Ac}$




2,3,5,4'-tetrahydroxystilbene-2-O- β -D-glucoside




2''-O-(3,4,5-trihydroxybenzoyl) quercitrin (galloyl quercitrin)




The stilbene glycoside




Gallic acid




Quercetin-3-O- β -D-glucuronide




Polydatin




Resveratrol




Anthraglycoside B




Emodin




Polygodial




Polygosumic acid




Quercetin-3-O- β -glucuronide




Quercetin-3-O- α -rhamnosyl-(1 \rightarrow 2)- β -glucuronide




Vanicoside B R=H
Lapathoside A R=OMe




R=(6''-feruloyl)- β -D-galactopyranosyl
Quercetin3-O-(6''-feruloyl)- β -D-galactopyranoside



Emodin



Emodin-8-O- β -D-glucopyranoside



Emodin-8-O- β -D-glucoside

Fig 1: Some compounds found in the genus *Polygonum*

CONCLUSION

To the best of our knowledge, the chemical composition of the genus *Polygonum* is rich, and a variety of biological activity have also been reported have been studies by modern investigations, such as antioxidation, antimicrobial, anticancer, antivirus, etc (Table 1). While, there are not enough systemic data for the pharmacokinetics and toxicity of the genus *Polygonum*.

This paper dedicated to compiling many active ingredients and intricate mechanisms of genus *Polygonum*. However, many comprehensive mechanisms still need to be investigated. In order to clarify the structure-activity relationship of the genus *Polygonum*. The main chemical constituents and their structures are shown in Fig 1). Considering the fact that the genus *Polygonum* spreading the world, only 27 % of them have been widely investigated, the genus *Polygonum* still remains to be a potential resource to research.

ACKNOWLEDGEMENT

This part of work is supported by the National Natural Science Foundation of China (no. 81173563), and the Compound Chinese Pharmaceutical Innovation Team of Beijing University of Chinese Medicine (no. 2011-CXTD-13). The first two named authors contributed equally to this work.

REFERENCES

1. Wang KJ, Zhang YJ, Yang CR. Recent Advance on the Chemistry and Bioactivity of Genus *Polygonum*. *Nat Prod Res Dev* 2006; 18: 151-164.

2. Yang XQ, Zou YM, Ye JR, Bao ZJ, Ding ZT. Chemical studies on the plants of Polygonum. Yun nan Chemical Technology 2003; 30: 31-33.
3. Mudasir AT, Mohamed MR, Seema A, Ikhlas AK. 5,6-Dihydropyranobenzopyrone: a previously undetermined antioxidant isolated from Polygonum amplexicaule. CJNM 2012; 10: 28-31.
4. Kiem PV, Nhiem NX, Cuong NX, Hoa TQ, Huong HT, Huong LM, Minh CV, Kim YH. New Phenylpropanoid Esters of Sucrose from Polygonum hydropiper and Their Antioxidant Activity. Arch Pharm Res 2008; 31: 1477-1482.
5. Zhao FP, Dieter S, Alfred B, Ramanathan S, Ngoh KG, Tet FC, Tan SN, Chia LS. Antioxidant flavonoids from leaves of Polygonum hydropiper L. Phytochemistry 2003; 62: 219-228.
6. Haraguchi H, Hashimoto K, Yagi A. Antioxidative substances in leaves of Polygonum hydropiper. J Agr Food Chem 1992; 40: 1349-1351.
7. Yagi A, Uemura T, Okamura N, Haraguchi H, Imoto T, Hashimoto K. Antioxidative sulphated flavonoids in leaves of Polygonum hydropiper. Phytochemistry 1994; 35: 885-887.
8. Hsu CY. Antioxidant activity of extract from Polygonum aviculare L. Biol Res 2006; 39: 281-288.
9. Qader SW, Abdulla MA, Chua LS, Sirat HM, Hamdan S. Pharmacological Mechanisms Underlying Gastro protective Activities of the Fractions Obtained from Polygonum minus in Sprague Dawley Rats. Int J Mol Sci 2012; 13: 1481-1496.
10. Chen Y, Wang MF, Robert TR, Ho CT. 2,2-Diphenyl-1-picrylhydrazyl Radical-Scavenging Active Components from Polygonum multiflorum Thunb. J Agr Food Chem 1999; 47: 2226-2228.
11. Lv LS, Gu XH, Tang J, Ho CT. Antioxidant activity of stilbene glycoside from Polygonum multiflorum Thunb in vivo. Food Chem, 2007, 104(4): 1678-1681.
12. Liu LP, Liao ZP, Yin D, Li WD, Liu D, Li Q, Huang QR, Yang YF, He M. The protective effects of Polygonum multiflorum stilbene glycoside preconditioning in an ischemia/reperfusion model of HUVECs. Acta Pharmacol Sin 2010; 31: 405-412.
13. Lee MH, Kao L, Lin CC. Comparison of the Antioxidant and Transmembrane Permeative Activities of the Different Polygonum cuspidatum Extracts in Phospholipid-Based Microemulsions. J Agr Food Chem 2011; 59: 9135-9141.
14. Wang Y, Xu H, Fu Q, Ma R, Xiang JZ. Protective effect of resveratrol derived from Polygonum cuspidatum and its liposomal form on nigral cells in Parkinsonian rats. Neurol Sci 2011; 304: 29-34.
15. Jang HG, Heo BG, Park YS, Namiesnik J, Barasch D, Katrich E, Vearasilp K, Trakhtenberg S, Gorinstein S. Chemical Composition, Antioxidant and Anticancer Effects of the Seeds and Leaves of Indigo (Polygonum tinctorium Ait.) Plant. Appl Biochem Biotechnol 2012; 167: 1986-2004.
16. Imad AE, Fawzia AB, Wissame M, Faycal H, Mohamed AD. Screening of biological activities of Polygonum maritimum L. From Algerian coast. Asian Pacific Journal of Tropical Biomedicine 2013; 3: 611-616.
17. Fan P, Terrier L, Hay AE, Marston A, Hostettmann K. Antioxidant and enzyme inhibition activities and chemical profiles of Polygonum sachalinensis F. Schmidt ex Maxim (Polygonaceae). Fitoterapia 2010; 81: 124-131.
18. Bralley EE, Greenspan P, Hargrove JL, Wicker L, Hartle DK. Topical anti-inflammatory activity of Polygonum cuspidatum extract in the TPA model of mouse ear inflammation. J Inflamm 2008; 5: 1-7.
19. Ghanim H, Sia CL, Abuaysheh S, Korzeniewski K, Patnaik P, Marumqanti A, Chaudhuri A, Dandona P. An Antiinflammatory and Reactive Oxygen Species Suppressive Effects of an Extract of Polygonum Cuspidatum Containing Resveratrol. J Clin Endocr Metab 2010; 95: 1-8.
20. Wu XB, Luo XQ, Gu SY, Xu JH. The effects of Polygonum cuspidatum extract on wound healing in rats. J Ethnopharmacol 2012; 141: 934-937.
21. Han JH, Koh W, Lee HJ, Lee EO, Lee SJ, Kim JT, Khil JH, Jeong SJ, Kim SH. Analgesic and anti-inflammatory effects of ethyl acetate fraction of Polygonum cuspidatum in experimental animals. Immunopharm Immunot 2012; 34: 191-195.
22. Cheng HW, Lee KC, Cheah KP, Chang ML, Lin CW, Li JS, Yu WY, Liu ET, Hu CM. Polygonum viviparum L. inhibits the lipopolysaccharide-induced inflammatory response in RAW 264.7 macrophages through haem oxygenase-1 induction and activation of the Nrf2 pathway. J Sci Food Agr 2013; 93: 491-497.
23. Datta BK, Datta SK, Chowdhury MM, Khan TH, Kundu JK, Rashid MA, Nahar L, Sarker SD. Analgesic, anti-inflammatory and CNS depressant activities of sesquiterpenes and a flavonoid glycoside from Polygonum viscosum. Die Pharm 2004; 59: 222-225.
24. Fan D, Zhou X, Zhao C, Chen HG, Zhao Y, Gong XJ. Anti-inflammatory, antiviral and quantitative study of quercetin-3-O-β-D-glucuronide in Polygonum perfoliatum L. Fitoterapia 2011; 82: 805-810.
25. Duwiejua M, Zeitlin IJ, Gray AL, Waterman PG. The anti-inflammatory compounds of Polygonum bistorta: isolation and characterisation. Planta medica 1999; 65: 371-374.
26. Pandit S, Kim HJ, Park SH, Jeon JG. Enhancement of fluoride activity against Streptococcus mutans biofilms by a substance separated from Polygonum cuspidatum. Biofouling 2012; 28: 279-287.
27. Ban SH, Kwon YR, Pandit S, Lee YS, Yi HK, Jeon JG. Effects of a bio-assay guided fraction from Polygonum cuspidatum root on the viability, acid production and glucosyltransferase of mutans streptococci. Fitoterapia 2010; 81: 30-34.
28. Shan B, Cai YZ, Brooks JD, Corke H. Antibacterial properties of Polygonum cuspidatum roots and their

- major bioactive constituents. *Food Chem* 2008; 109: 530–537.
29. Cecotti R, Carpana E, Falchero L, Paoletti R, Tava A. Determination of the Volatile Fraction of *Polygonum bistorta* L. at Different Growing Stages and Evaluation of Its Antimicrobial Activity against Two Major Honeybee (*Apis mellifera*) Pathogens. *Chem Biodivers* 2012; 9: 359-369.
 30. Alves TMA, Ribeiro FL, Kloos H, Zani CL. Polygodial, the Fungitoxic Component from the Brazilian Medicinal Plant *Polygonum punctatum*. *Mem I Oswaldo Cruz* 2001; 96: 831-833.
 31. Salama HMM, Marraiki N. Antimicrobial activity and phytochemical analyses of *Polygonum aviculare* L. (*Polygonaceae*), naturally growing in Egypt. *Saudi J Biol Sci* 2010; 17: 57-63.
 32. Xv FQ, Liu HB, Luo JG, Zhang JL, Guan HS. Chemical compositions and potency Preliminary Meridian. *Periodical of Ocean University of China* 2010; 40: 101-104.
 33. Begne MG, Yslas N, Reyes E, Quiroz V, Santana J, Jimenez G. Clinical effect of a Mexican *Sanguinaria* extract (*Polygonum aviculare* L.) on gingivitis. *J Ethnopharmacol* 2001; 74: 45-51.
 34. Cai J, Xie S, Feng J, Wang FP, Xu QF. Protective Effect of *Polygonum orientale* L. Extracts against *Clavibacter michiganense* subsp. *sepedonicum*, the Causal Agent of Bacterial Ring Rot of Potato. *PLOS ONE* 2013; 8: 68480.
 35. Liao SG, Zhang LJ, Sun F, Zhang JJ, Chen AY, Lan YY, Li YJ, Wang AM, He X, Xiong Y, et al. Antibacterial and anti-inflammatory effects and fractions from *Polygonum capitatum*. *J Ethnopharmacol* 2011; 134:1 006-1009.
 36. Kataoka M, Hirata K, Kunikata T, Ushio S, Iwaki K, Ohashi K, Ikeda M, Kurimoto M. Antibacterial action of tryptanthrin and kaempferol, isolated from the indigo plant (*Polygonum tinctorium* Lour.), against *Helicobacter pylori*-infected Mongolian gerbils. *J Gastroenterol* 2001; 36: 5-9.
 37. Datta BK, Rahman MM, Gray AI, Nahar L, Hossein SA, Auzi AA, Sarker SD. Polygodosomic acid, a new cadinane sesquiterpene from *Polygonum viscosum*, inhibits the growth of drug-resistant *Escherichia coli* and *Staphylococcus aureus* (MRSA) *in vitro*. *J Nat Med* 2007; 61: 391-396.
 38. Kuo YC, Sun CM, Ou JC, Tsai WJ. A tumor cell growth inhibitor from *Polygonum hypoleucum* Ohwi. *Life Sci* 1997; 61: 2335-2344.
 39. Lin YW, Yang FJ, Chen CL, Lee WT, Chen RS. Free radical scavenging activity and antiproliferative potential of *Polygonum cuspidatum* root extracts. *J Nat Med* 2010; 64: 146-152.
 40. Kimura Y, Okuda H. Resveratrol isolated from *Polygonum cuspidatum* root prevents tumor growth and metastasis to lung and tumor-induced neovascularization in Lewis lung carcinoma-bearing mice[J]. *J Nutr* 2001, 131(6): 1844-1849.
 41. Hu B, An HM, Shen KP, Song HY, Deng S. *Polygonum cuspidatum* Extract Induces Anoikis in Hepatocarcinoma Cells Associated with Generation of Reactive Oxygen Species and Down regulation of Focal Adhesion Kinase. *Evid-Based Compl Alt* 2012; 2012: 1-9.
 42. Shin JA, Shim JH, Jeon JG, Choi KH, Choi ES, Cho NP, Cho SD. Apoptotic effect of *Polygonum Cuspidatum* in oral cancer cells through the regulation of specificity protein 1. *Oral Dis* 2011; 17: 162-170.
 43. Smolarz HD, Budzianowski J, Bogucka-Kocka A, Kocki J, Mendyk E. Flavonoid Glucuronides with Anti-leukaemic Activity from *Polygonum amphibium* L. *Phytochemical Analysis* 2008; 19: 506-513.
 44. Wang KW, Zhu JR, Shen LQ. A new lignan with anti-tumor activity from *Polygonum perfoliatum* L. *Nat Prod Res* 2013; 27: 568-573.
 45. Roudkenar MH, Roushandeh AM, Rad JS. Effects of *polygonum aviculare* herbal extract on proliferation and apoptotic gene expression of MCF-7. *DARU J Pharm Sci* 2011; 19: 326-331.
 46. Takasaki M, Konoshima T, Kuroki S, Tokuda H, Nishino H. Cancer chemopreventive activity of phenylpropanoid esters of sucrose, vanicoside B and lapathoside A, from *Polygonum lapathifolium*. *Cancer Lett* 2001; 173: 133-138.
 47. Koya-Miyata S, Kimoto T, Micallef MJ, Hino K, Taniguchi M, Ushio S, Iwaki K, Ikeda M, Kurimoto. Prevention of azoxymethane-induced intestinal tumors by a crude ethyl acetate-extract and tryptanthrin extracted from *Polygonum tinctorium* Lour. *Anticancer Res* 2001; 21: 3295-3300.
 48. Chang JS, Liu HW, Wang KC, Chen MC, Chiang LC, Hua YC, Lin CC. Ethanol extract of *Polygonum cuspidatum* inhibits hepatitis B virus in a stable HBV-producing cell line. *Antivir Res* 2005; 66: 29-34.
 49. Liu Z, Wei F, Chen LJ, Xiong HR, Liu YY, Luo F, Hou W, Xiao H, Yang ZQ. In Vitro and in Vivo Studies of the Inhibitory Effects of Emodin Isolated from *Polygonum cuspidatum* on Coxsackievirus B4. *Molecules* 2013; 18: 11842-11858.
 50. Zhong Y, Yoshinaka Y, Takeda T, Shimizu N, Yoshizaki S, Inagaki Y. Highly potent anti-HIV-1 activity isolated from fermented *Polygonum tinctorium* Aiton. *Antivir Res* 2005; 66: 119-128.
 51. Datta B, Datta S, Khan T, Kundu J, Rashid M, Nahar L, Sarker S. Anti-cholinergic, cytotoxic and anti-HIV-1 activities of sesquiterpenes and a flavonoid glycoside from the aerial parts of *Polygonum viscosum*. *Pharm Biol* 2004; 42: 18-23.
 52. Sung YY, Yoon T, Yang WK, Kim SJ, Kim DS, Kim HK. The Antiobesity Effect of *Polygonum aviculare* L. Ethanol Extract in High-Fat Evid-based Compl Alt 2013; 2013: 626397.
 53. Chao PM, Kuo YH, Lin YS, Chen CH, Chen SW. The Metabolic Benefits of *Polygonum hypoleucum* Ohwi in HepG2 Cells and Wistar Rats under Lipogenic Stress. *J Agr Food Chem* 2010; 58: 5174-5180.

54. Yang PY, Almofti MR, Lu L, Kang H, Zhang J, Li TJ, Rui YC, Sun LN, Chen WS. Reduction of atherosclerosis in cholesterol-fed rabbits and decrease of expressions of intracellular adhesion molecule-1 and vascular endothelial growth factor in foam cells by a water-soluble fraction of *Polygonum multiflorum*. *J Pharmacol Sci* 2005; 99: 294-300.
55. Xing WW, Wu JZ, Jia M, Du J, Zhang H, Qin LP. Effects of polydatin from *Polygonum cuspidatum* on lipid profile in hyperlipidemic rabbits. *Biomed Pharmacother* 2009; 63: 457-462.
56. Park CS, Lee YC, Kim JD, Kim HM, Kim CH. Inhibitory effects of *Polygonum cuspidatum* water extract (PCWE) and its component resveratrol on acyl-coenzyme A-cholesterol acyltransferase activity for cholesteryl ester synthesis in HepG2 cells. *Vasc Pharmacol* 2004; 40: 279-284.
57. Li X, Matsumoto K, Murakami Y, Tezuka Y, Wu Y, Kadota S. Neuroprotective effects of *Polygonum multiflorum* on nigrostriatal dopaminergic degeneration induced by paraquat and maneb in mice. *Pharmacol, Biochem Be* 2005; 82: 345-352.
58. Jang JY, Kim HN, Kim YR, Choi YW, Choi YH, Lee JH, Shin HK, Choi BT. Hexane extract from *Polygonum multiflorum* attenuates glutamate-induced apoptosis in primary cultured cortical neurons. *J Ethnopharmacol* 2013; 145: 261-268.
59. Kim HN, Kim YR, Jang JY, Choi YW, Baek JU, Hong JW, Choi YH, Shin HK, Choi BT. Neuroprotective effects of *Polygonum multiflorum* extract against glutamate-induced oxidative toxicity in HT22 hippocampal cells. *J Ethnopharmacol* 2013; 150: 108-115.
60. Kim J, Kim MY, Leem KH, Moon S, Jamakattel-Pandit N, Choi H, Kim H, Bu YM. Key compound groups for the neuroprotective effect of roots of *Polygonum cuspidatum* on transient middle cerebral artery occlusion in Sprague-Dawley rats. *Nat Prod Res* 2010; 24: 1214-1226.
61. Matsuda H, Shimoda H, Morikawa T, Yoshikawa M. Phytoestrogens from the Roots of *Polygonum cuspidatum* (Polygonaceae): Structure-Requirement of Hydroxyanthraquinones for Estrogenic Activity. *Bioorg Med Chem Lett* 2001; 11: 1839-1842.
62. Hazarika A, Sarma HN. The estrogenic effects of *Polygonum hydropiper* root extract induce follicular recruitment and endometrial hyperplasia in female albino rats. *Contraception* 2006; 74: 426-434.
63. Rahman E, Goni SA, Rahman, Ahmed M. Antinociceptive activity of *Polygonum hydropiper*. *Fitoterapia* 2002; 73: 704-706.
64. Miyazawa M, Tamura N. Inhibitory Compound of Tyrosinase Activity from the Sprout of *Polygonum hydropiper* L. (Benitade). *Biological and Pharmaceutical Bulletin* 2007; 30: 595-597.
65. Xiang MX, Xu Z, Su HW, Hu J, Yan YJ. Emodin-8-O- β -D-Glucoside from *Polygonum Amplexicaule* D. Don var. *Sinense* Forb. Promotes Proliferation and Differentiation of Osteoblastic MC3T3-E1 Cells. *Molecules* 2011; 16: 728-737.
66. Leu YL, Hwang TL, Hu JW, Fang JY. Anthraquinones from *Polygonum cuspidatum* as tyrosinase inhibitors for demal use. *Phytother Res* 2008; 22: 552-556.