

Phosphate removal from the returned liquor of municipal wastewater treatment plant using iron-reducing bacteria

V. Ivanov¹, V. Stabnikov², W.Q. Zhuang¹, J.H. Tay¹ and S.T.L. Tay¹

¹School of Civil and Environmental Engineering, Nanyang Technological University, Singapore, and ²Department of Microbiology and Biotechnology, National University of Food Technologies, Kiev, Ukraine

2004/0884: received 30 July 2004, revised 11 November 2004 and accepted 12 November 2004

ABSTRACT

V. IVANOV, V. STABNIKOV, W.Q. ZHUANG, J.H. TAY AND S.T.L. TAY. 2005.

Aim: The application of iron-reducing bacteria (IRB) to phosphate removal from returned liquor (liquid fraction after activated sludge digestion and anaerobic sludge dewatering) of municipal wastewater treatment plant (WWTP) was studied.

Methods and Results: An enrichment culture and two pure cultures of IRB, *Stenotrophomonas maltophilia* BK and *Brachymonas denitrificans* MK identified by 16S rRNA gene sequencing, were produced using returned liquor from a municipal WWTP as carbon and energy source, and iron hydroxide as oxidant. The final concentration of phosphate increased from 70 to 90 mg l⁻¹ in the control and decreased from 70 to 1 mg l⁻¹ in the experiment. The mass ratio of removed P to produced Fe(II) was 0.17 g P g⁻¹ Fe(II). The strain *S. maltophilia* BK showed the ability to reduce Fe(III) using such xenobiotics as diphenylamine, m-cresol, 2,4-dichlorophenol and p-phenylphenol as sole sources of carbon under anaerobic conditions.

Conclusions: Bacterial reduction of ferric hydroxide enhanced the phosphate removal from the returned liquor.

Significance and Impact of the Study: The ability of the facultative anaerobes *S. maltophilia* BK and *B. denitrificans* MK to reduce Fe(III) was shown. These micro-organisms can be used for anaerobic removal of phosphate and xenobiotics by bacterial reduction of ferric ions.

Keywords: iron-reducing bacteria, municipal wastewater, phosphate, returned liquor.

ACKNOWLEDGEMENTS

The research was supported by a grant from the Nanyang Technological University, Republic of Singapore. The authors thank Mr Z. Xing for the technical help in the performance of this research.

REFERENCES

- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25, 3389-3402.
- American Public Health Association. (1998a) 3500-Fe B. Phenanthroline method. In *Standard Methods for the examination of Water and Wastewater*, 21st edn ed. Clesceri, L.S., Greenberg, A. A. and Eaton, A.D. pp. 3-76-3-78. Washington, DC: American Public Health Association/American Water Works Association/Water Environmental Federation.
- American Public Health Association. (1998b) 4500-P C. Vanadomolybdophosphoric acid colorimetric method. In *Standard Methods for the examination of Water and Wastewater*, 21st edn ed. Clesceri, L.S., Greenberg, A.A. and Eaton, A.D. pp. 4-412-4-145. Washington, DC: American Public Health Association/American Water Works Association/Water Environmental Federation.
- American Public Health Association. (1998c) 2540 B. Total solids dried at 103–105°C. In *Standard Methods for the examination of Water and Wastewater*, 21st edn ed. Clesceri, L.S., Greenberg, A.A. and Eaton, A.D. pp. 2-55-2-56. Washington, DC: American Public Health Association/American Water Works Association/Water Environmental Federation.
- American Public Health Association. (1998d) 2540 E. Fixed and volatile solids ignited at 550°C. In *Standard Methods for the examination of Water and Wastewater*, 21st edn ed. Clesceri, L.S., Greenberg, A.A. and Eaton, A.D. pp. 2-58-2-59. Washington, DC: American Public Health Association/American Water Works Association/Water Environmental Federation.
- Bink, P.R., Nickin, S. and Bruce, N.C. (1995) Degradation of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) by *Stenotrophomonas maltophilia* PBI. *Appl Environ Microbiol* 61, 1318–1322.
- Boulos, L., Prevost, M., Barbeau, B., Coallier, J. and Desjardins, R. (1999) LIVE/DEAD BacLight: application of a new rapid staining method for direct enumeration of viable and total bacteria in drinking water. *J Microbiol Methods* 37, 77–86.
- Bramucci, M.G. and Nagarajan, V. (2000) Industrial wastewater bioreactors: sources of novel microorganisms for biotechnology. *Trends Biotechnol* 18, 501–505.
- Caccavo, F., Jr (1999) Protein-mediated adhesion of the dissimilatory Fe(III)-reducing bacterium *Shewanella alga* BrY to hydrous ferric oxide. *Appl Environ Microbiol* 65, 5017-5022.
- Cole, J.R., Chai, B., Marsh, T.L., Farris, R.J., Wang, Q., Kulam, S.A., Chandra, S., McGarrell, D.M. et al. (2003) The Ribosomal Database Project (RDP-II): previewing a new autoaligner that allows regular updates and the new prokaryotic taxonomy. *Nucleic Acids Res* 31, 442-443.
- Coleman, M.L., Hedfick, D.B., Lovley, D.R., White, D.C. and Pye, K. (1993) Reduction of Fe(III) in sediments by sulphate-reducing bacteria. *Nature* 361, 436-438.
- Das, A. and Caccavo, F., Jr (2000) Dissimilatory Fe(III) oxide reduction by *Shewanella alga* BrY requires adhesion. *Curr Microbiol* 40, 344-347.
- Drzyzga, O. (2003) Diphenylamine and derivatives in the environment: a review. *Chemosphere* 53, 809-818.
- Eckenfelder, W.W. and Argaman, Y. (1991) Principles of biological and physical/chemical nitrogen removal. In *Phosphorus and Nitrogen Removal from Municipal Wastewater* ed. Sedlak, R. pp. 3-42. New York: Lewis Publishers.
- Felsenstein, J. (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17, 368-376.
- Francis, C.A., Obratsova, A.Y. and Tebo, B.M. (2000) Dissimilatory metal reduction by the facultative anaerobe *Pantoea agglomerans* SP1. *Appl Environ Microbiol* 66, 543-548.
- Fredrickson, J.K. and Gorby, Y.A. (1996) Environmental processes mediated by iron-reducing bacteria. *Curr Opin Biotechnol* 7, 287–294.
- Fytianos, K., Voudrias, E. and Raikos, N. (1998) Modeling of phosphorus removal from aqueous and wastewater sample using ferric iron. *Environ Pollut* 101, 123-130.
- Hall, T.A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41, 95-98.
- Hiraishi, A., Shin, Y.K. and Sugiyama, J. (1995) *Brachymonas denitrificans* gen. nov., sp. nov., an aerobic chemoorganotrophic bacterium which contains ridoquinones, and evolutionary relationships of ridoquinone producers to bacterial species with various quinone classes. *J Gen Appl Microbiol* 41, 99-117.
- Ivanov, V.N., Stabnikova, E.V., Stabnikov, V.P., Kim, I.S. and Zubair, A. (2002) Influence of iron compounds on the treatment of fat-containing wastewaters. *Appl Biochem Microbiol* 38, 255-258.
- Ivanov, V., Wang, J.Y., Stabnikova, O., Krasinko, V., Stabnikov, V., Tay, S.T.L. and Tay, J.H. (2004) Iron-mediated removal of ammonia from strong nitrogenous wastewater of food processing. *Water Sci Technol* 49, 421-431.
- Juhasz, A.L. and Naidu, R. (2000) Enrichment and isolation of non-specific aromatic degraders from unique uncontaminated (plant and faecal material) sources and contaminated soils. *J Appl Microbiol* 89, 642-650.
- Kazumi, J., Haggblom, M.M. and Young, L.Y. (1995) Degradation of monochlorinated and nonchlorinated aromatic compounds under iron-reducing conditions. *Appl Environ Microbiol* 61, 4069–4073.
- Kostka, J.E., Dalton, D.D., Skelton, H., Dollhopf, S. and Stucki, J.W. (2002) Growth of iron (III)-reducing bacteria on clay minerals as the sole electron acceptor and comparison of growth yield on a variety of oxidized iron forms. *Appl Environ Microbiol* 68, 6256-6262.

- Lane, D.J. (1991) 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial Systematics* ed. Stackebrandt, E. and Goodfellow, M. pp. 115-175. Chichester: John Wiley and Sons.
- Lens, P.N.L., O'Flaherty, V., Dijkema, C., Colleran, E. and Stams, A.J.M. (1996) Propionate degradation by mesophilic anaerobic sludge: degradation pathways and effects of other volatile fatty acids. *J Ferment Bioeng* 82, 387-391.
- Liu, W.-T., Chan, O.-C. and Fang, H.H.P. (2002) Characterization of microbial community in granular sludge treating brewery wastewater. *Water Res* 36, 1767-1775.
- Lovley, D.R. (1991) Dissimilatory Fe(III) and Mn(IV) reduction. *Microbiol Rev* 55, 259-287.
- Lovley, D.R. (1993) Dissimilatory metal reduction. *Annu Rev Microbiol* 47, 263-290.
- Lovley, D.R. (2000) Fe(III) and Mn(IV) reduction. In *Environmental Microbe-Metal Interactions*, Vol. 47 ed. Lovley, D.R. pp. 3-30. Washington, DC: ASM Press.
- Lovley, D.R. and Phillips, E.J.P. (1986) Organic matter mineralization with the reduction of ferric iron in anaerobic sediments. *Appl Environ Microbiol* 52, 683-689.
- Lovley, D.R., Phillips, E.J.P. and Lonergan, D.J. (1991) Enzymatic versus nonenzymatic mechanisms for Fe(III) reduction in aquatic sediments. *Environ Sci Technol* 25, 1062-1067.
- Lovley, D.R., Roden, E.E., Phillips, E.J.P. and Woodward, J.C. (1993) Enzymatic iron and uranium reduction by sulfate-reducing bacteria. *Mar Geol* 113, 41-53.
- Luedecke, D., Hermanowicz, S.W. and Jenkins, D. (1989) Precipitation of ferric phosphate in activated sludge: a chemical model and its verification. *Water Sci Technol* 21, 325-337.
- Maidak, B.L., Cole, J.R., Lilburn, T.G., Parker, C.T., Saxman, P.R., Jr, Farris, R.J., Garrity, G.M., Olsen, G.J. et al. (2001) The RDP-II (Ribosomal Database Project). *Nucleic Acids Res* 29, 173-174.
- Maszenan, A.M., Seviour, R.J., Patel, B.K.C., Rees, G.N. and McDougall, B.M. (1997) *Amaricoccus* gen. nov., a Gram-negative coccus occurring in regular packages or tetrads, isolated from activated sludge biomass, and descriptions of *Amaricoccus veronensis* sp. nov., *Amaricoccus tamworthensis* sp. nov., *Amaricoccus macauensis* sp. nov., and *Amaricoccus kaplicensis* sp. nov. *Int J Syst Bacteriol* 47, 727-734.
- Meganck, M.T.L. and Faup, G.M. (1988) Enhanced biological phosphorus removal from wastewaters. In *Biotreatment Systems*, Vol. 3 ed. Wise, D.L. pp. 111-203. Boca Raton, FL: CRC Press.
- Nealson, K.H. and Saffarini, D. (1994) Iron and manganese in anaerobic respiration: environmental significance, physiology, and regulation. *Annu Rev Microbiol* 48, 311-343.
- Nevin, K.P. and Lovley, D.R. (2000) Lack of production of electron-shuttling compounds or solubilization of Fe(III) during reduction of insoluble Fe(III) oxide by *Geobacter metallireducens*. *Appl Environ Microbiol* 66, 2248-2251.
- Nielsen, P.H. (1996) The significance of microbial Fe(III) reduction in the activated sludge process. *Water Sci Technol* 34, 129-136.
- Nielsen, J.L., Juretschko, S., Wagner, M. and Nielsen, P.H. (2002) Abundance and phylogenetic affiliation of iron reducers in activated sludge as assessed by fluorescence *in situ* hybridization and microautoradiography. *Appl Environ Microbiol* 68, 4629-4636.
- Olsen, G.J., Matsuda, H., Hagstrom, R. and Overbeek, R. (1994) FastDNAm1: a tool for construction of phylogenetic trees of DNA sequences using maximum likelihood. *Comput Appl Biosci* 10, 41-48.
- Phelps, C.D. and Young, L.Y. (1999) Anaerobic biodegradation of BTEX and gasoline in various aquatic sediments. *Biodegradation* 10, 15-25.
- Radianingtyas, H., Robinson, G.K. and Bull, A.T. (2003) Characterization of a soil-derived bacterial consortium degrading 4-chloroaniline. *Microbiology* 149, 3279-3287.
- Rousseaux, S., Hartmann, A. and Soulas, G. (2001) Isolation and characterisation of new Gram-negative and Gram-positive atrazine degrading bacteria from different French soils. *FEMS Microbiol Ecol* 34, 211-222.
- Schlöter, M., Lebuhn, M., Heulin, T. and Hartmann, A. (2000) Ecology and evolution of bacterial microdiversity. *FEMS Microbiol Rev* 24, 647-660.
- Stabnikov, V.P., Tay, S.T.L., Tay, J.H. and Ivanov, V.N. (2004) Effect of iron hydroxide on phosphate removal during anaerobic digestion of activated sludge. *Appl Biochem Microbiol* 40, 376-380.
- Taylor, K.G. (1998) Spatial and temporal variations in early diagenetic organic matter oxidation pathways in Lower Jurassic mudstones of eastern England. *Chem Geol* 145, 47-60.
- Tebo, B.M. and Obraztsova, A.Ya. (1998) Sulfate-reducing bacterium grows with Cr(VI), U(VI), Mn(IV), and Fe(III) as electron acceptors. *FEMS Microbiol Lett* 162, 193-198.
- US Environmental Protection Agency (1986) *Quality Criteria for Water 1986*. Washington, DC: US Environmental Protection Agency. Report 440/5-86-001. 453 pp.
- Wang, Q., Kuninobu, M., Ogawa, H.I. and Kato, Y. (1999) Degradation of volatile fatty acids in highly efficient anaerobic digestion. *Biomass Bioenerg* 16, 407-416.
- Weisberg, W., Barns, S., Pelletier, D. and Lane, D. (1991) 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* 173, 697-703.
- Yamamoto-Ikemoto, R., Matsui, S., Komori, T. and Bosque-Hamilton, E.K. (1998) Control of filamentous bulking and interactions among sulfur oxidation-reduction and iron oxidation-reduction in activated sludge using an iron coagulant. *Water Sci Technol* 38, 9-17.
- Zissi, U.S. and Lyberatos, G.C. (1999) Kinetics of growth and aniline degradation by *Stenotrophomonas maltophilia*. *Water Environ Res* 71, 43-49.