



Photo-induced proton gradients for the *in vitro* investigation of bacterial efflux pumps

Alice Verchère, Isabelle Broutin & Martin Picard

Laboratoire de Cristallographie et RMN Biologiques (CNRS UMR8015), Faculté de Pharmacie, Université Paris Descartes, Sorbonne Paris Cité, 4 Avenue de l'Observatoire, 75006 Paris, France.

SUBJECT AREAS:

BIOPHYSICS

BIOCHEMISTRY

MICROBIOLOGY

BIOLOGICAL MODELS

Received

19 January 2012

Accepted

16 February 2012

Published

7 March 2012

Correspondence and requests for materials should be addressed to I.B. (isabelle.broutin@parisdescartes.fr) or M.P. (martin.picard@parisdescartes.fr)

We describe an original activity assay for membrane transport that uses the proton motive force-dependent efflux pump MexAB from *Pseudomonas aeruginosa*. This pump is co-reconstituted into proteoliposomes together with bacteriorhodopsin (BR), a light-activated proton pump. In this system, upon illumination with visible light, the photo-induced proton gradient created by the BR is shown to be coupled to the active transport of substrates through the pump.

Multidrug efflux pumps from gram-negative bacteria are the focus of increasing interest because of their prominent role in the active efflux of xenobiotics out of the bacteria and, by extension, their prominent role in the worrisome problem of bacterial resistance. Efflux transporters are organised as multicomponent systems in which the RND pump (of the so-called Resistance, Nodulation, cell Division family) located in the inner membrane works in conjunction with a periplasmic protein (MFP, Membrane Fusion Protein) and an outer membrane protein (OMP)¹. The cytoplasmic inner membrane protein, which has broad substrate specificity, acts as an energy-dependent pump that is driven by the proton-motive force (pmf). We focus on the OprM/MexA/MexB system from the gram-negative bacterium *Pseudomonas aeruginosa* (Figure 1a).

Although it has recently benefited from significant structural breakthroughs^{2,3}, the efflux pump field still suffers from a critical lack of tools that allow the *in vitro* investigation of transport. This lack of tools arises from the difficulty of generating robust, quantitatively measurable and reproducible proton gradients across the membrane, a pre-requisite for the transporter to function. Initial attempts have been described that involve the use of valinomycin, a potassium-selective ionophore, which makes it possible to generate proton movement within liposomes that are first filled with KCl and subsequently diluted in a KCl-free buffer^{4,5}. We have made recommendations that will allow this procedure to become unambiguous and reproducible⁶, but we have to acknowledge that even with these caveats in mind, the valinomycin assay remains tedious. Hence, we have decided to improve this protocol by adapting a classic of the membrane bioenergetics field, namely the work of Ephraïm Racker and Walther Stoeckenius in which they used the bacteriorhodopsin from the archaeal organism *Halobacter halobium* to generate the photo-induced proton gradients needed to allow for ATP synthesis through the pH-dependent ATP synthase⁷. We have decided to adapt this pioneering work to convert light into the energy needed to activate the RND pump MexB.

Results

Rationale for the assay. In our system, the membrane proteins are reconstituted into liposomes containing 8-hydroxypyrene-1, 3, 6-trisulphonic acid (pyranine), which provides an optical read-out of the pH changes occurring within the vesicles^{8,9}. In Figure 1c, we show the fluorescence spectra of pyranine at different pHs. The pH dependence of the fluorescence is plotted and is shown to be linear over 1 pH unit. The proton gradient photo-induced by the BR is used by MexB to transport Hoechst 33342, a known substrate of the pump. Given that BR incorporation into liposomes favours one direction¹⁰ (proton pumping into the liposome lumen), only those MexB membrane proteins oriented inside-out will be potentially energised by the pH gradient (see Figure 1b).

The rationale for our assay is the following: In a control experiment in which only BR is present in the liposome membrane, the pH is supposed to decrease upon illumination, and hence, the pyranine fluorescence drops. In contrast, when BR is present together with an active and functional pmf-dependent transporter, the protons




Figure 1 | Rationale for the assay. A. Schematic representation of the MexA-MexB-OprM pump from *Pseudomonas aeruginosa*. B. Cartoon representation of liposomes reconstituted with BR (upper scheme) or with both BR and a functional and active membrane transporter (lower scheme). C. Pyranine fluorescence as a function of pH. *left*, excitation spectrum (emission wavelength set to 509 nm) of pyranine measured at pHs ranging from 6.6 to 8.5; *right*, pyranine fluorescence (λ_{ex} 455 nm, λ_{em} 509 nm) as a function of pH. inset: linear correlation between fluorescence and pH between pH 6 and pH 7. The following linear regression was calculated: Δ fluorescence (%) = $65.9 \times \Delta$ pH - 368 ($R^2=0.9946$).

pumped inside the vesicle by BR will be counter-transported, thereby leading to compensation for the liposomal acidification and hence a steady fluorescence signal.

Mandatory controls. Of foremost importance regarding the rationale of our assay is that the proteoliposomes must be as tight as possible. Indeed, one should make sure that the above-mentioned steady signal is due to counter-transport by the pump and not to the passive leakage of protons out of the liposome. We have taken extreme care in the optimisation of the preparation of the liposomes^{6,10}, and we systematically checked that the liposomes reconstituted with both membrane proteins were not significantly leakier than the control liposomes (see *Supplementary information, Figure S1*). Note that the proton gradients generated by the BR dissipate, leading to the eventual restoration of the initial fluorescence level after the liposomes are incubated in the dark; acidification can then be triggered from that point simply by illuminating again the suspension. Provided that the above prerequisites are fulfilled, the activity of the RND pump MexB can be monitored *in vitro*. In Figure 2, we present the averages of the results for six independent measurements.

Investigation of transport. One can but appreciate the sensitivity of the BR-mediated proton pumping procedure. We have defined the reconstitution and measurement conditions such that a pH decrease




Figure 2 | Monitoring of the liposomal acidification upon illumination. Pyranine fluorescence variations, measured as a function of time, were normalised to the corresponding pH variations based on the titration presented Figure 1c.



of approximately 0.3 units occurs upon light-induced proton pumping¹¹, thereby matching the ΔpH postulated to be required for activating the pump⁵. Surprisingly, when the measurement is performed with proteoliposomes containing BR and MexB, the pump seems to provide suboptimal compensation for the BR-mediated acidification: the fluorescence signal ends between that obtained with control liposomes devoid of any protein and that obtained with liposomes reconstituted with BR alone (see Figure 2, trace c versus trace a and trace b). Note that this behaviour is observed irrespective of the presence of substrate, as if MexB were able to pump protons even in the absence of any substrate (a so-called basal, substrate-independent, proton pumping activity), though at a rate seemingly too low to efficiently compensate for the gradient created by the BR. By contrast, when the measurement is performed with liposomes containing BR, the substrate, and both MexB and MexA, one now clearly observe the fluorescence signal expected for coupled proton counter transport: for one proton entering the liposome, one is transported out by the pump (see Figure 2, trace e). This observation is in line with the results of work of Pr Nikaido, who has shown that the MFP was mandatory for transport^{4,5}. Interestingly, when proteoliposomes containing BR, MexA and MexB are prepared beforehand without substrate and subjected to our assay, no coupled activity is observed (see Figure 2, trace f). However, when the same proteoliposomes are subsequently incubated with the substrate in the dark for a sufficient period of time, proton counter transport is detected. It is likely that the incubation makes it possible for the substrate to partition into the membranes of the liposome and then diffuse laterally to eventually reach the pump via its vestibule; hence, transport is made possible.

Discussion

We have designed an activity assay that allows the monitoring of the activity of RND efflux pumps. Our results might well shed some light on the controversy regarding the possible implication of the MFP in transport. In agreement with the results of previous work by Nikaido and Aires, we show here that MexA is a key element of the actual coupled transport through the efflux pump. In contrast, the Venter's group showed that MexB is able to transport Hoechst 33342 alone, in the absence of MexA¹². The difference between our respective results might very well arise from the different experimental procedures used in our respective laboratories. Indeed, in our protocol, inside-out reconstitution entraps the whole functional complex, and the resulting confinement increases the probability of interaction between MexB and its MFP partner. In contrast, Venter and colleagues performed a so-called "direct incorporation" experiment in which MexB is inserted unidirectionally, right-side out. Under such conditions, functional assembly will be more difficult to achieve because the MFP will have to explore a much larger volume in that case than in our inside-out liposomes.

We show that MexB behaves as if it were able to sustain a basal proton pumping activity that is potentiated by the presence of substrate. This behaviour is reminiscent of that of proteins in the functionally related ABC protein family, which exhibit a basal ATPase activity that is enhanced by the presence of substrates¹³. Interestingly, it has been shown recently that MacB, an ABC transporter from *E. coli* that works in conjunction with a MFP and an OMP, exhibits basal ATPase activity that is enhanced in the presence of its cognate MFP protein MacA, which is closely related to MexA^{14,15}. Note that we cannot exclude the possibility that what we believe to be substrate-free conditions are in fact not; it is possible that the system is contaminated by some endogenous compound (e.g., lipid, left-over from the purification) that can be transported by the pump.

Our protocol opens the way towards a better understanding of the activity of the RND efflux pump MexB and, more generally, of all

pmf-activated membrane proteins. However, we also envisage using this protocol for screening Efflux Pump Inhibitors (EPIs) because our protocol can be easily scaled down, automatised (e.g., in a 96-well plate reader equipped with relevant optical filters) and parallelised: in this context, the use of BR as a switch for triggering the measurement is certainly an asset because it eliminates any second-hand manipulation of the suspension.

Methods

Materials. 1,2-Dioleoyl-sn-glycero-3-phosphocholine (DOPC) was purchased from Avanti Polar Lipids. Cholesterol, 8-hydroxypyrene-1,3,6-trisulphonic acid (pyranine), Triton X-100, and Hoechst 33342 were obtained from Sigma Aldrich. SM₂ Bio-beads were obtained from Bio-Rad. n-Doodecyl- β -D-maltopyranoside (DDM) was purchased from Affymetrix (reference D310LA).

Preparation of the proteins. Expression and purification of the MexA, MexB and BR proteins were performed as previously described^{16,17,18}. Briefly, MexA and MexB were heterologously expressed from pBAD-mexA or pET-mexB (pBAD33-GFPuv, Invitrogen; pET22b vector, Novagen) in *E. coli* cells (strain C43 DE3) and purified as His-tagged proteins from the membranes solubilised with DDM. The yields of homogeneously purified MexA and MexB were approximately 6–7 mg and 0.5 mg per litre of culture, respectively. The purified proteins were concentrated using Vivaspin (VivaScience) concentrators with cutoffs of 30 kDa and 100 kDa for MexA and MexB, respectively.

Halobacter halobium cells were grown under illumination at 37°C in a liquid growth medium containing NaCl 4 M, MgSO₄ 150 mM, trisodium citrate 10 mM, KCl 30 mM, yeast extract 5 g/L, and peptone 5 g/L. Purple membrane was isolated using a sucrose gradient and solubilised as follows: a membrane suspension containing 15–20 mg BR at 7 g/L was sonicated for 5 min and incubated overnight at room temperature with Triton X-100 at a 1:5 protein-to-detergent ratio (w/w). The concentration of solubilised BR was estimated using ϵ (570 nm) = 54.000 M⁻¹.cm⁻¹ and ϵ (280 nm) = 1.080.000 M⁻¹.cm⁻¹.

Preparation of unilamellar vesicles. DOPC and cholesterol were dissolved in HEPES 25 mM pH 7, K₂SO₄ 100 mM and MgCl₂ 15 mM to final concentrations of 10 mg/mL and 1.5 mg/mL, respectively. The suspension was then heated for 10 min at 37°C. Then, pyranine and Hoechst 33342 were added to final concentrations of 2 mM and 140 μ M, respectively. This solution was then sonicated for 10 minutes with 30'' pulse/30'' pause cycles. The liposomes were subsequently extruded through 200-nm membranes and then through 100-nm membranes (20 passes for each type of membrane). Triton X-100 was then added to reach a 2.8 detergent/lipid ratio (w/w). This suspension was then incubated overnight at 4°C with gentle stirring. Proteins were added to the solubilised liposome suspension at the following ratios (w/w): lipids/MexB=20, MexB/MexA=2.5, and lipids/BR=30. Detergent removal was achieved upon addition of Bio-beads (previously washed with methanol, ethanol and water) at a Bio-bead/detergent ratio of 60 (w/w) for at least 3 h at 4°C in the dark. The proteoliposomes were then purified using a PD-10 desalting column (GE Healthcare) equilibrated with HEPES 25 mM pH 7, K₂SO₄ 100 mM and MgCl₂ 15 mM and kept in the dark at 18°C. The proteoliposomes were stored for up to 2–3 weeks under these conditions.

Fluorescence assay. Fluorescence measurements were conducted at 25.0°C using a JASCO FP-6200 spectrofluorimeter. The measurements were performed using the dual-wavelength mode, taking measurements with excitation/emission wavelengths of 550 nm/550 nm for 18'' (to activate BR) and then taking measurements with excitation/emission wavelengths of 455 nm/509 nm for 2'' (to measure the pyranine fluorescence). The excitation and emission bandwidths were 5 nm. Note that the measurements were performed in the presence of 50 nM valinomycin to prevent the formation of a reverse $\Delta\Psi$.

1. Nikaido, H. Multidrug Resistance in Bacteria. *Annu. Rev. Biochem.* **78**, 119–146 (2009).
2. Seeger, M. A. *et al.* Structural Asymmetry of AcrB Trimer Suggests a Peristaltic Pump Mechanism. *Science*. **313**, 1295–1298 (2006).
3. Murakami, S., Nakashima, R., Yamashita, E., Matsumoto, T. & Yamaguchi, A. Crystal structures of a multidrug transporter reveal a functionally rotating mechanism. *Nature*. **443**, 173–179 (2006).
4. Zgurskaya, H. I. & Nikaido, H. Bypassing the periplasm: reconstitution of the AcrAB multidrug efflux pump of *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 7190–7195 (1999).
5. Aires, J. R. & H. Nikaido. Aminoglycosides are captured from both periplasm and cytoplasm by the AcrD multidrug efflux transporter of *Escherichia coli*. *Journal of Bacteriology*. **187**, 1923–1929 (2005).
6. Picard, M., Verchère, A. & Broutin, I. Monitoring the active transport of efflux pumps after their reconstitution into proteoliposomes: caveats and keys. *Analytical Biochemistry* **420**, 194–196 (2012).
7. Racker, E. & Stoerkenius, W. Reconstitution of purple membrane vesicles catalyzing light-driven proton uptake and adenosine triphosphate formation. *J. Biol. Chem.* **249**, 662–663 (1974).



8. Clement, N. R. & J. M. Gould. Pyranine (8-hydroxy-1,3,6-pyrenetrisulfonate) as a probe of internal aqueous hydrogen ion concentration in phospholipid vesicles. *Biochemistry* **20**, 1534–1538 (1981).
9. Sedgwick, E. G. & P. D. Bragg. Differential movement of ions in artificial phospholipid vesicles. *FEBS Letters*. **272**, 81–84 (1990).
10. Rigaud, J.-L. & D. Lévy. Reconstitution of membrane proteins into liposomes. *Meth. Enzymol.* **372**, 65–86 (2003).
11. Seigneuret, M. & Rigaud J.-L. Analysis of Passive and Light-Driven Ion Movements in Large Bacteriorhodopsin Liposomes Reconstituted by Reverse-Phase Evaporation. 1. Factors Governing the passive proton permeability of the membrane. *Biochemistry* **25**, 6716–6722 (1986).
12. Welch, A., Awah, C. U., Jing, S., Van Veen, H. & Venter, H. Promiscuous partnering and independent activity of MexB, the multidrug transporter protein from *Pseudomonas aeruginosa*. *Biochem. J.* **430**, 355–364 (2010).
13. Al-Shawi MK. Catalytic and transport cycles of ABC exporters. *Essays Biochem.* **50**, 63–83 (2011).
14. Tikhonova, E. B., Devroy, V. K., Lau, S. Y. & Zgurskaya, H. I. Reconstitution of the *Escherichia coli* macrolide transporter: the periplasmic membrane fusion protein MacA stimulates the ATPase activity of MacB. *Mol Microbiol.* **63**, 895–910 (2007).
15. Modali, S. D. & Zgurskaya, H. I. The periplasmic membrane proximal domain of MacA acts as a switch in stimulation of ATP hydrolysis by MacB transporter. *Mol Microbiol.* **81**, 937–951 (2011).
16. Akama, H. Crystal Structure of the Membrane Fusion Protein, MexA, of the Multidrug Transporter in *Pseudomonas aeruginosa*. *J. Biol. Chem.* **279**, 25939–25942 (2004).
17. Mokhonov, V. *et al.* Multidrug transporter MexB of : overexpression, purification, and initial structural characterization. *Protein Expression and Purification.* **40**, 91–100 (2005).
18. Oesterhelt, D. & Stoekenius, W. Isolation of the cell membrane of *Halobacterium halobium* and its fractionation into red and purple membrane. *Meth. Enzymol.* **31**, 667–678 (1974).

Acknowledgments

We thank Manuela Dezi and Daniel Lévy for helpful discussions, and Clara Jaquet for the initial fluorescence measurements. This work was supported by a grant from Région Ile-de-France (DIM-Malinf) and by the Agence Nationale de la Recherche (project ANR-07-BLAN-0263).

Author contributions:

MP conceived the experiments. AV, IB and MP wrote the manuscript. AV performed the experiments.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

License: This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

How to cite this article: Verchère, A., Broutin, I. & Picard, M. Photo-induced proton gradients for the *in vitro* investigation of bacterial efflux pumps. *Sci. Rep.* **2**, 306; DOI:10.1038/srep00306 (2012).