

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

X-641-68-480

PREPRINT

NASA TM X-63411

**PHOTODISINTEGRATION OF ULTRAHIGH  
ENERGY COSMIC-RAYS  
BY  
THE UNIVERSAL RADIATION FIELD**

**F. W. STECKER**

**DECEMBER 1968**



**GODDARD SPACE FLIGHT CENTER  
GREENBELT, MARYLAND**

**N 69-17996**

(ACCESSION NUMBER)

(THRU)

11

(PAGES)


(CODE)

29

(CATEGORY)

FACILITY FORM 602

**NASA-TMX-63411**  
(NASA CR OR TMX OR AD NUMBER)



X-641-68-480  
Preprint

PHOTODISINTEGRATION OF ULTRAHIGH ENERGY  
COSMIC-RAYS BY THE UNIVERSAL  
RADIATION FIELD

F. W. Stecker  
Theoretical Studies Branch

December 1968

NASA GODDARD SPACE FLIGHT CENTER  
Greenbelt, Maryland

PHOTODISINTEGRATION OF ULTRAHIGH ENERGY  
COSMIC-RAYS BY THE UNIVERSAL  
RADIATION FIELD

F. W. Stecker  
Theoretical Studies Branch  
NASA Goddard Space Flight Center  
Greenbelt, Maryland

ABSTRACT

We have carried out a detailed calculation to determine the effectiveness of the universal radiation field in disintegrating ultrahigh energy cosmic-ray nuclei. We conclude that this process cannot explain the air-shower observations above  $10^{17}$  eV unless there exists a large metagalactic flux of infrared photons as may have recently been detected. Other implications of our results are discussed.

PHOTODISINTEGRATION OF ULTRAHIGH ENERGY  
COSMIC-RAYS BY THE UNIVERSAL  
RADIATION FIELD

In a recent paper<sup>1</sup>, we made a detailed study of the effect of photomeson production by the universal radiation field on ultrahigh energy cosmic-rays. Greisen<sup>2</sup> and Zatsepin and Kuz'min<sup>3</sup>, who first proposed the significance of this effect, also pointed out that the universal radiation field can disintegrate cosmic-ray nuclei of  $10^{19}$  eV energy and above. The purpose of this letter is to point out that by taking account of the details of the photodisintegration process, the recent air-shower data can be examined for implications on the origin and propagation time of the ultrahigh energy cosmic-rays.

In Reference 1, we showed in a detailed calculation that cosmic-ray protons of energies less than or equal to  $6 \times 10^{19}$  eV can exist for  $10^{10}$  years (the age of the universe) against attenuation by photomeson production. It was also shown that the lifetime of a  $10^{20}$  eV proton is of the order of  $10^9$  years and that protons of all energies have lifetimes of at least  $5 \times 10^7$  years, long enough to reach us if produced within the local supercluster region.

A proton of  $10^{20}$  eV energy was detected by Linsley<sup>4</sup> along with the detection of six other protons having energies greater than or equal to  $2 \times 10^{19}$  eV<sup>5</sup>. Recently, Andrews, et al.<sup>6</sup> have detected a proton of energy  $\geq 5 \times 10^{19}$  eV. As

we concluded in Reference 1, such observations are not incompatible with the existence of the universal blackbody radiation field, but only with the implicit assumption that ultrahigh energy cosmic-rays are primordial. These assumptions are compatible with the existence of these cosmic-rays with ages up to  $10^9$  years, and which may be reaching us from distances as great as 300 Mpc.

We now present a similar discussion of the implications of photodisintegration by the universal radiation field. Letting  $\epsilon'$  denote the energy of a blackbody photon in the rest frame of an ultrahigh energy cosmic-ray nucleus, and letting  $\sigma(\epsilon')$  denote the cross section of a nucleus of type i for photodisintegration, the lifetime of the nucleus is given by (cf. reference 1)

$$\tau_i(E_i) = 2\gamma_i^2 \hbar^3 \pi^2 c^2 \left[ \int_{\epsilon'_{th}/2\gamma_i}^{\infty} \frac{d\epsilon}{e^{\epsilon/kT} - 1} \int_{\epsilon'_{th}}^{2\gamma\epsilon} d\epsilon' \epsilon' \sigma_i(\epsilon') \right]^{-1} \quad (1)$$

In previous discussions<sup>2, 3, 7</sup>, photodisintegration has been stressed for iron nuclei where the cross section is large. However, recent observations<sup>5, 8</sup> indicate that cosmic-rays of energies  $\geq 10^{17}$  eV are purely protons. Such observations can be explained by photodisintegration only if all heavier nuclei can be broken down into individual nucleons. As one considers lighter and lighter nuclei, the photodisintegration cross section becomes smaller and smaller. Indeed<sup>9</sup>, the photodisintegration cross section of a nucleus of atomic number A is proportional to  $A^{4/3}$ . Therefore, the question posed here is whether even the

lightest nuclei can be broken down completely into their constituent nucleons. We thus consider here in detail, the photodisintegration of helium nuclei. We consider in particular, the well studied processes



and



keeping in mind the fact that reactions (2) and (3) must be followed by the further breakup of  $\text{He}^3$ . Thus, lifetimes for the complete breakup of He nuclei are at least twice as long as those calculated here.

The data on reactions (2) and (3) have been compiled by Gorbunov<sup>10</sup>, as shown in Figure 1. Equation (1) was solved numerically making use of these cross sections and the results are given in Figure 2 along with the lifetimes of protons against attenuation by photomeson production as given in Reference 1. We have also calculated the lifetimes for processes (2) and (3) as generated by dilute metagalactic starlight photons with a mean temperature of 5000°K and a dilution factor of  $2.1 \times 10^{-15}$  as given by Allen<sup>11</sup>. Recent calculations by Garmire<sup>12</sup> have yielded even lower estimates of the metagalactic starlight photon density. Such optical photon densities cannot produce complete photodisintegration even in  $10^{10}$  years.

One can see from Figure 2 that complete photodisintegration by the universal radiation field is effective only above  $10^{19}$  eV even for propagation times of the order of  $10^{10}$  years. Considering more likely propagation times as discussed in Reference 1, we find complete photodisintegration effective only above energies of  $2-3 \times 10^{19}$  eV. Thus, photodisintegration by microwave and dilute optical metagalactic photons cannot account for the absence of multinucleon nuclei between  $10^{17}$  and  $10^{19}$  eV.

It may still be possible to obtain adequate photodisintegration for complete breakup if a large flux of infrared photons exists in metagalactic space. Recent preliminary observations between 0.4 and 1.3 mm wavelengths by Shivanandan, et al.<sup>13</sup> indicate the presence of a photon flux 2 orders of magnitude higher than to be expected from a 3°K blackbody background. If this flux is universal, it may help account for the absence of multinucleon nuclei above  $10^{17}$  eV.

If the extragalactic ultrahigh energy cosmic-rays cannot undergo complete photodisintegration during propagation, then we must look for an alternative explanation. The most natural alternative which comes to mind, is the possibility that these cosmic-rays were exposed to a strong photon field at their source. This may have been a likely possibility during the length of time when they underwent acceleration. Indeed, powerful radiation fields may have been associated with, or even caused the acceleration process. The possibility of acceleration of cosmic-rays by radiation has been suggested by Tsytoovich.<sup>14</sup>



## ACKNOWLEDGMENTS

The author would like to acknowledge and thank Dr. Maurice M. Shapiro for suggesting this topic for investigation. Programming of the numerical computations was performed by Mr. Joseph Bredekamp.

## REFERENCES

1. Stecker, F. W., Phys. Rev. Letters 21, 1016 (1968).
2. Greisen, K., Phys. Rev. Letters 16, 748 (1966).
3. Zatsepin, G. T., and V. A. Kuz'min, JETP Letters 4, 114 (1966).
4. Linsley, J., Phys. Rev. Letters 10, 146 (1963).
5. Linsley, J., Phys. Rev. Letters 9, 123 (1962).  
Proc. Int. Conf. Cos. Rays, Jaipur 4, 77 (1963).
6. Andrews, D., A. C. Evans, R. J. O. Reid, R. M. Tennent, A. A. Watson and  
J. G. Wilson, Nature 219, 343 (1968).
7. Gerasimova, N. M., and I. L. Rozental', Sov. Phys. JETP 14, 350 (1962).
8. Cowsik, R., Can. Jour. Phys. 46, S142 (1968).
9. Blatt, J. M., and V. F. Weisskopf, Theoretical Nuclear Physics, Wiley, New  
York, 1952.
10. Gorbunov, A. N., Phys. Letters 27B, 436 (1968).
11. Allen, C. W., Astrophysical Quantities, 2nd ed., Althone Press, London,  
1963.

12. Garmire, G., Proc. Int. Conf. Cos. Rays, London 1, 315 (1965).
13. Shivanandan, K., J. R. Houck and M. O. Harwit, Phys. Rev. Letters 21,  
1460 (1968).
14. Tsytovich, V. N., Sov. Phys. Doklady 7, 43 (1962).




Figure 1. Total cross section for the processes  $\text{He}^4(\gamma, p)\text{H}^3$  and  $\text{He}^4(\gamma, n)\text{He}^3$  as a function of gamma-ray energy in the helium rest system.




Figure 2. Characteristic lifetimes for helium nuclei against photodisintegration and for protons against photomeson production as a function of energy.

## FIGURE CAPTIONS

- Figure 1.** Total cross section for the processes  $\text{He}^4(\gamma, p)\text{H}^3$  and  $\text{He}^4(\gamma, n)\text{He}^3$  as a function of gamma-ray energy in the helium rest system.
- Figure 2.** Characteristic lifetimes for helium nuclei against photodisintegration and for protons against photomeson production as a function of energy.