



King's Research Portal

DOI:

[10.1080/09638288.2017.1306587](https://doi.org/10.1080/09638288.2017.1306587)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Vancampfort, D., Mugisha, J., Richards, J., De Hert, M., Probst, M., & Stubbs, B. (2017). Physical activity correlates in people living with HIV/AIDS: a systematic review of 45 studies. *Disability and Rehabilitation*, 1-12. <https://doi.org/10.1080/09638288.2017.1306587>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

DISABILITY & REHABILITATION

Physical Activity Correlates in People Living with HIV/AIDS: A Systematic Review of 45 Studies

Davy Vancampfort^{a,b}, James Mugisha^{c,d}, Justin Richards^e, Marc De Hert^b, Michel Probst^a, Brendon Stubbs^{f,g}

^aKU Leuven – University of Leuven, Department of Rehabilitation Sciences, Leuven, Belgium

^bKU Leuven – University of Leuven, University Psychiatric Center KU Leuven, Leuven-Kortenberg Belgium

^cButabika National Referral and Mental Health Hospital, Kampala, Uganda

^dKyambogo University, Kampala, Uganda

^eSchool of Public Health & Charles Perkins Centre, University of Sydney, Sydney, NSW, Australia ^fPhysiotherapy Department, South London and Maudsley

NHS Foundation Trust, London, UK

^gHealth Service and Population Research Department, King's College London, De Crespigny Park, London, UK

Running title: HIV and physical activity

*Corresponding author: Davy Vancampfort. UPC KU Leuven, campus Kortenberg, KU Leuven – University of Leuven, Leuvensesteenweg 517, 3070 Kortenberg, Belgium; Tel.: +32 2 758 05 11; Fax: +32 2 759 9879. E-mail address: davy.vancampfort@kuleuven.be

Abstract

Purpose: Understanding barriers and facilitators of physical activity participation in persons living with HIV/AIDS is an essential first step in order to devise effective interventions. The present review provides a systematic quantitative review of the physical activity correlates in people with HIV/AIDS.

Methods: Major electronic databases were searched till August 2016. Keywords included 'physical activity' or 'exercise' or 'sports' and 'AIDS' or 'HIV'.

Results: Out of 55 correlates from 45 studies (N=13,167; mean age range=30.5-58.3years; 63.2% male) 5 consistent (i.e. reported in 4 or more studies) correlates were identified. Lower levels of physical activity were consistently associated with older age (6/10 studies), a lower educational level (6/7), a lower number of CD4 cells/ μ l (7/11), exposure to antiviral therapy (4/6) and the presence of lipodystrophy (4/4). Other important barriers were the presence of bodily pain (2/2), depression (3/3) and opportunistic infections (3/4). Facilitators were a higher cardiorespiratory fitness level (3/3), a higher self-efficacy (2/2), more perceived benefits (2/2) and a better health motivation (3/3).

Conclusions: The current review has elucidated that participation in physical activity by people with HIV/AIDS is associated with a range of complex factors which should be considered in rehabilitation programs.

Keywords: physical activity; exercise; physiotherapy; AIDS; HIV

Introduction

The Global Burden of Disease Expert Group estimated that approximately 30 million people were infected with HIV worldwide in 2013, the majority of whom reside in sub-Saharan Africa [1]. Life expectancy in those infected with HIV has improved significantly since the introduction of effective antiretroviral therapy. Between 1990 and 2013, antiviral therapy saved approximately 19 million life-years in the HIV-infected population [1]. Whilst the longevity in life expectancy has increased among people living with HIV/AIDS, people are tending to live with many more chronic diseases [2]. In people living with HIV/AIDS these chronic comorbidities develop earlier and more frequently than in non-HIV-infected counterparts [3]. The most prevalent comorbidities are cardiovascular diseases, chronic obstructive pulmonary disease, cancers, arthritis, osteoporosis, and liver disease [4]. Current dominant hypotheses link the increased incidence of these comorbidities to HIV-related chronic inflammation [5], adipose tissue abnormalities [6], side-effects of antiviral treatment [7, 8] and modifiable lifestyle factors, such as increased tobacco use [9] and decreased physical activity [10]. The disability caused by these chronic conditions may be exacerbated or alleviated by

intrinsic (personal attributes) and extrinsic (social support, stigma) contextual factors and impacts also the mental health and quality of life of people living with HIV/AIDS [11].

Self-management strategies, such as physical activity and exercise can address disability and optimize mental and physical health outcomes for people living with HIV/AIDS. While physical activity can be defined here as any bodily movement produced by skeletal muscles that results in energy expenditure, exercise is a planned, structured, repetitive and purposive form of physical activity that aims to improve or maintain one or more components of physical fitness [12]. Regular physical activity is considered to contribute to an improved body composition, cardiorespiratory and muscular fitness, lower levels of depressive symptoms and an improved physical and mental health related quality of life [13, 14, 15, 16, 17, 18]. It is therefore strongly recommended that people living with HIV/AIDS augment their physical activity levels, while adequate information and awareness is spread among service users and health care professionals, and corresponding opportunities for physical activity are built and maintained [19].

Despite the observed physical and mental benefits, a large proportion of people living with HIV/AIDS are still not engaging in physical activity on a regular basis [10]. Understanding barriers and facilitators of participation in physical activity in people living with HIV/AIDS is an essential first step in order to devise effective physical activity interventions. Behavioral theories, such as the socio-ecological model [20] have shown to be useful in attempting to understand the factors which influence physical activity behavior in vulnerable populations [21, 22, 23, 24, 25]. Socio-ecological models posit that multiple relevant attributes influence health behavior. These include intrapersonal (demographic, biological, psychological, emotional and cognitive), interpersonal/cultural (e.g., social support), physical environment (e.g., distance to the facilities, financial costs, enjoyable scenery), and policy (laws, rules, regulations, codes) factors [20]. Qualitative research in people living with HIV/AIDS exploring the perceived barriers for physical activity participation indicated that physical exertion, lack of social support, time and financial constraints and the distance to facilities are considered the most important and reported barriers [26, 27, 28]. Also uncertainty about the future [28], avoiding stigma [29], adverse weather conditions and domestic abuse and crime [27] are reported in this population. People living with HIV/AIDS participating in physical activity programs recognized as well that engaging in regular physical activity could provide energy, improve the self-concept and sleeping quality, and increase physical fitness and mental health [27, 29]. In order to elaborate and confirm such qualitative findings, quantitative research

which is able to identify potential correlates of actual physical activity levels in people living with HIV/AIDS is also needed. This information can then be used to target future physical activity interventions for people living with HIV/AIDS. A systematic review on physical activity correlates in people living with HIV/AIDS is however currently lacking. Systematic quantitative research of potential negative and positive correlates of physical activity will provide valuable information to implement physical activity in clinical practice and will inform future research. The present review therefore systematically evaluates published quantitative studies on correlates of physical activity in people living with HIV/AIDS. In addition to summarizing methods and results of these studies, gaps in the literature are identified and directions for future research are proposed.

Methods

This systematic review was conducted in accordance with the “ Meta-Analyses and Systematic Reviews of Observational Studies” - guidelines [30].

Data sources and searches

Two independent reviewers performed an electronic search of the health-related databases PubMed, CINAHL and Embase until August 1st 2016. Manual searches were also conducted using the reference lists from identified articles. The medical subject headings used were ‘physical activity’ OR ‘exercise’ OR ‘sports’ AND ‘HIV’ OR ‘AIDS’ in the title, abstract or index term fields.

Eligibility Criteria

Inclusion criteria were as follows: (a) a diagnosis of HIV or AIDS irrespective of the assessment method used, (b) participants were at least 18 years of age, (c) studies contained quantitative research and had been published in a peer-reviewed journal, (d) the dependent variable was a measure of physical activity participation. No restriction was placed on the selection of the outcome measure or the language of the article. For cohort or intervention studies, only associations of physical activity participation with baseline data were included. We excluded articles if the dependent variable was aerobic fitness, physical activity intention, self-efficacy, or other intermediate (non-behavioral) measures because these variables are less direct indicators of actual physical activity behavior [12]. Also, case reports and expert opinions were excluded.

Data Collection

Two reviewers independently extracted data from the included studies using a predetermined form. The form captured data in 6 domains including (a) gender, (b) age (mean), (c) ethnicity (%White), (d) the quality of the physical activity measure, and (e) physical activity correlates. The following categories were used to code the quality of the physical activity measure: (a) self-report with poor, unknown or not reported reliability/validity in people with HIV, (b) self-report with reported and acceptable reliability/validity in persons with HIV, and (c) acceptable objective measurements for people with HIV. Objective measurements included motion sensors such as accelerometers and pedometers, combined heart rate and accelerometer devices and the doubly labeled water method [20]. The acceptability of the psychometric properties of measurement tools was assessed according to previous recommendations [31]. In accordance with previous physical activity correlates reviews [21, 22, 23, 32, 33, 34] the following potential physical activity correlate categories were included: (a) demographic, (b) biological, (c) psychological / cognitive / emotional, (d) behavioral attributes/skills, (e) social/cultural factors, (f) physical environment, and (g) policy factors. Variables were classified as 'related' or 'not related' to physical activity based on statistical significance, and the direction of association for related variables

was identified. The detailed data tables were further analyzed (see § 2.4. and 2.5) to create tables that summarized the state of the literature on different variables.

Coding Associations with Physical Activity

A variety of statistical techniques were used to evaluate correlates, including uni-/ bivariate analyses, correlations, t-tests, and ANOVA. If both uni-/bivariate and multivariate tests were conducted, uni-/bivariate tests were reported for consistency across studies. The column 'related to physical activity' indicates, which studies reported significant associations between the variable and the physical activity measure. Direction of association is indicated with a '+' or '-'. The column 'unrelated to physical activity' indicates which studies reported non-significant associations between the variable and physical activity.

Summary Codes

A summary code for each variable was presented and calculated following previous recommendations [35, 36]. The summary code column contains a code to summarize the state of the literature for that variable. The percentages refer to the number of associations supporting the expected association divided by the total number of associations for the variable. In accordance with previous physical activity correlates reviews [21, 22, 23, 32, 33, 34], associations were coded with: '0' (0-33% of studies supporting association); '?' (34%-59% of studies supporting an association); or '+' or '-' (60%-100% of studies supporting an association). When correlates were reported in 4 or more studies the summary code for these correlates were considered as 'consistent' and coded with '++', '- -' or '??' .

Differences in Number of Significant Correlates


Using Fisher's exact tests, we explored differences in the number of significant correlates versus unrelated variables obtained via valid physical activity assessments versus assessments with unknown validity versus objective tools and between associations explored in studies with a sample size lower than versus equal to or larger than the median sample size.

Results

Study Selection

Out of 84 potentially eligible studies, 45 were included in this review [37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80]. The search strategy and reasons for exclusion are shown in Figure 1.

Figure 1. Flow Diagram of the Included and Excluded Studies



Participant and Study Characteristics

Across all 45 studies, a total of 13,176 people living with HIV/AIDS (mean age range=30.5-58.2years) were included in the analyses. Thirty-four studies had a cross-sectional design, ten were cohort studies and one was a randomized controlled trial. In 42 studies reporting on the gender distribution, the percentage of male participants was 63.2% (n=8,268/13,080). The sample size ranged from 21 to 2,864. The median sample size was 107. The majority of the studies were executed in the North-American (N=26), followed by South-America (N=6). There were 5 African and 5 European studies and one study was executed in Asia (China) and one in Australia. Concerning the quality of the physical activity measure, 27 studies were based on un-validated or unreliable self-report measures of physical activity or measures with unknown psychometric properties, 11 on validated and reliable self-report measures for people with HIV/AIDS and 7 studies used an objective measure of physical activity. Table 1 presents the characteristics of the included participants, the quality of physical activity assessments and the statistical analyses undertaken.

Table 1. Study Characteristics.

Nr	First author / year	Design	Country	Participants	PA measurement	Quality of PA Measurement	Statistical Analysis
1	Dirajlal-Fargo 2015 [80]	RCT	USA	72 (58♂); median age=45; 69% African American in the Rosuvastin group and 75 (57♂); median age=47; 67% African American in the placebo group	Adult AIDS Clinical Trials Group PA assessment	A	Regression analysis
2	Fazeli 2015 [79]	Cross-sectional	USA	100(88♂); mean age=58.2±6.5years; 82% White	IPAQ	B	Chi square analyses
3	Monroe 2015 [78]	Cross-sectional	USA	596♂; mean age=51years; 54% White, 17% African American, 29% Hispanic/other ; BMI=25.3	IPAQ	B	Adjusted quantile regression modeling
4	Olsen 2015 [77]	Cross-sectional	Ethiopia	116♂; mean age=37.6±8.6years; BMI=18.8±2.1; 232♂; mean age=30.5±7.8years; BMI=19.2±2.8	Accelerometer	C	Linear regression models
5	Ortega 2015 [76]	Cross-sectional	USA	21 (5♂) physically active with a mean age=44.1±17.2years, 73% African American and 27	self-reported historical PAQ	A	t-tests and multiple

6	Webel 2015 [106]	Cross-sectional	USA	%White and 48(♂) sedentary with a mean age=40.1±15.6years, 75% African American and 25% White 102(54♂); mean age=48±6.5years; 10% White, 84% African American, 6% Hispanic/other	(previous year) Diary	A	linear regression models t-tests
7	Erlandson 2014 [75]	Cross-sectional	USA	359 (305♂); mean age=52.0±5.2years; 74% White, 18% Hispanic	MLTPA	A	Spearman coefficients
8	Hsieh 2014 [74]	Cross-sectional	China	263 (200♂); mean age=38.4±9.8years; BMI=21.6±2.6; 94% Han Chinese	IPAQ	B	Logistic regression models
9	Justina 2014 [73]	Cross-sectional	Brazil	74 (45♂); mean age=44.3±9.2ears; 92% White	IPAQ	B	Regression analysis
10	Jaggers 2014 [72]	RCT	USA	63(31♂); mean age=48±11years	Accelerometer	C	ANOVA
11	Ramírez-Marrero 2014 [71]	Cross-sectional	Puerto Rico, USA	32 (13♂) Hispanics with lipodystrophy mean age=50.3±1.2 years; BMI=29.0±0.8 and 28 (15♂) Hispanics without lipodystrophy mean age=48.1±1.3years; BMI=26.4±1.4	Accelerometer	C	ANOVA

Table 1. Continued.

Nr	First author / year	Design	Country	Participants	PA Measurement	Quality of PA Measurement	Statistical Analysis
12	Raso 2014 [70]	Cross-sectional	Brazil	39♂; mean age=40.6±1.4years; BMI=24.8±0.6	IPAQ	B	Univariate regression models
13	Blashill 2013 [69]	Cohort	USA	860♂; mean age=43.8±9.9years; 80% White, 10% African American, 2% Asian, 8% other	Lipid Research Clinics PA questionnaire	A	Linear mixed effects modeling
14	Dufour 2013 [68]	Cross-sectional	USA	335(249♂); mean age=47.7±10.5 years; 51.3% White	Self-report exercise in the past 72h	A	T-tests and chi-square tests
15	Edward 2013 [67]	Cross-sectional	Nigeria	265(86♂); mean age=38.7±8.7 years	STEP-wise approach to Surveillance	A	Chi-square tests
16	Frantz 2013 [66]	Cross-sectional	South-Africa	407 (93♂); mean age=38.8±8.9years	Sub-Saharan African Activity	A	Chi-square tests

17	de Bruin 2012 [65]	Cohort	The Netherlands	499 (245♂); mean age=44.3±10.0years; BMI=26.5±4.9	Questionnaire Having 30 min exercise 2 days / week. IPAQ	A	Logistic regression models
18	Fillipas 2012 [64]	Cohort	Australia	80 outpatients, aged ≥18years	IPAQ	B	
19	Alencastro 2011 [63]	Cross-sectional	Brazil	1,240 (628♂); mean age=38.6±10.1years; BMI=24.9±4.4; 57.3% White	IPAQ	B	ANOVA
20	Kyser 2011 [62]	Cohort	USA	528 (410♂); median age=41years; 60% White, 28% African American, 12% Hispanic / Other	computer-assisted self-interview IPAQ	A	
21	Segatto 2011 [61]	Cross-sectional	Brazil	42 (23♂); age range=31–59 years	IPAQ	B	Chi-square tests
22	Muronya 2011 [60]	Cross-sectional	Malawi	174 (67♂); mean age=40.8years	STEP-wise approach to Surveillance	A	Odds ratios
23	Bonfanti 2010 [59]	Cohort	Italy	292 (219♂); median age=37years; 86.2% White	Unknown: yes versus no	A	Univariate analysis

Table 1. Continued.

Nr	First author / year [Ref nr]	Design	Country	Participants	PA Measurement	Quality of PA Measurement	Statistical Analysis
24	Lo Re 2009 [58]	Cross-sectional	USA	1,237 (765♂); median age=43years	mild [<4 h/week, intensive [\geq h /week]	A	Chi-square tests
25	Allard 2008 [57]	Cross-sectional	Canada	65♂; mean age=47.0±0.9years; BMI=26.5±0.4; 81.5% White, 10.8% African descents, 7.7% Asian descents	Activity logs	A	Mann-Whitney U-test
26	Basta 2008 [56]	Cross-sectional	USA	208 (181♂); mean age=42.6±7.1years; 84.6% African American or Black	IPAQ	B	
27	Kinsey 2008 [55]	Cross-sectional	South-Africa	186 (46♂ with a mean age=36±7years and 140♀ with a mean age=35±8years)	Combination MAQ, MLTPA and Baecke PAQ	A	Spearman's correlations / Mann-Whitney U-test

28	Kowal 2008 [54]	Cross-sectional	Canada	97 (81♂); mean age=39.4±8.7years; 71% White, 12% African, 4% Hispanic, 2% Haitian, 2% Aboriginal, 8% Other	Self-reported frequencies of exercising	A	Regression analysis
29	Littlewood 2008 [53]	Cross-sectional	USA	221 (124♂); mean age=40 years; 42% African-American, 46% White, 4% Native American, 4% Asian/Pacific Islander, 4% Other	Self-reported frequencies of exercising	A	Pearson correlation coefficients, ANOVA
30	Ramírez-Marrero 2008 [52]	Cross-sectional	Puerto Rico, USA	58 (35♂); mean age=46.5±8.8years; BMI=26.8±5.3	Accelerometer	C	Independent t-test
31	Florindo 2007 [51]	Cross-sectional	Brazil	230 (169♂); age range=20-59 years	Baecke PA questionnaire	B	Regression analysis
32	Florindo 2006 [50]	Cross-sectional	Brazil	30; mean age=37.2 years	Baecke PA questionnaire	B	Pearson correlation coefficients
33	Howard 2006 [49]	Cohort	USA	364♂; median age=54years; 61% African American, 24% Hispanic, 12% White, 4% Other	At least moderate PA ≥20 min >1 day per week	A	Regression analysis

Table 1. Continued.

Nr	First author / year [Ref nr]	Design	Country	Participants	PA Measurement	Quality of PA Measurement	Statistical Analysis
34	Jacobson 2006 [48]	Cohort	USA	342 (249♂) highly active antiviral therapy users with a mean age=42±7years, with 58% White, 31% African American, 11% Hispanic and 135 (97♂) non-highly active antiviral therapy users with a mean age=44±7years, with 44% White, 44% African American, 12% Hispanic	PA recall last 7 days	A	Cox proportional hazards
35	Salyer 2006 [47]	Cross-sectional	USA	95 (79♂); mean age=41.3±8.4years	Self-report days / week 30 min moderate PA	A	Factorial between-subjects ANOVA
36	Smit 2006 [46]	Cohort	USA	79 (57♂) with no treatment; mean age=40.7±0.7years; 96.1% African American, 1.3% Hispanic, 2.6% White and 134 (91♂) with treatment; mean age=44.7±0.5years; 94% African American, 3% Hispanic, 3% White	modified Paffenbarger questionnaire	A	Chi-square tests

37	Shah 2005 [45]	Cross-sectional	USA	45♂ all White with a mean age=47.1±8.4years versus 6♀ (5 White, 1 African American) with a mean age=42.5±5.9years	Self-report	A	Wilcoxon rank sum tests and Spearman's correlation coefficients
38	Bopp 2004 [44]	Cross-sectional	USA	66; mean age=39±8.years; 92% African American, 5% White, 3% Hispanic	Accelerometer	C	Pearson correlation coefficients
39	Clingerman 2003 [43]	Cross-sectional	USA	78 (70♂); mean age=40.4±8.3years; 59% African American	PAQ	A	Pearson correlation coefficients
40	Domingo 2003 [42]	Cross-sectional	Spain	75 (56♂) on stavudine with mean age=38.8±8.6years and 75 (62♂) on zidovudine with mean age=39.3±9.5years	MLTPA	A	Kruskal-Wallis test
41	Gavrila 2003 [41]	Cross-sectional	USA	120 (107♂); mean age=43.7±8.0years; 86.7% White, 7.5% Black, 5.8% Hispanic	Self-report	A	Logistic regression models

Table 1. Continued.

Nr	First author / year [Ref nr]	Design	Country	Participants	PA measurement	Quality of PA Measurement	Statistical Analysis
42	Collins 2001 [40]	Cohort	USA	2,864 (1403♂); mean age=38years; 49% White, 33% African American, 13% Hispanic	Self-report regular exercise	A	Logistic regression models
43	Sheehan 2000 [39]	Cohort	UK	33♂; median age=35years; mean BMI=20.8	Doubly labeled water	C	Linear regression models
44	Mustafa 1999 [38]	Cohort	USA	156♂; mean age=35.0±6.4years;	How many times a week do you engage in physical exercise?	A	Cox propotional hazards model
45	Macallan 1995 [37]	Cross-sectional	UK	27♂; median age=35years	Doubly labeled water	C	ANOVA

A=self-report of poor or unknown reliability/validity in persons living with HIV, B=self-report with acceptable reliability/validity in persons living with HIV, c=objective PA assessment, RCT= randomized controlled trial, PA=physical activity, BMI=body mass index, IPAQ= International Physical Activity Questionnaire; MAQ= Modifiable Activity Questionnaire, MLTPA=Minnesota Leisure Time Physical Activity, PAQ=Physical Activity Questionnaire.

Correlates of Physical Activity in People Living with HIV/AIDS

Table 2 summarizes associations between 55 potential correlates and the physical activity participation in people with HIV/AIDS.

Demographic Correlates

For 3 of the 8 demographic variables there was a consistent (reported in 4 or more studies) finding. Older age (6/10 studies; 60%) was consistently associated with lower and a higher educational level with higher physical activity levels (6/7; 86%). Gender differences were inconsistently reported, i.e. while 6 studies indicated men engaged in more physical activity than women, another reported the opposite, while 8 other studies showed no difference between genders. While one study reported a higher physical activity levels in the non-white population, another reported lower levels and 4 studies reported no associations. Being married, having a manual labor versus non-manual labor job and a higher annual income were all in one study significantly associated with a higher physical activity level while only 1 of 2 studies found that having a job was associated with more physical activity.

Biological Correlates

Seventeen biological correlates were included. There was consistent evidence for 4 correlates. Exposure to antiviral therapy (4/6, 67%), a lower number of CD4 cells/ μ l (7/11, 64%) and the presence of lipodystrophy (4/4, 100%) were all consistently associated with lower levels of physical activity. A potential biological correlate of interest, which across three studies was associated with higher levels of physical activity was a higher cardiorespiratory fitness level. The presence of opportunistic infections was negatively associated with physical activity in 3 studies. The body mass index (3/8; 37.5%) and a higher viral load (4/8; 50%)

were inconsistently associated with physical activity levels. The presence of cardio-metabolic comorbidity is not associated with lower physical activity levels (3/10; 30%). Other biological correlates of interest are summarized in Table 2.

Behavioral Attributes/Skills

Eight behavioral attributes were examined (see Table 2), but due to limited data none were considered consistently related to physical activity. Current alcohol use (1/3; 33%) and illicit drugs (1/5; 20%) use seem not to be associated with current physical activity behavior. The association with adherence to antiviral therapy was inconsistently associated with physical activity levels (1/2; 50%). Findings reported in only one study are presented in Table 2.

Psychological, Cognitive and Emotional Correlates

Twenty correlates were included. No correlate was consistently reported in 4 or more studies. A more positive health attitude was in 3 studies associated with more physical activity, while a higher self-efficacy, more perceived benefits and a better perceived physical functioning and general health were in 2 studies associated with more physical activity. Higher levels of depressive symptoms were in 3 studies, and more perceived bodily pain in 2 studies reported as a negative correlate, i.e. higher levels of depression and more bodily pain are associated with lower physical activity levels. A better emotional functioning was in 3 studies unrelated to the physical activity levels. Details and findings reported in only one study are presented in Table 2.

Social/Cultural Factors

Two of 3 studies (67%) reported on social support as a potential positive correlate.

Environmental Factors

The presence of food insecurity was in Ethiopian men but not in women associated with more physical activity.

Policy Factors

No policy-level correlates were located in the systematic review of the literature.

Table 2. Summary of the Physical Activity Correlates in People Living with HIV

Determinant Variable	Significantly Related to PA		Unrelated to PA		Summary Code ^a	
	Study*	Assoc.	Study*	Assoc.	% Studies Reporting Assoc.	
Demographic Factors						
Age (older)	[78]M [106] [74] [66] [60] [40]	-	[76] [72] [68] [43]	--	60% (6/10)	
Gender (female)	[77] [106] [66] [65] [46] [40] / [63]	-/+	[76] [74] [67] [60] [55] [52] [45] [43]	??	43% (6/14 for -)	
Race (non-White)	[78]M / [40]	-/+	[76] [68] [47] [43]	0	20% (1/5 for -)	
Education (higher)	[78]M [76] [74] [68] [65] [40]	+	[43]	++	86% (6/7)	
Marital status (married)	[40]	+		+	100% (1/1)	
Employment (yes)	[46]	+	[43]	?	50% (0/1)	
Employment (manual labor vs. non-manual)	[74]	+		+	100% (1/1)	
Annual income (higher)	[43]	+		+	100% (1/1)	
Biological Factors						
Illness duration (longer)			[76] [74] [68]	0	0% (0/3)	
AIDS diagnosis (present)	[68]	-		0	0% (0/1)	
Exposure to antiviral therapy (yes)	[68] [60] [48] [46]	-	[66] [55]	--	67% (4/6)	
Duration antiviral treatment (longer)			[60] [46]	0	0% (0/2)	
HIV status (WHO clinical stage higher)	[77]	-		-	100% (1/1)	
HIV viral load (higher)	[77] [69]M [44] [43]	-	[76] [68] [61] [46]	??	50% (4/8)	
Number of CD4 cells/ul (lower)	[77] [68] [61] [55] [40] [39]M [38]M	-	[76] [74] [44] [43]	--	64% (7/11)	
Body mass index (higher)	[78] [77] [72]	-	[80] [76] [74] [68] [57]M	??	37.5% (3/8)	
Waist circumference (higher)	[72]		[47]	?	50% (1/2)	
Central obesity (present)	[51]	-		-	100% (1/1)	
Lipodystrophy (present)	[73] [71] [61] [42]	-		--	100% (4/4)	
Wasting (more weight loss)	[37]M	-		-	100% (1/1)	
Cardiorespiratory fitness (higher)	[50] [72] [47]	+		+	100% (3/3)	
Neurocognitive impairments (more)	[76] [68]	-		-	100% (2/2)	
Muscular fitness (higher)			[70]M	0	100% (1/1)	
Cardio-metabolic comorbidity (present)	[63]M [59] [49]M		[79] [72] [71] [64] [63]F [48] [47] [45] [42] [41]	0	30% (3/10)	
Opportunistic infections (present)	[77] [58] [39]M	-	[68] [40]	-	75% (3/4)	

Table 2. Continued.

Determinant Variable	Significantly Related to PA		Unrelated to PA		Summary Code ^a	
	Study*	Assoc.	Study*	Assoc.	% Studies Reporting Assoc.	
Behavioral Attributes /Skills						
Adherence to antiviral therapy (yes)	[62]	+	[53]	?	50% (1/2)	
Smoking (more cigarettes per day)	[53]	-		-	100% (1/1)	
Smoking history (present)			[74]	0	0% (0/1)	
Physical activity history (active)	[65]	+		+	100% (1/1)	
Sedentary behavior (more television watching)	[43]	-		-	100% (1/1)	
Eating habits (healthier)	[47]	+		+	100% (1/1)	
Alcohol abuse (current)	[53]	-	[76] [74]	0	33% (1/3)	
Illicit drug use (present)	[53]	-	[76] [68] [46] [40]	0	20% (1/5)	
Psychological, Cognitive and Emotional Factors						
Exercise knowledge (present)			[74]	0	0% (0/1)	
Self-efficacy (higher)	[74] [65]	+		+	100% (2/2)	
Body image (better)	[64]	+		+	100% (1/1)	
Health motivation /attitude (higher / better)	[74] [65] [40]	+		+	100% (3/3)	
Depression (present)	[69] [68] [53]	-		-	100% (3/3)	
Perceived benefits (more)	[74] [53]	+		+	100% (2/2)	
Perceived barriers (more)	[74]	-		-	100% (1/1)	
Perceived physical functioning (better)	[75] [68] [40]	+		+	100% (3/3)	
Perceived physical limitations (more)	[75]	-		-	100% (1/1)	
Perceived bodily pain (worse)	[75] [54]	-		-	100% (2/2)	
Perceived general health (better)	[75] [43]	+		+	100% (2/2)	
Perceived vitality (better)	[75]	+	[44]	?	50% (1/2)	
Perceived social functioning (better)	[75]	+		+	100% (1/1)	
Perceived emotional functioning (better)			[75] [68] [40]	0	0% (0/3)	
Perceived emotional limitations (more)	[75]	-		-	100% (1/1)	
Perceived stress (higher)			[44]	0	0% (0/1)	
Perceived appetite (better)	[39]M	+		+	100% (1/1)	
Sleep quality (higher)			[44]	0	0% (0/1)	
State and trait anxiety (higher)			[44]	0	0% (0/1)	
Trans-theoretical model (higher stage)	[56]	+		+	100% (1/1)	
Social/Cultural Factors						
Social support (more)	[64] [43]	+	[53]	+	67% (2/3)	
Physical Environment						
Food insecurity (present)	[77]M	+	[77]F	?	50% (1/2)	

Policy Factors					
-----------------------	--	--	--	--	--

PA=physical activity; *Reference numbers; °The percentages in parentheses refer to the number of associations supporting the expected association divided by the total number of associations for the variable. Associations are coded with: "0" (0-33% of studies supporting association); "?" (34%-59% of studies supporting an association); or "+" or "-" (60%-100% of studies supporting an association). When four or more studies support an association or no association, it is coded as "--", or "++" indicating that there is consistent evidence for that correlate. The "??" code indicates a variable that was investigated four or more times studied with considerable lack of consistency in the findings. M=only valid for men, F=only valid for women, SDT= self-determination theory.

Differences in Number of Significant Correlates

Fisher's exact tests showed there were no differences between un-validated and validated self-report instruments (P=0.49) or between self-report and objective tools (P=0.81). In contrast, significantly (P=0.007) more correlates and less unrelated variables were obtained in studies with a sample size equal to or larger than the median sample size.

Discussion

General Findings

To the best of the authors' knowledge, the present review is the first to systematically document the correlates of physical activity in persons with HIV/AIDS. Out of 55 correlates from 45 studies, 5 consistent (i.e. reported in 4 or more studies and consistently associated in at least 60% of the cases) correlates were obtained. Lower levels of physical activity were consistently associated with older age, a lower number of CD4 cells/ μ l, exposure to antiviral therapy and the presence of lipodystrophy while higher levels were found in those with a higher educational level. One should take into account that several of these correlates may be interrelated with each other. For example, the presence of lipodystrophy is a known side-effect of exposure to antiviral therapy (stavudine which the World Health Organization recommends to be phased out due to its side-effects such as lipodystrophy [81]), and the number of CD4 cells/ μ l is also associated with exposure to antiviral therapy.

For gender, the body mass index and HIV viral load, findings were consistently conflicting. We believe that the limited number of consistent correlates observed in the present review might be due to differences in study design, sample characteristics, choice of assessments/correlates, statistical analyses, and differences in sample sizes. More significant correlates were obtained in studies with a sample size equal to or larger than the median sample size. Nevertheless, although there was a large heterogeneity, our varied findings clearly illustrate that participation in physical activity by people living with HIV/AIDS is associated with a range of complex factors.

Knowledge about demographic correlates of physical activity behavior will help to identify these high-risk persons in whom physical activity is low or, even, likely to be reduced and who may require intensified and targeted interventions. The current review shows that older patients with HIV/AIDS, those with a lower educational level and those on antiviral therapy are the most vulnerable patients in need for intensified care. The observation that older age was associated with a lower physical activity participation is in agreement with findings in the general population [36] and seems not to be due to a longer illness duration which was in none of 3 studies a significant correlate. Special attention should also be given to people with lipodystrophy. There are several pathways through which the presence of lipodystrophy might be a barrier to adopt and maintain an active lifestyle. Its presence has important psychological effects, ranging from bodily discomfort to low self-esteem and depression [82]. The observation in one study [64] that a better body image in people living with HIV/AIDS is associated with higher physical activity levels might confirm this hypothesis. Owing to its physical manifestations, lipodystrophy is viewed as a visible marker of the HIV disease as well which might on its turn cause stigmatization and social isolation [83]. Patients and healthcare professionals should be informed that positive experiences when being physically active can improve the physical comfort and body satisfaction of people living with HIV/AIDS [84]. This finding is also relevant in resource-limited areas of sub-Saharan Africa [85]. Researchers and clinicians should explore in more detail what techniques can stimulate these positive experiences in people with HIV/AIDS who are confronted with lipodystrophy.

The fact that immunological parameters such as a lower number of CD4 cells/ μ l are associated with less physical activity might be due to the fact that this a measure for the disease severity and the need for treatment and might be associated with symptoms as nausea [86], depression [87] and pain [88]. Two different studies indicated that respectively the presence of depression and bodily pain are negatively associated with physical activity among people with people

with HIV/AIDS, a finding which has been observed in vulnerable mental health populations as well [34, 89]. A systematic review of 61 studies in people living with HIV/AIDS demonstrated that the prevalence of pain ranged from a point prevalence of 54% (95%CI=51%–56%) to 83% (95%CI=76%–88%) using a three-month recall period [88]. The types of pain experienced by people with HIV/AIDS and the aetiology appear to vary. As indicated, people living with HIV/AIDS may experience pain as a direct result of the virus on the peripheral or central nervous systems. Pain may be due to resultant opportunistic infections as well, which was reported as a negative correlate in 3 of 4 studies, or pain may arise as a result of the side effects of anti-retroviral treatment [88] [90]. Despite an increasing awareness of pain as a significant contributor to the disability and impaired health related quality of life, the problem of its under-management persists [88]. Although there is preliminary evidence [91, 92] that inclusion of physiotherapy as a complementary treatment for pain-management and simultaneously improving mental and physical health outcomes and reducing disability is promising, also in resource-limited areas of sub-Saharan Africa [93], more research is needed.

Our data suggest that higher cardiorespiratory fitness is positively associated with physical activity, an observation made in the general population as well [36]. Therefore, physiotherapy should focus on cardiorespiratory fitness parameters. Cardiorespiratory fitness is the ability of the circulatory and respiratory systems to supply oxygen to working muscles during sustained physical activity [12] and is a strong and independent predictor of disability and all-cause mortality [94]. Research in a disadvantaged setting in South-Africa demonstrated that exercise might improve cardiorespiratory fitness, although follow-up is needed [95].

Finally, our data show that more research regarding the association between higher HIV loads and physical activity participation is needed. Until clear clinical practice guidelines become available, patients with high HIV viral load levels should be recommended to perform moderate instead of high intensity physical activity. While moderate intensity physical activity improves the immune function in people with HIV/AIDS [96, 97], high-intensity exercise is known to have immunosuppressive effects in the general population [98, 99] and in people living with HIV/AIDS [100].

Limitations and Recommendations for Future Research

There are several limitations to this review, which should be acknowledged.

First of all, the diversity of physical activity measures prevented us from performing a formal meta-analysis. Self-report questionnaires are, for example, known to require motivation to complete all of the questions and often the detail regarding the level (frequency, duration and intensity) and type of physical activity is not consistently evaluated. To the best of our knowledge only the International Physical Activity Questionnaire [101] and the Baecke Physical Activity Questionnaire [50] have been validated in people with HIV/AIDS. Fewer significant associations would be expected in studies that relied on un-validated self-report measures and in self-report versus objective assessments. However, Fisher's exact tests showed there were no differences between un-validated and validated self-report instruments or between self-report and objective tools. Considering the wide diversity in physical activity assessments, our findings do reveal that there is a high need for researchers to adopt a clear consensus on which assessment tools should be recommended in people living with HIV/AIDS.

Second, all correlates investigated were only documented in a small number of studies. Examination of the same, standardized variables in different studies is necessary in order to build a consistent body of evidence that can support or refute the potential influence of individual variables.

Third, the majority of the studies investigated physical activity correlates at only one level of the socio-ecological model, with very limited data about the potential role of social and environmental variables and no studies on policy factors. Future studies should attempt to analyze the role of multiple correlates of physical activity from a broad socio-ecological perspective. Given that research suggests that maintaining changes in physical activity requires a multilevel approach [20], exploring these interactions in people living with HIV/AIDS is highly important. At the social level of the socio-ecological model, future research could focus on the amount and type of social support necessary to begin and maintain physical activity behavior in people living with HIV/AIDS. This kind of research could explore whether: (a) the relationship between physical activity participation and professional support is a dynamic process in which the sources of support or need for support change over time, and (b) any social barriers for people living with HIV/AIDS can be identified and addressed by involving significant others in the rehabilitation process. Data on the role of social support in facilitating physical activity behavior in people living with HIV/AIDS are currently inconsistent. At the environmental level of the socio-ecological model, the role of the built environment on a person's physical activity behavior should be evaluated. Built environments are the totality of places built or designed, including buildings, grounds around buildings, layout of communities, transportation infrastructure and parks and trails [102]. There are self-report tools available for assessing the built environment [103], which have been used in sub-Saharan

African countries where the majority of people with HIV/AIDS live [104]. Correlates at the policy level of the socio-ecological model are likely to be best initially explored using a qualitative approach [31]. Researchers should examine, which policies are currently in place to motivate people living with HIV/AIDS to engage in an active and healthy lifestyle. Interviews of people living with HIV/AIDS but also healthcare professionals and policy makers may provide further insight as to what is needed to stimulate an active lifestyle. If the purpose is to inform and motivate policy changes that will improve the quality of life and reduce the disability in people living with HIV/AIDS merely documenting the relationship between for example environmental changes and physical activity behavior is likely to be insufficient. At some point, environmental and policy change research will need to include assessments of broader health outcomes in people living with HIV/AIDS, such as changes in the prevalence of chronic co-morbidities, health care service utilization, as well as the economic costs and benefits of proposed policy changes.

Fourth, while the majority of people with HIV/AIDS live in Sub-Saharan Africa [1], only a limited number of studies in this review (5/45) were executed in this region. This lack of studies from Sub-Saharan African countries highlights the gap between where most research is done and where the largest public health impacts of physical inactivity for people with HIV/AIDS are probably located. The effect of HIV/AIDS on African people has larger consequences because if their productivity is affected, this directly affects family welfare and increases the scale of both family and community poverty. In Sub-Saharan Africa, many HIV patients are relying on labor-demanding jobs in the informal sector with no job security or compensation for lost income. Maintaining physical strength and an adequate activity level is thus of crucial importance for their livelihoods [105]. Therefore, there is a high need for exploring in particular physical activity participation correlates in Sub-Saharan Africa. Future physical activity research in Sub-Saharan Africa should for example focus on the socio-cultural context in which people with HIV/AIDS are physically active and which is often different than in Western countries. For instance, in most Sub-Saharan African countries, if women are thin, they may be stigmatized for being poor or for having HIV [106]. Such prejudices might prevent women from being physically active as they are afraid of losing weight. Finally, future studies in Sub-Saharan Africa would benefit from assessing to what extent the lack of a comprehensive health policy in most of these countries and specific environmental factors such as lack access to physical activity related treatment and recreation facilities, unsafety due to

crime or dangerous traffic in urban settings, food insecurity, civil conflicts, and extreme weather conditions are all linked to physical inactivity in people with HIV/AIDS.

Fifth, the finding that people taking antiviral therapy may have lower levels of physical activity is of interest and perhaps differs from expert opinion and anecdotal clinical observations. More research is required to confirm or refute this finding and explore why this relationship may have been observed.

In conclusion, our results demonstrate that participation in physical activity by people with HIV/AIDS is determined by a range of complex factors. Special attention should be given to older patients, those with a lower educational level, those presenting with lipodystrophy and those on antiviral therapy. We recommend that health care professionals, and in particular physiotherapists, should also consider the presence of pain and depression when encouraging people with HIV/AIDS to engage in physical activity. This will be particularly pertinent given that previous literature has demonstrated the positive effects of physical activity on the physical and mental health comorbidities seen more recently in this group.

Acknowledgements

None.

Conflict of Interests

All authors declare that they have no conflicts of interest relevant to the content of this review.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. Brendon Stubbs was supported by the Collaboration for Leadership in Applied Health Research and Care South London theme. No sources of funding were used to assist in the preparation of this article.

References

1. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The lancet*. 2013;380:2197-223.
2. Morris A, Gingo MR, George MP, Lucht L, Kessinger C, Singh V, Hillenbrand M, Busch M, McMahon D, Norris KA. Cardiopulmonary function in individuals with HIV infection in the antiretroviral therapy era. *AIDS (London, England)*. 2012;26:731.
3. Schouten J, Wit FW, Stolte IG, Kootstra N, van der Valk M, Geerlings SG, Prins M, Reiss P, Reiss P, Wit F. Cross-sectional comparison of the prevalence of age-associated comorbidities and their risk factors between HIV-infected and uninfected individuals: the AGEHIV cohort study. *Clinical Infectious Diseases*. 2014:ciu701.
4. Deeks SG, Phillips AN. Clinical review: HIV infection, antiretroviral treatment, ageing, and non-AIDS related morbidity. *Bmj*. 2009;338:288-92.
5. Hearps AC, Martin GE, Rajasuriar R, Crowe SM. Inflammatory co-morbidities in HIV+ individuals: learning lessons from healthy ageing. *Current HIV/AIDS Reports*. 2014;11:20-34.
6. Grinspoon S, Carr A. Cardiovascular risk and body-fat abnormalities in HIV-infected adults. *New England Journal of Medicine*. 2005;352:48-62.
7. El-Sadr WM. Class of antiretroviral drugs and the risk of myocardial infarction. *New England Journal of Medicine*. 2007;356:1723-35.
8. Dillon DG, Gurdasani D, Riha J, Ekoru K, Asiki G, Mayanja BN, Levitt NS, Crowther NJ, Nyirenda M, Njelekela M. Association of HIV and ART with cardiometabolic traits in sub-Saharan Africa: a systematic review and meta-analysis. *International journal of epidemiology*. 2013;42:1754-71.
9. Lifson AR, Lando HA. Smoking and HIV: prevalence, health risks, and cessation strategies. *Current HIV/AIDS Reports*. 2012;9:223-30.
10. Schuelter-Trevisol F, H Wolff F, R Alencastro P, Grigoletti S, L Ikeda M, BM Brandao A, T Barcellos N, C Fuchs S. Physical activity: do patients infected with HIV practice? How much? A systematic review. *Current HIV research*. 2012;10:487-97.
11. O'Brien KK, Davis AM, Strike C, Young NL, Bayoumi AM. Putting episodic disability into context: a qualitative study exploring factors that influence disability experienced by adults living with HIV/AIDS. *Journal of the International AIDS Society*. 2009;12:30.

12. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Reports*. 1985;100:126.
13. O'Brien K, Nixon S, Tynan AM, Glazier R. Aerobic exercise interventions for adults living with HIV/AIDS. *Cochrane Database Syst Rev*. 2010;8.
14. Leach L, Bassett S, Smithdorf G, Andrews B, Travill A. Suppl 1: M3: A Systematic Review of the Effects of Exercise Interventions on Body Composition in HIV+ Adults. *The open AIDS journal*. 2015;9:66.
15. Neto MG, Conceição CS, Carvalho VO, Brites C. Effects of Combined Aerobic and Resistance Exercise on Exercise Capacity, Muscle Strength and Quality of Life in HIV-Infected Patients: A Systematic Review and Meta-Analysis. *PLoS one*. 2015;10:e0138066.
16. Fillipas S, Cherry C, Cicuttini F, Smirneos L, Holland A. The effects of exercise training on metabolic and morphological outcomes for people living with HIV: a systematic review of randomised controlled trials. *HIV clinical trials*. 2010;11:270-82.
17. Jagers J, Hand G, Dudgeon W, Burgess S, Phillips K, Durstine J, Blair S. Aerobic and resistance training improves mood state among adults living with HIV. *International journal of sports medicine*. 2014.
18. O'Brien, Tynan A-M, Nixon SA, Glazier RH. Effectiveness of aerobic exercise for adults living with HIV: systematic review and meta-analysis using the Cochrane Collaboration protocol. *BMC infectious diseases*. 2016;16:1.
19. Ley C, Prista A. Suppl 1: M1: Editorial Physical Activity and HIV in Africa. *The open AIDS journal*. 2015;9:60.
20. Sallis JF, Cervero RB, Ascher W, Henderson KA, Kraft MK, Kerr J. An ecological approach to creating active living communities. *Annu Rev Public Health*. 2006;27:297-322.
21. Stubbs B, Eggermont L, Soundy A, Probst M, Vandenbulcke M, Vancampfort D. What are the factors associated with physical activity (PA) participation in community dwelling adults with dementia? A systematic review of PA correlates. *Archives of gerontology and geriatrics*. 2014;59:195-203.

22. Stubbs B, Hurley M, Smith T. What are the factors that influence physical activity participation in adults with knee and hip osteoarthritis? A systematic review of physical activity correlates. *Clinical rehabilitation*. 2015;29:80-94.
23. Vancampfort D, Vanderlinden J, Stubbs B, Soundy A, Pieters G, Hert MD, Probst M. Physical activity correlates in persons with binge eating disorder: A systematic review. *European Eating Disorders Review*. 2014;22:1-8.
24. Vancampfort D, Knapen J, Probst M, Scheewe T, Remans S, De Hert M. A systematic review of correlates of physical activity in patients with schizophrenia. *Acta Psychiatrica Scandinavica*. 2012;125:352-62.
25. Vancampfort D, De Hert M, Stubbs B, Soundy A, De Herdt A, Detraux J, Probst M. A systematic review of physical activity correlates in alcohol use disorders. *Archives of psychiatric nursing*. 2015;29:196-201.
26. Rehm K, D K-P. Physical activity levels and perceived benefits and barriers to physical activity in HIV-infected women living in the deep south of the United States. *AIDS care*. 2016.
27. Roos R, Myezwa H, van Aswegen H. "Not easy at all but I am trying": barriers and facilitators to physical activity in a South African cohort of people living with HIV participating in a home-based pedometer walking programme. *AIDS care*. 2015;27:235-9.
28. Simonik A, Vader K, Ellis D, Kesbian D, Leung P, Jachyra P, Carusone SC, O'Brien KK. Are you ready? Exploring readiness to engage in exercise among people living with HIV and multimorbidity in Toronto, Canada: a qualitative study. *BMJ open*. 2016;6:e010029.
29. Ley C, Barrio MR, Leach L. Suppl 1: M6: Social-Ecological, Motivational and Volitional Factors for Initiating and Maintaining Physical Activity in the Context of HIV. *The open AIDS journal*. 2015;9:96.
30. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA, Thacker SB. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA*. 2000;283:2008-12.
31. DeVon HA, Block ME, Moyle-Wright P, Ernst DM, Hayden SJ, Lazzara DJ, Savoy SM, Kostas-Polston E. A psychometric toolbox for testing validity and reliability. *Journal of Nursing scholarship*. 2007;39:155-64.

32. Vancampfort D, Correll CU, Probst M, Sienaert P, Wyckaert S, De Herdt A, Knapen J, De Wachter D, De Hert M. A review of physical activity correlates in patients with bipolar disorder. *Journal of Affective Disorders*. 2013b;145:285-91.
33. Vancampfort D, De Hert M, Stubbs B, Soundy A, De Herdt A, Detraux J, Probst M. A systematic review of physical activity correlates in alcohol use disorders. *Archives of Psychiatric Nursing*.
34. Vancampfort D, Stubbs B, Sienaert P, Wyckaert S, De Hert M, Rosenbaum S, Probst M. What are the factors that influence physical activity participation in individuals with depression? A review of physical activity correlates from 59 studies. *Psychiatria Danubina*. 2015;27:210.
35. Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Medicine and science in sports and exercise*. 2000;32:963-75.
36. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: review and update. *Medicine & Science in Sports & Exercise*. 2002.
37. Macallan DC, Noble C, Baldwin C, Jebb SA, Prentice AM, Coward WA, Sawyer MB, McManus TJ, Griffin GE. Energy expenditure and wasting in human immunodeficiency virus infection. *New England Journal of Medicine*. 1995;333:83-8.
38. Mustafa T, Sy FS, Macera CA, Thompson SJ, Jackson KL, Selassie A, Dean LL. Association between exercise and HIV disease progression in a cohort of homosexual men. *Annals of Epidemiology*. 1999;9:127-31.
39. Sheehan LA, Macallan DC. Determinants of energy intake and energy expenditure in HIV and AIDS. *Nutrition*. 2000;16:101-6.
40. Collins RL, Kanouse DE, Gifford AL, Senterfitt JW, Schuster MA, McCaffrey DF, Shapiro MF, Wenger NS. Changes in health-promoting behavior following diagnosis with HIV: prevalence and correlates in a national probability sample. *Health Psychology*. 2001;20:351.
41. Gavrila A, Tsiodras S, Doweiko J, Nagy GS, Brodovicz K, Hsu W, Karchmer AW, Mantzoros CS. Exercise and vitamin E intake are independently associated with metabolic abnormalities in human immunodeficiency virus—positive subjects: a cross-sectional study. *Clinical infectious diseases*. 2003;36:1593-601.

42. Domingo P, Sambeat MA, Pérez A, Ordoñez J, Rodríguez J, Vázquez G. Fat distribution and metabolic abnormalities in HIV-infected patients on first combination antiretroviral therapy including stavudine or zidovudine: role of physical activity as a protective factor. *Antiviral therapy*. 2003;8:223-32.
43. Clingerman EM. Participation in physical activity by persons living with HIV disease. *Journal of the Association of Nurses in AIDS Care*. 2003;14:59-70.
44. Bopp CM, Phillips KD, Fulk LJ, Dudgeon WD, Sowell R, Hand GA. Physical activity and immunity in HIV-infected individuals. *AIDS care*. 2004;16:387-93.
45. Shah M, Tierney K, Adams-Huet B, Boonyavarakul A, Jacob K, Quittner C, Dinges W, Peterson D, Garg A. The role of diet, exercise and smoking in dyslipidaemia in HIV-infected patients with lipodystrophy. *HIV medicine*. 2005;6:291-8.
46. Smit E, Crespo C, Semba R, Jaworowicz D, Vlahov D, Ricketts E, Ramirez-Marrero F, Tang A. Physical activity in a cohort of HIV-positive and HIV-negative injection drug users. *AIDS care*. 2006;18:1040-5.
47. Salyer J, Lyon DE, Settle J, Elswick R, Rackley D. Coronary heart disease risks and lifestyle behaviors in persons with HIV infection. *Journal of the Association of Nurses in AIDS Care*. 2006;17:3-17.
48. Jacobson DL, Tang AM, Spiegelman D, Thomas AM, Skinner S, Gorbach SL, Wanke C. Incidence of metabolic syndrome in a cohort of HIV-infected adults and prevalence relative to the US population (National Health and Nutrition Examination Survey). *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2006;43:458-66.
49. Howard A, Floris-Moore M, Lo Y, Arnsten J, Fleischer N, Klein R. Abnormal glucose metabolism among older men with or at risk of HIV infection. *HIV medicine*. 2006;7:389-96.
50. Florindo AA, Latorre MdRDd, Santos ECMd, Negrão CE, Azevedo LF, Segurado AAC. Validity and reliability of the Baecke questionnaire for the evaluation of habitual physical activity among people living with HIV/AIDS. *Cadernos de saúde pública*. 2006;22:535-41.
51. Florindo AA, de Oliveira MdRD, Jaime PC, Segurado AAC. Leisure time physical activity prevents accumulation of central fat in HIV/AIDS subjects on highly active antiretroviral therapy. *International journal of STD & AIDS*. 2007;18:692-6.

52. Ramírez-Marrero FA, Rivera-Brown AM, Nazario CM, Rodríguez-Orengo JF, Smit E, Smith BA. Self-reported physical activity in Hispanic adults living with HIV: comparison with accelerometer and pedometer. *Journal of the Association of Nurses in AIDS Care*. 2008;19:283-94.
53. Littlewood RA, Vanable PA, Carey MP, Blair DC. The association of benefit finding to psychosocial and health behavior adaptation among HIV+ men and women. *Journal of Behavioral Medicine*. 2008;31:145-55.
54. Kowal J, Overduin LY, Balfour L, Tasca GA, Corace K, Cameron DW. The role of psychological and behavioral variables in quality of life and the experience of bodily pain among persons living with HIV. *Journal of pain and symptom management*. 2008;36:247-58.
55. Kinsey K, McVeigh J, Chantler I. Habitual physical activity levels are positively correlated with CD4 counts in an HIV-positive South African population. *African Journal of AIDS Research*. 2008;7:237-42.
56. Basta TB, Reece M, Wilson MG. The transtheoretical model and exercise among individuals living with HIV. *American journal of health behavior*. 2008;32:356-67.
57. Allard JP, Arendt BM, Aghdassi E, Mohammed SS, Fung LY, Jalali P, Salit IE. Dietary intake and physical activity in a Canadian population sample of male patients with HIV infection and metabolic abnormalities. *Current HIV research*. 2008;6:82-90.
58. Re III VL, Guaraldi G, Leonard MB, Localio AR, Lin J, Orlando G, Zirilli L, Rochira V, Kostman JR, Tebas P. Viral hepatitis is associated with reduced bone mineral density in HIV-infected women but not men. *AIDS (London, England)*. 2009;23:2191.
59. Bonfanti P, De Socio GL, Marconi P, Franzetti M, Martinelli C, Vichi F, Penco G, Madeddu G, Orofino G, Valsecchi L. Is metabolic syndrome associated to HIV infection per se? Results from the HERMES study. *Current HIV research*. 2010;8:165-71.
60. Muronya W, Sanga E, Talama G, Kumwenda JJ, van Oosterhout JJ. Cardiovascular risk factors in adult Malawians on long-term antiretroviral therapy. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2011;105:644-9.
61. Segatto AFM, Freitas Junior IF, Santos VRd, Alves KCP, Barbosa DA, Portelinha Filho AM, Monteiro HL. Lipodystrophy in HIV/AIDS patients with different levels of physical activity while on antiretroviral therapy. *Revista da Sociedade Brasileira de Medicina Tropical*. 2011;44:420-4.

62. Kyser M, Buchacz K, Bush TJ, Conley LJ, Hammer J, Henry K, Kojic EM, Milam J, Overton ET, Wood KC. Factors associated with non-adherence to antiretroviral therapy in the SUN study. *AIDS care*. 2011;23:601-11.
63. Alencastro PR, Fuchs SC, Wolff FH, Ikeda ML, Brandão AB, Barcellos NT. Independent predictors of metabolic syndrome in HIV-infected patients. *AIDS patient care and STDs*. 2011;25:627-34.
64. Fillipas S, Cicuttini F, Holland A, Cherry C. Physical activity participation and cardiovascular fitness in people living with HIV. A one-year longitudinal study. *Retrovirology*. 2012;9:P63.
65. de Bruin M, Sheeran P, Kok G, Hiemstra A, Prins JM, Hospers HJ, van Breukelen GJ. Self-regulatory processes mediate the intention-behavior relation for adherence and exercise behaviors. *Health Psychology*. 2012;31:695.
66. Frantz JM, Murenzi A. The physical activity levels among people living with human immunodeficiency virus/acquired immunodeficiency syndrome receiving high active antiretroviral therapy in Rwanda. *SAHARA-J*. 2013;10:113-8.
67. Edward AO, Oladayo AA, Omolola AS, Adetiloye AA, Adedayo PA. Prevalence of traditional cardiovascular risk factors and evaluation of cardiovascular risk using three risk equations in Nigerians living with human immunodeficiency virus. *North American journal of medical sciences*. 2013;5:680.
68. Dufour CA, Marquine MJ, Fazeli PL, Henry BL, Ellis RJ, Grant I, Moore DJ, Group H. Physical exercise is associated with less neurocognitive impairment among HIV-infected adults. *Journal of neurovirology*. 2013;19:410-7.
69. Blashill AJ, Mayer KH, Crane H, Magidson JF, Grasso C, Mathews WC, Saag MS, Safren SA. Physical activity and health outcomes among HIV-infected men who have sex with men: A longitudinal mediational analysis. *Annals of Behavioral Medicine*. 2013;46:149-56.
70. Raso V, Shephard RJ, Casseb J, Duarte AJdS, Silva PRS, Greve JMDA. Association between muscle strength and the cardiopulmonary status of individuals living with HIV/AIDS. *Clinics*. 2013;68:359-64.

71. Ramírez-Marrero FA, Santana-Bagur JL, Joyner MJ, Rodríguez-Zayas J, Frontera W. Metabolic syndrome in relation to cardiorespiratory fitness, active and sedentary behavior in HIV+ Hispanics with and without lipodystrophy: Fitness and metabolic syndrome in HIV. *Puerto Rico Health Sciences Journal*. 2014;33:163.
72. Jaggars JR, Prasad VK, Dudgeon WD, Blair SN, Sui X, Burgess S, Hand GA. Associations between physical activity and sedentary time on components of metabolic syndrome among adults with HIV. *AIDS care*. 2014;26:1387-92.
73. Della Justina LB, Luiz MC, Maurici R, Schuelter-Trevisol F. Prevalence and factors associated with lipodystrophy in AIDS patients. *Revista da Sociedade Brasileira de Medicina Tropical*. 2014;47:30-7.
74. Hsieh E, Fraenkel L, Bradley EH, Xia W, Insogna KL, Cui Q, Li K, Li T. Osteoporosis knowledge, self-efficacy, and health beliefs among Chinese individuals with HIV. *Archives of osteoporosis*. 2014;9:1-10.
75. Erlandson KM, Allshouse AA, Jankowski CM, Mawhinney S, Kohrt WM, Campbell TB. Relationship of physical function and quality of life among persons aging with HIV infection. *AIDS (London, England)*. 2014;28:1939-43.
76. Ortega M, Baker LM, Vaida F, Paul R, Basco B, Ances BM. Physical Activity Affects Brain Integrity in HIV+ Individuals. *Journal of the International Neuropsychological Society*. 2015;21:880-9.
77. Olsen MF, Kæstel P, Tesfaye M, Abdissa A, Yilma D, Girma T, Mølgaard C, Faurholt-Jepsen D, Christensen DL, Brage S. Physical activity and capacity at initiation of antiretroviral treatment in HIV patients in Ethiopia. *Epidemiology and infection*. 2015;143:1048.
78. Monroe AK, Brown TT, Cox C, Reynolds SM, Wiley DJ, Palella FJ, Kingsley LA, Plankey MW. Physical Activity and Its Association with Insulin Resistance in Multicenter AIDS Cohort Study Men. *AIDS research and human retroviruses*. 2015;31:1250-6.
79. Fazeli PL, Marquine MJ, Dufour C, Henry BL, Montoya J, Gouaux B, Moore RC, Letendre SL, Woods SP, Grant I. Physical activity is associated with better neurocognitive and everyday functioning among older adults with HIV disease. *AIDS and Behavior*. 2015;19:1470-7.

80. Dirajlal-Fargo S, Webel AR, Longenecker CT, Kinley B, Labbato D, Sattar A, McComsey GA. The effect of physical activity on cardiometabolic health and inflammation in treated HIV infection. *Antiviral therapy*. 2015.
81. World Health Organization. Antiretroviral therapy for HIV infection in adults and adolescents: recommendations for a public health approach-2010 revision. Geneva: World Health Organization 2010.
82. Power R, Tate H, McGill S, Taylor C. A qualitative study of the psychosocial implications of lipodystrophy syndrome on HIV positive individuals. *Sexually transmitted infections*. 2003;79:137-41.
83. Alexias G, Savvakis M, Stratopoulou I. Embodiment and biographical disruption in people living with HIV/AIDS (PLWHA). *AIDS care*. 2015:1-6.
84. Fillipas S, Oldmeadow LB, Bailey MJ, Cherry CL. A six-month, supervised, aerobic and resistance exercise program improves self-efficacy in people with human immunodeficiency virus: a randomised controlled trial. *Australian Journal of Physiotherapy*. 2006;52:185-90.
85. Mutimura E, Stewart A, Crowther NJ, Yarasheski KE, Cade WT. The effects of exercise training on quality of life in HAART-treated HIV-positive Rwandan subjects with body fat redistribution. *Quality of Life research*. 2008;17:377-85.
86. Chubineh S, McGowan J. Nausea and vomiting in HIV: a symptom review. *International journal of STD & AIDS*. 2008;19:723-8.
87. Pala AN, Steca P, Bagrodia R, Helpman L, Colangeli V, Viale P, Wainberg M. Subtypes of depressive symptoms and inflammatory biomarkers: An exploratory study on a sample of HIV-positive patients. *Brain, behavior, and immunity*. 2016.
88. Parker R, Stein DJ, Jelsma J. Pain in people living with HIV/AIDS: a systematic review. *Journal of the International AIDS Society*. 2014;17.
89. Stubbs B, Binnekade TT, Soundy A, Schofield P, Huijnen IP, Eggermont LH. Are Older Adults with Chronic Musculoskeletal Pain Less Active than Older Adults Without Pain? A Systematic Review and Meta-Analysis. *Pain Medicine*. 2013;14:1316-31.
90. Hewitt DJ, McDonald M, Portenoy RK, Rosenfeld B, Passik S, Breitbart W. Pain syndromes and etiologies in ambulatory AIDS patients. *Pain*. 1997;70:117-23.

91. Pullen SD, Chigbo NN, Nwigwe EC, Chukwuka CJ, Amah CC, Idu SC. Physiotherapy intervention as a complementary treatment for people living with HIV/AIDS. *HIV/AIDS (Auckland, NZ)*. 2014;6:99.
92. Mgbemena O, Westfall AO, Ritchie CS, Hicks J, Raper JL, Overton ET, Norton WE, Merlin JS. Preliminary outcomes of a pilot physical therapy program for HIV-infected patients with chronic pain. *AIDS care*. 2015;27:244-7.
93. Mkandla K, Myezwa H, Musenge E. The effects of progressive-resisted exercises on muscle strength and health-related quality of life in persons with HIV-related poly-neuropathy in Zimbabwe. *AIDS care*. 2015:1-5.
94. Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Asumi M, Sugawara A, Totsuka K, Shimano H, Ohashi Y. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*. 2009;301:2024-35.
95. Ley C, Leach L, Barrio MR, Bassett S. Effects of an exercise programme with people living with HIV: research in a disadvantaged setting. *African Journal of AIDS Research*. 2014;13:313-9.
96. Ezema C, Onwunali A, Lamina S, Ezugwu U, Amaeze A, Nwankwo M. Effect of aerobic exercise training on cardiovascular parameters and CD4 cell count of people living with human immunodeficiency virus/acquired immune deficiency syndrome: A randomized controlled trial. *Nigerian journal of clinical practice*. 2014;17:543-8.
97. Garcia A, Fraga GA, Vieira Jr RC, Silva CMS, Trombeta JCDS, Navalta JW, Prestes J, Voltarelli FA. Effects of combined exercise training on immunological, physical and biochemical parameters in individuals with HIV/AIDS. *Journal of sports sciences*. 2014;32:785-92.
98. Tuan T-C, Hsu T-G, Fong M-C, Hsu C-F, Tsai KK, Lee C-Y, Kong C-W. Deleterious effects of short-term, high-intensity exercise on immune function: evidence from leucocyte mitochondrial alterations and apoptosis. *British journal of sports medicine*. 2008;42:11-5.
99. Simpson RJ, Kunz H, Agha N, Graff R. Chapter Fifteen-Exercise and the Regulation of Immune Functions. *Progress in molecular biology and translational science*. 2015;135:355-80.

100. d'Ettorre G, Ceccarelli G, Giustini N, Mastroianni CM, Silvestri G, Vullo V. Taming HIV-related inflammation with physical activity: a matter of timing. *AIDS research and human retroviruses*. 2014;30:936-44.
101. Fillipas S, Cicuttini F, Holland AE, Cherry CL. The international physical activity questionnaire overestimates moderate and vigorous physical activity in HIV-infected individuals compared with accelerometry. *Journal of the Association of Nurses in AIDS Care*. 2010;21:173-81.
102. Handy S, Does the Built Environment Influence Physical Activity? Examining the Evidence. *Critical Assessment of the Literature of the Relationships Among Transportation, Land Use and*. 2005: Publisher.
103. Cerin E, Saelens BE, Sallis JF, Frank LD. Neighborhood Environment Walkability Scale: validity and development of a short form. *Medicine and Science in Sports and Exercise*. 2006;38:1682.
104. Oyeyemi AL, Kasoma SS, Onywera VO, Assah F, Adedoyin RA, Conway TL, Moss SJ, Ocansey R, Kolbe-Alexander TL, Akinroye KK. NEWS for Africa: adaptation and reliability of a built environment questionnaire for physical activity in seven African countries. *International Journal of Behavioral Nutrition and Physical Activity*. 2016;13:1.
105. Schatz E, Gilbert L. "My legs affect me a lot.... I can no longer walk to the forest to fetch firewood": Challenges Related to Health and the Performance of Daily Tasks for Older Women in a High HIV Context. *Health Care for Women International*. 2014;35:771-88.
106. Kinsman J, Norris SA, Kahn K, Twine R, Riggle K, Edin K, . . . Micklesfield LK. A model for promoting physical activity among rural South African adolescent girls. *Global Health Action*, 2014;8:28790.

