

Physiological uncoupling of mitochondrial oxidative phosphorylation. Studies in different yeast species

Sergio Guerrero-Castillo · Daniela Araiza-Olivera · Alfredo Cabrera-Orefice ·
Juan Espinasa-Jaramillo · Manuel Gutiérrez-Aguilar · Luis A. Luévano-Martínez ·
Armando Zepeda-Bastida · Salvador Uribe-Carvajal

© Springer Science+Business Media, LLC 2011

Abstract Under non-phosphorylating conditions a high proton transmembrane gradient inhibits the rate of oxygen consumption mediated by the mitochondrial respiratory chain (state IV). Slow electron transit leads to production of reactive oxygen species (ROS) capable of participating in deleterious side reactions. In order to avoid overproducing ROS, mitochondria maintain a high rate of O₂ consumption by activating different exquisitely controlled uncoupling pathways. Different yeast species possess one or more uncoupling systems that work through one of two possible mechanisms: i) Proton sinks and ii) Non-pumping redox enzymes. Proton sinks are exemplified by mitochondrial unspecific channels (MUC) and by uncoupling proteins (UCP). *Saccharomyces cerevisiae* and *Debaryomyces hansenii* express highly regulated MUCs. Also, a UCP was described in *Yarrowia lipolytica* which promotes uncoupled O₂ consumption. Non-pumping alternative oxido-reductases may substitute for a pump, as in *S. cerevisiae* or may coexist with a complete set of pumps as in the branched respiratory chains from *Y. lipolytica* or *D. hansenii*. In addition, pumps may suffer intrinsic uncoupling (slipping). Promising models for study are unicellular parasites which can turn off their aerobic metabolism completely. The variety of energy dissipating systems in eukaryote species is probably designed to control ROS production in the different environments where each species lives.

Keywords Uncoupling · Permeability transition · Uncoupling protein · Branched respiratory chain · *Saccharomyces cerevisiae* · *Yarrowia lipolytica* · *Debaryomyces hansenii*

Introduction

In mitochondria, oxidative phosphorylation results from the coupling between the redox-primary proton pumps in the respiratory chain and the F₁F₀-ATP synthase. The redox H⁺ pumps create a pH gradient (Δ pH) used by the F₁F₀-ATP synthase to phosphorylate ADP. The efficiency of this system varies when electrons enter or exit the respiratory chain at different enzymes or when the H⁺ gradient is used by secondary pumps for the active transport of proteins, ions and metabolites (Nicholls and Ferguson 2002) (Fig. 1).

Three of the four respiratory complexes in an orthodox respiratory chain are proton pumps. These enzymes oxidize substrates, transferring electron(s) to the next acceptor in the chain and expelling H⁺(s) to the intermembrane space. Recycling of the electron within a given pump often results in H⁺/e⁻ stoichiometries higher than 1 (Brandt 2006; Hosler et al. 2006; Trumpower 1990). This high efficiency comes at a price, as redox reactions involve several steps where incomplete reductions transiently convert coenzymes into reactive free radicals (Drose and Brandt 2008; Kushnareva et al. 2002). Therefore, when the mitochondrial ADP concentration drops, the rate of electron flux through the respiratory chain decreases (State IV respiration) and mitochondria become an important source of superoxide and other reactive oxygen species (ROS) (Chen et al. 2003). ROS production has diverse functions, such as signaling and apoptosis (Forman et al. 2010; Perrone et al. 2008).

S. Guerrero-Castillo · D. Araiza-Olivera · A. Cabrera-Orefice ·
J. Espinasa-Jaramillo · M. Gutiérrez-Aguilar ·
L. A. Luévano-Martínez · A. Zepeda-Bastida ·
S. Uribe-Carvajal (✉)
Departamento de Genética Molecular, Instituto de Fisiología
Celular, Universidad Nacional Autónoma de México,
Ciudad Universitaria, Apdo. Postal 70–242, Mexico City, México
DF, Mexico
e-mail: suribe@ifc.unam.mx





Fig. 1 Oxidative phosphorylation efficiency variations due to different systems that use protons. The proton gradient generated by the respiratory chain may be used by (a) the F_1F_0 -ATP synthase (F_0F_1) for ADP phosphorylation (b) the transport of ions or metabolites across the inner mitochondrial membrane (IMM) either as (c) antiporter or (c') symporter. (d) ion uniport. RC, respiratory chain

However, overproduction of ROS may lead to ageing and disease (Drakulic et al. 2005; Wilhelm et al. 2006).

The labile nature of the superoxide radical has made difficult the identification of all its mitochondrial sources. Still, it is known that the ubiquinone and flavin oxidation centers produce ROS (Chen et al. 2003; Starkov et al. 2004; Zundorf et al. 2009). During the redox ubiquinone/ubiquinol reaction, oxidized ubiquinone is partially reduced by one electron in the Q_0 site of the bc_1 complex becoming a potential superoxide source (Drose and Brandt 2008). At a high mitochondrial transmembrane potential, semiquinone accumulates participating in a side reactions that produce ROS (Koshkin et al. 2003; Rottenberg et al. 2009).

In cells and mitochondria there are different enzymes that eliminate ROS, such as Mn^{2+} SOD-dismutases, catalase and glutathione peroxidases. However, ROS overproduction may overwhelm these systems and thus different energy-dissipating uncoupling mechanisms may be activated to prevent such overproduction. These “physiological uncoupling” mechanisms would prevent ROS over-accumulation by inducing increased electron flux (Czarna and Jarmuszkiewicz 2005; Maxwell et al. 1999).

Fig. 2 Physiological uncoupling systems in yeast mitochondria. The rate of oxygen consumption may be accelerated independently of the synthesis of ATP by either depleting the transmembrane pH gradient or by reducing oxygen without contributing to the proton gradient. These mechanisms are present in different yeast species



Among plants, yeast and fungi, there are different strategies aimed at preventing ROS overproduction (Kowaltowski et al. 1998; Magnani et al. 2008). In different yeast species it has been observed that oxidative phosphorylation can be uncoupled by different mechanisms (Fig. 2). Oxidative phosphorylation may be uncoupled through dissipation of the H^+ gradient through proton sinks, also termed extrinsic uncouplers (Kadenbach 2003); these may be channels or transporters and are represented by two well studied systems. These are the yeast mitochondrial unspecific channel (MUC) (Manon et al. 1998), which in mammals is known as the permeability transition pore (PTP) (Haworth and Hunter 1979), and the uncoupling proteins (UCP) (Nicholls and Rial 1999) that specifically dissipate H^+ gradients (Fig. 3). The second respiratory chain uncoupling mechanism, also termed intrinsic uncoupling (Kadenbach 2003) is the catalysis of redox reactions without pumping protons. Non-pumping redox enzymes are widely represented in the branched mitochondrial respiratory chains observed in plants and unicellular organisms (Rasmusson et al. 2004; Umbach and Siedow 2000; Wagner and Moore 1997). Among these enzymes, there are type-II NADH dehydrogenases (NDH2) and alternative oxidases (AOX). In addition, the variations in H^+/e^- stoichiometry (slipping) are another source of uncoupling.

Proton dissipating pathways

The mitochondrial unselective channel

Mitochondrial unspecific channels (MUCs) have been detected in yeast such as *Saccharomyces cerevisiae* (S_c MUC) (Guerin et al. 1994; Prieto et al. 1992) and *Debaryomyces hansenii* (D_h MUC) (Cabrera-Orefice et al. 2010). MUC opening results in a mitochondrial permeability transition (PT) similar to that described in mammals, i.e. a large




Fig. 3 Proton Sinks. Two proton sink systems are exemplified. Once the (a) Respiratory chain establishes a proton gradient, protons may be returned to the matrix through (b) unspecific channels or through (c) uncoupling proteins that are specific protonophores

increase in conductivity that depletes electrochemical gradients (Azzolin et al. 2010).

The s_c MUC has been thoroughly characterized. s_c MUC opens in response to ATP, while it is closed by Pi or ADP (Prieto et al. 1995). This suggests that s_c MUC is controlled by the phosphorylation potential (Wallace et al. 1994). In addition, both the s_c MUC (Perez-Vazquez et al. 2003) and the D_h MUC (Cabrera-Orefice et al. 2010) are closed by Mg^{2+} and by Ca^{2+} . Furthermore, the s_c MUC seems to be controlled cooperatively by Ca^{2+} , Mg^{2+} and Pi [to be published]. In *S. cerevisiae*, a rise in cytoplasmic $[Ca^{2+}]$ precedes processes such as division, mating (Nakajima-Shimada et al. 2000; Ohya et al. 1991); or even a death program resembling apoptosis (Nakajima-Shimada et al. 2000; Ohya et al. 1991; Pozniakovskiy et al. 2005). That is, a rise in $[Ca^{2+}]_{cyt}$ indicates that the cell is about to spend a large amount of energy (Anraku et al. 1991; Manon and Guerin 1998). Both the s_c MUC and the D_h MUC close in response to low [ATP] or high [Pi] while in contrast, when there is a surplus of ATP and no signals this indicates an oncoming need for energy, yeast MUCs open, dissipating the transmembrane potential and thus allowing the rate of oxygen consumption to increase (Prieto et al. 1992) and the production of ROS to decrease (Korshunov et al. 1997).

In *S. cerevisiae*, Ca^{2+} closes MUC, probably through its interaction with the voltage-dependent anionic channel (VDAC) (Gutierrez-Aguilar et al. 2007). The Ca^{2+} -VDAC interaction has also been proposed for vertebrates (Gincel et al. 2001). In both cases, the possibility that VDAC is a regulatory pore component has been suggested (Baines et al. 2007; Gutierrez-Aguilar et al. 2007). In regard to the possible component of MUC in the IMM, in *S. cerevisiae* it has long been evident that Pi is a strong MUC regulator (Azzolin et al. 2010; Cortes et al. 2000; Jung et al. 1997; Manon and Guerin 1997; Prieto et al. 1992; Velours et al. 1977). From this, it should not be surprising that recent evidence suggests that the mitochondrial phosphate carrier (PiC) is a constituent of the s_c MUC: in the absence of PiC, s_c MUC changes its solute size exclusion size and Pi sensitivity (Gutierrez-Aguilar et al. 2010). In mammals,

PiC has also been proposed to be part of this channel (Leung et al. 2008).

Different modulators of MUCs have been reported depending on the species, strain or even tissue under study (Berman et al. 2000; Fortes et al. 2001; Friberg et al. 1999; Manon et al. 1998), suggesting that MUCs have evolved in response to selective pressure, e.g. in *D. hansenii*, the MUC is closed by monovalent cations (Cabrera-Orefice et al. 2010). This closure probably results in higher production of ATP, as it correlates with increased growth rate and mass yield (Gonzalez-Hernandez et al. 2004) and probably constitutes an adaptation to the high Na^+ contents of sea water (Gustafsson and Norkrans 1976).

Adding to the ongoing debate on the physiological role of MUCs, it is suggested that their role as physiological uncouplers should be considered; i.e. MUCs probably are highly regulated energy dissipative systems that decrease mitochondrial gradients when the demand for energy is low.

PT does not seem to be universal. *Yarrowia lipolytica* and *Endomyces magnusii* undergo PT only upon forced conditions which include incubation with the Ca^{2+} ionophore ETH129 (Kovaleva et al. 2009; Yamada et al. 2009). If MUC-mediated uncoupling is important to inhibit ROS production, and *Y. lipolytica* and *E. magnusii* seem to lack such a structure, then these yeast species should possess alternative uncoupling systems. Indeed, in *Y. lipolytica* mitochondria there are two such systems that might function as uncouplers: an uncoupling protein (Luevano-Martinez et al. 2010) and a branched respiratory chain (Guerrero-Castillo et al. 2009; Kerscher et al. 2002).

Uncoupling proteins

Uncoupling protein (UCP)-like activities have been detected in mitochondria from unicellular organisms, higher eukaryotes and plants (Jarmuszkiewicz et al. 2010). The physiological role of UCPs in unicellular organisms is still debated: the small size of unicellular eukaryotes makes a thermogenic role unlikely, as it is impossible to form a temperature gradient between the cell and the environment although, in *Acanthamoeba castellanii* UCP expression does increase in cells growing at 4 °C (Jarmuszkiewicz et al. 2004). Here, it is proposed that unicellular UCPs are capable of decreasing the mitochondrial $\Delta\Psi$ with the aim of decreasing production of ROS. Also, in unicellular organisms resistance to exogenous ROS is enhanced by UCP activity (Kowaltowski et al. 1998; Ricquier 2005), probably because UCP decreases endogenous ROS production (Krauss et al. 2005) and thus detoxifying enzymes are free to deal with the exogenous species: e.g. strains of *Candida albicans* devoid of UCP are less invasive than the wild type (Cavalheiro et al. 2004; Cheng et al. 2007).

In addition to the available functional evidence, recently a protein exhibiting UCP-like activity was identified in *Y. lipolytica* (Luevano-Martinez et al. 2010). The UCP activity was regulated similarly to the UCP1 from brown adipose tissue. After an extensive phylogenetic search for a UCP ortholog in this yeast, it was demonstrated that the mitochondrial oxaloacetate carrier (OAC) from *Y. lipolytica* is a *bona fide* UCP. The *Y. lipolytica* OAC displayed both, a sulfate/oxaloacetate transport and a UCP behavior. It is noteworthy that in the unicellular organisms where UCP activity has been reported, the green algae *Chlamydomonas reinhardtii*, the amoeba *Dictyostelium discoideum* (DictyBase) and the yeast *Candida albicans* (Cavalheiro et al. 2004; Jarmuszkiewicz et al. 2002) the only UCP-like proteins seem to be the mitochondrial oxaloacetate carriers (results not published). In regard to whether a UCP might prevent ROS overproduction, in *Y. lipolytica*, it has been demonstrated that this protein is over-expressed in the stationary phase, where a degree of uncoupling would be needed to maintain a high rate of oxygen consumption in the absence of ATP synthesis (Luevano-Martinez et al. 2010).

Redox enzymes that do not pump protons

Branched mitochondrial respiratory chains

Redox enzymes lacking pumping activity are constituted by a single protein subunit. These enzymes probably appeared early in the reducing world, before the appearance of oxidative phosphorylation, fulfilling the need to detoxify oxygen from the vicinity of enzymes and membranes. Some prokaryotes still use oxidoreductase-mediated detoxification of oxygen to protect their fragile nitrogen reducing enzymes (Flores-Encarnacion et al. 1999).

Alternative redox enzymes do not contribute to the proton gradient. Branched mitochondrial respiratory chains may contain a number of different enzymes that donate electrons to the quinone pool including complex I (the only proton pump), succinate dehydrogenase, glycerol phosphate dehydrogenase, dihydroorotate dehydrogenase and internal or external type II NADH dehydrogenases. Then the electrons in reduced ubiquinol follow two possible pathways reaching either the cytochrome pathway (complexes III and IV), or the alternative oxidase (AOX). In these respiratory chains, different electron pathways may be envisioned that bypass energy-conserving respiratory complexes I, III and/or IV, i.e. branched chains seem to be able to reduce oxygen while using 0, 1, 2 or 3 proton pumps (Fig. 4).

In mitochondria, the most widely distributed monosubunit redox enzymes are type II NADH dehydrogenases

(NDH2) and alternative oxidases (AOX). NDH2s may be located on either surface of the IMM. External NDH2s (NDH2e) oxidize cytosolic NADH, while internal NDH2s (NDH2i) oxidize NADH from the matrix in a rotenone-insensitive reaction. The structure (Fisher et al. 2007; Fisher et al. 2009; Gonzalez-Meler et al. 1999; Kerscher 2000; Melo et al. 2004; Schmid and Gerloff 2004) and kinetics (Fisher et al. 2009; Velazquez and Pardo 2001) of NDH2s from different organisms have been reported. AOX is a single subunit enzyme (Albury et al. 2002; Andersson and Nordlund 1999; Berthold et al. 2000; Moore and Siedow 1991). AOX activity is regulated by nucleotides, by dimerization and/or by α -ketoacids (Hoefnagel et al. 1995; Millar et al. 1993; Millenaar et al. 1998). Some yeast species contain two AOX isoforms, one being constitutively expressed and a second one induced by stress (Siedow and Umbach 2000). It is noteworthy that AOX is present only in fungi that express complex I, possibly because in a respiratory chain without Complex I, any electron reaching AOX would be totally unproductive (Joseph-Horne et al. 2001).

In mitochondria with alternative components, the pathway that electrons follow has to be strictly controlled. A direct reaction between NDH2, ubiquinone and AOX would result in a non-productive, uncoupled pathway, i.e. no protons would be pumped. Furthermore, at the external face of the inner membrane, NDH2 receives the hydride from NADH and takes one H^+ , transferring both hydrogen atoms to ubiquinone. Then ubiquinone is regenerated by AOX which in turn transfers its hydrogen atoms to oxygen producing water. This sequence of reactions results in the dissipation of a H^+ , i.e. it has a H^+/e^- pumping stoichiometry of -0.5 . Therefore, when energy is required, alternative redox enzymes need to be isolated from each other, probably by binding to the proton-pumping complexes. In contrast, when phosphorylation is not active, as in the stationary phase, the non-producing electron transfer between NDH2 and AOX would be useful to maintain a high rate of oxygen consumption at a high transmembrane potential, preventing semiquinone accumulation and decreasing ROS formation (Joseph-Horne et al. 2001).

Proton/electron stoichiometry variations. Slipping

Non-branched respiratory chains seem to use other mechanisms to regulate the efficiency of oxidative phosphorylation (van Dam et al. 1990). Uncoupling may result from increased proton conductance at the lipid bilayer (Luvisetto and Azzone 1989; Luvisetto et al. 1991). A second mechanism would be the decrease in the efficiency of a respiratory pump (slipping) (Pietrobon et al. 1981; Pietrobon et al. 1983). Intrinsic uncoupling or slipping is defined as a decrease in the efficiency of a




Fig. 4 In a branched respiratory chain the number of proton pumps participating in electron transfer may vary from three to zero. In branched respiratory chains electrons may follow different routes to reach oxygen. Thus the number of proton pumps involved may change: **a** complexes I, III and IV: three proton pumps are involved. **b** from succinate dehydrogenase through the cytochrome pathway, two proton pumps. **c** from NDH2e through complexes III-IV; two proton pumps, although H^+/e^- is 2.5 instead of 3 as in **(b)**. **d** from complex I though AOX; one pump. **e** from succinate dehydrogenase through AOX; No proton pumps

proton pump (decrease of the H^+/e^- or H^+/ATP stoichiometry) resulting in a diminished P/O ratio (Kadenbach 2003).

Slipping has been reported in cytochrome c oxidase (Azzone et al. 1985; Frank and Kadenbach 1996). The F_1F_0 -synthase can also undergo slipping, hydrolyzing ATP without pumping protons (Feniouk et al. 2005). In addition, protons can reenter the matrix through the pumps without moving electrons backwards or making ATP (Pietrobon et al. 1983).

Slipping accelerates the rate of oxygen consumption as more electrons are needed to maintain a high ΔpH . Normally in the proton pump the chemical reaction and the transport of protons are tightly coupled, while during slipping both processes become independent (Mourier et al. 2010). Upon slipping, the rate of electron flux increases while the proton motive force remains constant and energy is dissipated as heat (Kadenbach 2003).

In *S. cerevisiae* mitochondria, a remarkable change in the stoichiometry of proton pumping has been described. Feeding the respiratory chain with substrates for different quinone reductases leads to an increase in the rate of

participate. **f** NDH2e through AOX; zero proton pumps participate and in addition, the combined activity of NDH2e with AOX would consume a H^+ from the intermembrane space, yielding a negative stoichiometry of $-0.5 H^+/e^-$. Numbers I, II, III₂ and IV represent each of the four respiratory complexes; NDH2e, external NADH dehydrogenase; AOX, alternative oxidase; IMM, inner mitochondrial membrane. Protons in red are used for ubiquinone reduction in the intermembrane side of the IMM, i.e. they do not contribute **(c)** or contribute negatively to the H^+/e^- stoichiometry **(f)**

oxygen consumption without increasing the rate of ATP phosphorylation (Mourier et al. 2010). This phenomenon has been termed active leak and is probably due to slipping of an oxidative phosphorylation pump, although an increase in the proton conductance of the bilayer has not been ruled out.

Interactions between the cytoplasm and mitochondria regulate the efficiency of oxidative phosphorylation

At any given moment the cell's energy needs determine which metabolic pathways are activated or inhibited (Devin and Rigoulet 2007). The catabolism/anabolism activity ratio is determined by metabolic fluxes (Cascaete et al. 1994; Moreno-Sanchez et al. 2010; Ovadi and Saks 2004; Srere 1987). Upon oxygenation, the rate of glycolysis decreases. This may be explained by the allosteric regulation of glycolytic enzymes by ATP and fructose 2,6-bisphosphate and by the competition for ADP and for reducing equivalents observed between glycolysis and

oxidative phosphorylation (Beauvoit et al. 1993; Gosalvez et al. 1974).

In *Saccharomyces cerevisiae*, glycolysis is the main source of ATP; however, in the presence of non-fermentable substrates oxidative phosphorylation becomes the main energy source. During fermentation the genes that encode for oxidative metabolism enzymes stop their expression (Takeda 1981), e.g. glucose addition inhibits the expression of cytochrome c (Thevelein 1994; Zitomer and Nichols 1978), while glycolytic intermediates are accumulated to induce the expression of glycolytic enzymes (Boles et al. 1993).

In *S. cerevisiae* the addition of glucose induces the transition to fermentative metabolism, where glycolysis is increased and oxidative phosphorylation is decreased (den Hollander et al. 1986). This is the Crabtree effect. There are both Crabtree-positive and negative yeast species. Recent studies indicate that fructose 1,6-bisphosphate inhibits oxygen consumption through an interaction with complexes III and IV. In contrast, physiological concentrations of glucose 6-phosphate and fructose 6-phosphate stimulate the respiratory flux, possibly inducing slipping (Diaz-Ruiz et al. 2008).

Unicellular organisms other than yeast

Protists make up the bulk of the eukaryotes, while vertebrates and fungi represent only a small fraction. Protists present a wide variety of physiological properties. There are very few bioenergetics studies on these organisms. *Giardia lamblia* (Hashimoto et al. 1994) and *Entamoeba histolytica* (Tovar et al. 1999) have lost their mitochondria. Other protists, such as some Trichomonadidae and ciliates, have organelles called hydrogenosomes, which are related to mitochondria (de Souza et al. 2009; Mather and Vaidya 2008).

Unicellular parasites have evolved to adapt their metabolism for survival within the host. Depending on the environment and stage in their life cycle, *Plasmodium*, *Trypanosoma* and *Leishmania* can make a complete switch from a glycolytic to an aerobic metabolism and back, such that in *Plasmodium falciparum* the activities of complex III, IV and dihydroorotate dehydrogenase, are 10 times higher in the sexual than in the asexual stage (Monzote and Gille 2010). Likewise, mitochondria have adapted to the metabolic conditions found within the host, e.g. in the mosquito, *Plasmodium* gametocytes are aerobic and mitochondria are typical. In contrast, in the vertebrate host, sporozoites and merozoites are adapted to microaerophilia and contain few, underdeveloped mitochondria (Segura and Blair 2003).

Throughout the trypanosomatid life cycle, mitochondrial activity varies widely (Schneider 2001). In the bloodstream,

these protozoans are anaerobic while in the gut of the insect they perform oxidative phosphorylation. In *Toxoplasma* most energy is obtained from glycolysis, although the mitochondrial DNA sequence of these parasites shows significant differences from the mammalian host, suggesting possible drug targets (Monzote and Gille 2010). Remarkably, the mitochondrial DNAs from trypanosomatids and Apicomplexa lack genes for transfer RNA (Mather and Vaidya 2008).

Concluding remarks

Aerobic metabolism is at the same time highly efficient and very dangerous. The reactive oxygen species produced by the respiratory chain can react with, and damage different components of the cell. Diverse mechanisms have evolved to prevent the deleterious effect of ROS. There are many detoxifying enzymes such as glutathione reductase, superoxide dismutase or catalase. In addition, upriver from these reactions, there are diverse mitochondrial systems designed to prevent ROS overproduction. These systems promote physiological uncoupling to ensure that the redox enzymes in the respiratory chain work at a fast rate, thus preventing reactive intermediates from participating in collateral reactions.

There are two mitochondrial uncoupling mechanisms: a) Those that dissipate the pH gradient and b) Non-productive redox reactions. Both mechanisms are widely spread in nature. Physiological proton sinks are the uncoupling proteins and the mitochondrial unspecific channels, while non productive redox reactions are catalyzed by redox/non-pumping alternative dehydrogenases and by orthodox complexes that undergo slipping.

The relationship between the cytoplasmic and the mitochondrial metabolic pathways needs to be better understood. The ability of some products from glycolysis to regulate oxidative phosphorylation is illustrative. The comparison between Crabtree positive and Crabtree negative yeast species may help understand the mechanisms and consequences of these interactions.

Understanding the mechanisms underlying the control and production of ROS may help to select more resistant organisms for biotechnological applications. Also, various ROS-related diseases have to be understood in order to design better treatments. In this light, it seems useful to know that uncoupling prevents ROS production.

During evolution, each eukaryote species preserved one or more ROS overproduction-prevention mechanism(s). Yeast species are ideal to study each mechanism. Other unicellular organisms may be helpful to understand their ability to shut down aerobic metabolism without being overwhelmed by ROS production.

Acknowledgements Partially funded by grants from CONACYT 79989 and by DGAPA/UNAM, IN217109. SGC, DAO, ACO, MGA and LALM are CONACYT fellows enrolled in the Biochemistry Graduate Program at UNAM. JEJ has a SNI-III aid fellowship. The assistance of Dr Natalia Chiquete-Félix is acknowledged. We thank Dr Soledad Funes-Argüello for critically reading the manuscript.

References

- Albury MS, Affourtit C, Crichton PG, Moore AL (2002) Structure of the plant alternative oxidase. Site-directed mutagenesis provides new information on the active site and membrane topology. *J Biol Chem* 277:1190–1194
- Andersson ME, Nordlund P (1999) A revised model of the active site of alternative oxidase. *FEBS Lett* 449:17–22
- Anraku Y, Ohya Y, Iida H (1991) Cell cycle control by calcium and calmodulin in *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 1093:169–177
- Azzolin L, von Stockum S, Basso E, Petronilli V, Forte MA, Bernardi P (2010) The mitochondrial permeability transition from yeast to mammals. *FEBS Lett* 584:2504–2509
- Azzone GF, Zoratti M, Petronilli V, Pietrobon D (1985) The stoichiometry of H⁺ pumping in cytochrome oxidase and the mechanism of uncoupling. *J Inorg Biochem* 23:349–356
- Baines CP, Kaiser RA, Sheiko T, Craigen WJ, Molkenin JD (2007) Voltage-dependent anion channels are dispensable for mitochondrial-dependent cell death. *Nat Cell Biol* 9:550–555
- Beauvoit B, Rigoulet M, Bunoust O, Raffard G, Canioni P, Guerin B (1993) Interactions between glucose metabolism and oxidative phosphorylations on respiratory-competent *Saccharomyces cerevisiae* cells. *Eur J Biochem* 214:163–172
- Berman SB, Watkins SC, Hastings TG (2000) Quantitative biochemical and ultrastructural comparison of mitochondrial permeability transition in isolated brain and liver mitochondria: evidence for reduced sensitivity of brain mitochondria. *Exp Neurol* 164:415–425
- Berthold DA, Andersson ME, Nordlund P (2000) New insight into the structure and function of the alternative oxidase. *Biochim Biophys Acta* 1460:241–254
- Boles E, Heinisch J, Zimmermann FK (1993) Different signals control the activation of glycolysis in the yeast *Saccharomyces cerevisiae*. *Yeast* 9:761–770
- Brandt U (2006) Energy converting NADH:quinone oxidoreductase (complex I). *Annu Rev Biochem* 75:69–92
- Cabrera-Orefice A, Guerrero-Castillo S, Luevano-Martinez LA, Pena A, Uribe-Carvajal S (2010) Mitochondria from the salt-tolerant yeast *Debaryomyces hansenii* (halophilic organelles?). *J Bioenerg Biomembr* 42:11–19
- Cascante M, Sorribas A, Canela EI (1994) Enzyme-enzyme interactions and metabolite channelling: alternative mechanisms and their evolutionary significance. *Biochem J* 298(Pt 2):313–320
- Cavalheiro RA, Fortes F, Borecky J, Faustini VC, Schreiber AZ, Vercesi AE (2004) Respiration, oxidative phosphorylation, and uncoupling protein in *Candida albicans*. *Braz J Med Biol Res* 37:1455–1461
- Cortes P, Castrejon V, Sampedro JG, Uribe S (2000) Interactions of arsenate, sulfate and phosphate with yeast mitochondria. *Biochim Biophys Acta* 1456:67–76
- Chen Q, Vazquez EJ, Moghaddas S, Hoppel CL, Lesnfsky EJ (2003) Production of reactive oxygen species by mitochondria: central role of complex III. *J Biol Chem* 278:36027–36031
- Cheng S, Clancy CJ, Zhang Z, Hao B, Wang W, Iczkowski KA, Pfaller MA, Nguyen MH (2007) Uncoupling of oxidative phosphorylation enables *Candida albicans* to resist killing by phagocytes and persist in tissue. *Cell Microbiol* 9:492–501
- Czarna M, Jarmuszkiewicz W (2005) Activation of alternative oxidase and uncoupling protein lowers hydrogen peroxide formation in amoeba *Acanthamoeba castellanii* mitochondria. *FEBS Lett* 579:3136–3140
- de Souza W, Attias M, Rodrigues JC (2009) Particularities of mitochondrial structure in parasitic protists (Apicomplexa and Kinetoplastida). *Int J Biochem Cell Biol* 41:2069–2080
- den Hollander JA, Ugurbil K, Brown TR, Bednar M, Redfield C, Shulman RG (1986) Studies of anaerobic and aerobic glycolysis in *Saccharomyces cerevisiae*. *Biochemistry* 25:203–211
- Devin A, Rigoulet M (2007) Mechanisms of mitochondrial response to variations in energy demand in eukaryotic cells. *Am J Physiol Cell Physiol* 292:C52–C58
- Diaz-Ruiz R, Averet N, Araiza D, Pinson B, Uribe-Carvajal S, Devin A, Rigoulet M (2008) Mitochondrial oxidative phosphorylation is regulated by fructose 1,6-bisphosphate. A possible role in Crabtree effect induction? *J Biol Chem* 283:26948–26955
- Drakulic T, Temple MD, Guido R, Jarolim S, Breitenbach M, Attfield PV, Dawes IW (2005) Involvement of oxidative stress response genes in redox homeostasis, the level of reactive oxygen species, and ageing in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 5:1215–1228
- Drose S, Brandt U (2008) The mechanism of mitochondrial superoxide production by the cytochrome bc₁ complex. *J Biol Chem* 283:21649–21654
- Feniouk BA, Mulikidjanian AY, Junge W (2005) Proton slip in the ATP synthase of *Rhodobacter capsulatus*: induction, proton conduction, and nucleotide dependence. *Biochim Biophys Acta* 1706:184–194
- Fisher N, Bray PG, Ward SA, Biagini GA (2007) The malaria parasite type II NADH:quinone oxidoreductase: an alternative enzyme for an alternative lifestyle. *Trends Parasitol* 23:305–310
- Fisher N, Warman AJ, Ward SA, Biagini GA (2009) Chapter 17 Type II NADH: quinone oxidoreductases of *Plasmodium falciparum* and *Mycobacterium tuberculosis* kinetic and high-throughput assays. *Meth Enzymol* 456:303–320
- Flores-Encarnacion M, Contreras-Zentella M, Soto-Urzuu L, Aguilar GR, Baca BE, Escamilla JE (1999) The respiratory system and diazotrophic activity of *Acetobacter diazotrophicus* PAL5. *J Bacteriol* 181:6987–6995
- Forman HJ, Maiorino M, Ursini F (2010) Signaling functions of reactive oxygen species. *Biochemistry* 49:835–842
- Fortes F, Castilho RF, Catisti R, Camieri EG, Vercesi AE (2001) Ca²⁺ induces a cyclosporin A-insensitive permeability transition pore in isolated potato tuber mitochondria mediated by reactive oxygen species. *J Bioenerg Biomembr* 33:43–51
- Frank V, Kadenbach B (1996) Regulation of the H⁺/e⁻ stoichiometry of cytochrome c oxidase from bovine heart by intramitochondrial ATP/ADP ratios. *FEBS Lett* 382:121–124
- Friberg H, Connern C, Halestrap AP, Wieloch T (1999) Differences in the activation of the mitochondrial permeability transition among brain regions in the rat correlate with selective vulnerability. *J Neurochem* 72:2488–2497
- Gincel D, Zaid H, Shoshan-Barmatz V (2001) Calcium binding and translocation by the voltage-dependent anion channel: a possible regulatory mechanism in mitochondrial function. *Biochem J* 358:147–155
- Gonzalez-Hernandez JC, Cardenas-Monroy CA, Pena A (2004) Sodium and potassium transport in the halophilic yeast *Debaryomyces hansenii*. *Yeast* 21:403–412
- Gonzalez-Meler MA, Ribas-Carbo M, Giles L, Siedow JN (1999) The effect of growth and measurement temperature on the activity of the alternative respiratory pathway. *Plant Physiol* 120:765–772
- Gosalvez M, Perez-Garcia J, Weinhouse S (1974) Competition for ADP between pyruvate kinase and mitochondrial oxidative phosphorylation as a control mechanism in glycolysis. *Eur J Biochem* 46:133–140

- Guerin B, Bunoust O, Rouqueys V, Rigoulet M (1994) ATP-induced unspecific channel in yeast mitochondria. *J Biol Chem* 269:25406–25410
- Guerrero-Castillo S, Vazquez-Acevedo M, Gonzalez-Halphen D, Uribe-Carvajal S (2009) In *Yarrowia lipolytica* mitochondria, the alternative NADH dehydrogenase interacts specifically with the cytochrome complexes of the classic respiratory pathway. *Biochim Biophys Acta* 1787:75–85
- Gustafsson L, Norkrans B (1976) On the mechanism of salt tolerance. Production of glycerol and heat during growth of *Debaryomyces hansenii*. *Arch Microbiol* 110:177–183
- Gutierrez-Aguilar M, Perez-Vazquez V, Bunoust O, Manon S, Rigoulet M, Uribe S (2007) In yeast, Ca²⁺ and octylguanidine interact with porin (VDAC) preventing the mitochondrial permeability transition. *Biochim Biophys Acta* 1767:1245–1251
- Gutierrez-Aguilar M, Perez-Martinez X, Chavez E, Uribe-Carvajal S (2010) In *Saccharomyces cerevisiae*, the phosphate carrier is a component of the mitochondrial unselective channel. *Arch Biochem Biophys* 494:184–191
- Hashimoto T, Nakamura Y, Nakamura F, Shirakura T, Adachi J, Goto N, Okamoto K, Hasegawa M (1994) Protein phylogeny gives a robust estimation for early divergences of eukaryotes: phylogenetic place of a mitochondria-lacking protozoan, *Giardia lamblia*. *Mol Biol Evol* 11:65–71
- Haworth RA, Hunter DR (1979) The Ca²⁺-induced membrane transition in mitochondria. II. Nature of the Ca²⁺ trigger site. *Arch Biochem Biophys* 195:460–467
- Hoefnagel MH, Millar AH, Wiskich JT, Day DA (1995) Cytochrome and alternative respiratory pathways compete for electrons in the presence of pyruvate in soybean mitochondria. *Arch Biochem Biophys* 318:394–400
- Hosler JP, Ferguson-Miller S, Mills DA (2006) Energy transduction: proton transfer through the respiratory complexes. *Annu Rev Biochem* 75:165–187
- Jarmuszkiewicz W, Behrendt M, Navet R, Sluse FE (2002) Uncoupling protein and alternative oxidase of *Dictyostelium discoideum*: occurrence, properties and protein expression during vegetative life and starvation-induced early development. *FEBS Lett* 532:459–464
- Jarmuszkiewicz W, Antos N, Swida A, Czarna M, Sluse FE (2004) The effect of growth at low temperature on the activity and expression of the uncoupling protein in *Acanthamoeba castellanii* mitochondria. *FEBS Lett* 569:178–184
- Jarmuszkiewicz W, Woyda-Ploszczyca A, Antos-Krzeminska N, Sluse FE (2010) Mitochondrial uncoupling proteins in unicellular eukaryotes. *Biochim Biophys Acta* 1797:792–799
- Joseph-Home T, Hollomon DW, Wood PM (2001) Fungal respiration: a fusion of standard and alternative components. *Biochim Biophys Acta* 1504:179–195
- Jung DW, Bradshaw PC, Pfeiffer DR (1997) Properties of a cyclosporin-insensitive permeability transition pore in yeast mitochondria. *J Biol Chem* 272:21104–21112
- Kadenbach B (2003) Intrinsic and extrinsic uncoupling of oxidative phosphorylation. *Biochim Biophys Acta* 1604:77–94
- Kerscher SJ (2000) Diversity and origin of alternative NADH:ubiquinone oxidoreductases. *Biochim Biophys Acta* 1459:274–283
- Kerscher S, Drose S, Zwicker K, Zickermann V, Brandt U (2002) *Yarrowia lipolytica*, a yeast genetic system to study mitochondrial complex I. *Biochim Biophys Acta* 1555:83–91
- Korshunov SS, Skulachev VP, Starkov AA (1997) High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria. *FEBS Lett* 416:15–18
- Koshkin V, Wang X, Scherer PE, Chan CB, Wheeler MB (2003) Mitochondrial functional state in clonal pancreatic beta-cells exposed to free fatty acids. *J Biol Chem* 278:19709–19715
- Kovaleva MV, Sukhanova EI, Trendeleva TA, Zyl'kova MV, Ural'skaya LA, Popova KM, Saris NE, Zvyagil'skaya RA (2009) Induction of a non-specific permeability transition in mitochondria from *Yarrowia lipolytica* and *Dipodascus (Endomyces) magnusii* yeasts. *J Bioenerg Biomembr* 41:239–249
- Kowaltowski AJ, Costa AD, Vercesi AE (1998) Activation of the potato plant uncoupling mitochondrial protein inhibits reactive oxygen species generation by the respiratory chain. *FEBS Lett* 425:213–216
- Krauss S, Zhang CY, Lowell BB (2005) The mitochondrial uncoupling-protein homologues. *Nat Rev Mol Cell Biol* 6:248–261
- Kushnareva Y, Murphy AN, Andreyev A (2002) Complex I-mediated reactive oxygen species generation: modulation by cytochrome c and NAD(P)⁺ oxidation-reduction state. *Biochem J* 368:545–553
- Leung AW, Varanyuwatana P, Halestrap AP (2008) The mitochondrial phosphate carrier interacts with cyclophilin D and may play a key role in the permeability transition. *J Biol Chem* 283:26312–26323
- Luevano-Martinez LA, Moyano E, de Lacoba MG, Rial E, Uribe-Carvajal S (2010) Identification of the mitochondrial carrier that provides *Yarrowia lipolytica* with a fatty acid-induced and nucleotide-sensitive uncoupling protein-like activity. *Biochim Biophys Acta* 1797:81–88
- Luvisetto S, Azzone GF (1989) Local protons and uncoupling of aerobic and artificial delta muH-driven ATP synthesis. *Biochemistry* 28:1109–1116
- Luvisetto S, Conti E, Buso M, Azzone GF (1991) Flux ratios and pump stoichiometries at sites II and III in liver mitochondria. Effect of slips and leaks. *J Biol Chem* 266:1034–1042
- Magnani T, Soriani FM, Martins Vde P, Policarpo AC, Sorgi CA, Faccioli LH, Curti C, Uyemura SA (2008) Silencing of mitochondrial alternative oxidase gene of *Aspergillus fumigatus* enhances reactive oxygen species production and killing of the fungus by macrophages. *J Bioenerg Biomembr* 40:631–636
- Manon S, Guerin M (1997) The ATP-induced K⁽⁺⁾-transport pathway of yeast mitochondria may function as an uncoupling pathway. *Biochim Biophys Acta* 1318:317–321
- Manon S, Guerin M (1998) Investigation of the yeast mitochondrial unselective channel in intact and permeabilized spheroplasts. *Biochem Mol Biol Int* 44:565–575
- Manon S, Roucou X, Guerin M, Rigoulet M, Guerin B (1998) Characterization of the yeast mitochondria unselective channel: a counterpart to the mammalian permeability transition pore? *J Bioenerg Biomembr* 30:419–429
- Mather MW, Vaidya AB (2008) Mitochondria in malaria and related parasites: ancient, diverse and streamlined. *J Bioenerg Biomembr* 40:425–433
- Maxwell DP, Wang Y, McIntosh L (1999) The alternative oxidase lowers mitochondrial reactive oxygen production in plant cells. *Proc Natl Acad Sci USA* 96:8271–8276
- Melo AM, Bandejas TM, Teixeira M (2004) New insights into type II NAD(P)H:quinone oxidoreductases. *Microbiol Mol Biol Rev* 68:603–616
- Millar AH, Wiskich JT, Whelan J, Day DA (1993) Organic acid activation of the alternative oxidase of plant mitochondria. *FEBS Lett* 329:259–262
- Millenaar FF, Benschop JJ, Wagner AM, Lambers H (1998) The role of the alternative oxidase in stabilizing the in vivo reduction state of the ubiquinone pool and the activation state of the alternative oxidase. *Plant Physiol* 118:599–607
- Monzote L, Gille L (2010) Mitochondria as a promising antiparasitic target. *Curr Clin Pharmacol* 5:55–60
- Moore AL, Siedow JN (1991) The regulation and nature of the cyanide-resistant alternative oxidase of plant mitochondria. *Biochim Biophys Acta* 1059:121–140

- Moreno-Sanchez R, Saavedra E, Rodriguez-Enriquez S, Gallardo-Perez JC, Quezada H, Westerhoff HV (2010) Metabolic control analysis indicates a change of strategy in the treatment of cancer. *Mitochondrion* 10:626–639
- Mourier A, Devin A, Rigoulet M (2010) Active proton leak in mitochondria: a new way to regulate substrate oxidation. *Biochim Biophys Acta* 1797:255–261
- Nakajima-Shimada J, Sakaguchi S, Tsuji FI, Anraku Y, Iida H (2000) Ca²⁺ signal is generated only once in the mating pheromone response pathway in *Saccharomyces cerevisiae*. *Cell Struct Funct* 25:125–131
- Nicholls DG, Ferguson SJ (2002) *Bioenergetics* 3. Academic Press, London, ISBN 0125181213
- Nicholls DG, Rial E (1999) A history of the first uncoupling protein, UCP1. *J Bioenerg Biomembr* 31:399–406
- Ohya Y, Umemoto N, Tanida I, Ohta A, Iida H, Anraku Y (1991) Calcium-sensitive cts mutants of *Saccharomyces cerevisiae* showing a Pet⁻ phenotype are ascribable to defects of vacuolar membrane H⁽⁺⁾-ATPase activity. *J Biol Chem* 266:13971–13977
- Ovadi J, Saks V (2004) On the origin of intracellular compartmentation and organized metabolic systems. *Mol Cell Biochem* 256–257:5–12
- Perez-Vazquez V, Saavedra-Molina A, Uribe S (2003) In *Saccharomyces cerevisiae*, cations control the fate of the energy derived from oxidative metabolism through the opening and closing of the yeast mitochondrial unselective channel. *J Bioenerg Biomembr* 35:231–241
- Perrone GG, Tan SX, Dawes IW (2008) Reactive oxygen species and yeast apoptosis. *Biochim Biophys Acta* 1783:1354–1368
- Pietrobon D, Azzone GF, Walz D (1981) Effect of funiculosin and antimycin A on the redox-driven H⁺ pumps in mitochondria: on the nature of ‘leaks’. *Eur J Biochem* 117:389–394
- Pietrobon D, Zoratti M, Azzone GF (1983) Molecular slipping in redox and ATPase H⁺ pumps. *Biochim Biophys Acta* 723:317–321
- Pozniakovskiy AI, Knorre DA, Markova OV, Hyman AA, Skulachev VP, Severin FF (2005) Role of mitochondria in the pheromone- and amiodarone-induced programmed death of yeast. *J Cell Biol* 168:257–269
- Prieto S, Bouillaud F, Ricquier D, Rial E (1992) Activation by ATP of a proton-conducting pathway in yeast mitochondria. *Eur J Biochem* 208:487–491
- Prieto S, Bouillaud F, Rial E (1995) The mechanism for the ATP-induced uncoupling of respiration in mitochondria of the yeast *Saccharomyces cerevisiae*. *Biochem J* 307(Pt 3):657–661
- Rasmusson AG, Soole KL, Elthon TE (2004) Alternative NAD(P)H dehydrogenases of plant mitochondria. *Annu Rev Plant Biol* 55:23–39
- Ricquier D (2005) Respiration uncoupling and metabolism in the control of energy expenditure. *Proc Nutr Soc* 64:47–52
- Rottenberg H, Covian R, Trumppower BL (2009) Membrane potential greatly enhances superoxide generation by the cytochrome bcl complex reconstituted into phospholipid vesicles. *J Biol Chem* 284:19203–19210
- Schmid R, Gerloff DL (2004) Functional properties of the alternative NADH:ubiquinone oxidoreductase from *E. coli* through comparative 3-D modelling. *FEBS Lett* 578:163–168
- Schneider A (2001) Unique aspects of mitochondrial biogenesis in trypanosomatids. *Int J Parasitol* 31:1403–1415
- Segura C, Blair S (2003) Mitochondria in the Plasmodium genera. *Biomedica* 23:351–363
- Siedow JN, Umbach AL (2000) The mitochondrial cyanide-resistant oxidase: structural conservation amid regulatory diversity. *Biochim Biophys Acta* 1459:432–439
- Srere PA (1987) Complexes of sequential metabolic enzymes. *Annu Rev Biochem* 56:89–124
- Starkov AA, Fiskum G, Chinopoulos C, Lorenzo BJ, Browne SE, Patel MS, Beal MF (2004) Mitochondrial alpha-ketoglutarate dehydrogenase complex generates reactive oxygen species. *J Neurosci* 24:7779–7788
- Takeda M (1981) Glucose-induced inactivation of mitochondrial enzymes in the yeast *Saccharomyces cerevisiae*. *Biochem J* 198:281–287
- Thevelein JM (1994) Signal transduction in yeast. *Yeast* 10:1753–1790
- Tovar J, Fischer A, Clark CG (1999) The mitosome, a novel organelle related to mitochondria in the amitochondrial parasite *Entamoeba histolytica*. *Mol Microbiol* 32:1013–1021
- Trumppower BL (1990) Cytochrome bcl complexes of microorganisms. *Microbiol Rev* 54:101–129
- Umbach AL, Siedow JN (2000) The cyanide-resistant alternative oxidases from the fungi *Pichia stipitis* and *Neurospora crassa* are monomeric and lack regulatory features of the plant enzyme. *Arch Biochem Biophys* 378:234–245
- van Dam K, Shinohara Y, Unami A, Yoshida K, Terada H (1990) Slipping pumps or proton leaks in oxidative phosphorylation. The local anesthetic bupivacaine causes slip in cytochrome c oxidase of mitochondria. *FEBS Lett* 277:131–133
- Velazquez I, Pardo JP (2001) Kinetic characterization of the rotenone-insensitive internal NADH: ubiquinone oxidoreductase of mitochondria from *Saccharomyces cerevisiae*. *Arch Biochem Biophys* 389:7–14
- Velours J, Rigoulet M, Guerin B (1977) Protection of yeast mitochondrial structure by phosphate and other H⁺ donating anions. *FEBS Lett* 81:18–22
- Wagner AM, Moore AL (1997) Structure and function of the plant alternative oxidase: its putative role in the oxygen defence mechanism. *Biosci Rep* 17:319–333
- Wallace PG, Pedler SM, Wallace JC, Berry MN (1994) A method for the determination of the cellular phosphorylation potential and glycolytic intermediates in yeast. *Anal Biochem* 222:404–408
- Wilhelm J, Fuksova H, Schwippelova Z, Vytasek R, Pichova A (2006) The effects of reactive oxygen and nitrogen species during yeast replicative ageing. *Biofactors* 27:185–193
- Yamada A, Yamamoto T, Yoshimura Y, Gouda S, Kawashima S, Yamazaki N, Yamashita K, Kataoka M, Nagata T, Terada H et al (2009) Ca²⁺ induced permeability transition can be observed even in yeast mitochondria under optimized experimental conditions. *Biochim Biophys Acta* 1787:1486–1491
- Zitomer RS, Nichols DL (1978) Kinetics of glucose repression of yeast cytochrome c. *J Bacteriol* 135:39–44
- Zundorf G, Kahlert S, Bunik VI, Reiser G (2009) alpha-Ketoglutarate dehydrogenase contributes to production of reactive oxygen species in glutamate-stimulated hippocampal neurons in situ. *Neuroscience* 158:610–616