

## Phytochemistry, GC-MS Analysis, Antioxidant and Antimicrobial Potential of Essential Oil From Five *Citrus* Species

Shabnam Javed<sup>1</sup>, Ayesha Javaid<sup>2</sup>, Shaista Nawaz<sup>3</sup>, M. K. Saeed<sup>3</sup>, Zaid Mahmood<sup>2</sup>, S. Z. Siddiqui<sup>4</sup> & Rauf Ahmad<sup>5</sup>

<sup>1</sup> Centre for Undergraduate Studies, University of the Punjab, Quaid-e-Azam Campus, Lahore-54590, Pakistan

<sup>2</sup> Institute of Chemistry, University of the Punjab, Lahore, Pakistan

<sup>3</sup> Food and Biotechnology Research center, Pakistan Council of Scientific and Industrial Research Laboratories Complex, Ferozepur Road, Lahore, Pakistan

<sup>4</sup> Department of Chemistry, Government College University, Lahore, Pakistan

<sup>5</sup> Centre for environment protection studies, Pakistan Council of Scientific and Industrial Research Laboratories Complex, Ferozepur Road, Lahore, Pakistan

Correspondance: Shabnam Javed, Centre for Undergraduate Studies, University of the Punjab, Quaid-e-Azam Campus, Lahore-54590, Pakistan. Tel: 92-042-9923-2057. E-mail: shabnamjaved@yahoo.com

Received: November 13, 2013 Accepted: December 2, 2013 Online Published: February 15, 2014

doi:10.5539/jas.v6n3p201

URL: <http://dx.doi.org/10.5539/jas.v6n3p201>

### Abstract

*Citrus* essential oils were extraction from hydro distillation technique yielding *Citrus* oil with reasonable yield. Phytochemical screening of all five *Citrus* oils showed that alkaloids, tannins, sterols, terpenoids, saponins, flavonoids were present (50-80%). GC/MS analysis showed highest percentage of limonene (58-89%) in *Citrus* oils. Antioxidant study revealed that *Citrus* peel oils have strong scavenging activity (83%-91%). Antimicrobial activity was evaluated by agar well method against eight common pathogens depicted marked antimicrobial potential especially tangerine (4.9-1.9 cm inhibition zones) and grapefruit oil (4.5-1.2 cm) inhibition zones. The studies emphasized the therapeutic and commercial utilization of *Citrus* peel essential oils as food preservatives, phytomedicine and antioxidant agent.

**Keywords:** Citrus essential oils (EO), GC-MS analysis, phytochemicals, DPPH assay, antimicrobial activity

### 1. Introduction

All over the world *Citrus* is one of the widespread genus due to its prominent production. *Citrus* essential oils are naturally occurring, volatile and odoriferous oils synthesized by non woody parts of aromatic plants such as seeds, buds, leaves, flowers, stems, fruits, twigs and roots etc. and accumulated in secretory or epidermis cells and also sometimes in cavities (Ahmad, 2006). Essential oil from *Citrus* fruit peel is the fundamental product of genus *Citrus* and typically isolated by distillation or solvent extraction (Mondello et al., 2005). These are the complex mixtures of about 400 compounds of which 1-15% are non-volatile whereas 85-99% is the volatile constituents (Nannapaneni et al., 2009). Other organic compounds present in *Citrus* essential oils are aliphatic hydrocarbons, alcohols (linalool), aldehydes (citral), acids, esters and some aromatic compounds (Sharma & Tripathi, 2006). Svoboda & Greenaway (2003) reported the chief chemical constituent of *Citrus* essential oils is limonene and have a range of 32 to 98%. *Citrus* essential oils act as natural antioxidants because flavanone glycosides namely naringin, narirutin, hesperidin and neohesperidin are valuable phenolic compounds found in *Citrus* peel oil which make them liable to avert rancidity of food (Anagnostopoulou et al., 2006).

Essential oils of *Citrus* peels are medicinally very important and show variety of biological effects because they are rich in flavonoids (flavone, flavonol and flavanone), terpenes, carotenes and coumarines which are responsible for antimicrobial activity (Tepe et al., 2005). Consequently *Citrus* essential oils are extensively used in pharmaceuticals as an antimicrobial, anti-diabetic, antioxidant, insect repellent, carminative, larvicidal, antiviral, antihepatotoxic and antimutagenic agent (Kanaze et al., 2008).

The rapidly growing importance of *Citrus* based essential oils in food, pharmaceuticals, perfumes, flavor and fragrance has forced Pakistan to import increasing amounts of *Citrus* oils despite its rich variegated acreage of *Citrus* fruits and one of the largest *Citrus* fruits producing country of the world. These factors provide the

opportunity for the production of highest grade essential oils from *Citrus* fruit peel. So there was an urgent need to focus on the extraction of essential oils as solid waste management and to improve our economy. This study was aimed for the assessment of phytochemical constituents, antioxidant as well as antimicrobial activities of five *Citrus* species, *Citrus sinensis* (L.) var. Malta, *C. sinensis* (L.) var. Mousami, *C. reticulata* (L.) var. Tangerine, *C. reticulata* (L.) var. Mandarin and *C. paradisi* (L.) Grapefruit.

## 2. Materials and Methods

### 2.1 Collection and Identification of Citrus Fruit Peels

Peels of five varieties of *Citrus* fruits were collected from local *Citrus* juice shop near University of the Punjab, Quaid-e-Azam campus, Lahore, Pakistan during the month of January & February 2012. Voucher specimen number PU.HHC.901, PU.HHC.902, PU.HHC.903, PU.HHC.904 and PU.HHC.905 were assigned to Malta, Mousami, Tangerine, Mandarin and Grapefruit respectively.

### 2.2 Extraction of Citrus Essential Oil

Essential oils of selected *Citrus* species was extracted by hydro distillation unit for 3-4 hours extraction. Mixture of *Citrus* oils and water was incorporated which was separated, in two liquid layers which was isolated. Hydro-distilled pure oil obtained was stored in dark brown sealed vials at 4°C until analysis.

### 2.3 Physicochemical and Phytochemical Investigation of Citrus EO

Physicochemical characteristics of *Citrus* essential oils including refractive index, optical rotation, specific gravity, color, odor and solubility were analyzed by the method of AOAC (2005). The chemical tests were carried out for screening of bioactive compounds present in *Citrus* essential oils using standard methods (Sofowora, 1993; Trease & Evans, 1989).

### 2.4 GC-MS Analysis of Citrus EO

*Citrus* essential oils were analyzed for their chemical composition by GC-MS analysis. GC/MS JOEL model JMS-A × 5050 H mass spectrometer (JOEL, Japan) Hewlett Packard 5890 Gas Chromatograph (JOEL, Japan). Helium as carrier gas, split ratio 1:100, electrical energy 70 eV, ionization current 200 μA, ionization temperature 250°C, column temperature with 6°C/min rise to 230°C. The chemical constituents were identified by their retention time and compared with known spectrum deposited in the National Institute Standard and Technology (NIST) library (NIST147.LIB).

### 2.5 DPPH Assay

Antioxidant potential was assessed by evaluating scavenging effect of each of five varieties of *Citrus* peel oils on DPPH. 500 μL of each essential oil was added in 3 ml of 0.002% methanolic solution of DPPH and shaken well. Absorbance was noted at 517 nm for all sample solutions and blank (contain only DPPH) after a stay time of 30 min in dark (Amin et al., 2006). All measurements were performed in triplicates. Scavenging potential of *Citrus* peel oils was determined in terms of percentage inhibition (I %) of DPPH by given formula:

$$\text{Percent inhibition} = (A_{\text{blank}} - A_{\text{sample}} / A_{\text{blank}}) \times 100$$

Where  $A_{\text{blank}}$  represents absorbance of DPPH only at 517 nm and  $A_{\text{sample}}$  represents absorbance of sample under investigation at 517 nm.

### 2.6 Antimicrobial Activity of Citrus Oils

#### 2.6.1 Test Organisms

Antimicrobial activity of *Citrus* peel essential oils were studied against the two Gram positive bacteria *Listeria monocytogenes* and *Corynebacterium minutissimum* and three Gram negative bacteria *Escherichia coli*, *Yersinia* sp. and *Klebsiella planticola* whereas three fungal strains named as *Aspergillus flavus*, *A. fumigates* and *A. niger* were used. All microorganisms were obtained from First Fungal Culture Bank of Pakistan (FCBP), Institute of Agricultural Sciences, University of the Punjab, Lahore.

#### 2.6.2 Agar Well Diffusion Method

Antimicrobial potential of *Citrus* peel essential oils was assessed using agar well method (Kim et al., 1995). Each microbial concentration was made 10<sup>6</sup> CFU/ml. Wells (8 mm) were prepared in plates (single well in case of fungi whereas four wells in case of bacteria). About 60 μL of the essential oil was dripped into the wells. Water was used as control. The inoculated plates were incubated for 24 hours at 37°C for bacterial isolates and for 72 h at 27°C for fungi isolates. Study was conducted in triplicates. Biostatic efficacy against test organisms was investigated by measuring the inhibition zones in comparison to a control.

### 3. Results and Discussion

#### 3.1. Yield of Citrus EO by Hydro Distillation

For *Citrus* oil extraction by the hydro distillation, five batches of *Citrus* species were carried out to determine average productivity of *Citrus* oils (Table 1). *Citrus* oils yield were in the range of (0.28-0.45%) for 3-4 hours extraction, comparable and better in some varieties as compared to other extraction techniques reported in literature (Minh Tu et al., 2002; Lota et al., 2000). The highest yield among all *Citrus* essential oil was calculated for *C. paradise* Grapefruit 0.45% followed by *C. sinensis* var. Malta 0.37%, *C. reticulata* var. Mandarin 0.33%, *C. sinensis* var. Mousami 0.30%, *C. reticulata* var. Tangerine 0.28% (Table 1). According to the previous work of Minh Tu et al. (2002) the yield of orange essential oil was 0.13% and tangerine essential oil was 0.25%. The essential oil yielded from various cultivars of mandarin was reported, 0.1% to 0.45% (Lota et al., 2000). Varying yields of essential oil are due to different extraction methods, units, soil and climatic conditions (Huet, 1991).

Table 1. Percentage yield of *Citrus* essential oils by fabricated unit

<i>Citrus</i> species Essential oils (EO)	Raw Material Input (g)	Time extract (minutes)	Oil volume (ml)	Productivity (ml/2000g) %
<i>C. paradisi</i>	2000	215	9.0	0.45
<i>C. sinensis</i> var. Malta	2000	230	7.5	0.37
<i>C. reticulata</i> var. Mandarin	2000	210	6.6	0.33
<i>C. sinensis</i> var. Mousami	2000	235	6.0	0.30
<i>C. reticulata</i> var. Tangerine	2000	220	5.6	0.28

#### 3.2 Physicochemical Characterization

Specific gravity of *Citrus* essential oils in present work was ranged from 0.842-0.858, refractive indices between 1.465-1.476 and all essential oils were found optically active (Table 2). These results were in line with preceding work on essential oils of *Citrus* species (Guenther, 1948).

Table 2. Physicochemical properties of *C. reticulata* var. Mandarin, *C. sinensis* var. Mousami, *C. paradisi*, *C. sinensis* var. Malta, *C. reticulata* var. Tangerine

Physical Parameters	<i>Citrus</i> oils				
	Mandarin	Mousami	Grapefruit	Malta	Tangerine
Color	Yellow	Light yellow	Light yellow	Pale yellow	Light yellow
Odour	Pleasant, Intense	Pleasant, less Intense	Pleasant, less Intense	Pleasant, Intense	Pleasant, less Intense
Refractive index (25°C)	1.465	1.471	1.476	1.471	1.468
Optical rotation (25°C)	+86	+91	+93	+89	+88
Specific gravity (25°C)	0.844	0.849	0.858	0.847	0.842
Solubility					
Water	Insoluble	Insoluble	Insoluble	Insoluble	Insoluble
Ether	Soluble	Soluble	Soluble	Soluble	Soluble
Acetone	Fairly Soluble	Fairly Soluble	Fairly Soluble	Fairly Soluble	Fairly Soluble

### 3.3 Phytochemical Investigation

Phytochemical screening of all tested *Citrus* peel oils gave presence of alkaloids, tannins, terpenoids, saponins and combined anthraquinones in significant amounts (25-80%). Flavonoids and sterols were highly present in five *Citrus* essential oils (>80%). Anthraquinones were moderately present (50-80%) in essential oils of *C. sinensis* var. Mousami and Malta whereas highly depicted in *C. paradisi*, *C. reticulata* var. Tangerine and var. Mandarin (>80%). In all *Citrus* essential oils, alkaloids were in the range of 50-80%. Only *C. paradisi* peel oil depicted the presence of coumarin whereas all *Citrus* peel oils gave negative test for phlobatanins (Table 3). Mondello et al. (2005) explored flavonoids, terpenes, coumarins and carotenes as the major phytochemicals of *Citrus* essential oils. Likewise, Okwu et al. (2007) screened phytochemicals of five *Citrus* species and revealed the presence of saponins, tannins, flavonoids, alkaloids and phenols.

Table 3. Phytochemical Constituents of *C. reticulata* var. Mandarin, *C. sinensis* var. Mousami, *C. paradisi*, *C. sinensis* var. Malta, *C. reticulata* var. Tangerine

Phytochemicals	Citrus oils				
	Mandarin	Mosammi	Grapefruit	Malta	Tangerine
Alkaloids	++	++	++	++	++
Tannins	+	+	++	+	+++
Sterols	+++	+++	+++	+++	+++
Terpenoids	++	+	++	++	+++
Saponins	++	+	+	++	++
Flavonoids	+++	++	+++	+++	+++
Coumarins	-	-	+	-	-
Anthraquinones	+++	++	+++	++	+++
Combined anthraquinones	++	+	++	+	++
Phlobatanins	-	-	++	-	-

(-)=absent, (+) = present, > 50 %, compared with control, (++) = 50% < 80%, (+++) =>80%.

### 3.4 Chemical Composition by GC-MS Analysis

Limonene was identified as the key element of *Citrus* peel oils. GC-MS analysis of *Citrus* peel oils revealed that among five *Citrus* essential oils, Grapefruit essential oil displayed highest concentration of limonene (89.84%) followed by essential oils of Malta (88.57%), Mousami (87.84%), Mandarin (87.45%) and Tangerine (58.50%). Other chemical constituents identified were limonene oxide,  $\alpha$ -terpineol, carvone, carveol, eugenol, spathulenol and caryophyllene oxide.  $\alpha$ -Terpineol (12.55%) was the second major component of all five *Citrus* essential oils with Mandarin peel oil containing highest of it (Table 4). These results were in line with the former work on Brazilian tangerines in which limonene was the major component (Feger et al., 2003). Similarly, Espina et al., (2011) reported limonene (85.50%) and  $\alpha$ -terpineol (0.36%) as chief component of *Citrus* essential oils. In present study, apart from limonene (88.57%) and  $\alpha$ -terpineol (8.45%), Malta peel oil also contained eugenol (0.58%), spathulenol (0.55%), caryophyllene oxide (0.99%), n-hexadecanoic acid (0.86%). Tangerine peel oil showed presence of carvone (16.97%), carveol (8.77%), 3-cyclohexene-1-methanol (8.91%) and limonene oxide (6.85%) in addition to limonene (58.50%). In Mousami peel oil,  $\alpha$ -terpineol (12.16%) was major component after limonene (Table 4). Most of the compounds identified in *Citrus* peel oils were found to be hydrocarbons in nature (Ayoola et al., 2008). The major constituents of *Citrus* peel oils investigated by Vekiari et al., (2002) were limonene, neral, geranial,  $\beta$ -pinene,  $\beta$ -caryophyllene and neryl acetate. Preceding work stated limonene (86.27%),  $\gamma$ -terpinene (2.11%) and  $\alpha$ -pinene (1.26%) in Grapefruit peel essential oil and limonene (76.28%),  $\beta$ -pinene (5.45%), linalool (2.32%), citral (1.74%) and  $\alpha$ -pinene (1.26%) in Mousami peel oil (Ahmad et al., 2006). The different chemical constituents of different *Citrus* species are assumed may be due to different genetic characteristics.

Table 4. Chemical constituents of *Citrus* oils (GC-MS Analysis)

<i>Citrus</i> Essential oils (EO)	Retention time	Compounds	% Area
Mousami EO	7.215	Limonene	87.84
	8.867	$\alpha$ -terpineol	12.16
	7.210	Limonene	88.57
Malta EO	8.895	$\alpha$ -terpineol	8.45
	10.693	Eugenol	0.58
	11.417	Spathulenol	0.55
	11.486	Caryophyllene oxide	0.99
	13.935	n-Hexadecanoic acid	0.86
Grapefruit EO	7.215	Limonene	89.84
	8.871	$\alpha$ -terpineol	10.16
Mandarin EO	7.216	Limonene	87.45
	8.893	$\alpha$ -terpineol	12.55
	7.218	Limonene	58.50
	8.345	Limonene oxide	6.85
Tangerine EO	9.096	Carveol	8.77
	9.332	Carvone	16.97
	10.152	3-Cyclohexene-1-methanol	8.91

### 3.5 Antioxidant Activity

Antioxidant study revealed that all the *Citrus* peel oils have strong potential to reduce DPPH radical to DPPH-H (83%-91%). Highest antioxidant activity was shown by *C. reticulata* var. Mandarin (91.1%) followed by *C. reticulata* var. Tangerine (88.0%), *C. paradisi* (87.2%), *C. sinensis* var. Malta (86.0%) and *C. sinensis* var. Mousami 83.2% for 500  $\mu$ l/ml oil (Figure 1). Antioxidant efficacy of *Citrus* peel oils in decreasing order was as follows: Mandarin > Grapefruit > Tangerine > Malta > Mousami. These results were in accordance with work of Kamal et al. (2013) that *C. reticulata* var. Mandarin showed maximum antioxidant potential whereas *C. sinensis* var. Mousami showed minimum antioxidant potential. Correspondingly, Yang et al. (2010) reported limonene is a major constituent of *Citrus* peel oils having antioxidant potential equivalent to that of strong antioxidants.





Figure 1. Antioxidant activity of *Citrus* oils by DPPH assay using BHT as standard. Data expressed as  $\pm$  standard error bars

(A)



(B)




Figure 2. Antimicrobial activity of *Citrus* oils (A) Antibacterial activity of *Citrus* oils against five bacterial species (B) Antifungal activity of *Citrus* oils against three fungal pathogens. Error bars specify the mean  $\pm$  standard error

### 3.6 Antimicrobial Activity

The antimicrobial activity of *Citrus sinensis*(L.) var. Malta and Mousami, *Citrus reticulata* (L.) var. Tangerine and Mandarin and *Citrus paradise* (L.) grapefruit was examined on five bacterial strains *L. monocytogenes*, *C. minutissimum*, *E. coli*, *Y. sp.* and *K. planticola* (Figure 2A). The essential oils had various degrees of inhibition potential against the five bacterial strains. Tangerine oil depicted maximum bacteriostatic activity against all the test bacteria i.e., *E. coli*, *Y. sp.*, *K. planticola*, *L. monocytogenes* with inhibition zones (4.9 cm, 4.8 cm and 4.6 cm and 1.9 cm, respectively) except *C. minutissimum* var. Mandarin essential oil revealed second highest inhibitory potential against *E. coli* and *C. minutissimum* (4.7 cm and 2.8 cm) while Mousami oil against both *L. monocytogenes* (1.3 cm) and *Y. sp.* (4.7 cm). Grapefruit essential oil displayed marked bactericidal activity against *K. planticola* depicting the inhibition zone of 4.4cm. Least antibacterial effect was shown by Mousami in case of *C. minutissimum*, *E. coli* and *K. planticola* (1.6 cm, 3.7 cm and 3.2 cm, respectively), whereas Grapefruit and Mandarin against *Y. sp.* (4.5 cm each). The growth of *L. monocytogenes* was least affected by Malta oil (1.1 cm). *C. minutissimum* and *L. monocytogenes* were least susceptible to growth inhibition by *Citrus* essential oils. The antimicrobial activities of *Citrus* species are strongly related with chemical constituents like flavonoids and phenols (Viuda-Martos et al., 2008). The active ingredients responsible for antimicrobial potential of *Citrus* peel oils are monoterpene components (Pavithra et al., 2009). D-limonene, linalool or citral are major attributors for the antimicrobial capacity of *Citrus* peel oils. Previous work revealed that the inhibitory influence of *Citrus* peel essential oils is owed to the presence of linalool rather than limonene (Fisher & Phillips, 2006). However it has

been found that antimicrobial activity is not only produced by one particular major component but also due to the antagonistic and synergistic effects of variety of compounds (Deba et al., 2008).

The essential oils of Grapefruit, Mousami, Malta, Mandarin and Tangerine showed the tendency to impede the growth of molds *A. flavus*, *A. fumigates* and *A. niger* (Figure 2B). In all molds, Tangerine and Grapefruit oils revealed great antifungal potentials. In case of *A. flavus* and *A. fumigatus*, Tangerine peel oil was the best growth inhibitor (3.4 cm-4.0 cm). Followed by Grapefruit and Malta peel oils, which showed almost equal reduction in growths for these two molds. Mandarin and Mousami showed the lowest growth reductions of *A. flavus* and *A. fumigatus*. The mycelium growth of *A. niger* was most susceptible to grapefruit essential oil with inhibition zone (2.8 cm) while least affected by mousami peel oil with inhibition zone (0.7 cm). Viuda-Martos et al. (2008) studied inhibitory influence of mandarin, orange and grapefruit peel essential oils on four fungal pathogens. *A. flavus* growth was best prevented by Mandarin essential oil whereas *A. niger* growth was most susceptible to Orange peel oil. Grapefruit essential oil was the most efficient against *P. verrucosum* and *P. chrysogenum*.

#### 4. Conclusion

The results of present investigation are the basis for extraction of essential oil of *Citrus* by cheaper methods to design pilot plant to extract EO for industrial production. Some major restraints in viable industrial exploitation of medicinal plants are due to the poor agricultural practices, quality control trials, strain in marketing and dearth of research on process and product development. Coordination among various institutes and organizations of the country can lead for sustainable commercial utilization.

#### References

- A. O. A. C. (2005). *Official methods of Analysis* (15th ed.). Association of Official Analytical Chemists, Washington DC.
- Ahmad, M. M., Rehman, S., Iqbal, Z., Anjum, F. M., & Sultan, J. I. (2006). Genetic variability to essential oil composition in four *Citrus* fruit species. *Pakistan Journal of Botany*, *38*, 319-324.
- Amin, I., Norazaidah, Y., & Emmy Hainida, K. I. (2006). Antioxidant activity and phenolic content of raw and blanched *Amaranthus* species. *Food Chemistry*, *94*, 47-52. <http://dx.doi.org/10.1016/j.foodchem.2004.10.048>
- Anagnostopoulou, M. A., Kefalas, P., Papageorgiou, V. P., Assimopoulou, A. N., & Boskou, D. (2006). Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*). *Food Chemistry*, *94*, 19-25. <http://dx.doi.org/10.1016/j.foodchem.2004.09.047>
- Ayoola, G. A., Johnson, O. O., Adelowotan, T., Aibinu, I. E., Adenipekun, E., Adepoju-Bello, A. A., ... Odugbemi, T. O. (2008). Evaluation of the chemical constituents and the antimicrobial activity of the volatile oil of *Citrus reticulata* fruit (Tangerine fruit peel) from South West Nigeria. *African Journal of Biotechnology*, *7*(13), 2227-2231.
- Deba, F., Xuan, T. D., Yasuda, M., & Tawata, S. (2008). Chemical composition and antioxidant, antibacterial and antifungal activities of the essential oils from *Bidens pilosa* Linn. Var. *Radiata*. *Food Control*, *19*, 346-352. <http://dx.doi.org/10.1016/j.lwt.2007.04.011>
- Espina, L., Somolinos, M., Lorán, S., Conchello, P., García, D., & Pagán, R. (2011). Chemical composition of commercial *Citrus* fruit essential oils and evaluation of their antimicrobial activity acting alone or in combined processes. *Food Control*, *22*, 896-902. <http://dx.doi.org/10.1016/j.foodcont.2010.11.021>
- Feger, W., Brandauer, H., & Ziegler, H. (2003). Analytical investigation of Murcott (Honey) tangerine peels oil. *Journal of Essential Oil Research*, *15*(3), 143-147. <http://dx.doi.org/10.1080/10412905.2003.9712097>
- Fisher, K., & Phillips, C. A. (2006). The effect of lemon, orange and bergamot essential oils and their components on the survival of *Campylobacter jejuni*, *Escherichia coli* O157, *Listeria monocytogenes*, *Bacillus cereus* and *Staphylococcus aureus* *in vitro* and in food systems. *Journal of Applied Microbiology*, *101*, 1232-1240. <http://dx.doi.org/10.1111/j.1365-2672.2006.03035.x>
- Guenther, E. (1948). *The Essential Oils. The examination and analysis of essential oils, synthetics, and isolates* (pp. 229-367). New York: Van D. Nostrand Company Inc.
- Huet, R. (1991). Les huiles essentielles d'agrumes. *Fruits*, *46*(5), 551-576.
- Kamal, G. M., Ashraf, M. Y., Hussain, A. I., Shahzadi, A., & Chughtai, M. I. (2013). Antioxidant potential of peel essential oils of three Pakistani *Citrus* species: *Citrus reticulata*, *Citrus sinensis* and *Citrus paradisi*. *Pakistan Journal of Botany*, *45*(4), 1449-1454

- Kanaze, F. I., Termentzi, A., Gabrieli, C., Niopas, I., Georgarakis, M., & Kokkalou, E. (2008). The phytochemical analysis and antioxidant activity assessment of orange peel (*Citrus sinensis*) cultivated in Greece-Crete indicates a new commercial source of hesperidin. *Biomedical Chromatography*, *23*, 239-249. <http://dx.doi.org/10.1002/bmc.1090>
- Kim, J., Marshall, M. R., & Wei, C. I. (1995). Antibacterial activity of some essential oil components against five food borne pathogens. *Journal of Agriculture and Food Chemistry*, *43*, 2839-2845. <http://dx.doi.org/10.1021/jf00059a013>
- Lota, M. L., De Rocca Serra, D., Tomi, F., & Casanova, J. (2000). Chemical variability of peel and leaf essential oils of mandarins from *Citrus reticulata* Blanco. *Biochemical Systematics and Ecology*, *28*, 61-78. [http://dx.doi.org/10.1016/S0305-1978\(99\)00036-8](http://dx.doi.org/10.1016/S0305-1978(99)00036-8)
- Minh Tu, N. T., Thanh, L. X., Une, A., Ukeda, H., & Sawamura, M. (2002). Volatile constituents of Vietnamese pummelo, orange, tangerine and lime peel oils. *Flavour and Fragrance Journal*, *17*, 169-174. <http://dx.doi.org/10.1002/ffj.1076>
- Mondello, L., Casilli, A., Tranchida, P. Q., & Dugo, P. (2005). Comprehensive two-dimensional GC for the analysis of *Citrus* essential oils. *Flavour and Fragrance Journal*, *20*, 136-140. <http://dx.doi.org/10.1002/ffj.1506>
- Nannapaneni, R., Chalova, V. I., Crandall, P. G., Ricke, S. C., Johnson, M. G., & O'Bryan, C. A. (2009). *Campylobacter* and *Arcobacter* species sensitivity to commercial orange oil fractions. *International Journal of Food Microbiology*, *129*(1), 43-49. <http://dx.doi.org/10.1016/j.ijfoodmicro.2008.11.008>
- Okwu, D. E., Awurum, A. N., & Okoronkwo, J. I. (2007). Phytochemical composition and *in vitro* antifungal activity screening of extracts from *Citrus* plants against *Fusarium Oxysporum* of Okra plant (*Hibiscus esculentus*). *African Crop Science Conference Proceedings*, *8*, 1755-1758.
- Pavithra, P. S., Sreevidya, N., & Verma, R. S. (2009). Antibacterial activity and chemical composition of essential oil of *Pamburusmissionis*. *Journal of Ethnopharmacology*, *124*, 151-153. <http://dx.doi.org/10.1016/j.jep.2009.04.016>
- Sharma, N., & Tripathi, A. (2006). Effects of *Citrus sinensis* epicarp essential oil on growth and morphogenesis of *Aspergillus niger* Van Tieghem. *Microbiological Research*, *10*, 1016-1020.
- Sofowora, A. E. (1993). *Medicinal Plants and Traditional Medicine in Africa* (p. 288). Ibadan, Nigeria: Spectrum Books Ltd.
- Svoboda, K. P., & Greenaway, R. I. (2003). Lemon scented plants. *International Journal of Aromatherapy*, *13*(1), 23-32. [http://dx.doi.org/10.1016/S0962-4562\(03\)00048-1](http://dx.doi.org/10.1016/S0962-4562(03)00048-1)
- Tepe, B., Daferera, D., Sokmen, A., Sokmen, M., & Polissiou, M. (2005). Antimicrobial and antioxidant activities of the essential oils and various extracts of *Salvia tomentosa* Miller (Lamiaceae). *Food Chemistry*, *90*, 333-340. <http://dx.doi.org/10.1016/j.foodchem.2003.09.013>
- Trease, G. E., & Evans, W. C. (1989). *Pharmacognosy* (2nd ed.). Braille Tiridel and Macmillan Publishers.
- Vekiari, S. A., Protopapadakis, E. E., Parthena, P., Dimitrios, P., Panou, C., & Vamvakias, M. (2002). Composition and seasonal variation of the essential oil from leaves and peel of a Cretan lemon variety. *Journal of Agricultural Food Chemistry*, *50*(1), 147-153. <http://dx.doi.org/10.1021/jf001369a>
- Viuda-Martos, M., Ruiz-Navajas, Y., Fernandez-Lo'pez, J., & Perez-Alvarez, J. (2008). Antifungal activity of lemon (*Citrus lemon* L.), mandarin (*Citrus reticulata* L.), grapefruit (*Citrus paradisi* L.) and orange (*Citrus sinensis* L.) essential oils. *Food Control*, *19*, 1130-1138. <http://dx.doi.org/10.1016/j.foodcont.2007.12.003>
- Yang, S. A., Jeon, S. K., Lee, E. J, Shim, C. H., & Lee, I. S. (2010). Comparative study of the chemical composition and antioxidant activity of six essential oils and their components. *Natural Product Research*, *24*, 140-151. <http://dx.doi.org/10.1080/14786410802496598>

## Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).