

ARTICLE

Received 18 Feb 2014 | Accepted 8 Jul 2014 | Published 14 Aug 2014

DOI: 10.1038/ncomms5636

Phytochrome-interacting transcription factors PIF4 and PIF5 induce leaf senescence in *Arabidopsis*

Yasuhito Sakuraba^{1,*}, Jinkil Jeong^{2,*}, Min-Young Kang¹, Junghyun Kim², Nam-Chon Paek¹ & Giltso Choi²

Plants initiate senescence to shed photosynthetically inefficient leaves. Light deprivation induces leaf senescence, which involves massive transcriptional reprogramming to dismantle cellular components and remobilize nutrients. In darkness, intermittent pulses of red light can inhibit senescence, likely via phytochromes. However, the precise molecular mechanisms transducing the signals from light perception to the inhibition of senescence remain elusive. Here, we show that in *Arabidopsis*, dark-induced senescence requires phytochrome-interacting transcription factors PIF4 and PIF5 (PIF4/PIF5). ELF3 and phytochrome B inhibit senescence by repressing PIF4/PIF5 at the transcriptional and post-translational levels, respectively. PIF4/PIF5 act in the signalling pathways of two senescence-promoting hormones, ethylene and abscisic acid, by directly activating expression of *EIN3*, *ABI5* and *EEL*. In turn, PIF4, PIF5, *EIN3*, *ABI5* and *EEL* directly activate the expression of the major senescence-promoting NAC transcription factor *ORESARA1*, thus forming multiple, coherent feed-forward loops. Our results reveal how classical light signalling connects to senescence in *Arabidopsis*.

¹Department of Plant Science, Plant Genomics and Breeding Institute, Research Institute of Agriculture and Life Sciences, Seoul National University, Seoul 151-921, Korea. ²Department of Biological Sciences, KAIST, Daejeon 305-701, Korea. * These authors contributed equally to this work. Correspondence and requests for materials should be addressed to N.-C.P. (email: ncpaek@snu.ac.kr) or to G.C. (email: gchoi@kaist.edu).

Leaf senescence, the final stage of leaf development, actively destabilizes intracellular organelles, including decomposition of macromolecules, to relocate nutrients into developing tissues or storage organs. Senescence occurs autonomously in an age-dependent manner, controlled by an innate genetic programme. However, unfavorable environmental stresses, such as darkness, drought, high temperature, high salinity and pathogen attacks, can also trigger senescence during leaf development¹. For example, light deprivation induces senescence in lower leaves that are shaded by upper leaves².

Molecular genetic studies using *Arabidopsis thaliana* have identified many senescence-associated genes (SAGs), including genes involved in hormone signalling, transcriptional regulation and chlorophyll (Chl) catabolism^{1,3}. The developmental and environmental signals of senescence commonly activate senescence-associated transcriptional factors (senTFs), which modulate the expression of SAGs⁴. To date, transcriptome analyses and molecular genetic studies, in *Arabidopsis* and other plants, have identified many senTFs^{3,5}. In particular, the senTFs include a few plant-specific NAC genes, such as *ORESAR1* (*ORE1*, also known as *NAC092*)⁵, indicating that the senescence-induced NAC transcription factors have important roles in modulating senescence. *ORE1* promotes senescence by regulating the expression of hundreds of SAGs that function in the breakdown of nucleic acids and proteins, and the transport of sugar^{6,7}. Treatment with ethylene or abscisic acid (ABA) upregulates *ORE1* expression⁸, suggesting a close relationship between hormone signalling and *ORE1*. Among the senescence-induced NACs, *NAC-LIKE*, *ACTIVATED BY AP3/PI* (*NAP/NAC029*) and *NAC016* also promote senescence, as demonstrated by the delayed senescence phenotype of *nap* and *nac016* mutants^{9,10}. In contrast, *JUNGBRUNNEN 1* (*JUB1/NAC042*) and *VND-INTERACTING 2* (*VNI2/NAC083*) inhibit leaf senescence^{11,12}. Thus, these observations show that the senescence-promoting and senescence-inhibiting NACs finely tune the expression of SAGs in response to developmental and environmental cues.

Plant hormones coordinate senescence in connection with senTFs, and ensure that senescence proceeds in an orderly fashion at the organismal level. Among plant hormones, ethylene promotes senescence, as demonstrated by accelerated senescence in ethylene-treated plants¹³ and delayed senescence in mutants impaired in ethylene signalling, including mutants in the ethylene signalling component *ETHYLENE INSENSITIVE 2* (*EIN2*), the transcription factor *EIN3* and the ethylene receptor *ETHYLENE INSENSITIVE 1* (*ETR1*) (refs 14–16). ABA also induces senescence rapidly¹⁷. Mutants of *RECEPTOR-LIKE PROTEIN KINASE 1* (*RPK1*) and *SENESCENCE-ASSOCIATED GENE 113* (*SAG113*), which are insensitive to ABA, display delayed senescence phenotypes^{18,19}. Senescence is rapidly induced by treatment with salicylic or jasmonic acids^{20,21}, but is extremely delayed by cytokinin treatment²². The exact molecular mechanisms of how these hormone signals coordinate the ageing or stress signals for leaf senescence are not fully understood.

Plant growth and development requires light²³, and plants require photoreceptors to adapt to ambient light conditions throughout development. Red light has long been considered to be a key negative signal for plant senescence, because pulses of red light substantially delay dark-induced senescence (DIS) in a few plant species, supposedly via phytochromes²⁴ and a low ratio of red/far-red light causes senescence in tobacco leaves²⁵. Red light activates and far-red light inactivates phytochromes; darkness also causes slow inactivation of phytochromes²³. Photoactivated phytochromes move from the cytosol to the nucleus, where they interact with and inactivate the negative regulators of light signalling, resulting in massive transcriptome

changes²³. In *Arabidopsis*, *phyA* and *phyB* regulate a wide range of light responses such as seed germination, seedling photomorphogenesis, hypocotyl gravitropism, shade avoidance and flowering time²³. Phytochromes inactivate phytochrome-interacting bHLH transcription factors (PIFs), which inhibit light responses in the dark²⁶. At the molecular level, active phytochromes directly interact with PIFs to induce their detachment from DNA²⁷, and promote their degradation by the 26S proteasome^{28,29}. PIFs are also regulated at the transcriptional level; the EARLY FLOWERING 3 (ELF3)-ELF4-LUX ARRHYTHMO (LUX) complex directly associates with the promoters of *PIF4* and *PIF5* (*PIF4/PIF5*) and suppresses their expression in the regulation of circadian responses³⁰.

Although the regulatory function of the phytochrome-PIF signalling module has been intensively investigated in the light-responsive development of *Arabidopsis* seedlings, it is unknown if this signalling module has a role in plant senescence. Here, we show the molecular mechanism of how the phytochrome-PIF signalling module activates leaf senescence in *Arabidopsis*. We found that among the four PIFs, *PIF4/PIF5* function as the essential, central transcriptional activators of DIS and that ELF3 and red light-activated *phyB* repress the senescence-promoting activity of *PIF4/PIF5* at the transcriptional and post-translational levels, respectively. The two PIFs activate the expression of *ORE1* by forming multiple coherent feed-forward loops, together with two distinct classes of transcription factors, EIN3 in ethylene signalling and Group A bZIPs (ABA INSENSITIVE 5 (ABI5) and ENHANCED EM LEVEL (EEL)) in ABA signalling.

Results

The Pfr form of *phyB* inhibits leaf senescence. Phytochromes are red/far-red photoreceptors that can be interconverted to active Pfr and inactive Pr forms by red and far-red light irradiation, respectively³¹. To examine the effect of phytochrome-mediated light signalling on leaf senescence, 7-day-old *Arabidopsis* wild-type (WT) seedlings were incubated for 10-day in darkness (control), and in darkness with intermittent pulses of red light (*phyB_{on}*) or with pulses of red light followed by far-red light (*phyB_{off}*) (Fig. 1a, left panel). WT cotyledons turned yellow during dark incubation, a typical symptom of senescence in plants. When the light pulses were given, we found that the cotyledons stayed green in *phyB_{on}* conditions but turned yellow in *phyB_{off}* conditions (Fig. 1a, right panel), indicating that the Pfr form of *phyB* inhibits senescence. To investigate the role of *phyB* in senescence, we examined the phenotypes of 7-day-old seedlings of *phyB* mutants and *PHYB*-overexpressing (*PHYB-OX*) plants after 8 days of dark incubation (DDI). *phyB* mutants senesced faster and *PHYB-OX* plants senesced slower than WT (Fig. 1b,e,f). Similar to the seedling phenotypes, when attached or detached leaves of 3-week-old plants were incubated in darkness or individual leaves were covered with aluminium foil under long day conditions, *phyB* mutants senesced faster and *PHYB-OX* plants senesced slower than WT (Fig. 1c,d; Supplementary Fig. 1a and Supplementary Methods). Consistent with their visible phenotypes, *phyB* mutants had lower Chl levels, higher ion leakage rates, higher expression of two senescence marker genes (*SENESCENCE 4* (*SEN4*) and *SAG12*), and lower expression of a photosynthetic gene (*LIGHT HARVESTING CHLOROPHYLL A/B BINDING PROTEIN 1* (*Lhcb1*)) than WT seedlings or individually covered leaves of adult plants during dark incubation, while *PHYB-OX* had the opposite phenotypes, with the exception of ion leakage rate (Fig. 1e–g, Supplementary Fig. 1b–d). It appears that *PHYB* overexpression is not sufficient to retard the breakdown of membrane integrity of seedlings during DIS. In contrast, *phyA* mutants and *PHYA-OX* plants




Figure 1 | PhyB suppresses senescence. (a) Suppression of senescence by treatment with intermittent red light pulses during dark incubation. Light-grown 7-day-old WT (Col-0 ecotype) seedlings were transferred to darkness for 10 days without any light (continuous darkness), with intermittent red light pulses (*phyB_{on}*) or with red light pulses immediately followed by far-red light pulses (*phyB_{off}*). R and FR indicate red light and far-red light pulses, respectively. (b–f) Suppression of senescence by *phyB* as judged by changes in leaf colour. Light-grown 7-day-old WT (Col-0), *phyB* (*phyB-9*) and *PHYB-OX* seedlings (b), and attached (c) or detached leaves (d) of 3-week-old WT, *phyB* and *PHYB-OX* plants were transferred to darkness for the indicated number of days. Total Chl levels (e) and ion leakage rates (f) in (b–d) were determined before and after dark treatment. (g) Expression levels of *SEN4*, *SAG12* and *Lhcb1* of WT, *phyB* and *PHYB-OX* in 7-day-old seedlings or 3-week-old plants during dark incubation. For quantitative real-time PCR, the relative expression levels were determined by normalizing to the transcript levels of *GAPDH*. *PHYB-OX* indicates the plants (Col) overexpressing *PHYB* under the control of 35S promoter. Mutant information is listed in Supplementary Table 3. Data are means \pm s.d. of more than four independent biological replicates. * $P < 0.05$, ** $P < 0.01$ (Student's *t*-test). DAG, day(s) after germination.

senesced at the same rate as WT (Supplementary Fig. 2). The results indicate that *phyB* is the major photoreceptor signalling to suppress senescence in *Arabidopsis*.

Mutations in *PIF4* and *PIF5* delay leaf senescence. We next investigated which *phyB* signalling components regulate DIS in *Arabidopsis* leaves. Previous work showed that *pif* quadruple (*pifQ*) mutants (*pif1 pif3 pif4 pif5*) show a constitutive photomorphogenic phenotype when grown in darkness, and *phyB* promotes light responses by repressing the activity of PIFs^{32,33}. We found that *pifQ* mutants maintained green cotyledons at 10 DDI, while WT cotyledons turned completely yellow (Fig. 2a), indicating that PIFs promote senescence in light-deprived conditions. To determine which PIFs promote senescence, we examined the seedling phenotypes of four *pif* single mutants during dark incubation. The *pif1* and *pif3* mutants turned yellow

at the same rate as WT, but *pif4* and *pif5* mutants, similar to *pifQ* mutants, senesced slower than WT (Fig. 2b). We also observed a similar delayed senescence pattern in the attached, detached or individually covered leaves of *pif4*, *pif5* and *pifQ* mutants (Fig. 2c,d; Supplementary Fig. 3a). Consistent with their visible phenotypes, the mutants had higher Chl levels, lower ion leakage rates, lower expression of *SEN4* and higher expression of *Lhcb1* than WT (Supplementary Figs 3b,c, 4 and 5). In addition, *pifQ* mutants maintained better-preserved chloroplast structure and *pif4* and *pif5* mutants retained higher levels of photosystem proteins (Supplementary Fig. 6 and Supplementary Methods). Moreover, *PIF4/PIF5* also promote age-dependent leaf senescence, as *pif4* and *pif5* mutant leaves senesced slower than WT leaves under long days (Supplementary Fig. 7), suggesting that *PIF4/PIF5* are the essential transcription factors that promote leaf senescence under not only dark-induced, but also under natural senescence conditions.




Figure 2 | PIF4 and PIF5 (PIF4/PIF5) promote senescence. (a-d) Delayed senescence of *pif4* and *pif5* mutants in darkness. WT and *pifQ* indicate wild-type (Col-0) and *pif1 pif3 pif4 pif5* quadruple mutants, respectively. For the assay, 7-day-old seedlings were grown in continuous light (a,b), and attached (c) or detached (4th or 5th) leaves (d) of 3-week-old plants grown under long days were incubated in darkness and photographed. Mutant information is listed in Supplementary Table 3. (e) *PIL1* expression in WT (black line), *phyB* mutants (*phyB-9*; red line), and *pifQ* mutants (blue line) during dark incubation. Means \pm s.d. were calculated from at least four independent replicates. (f) *PIF4/PIF5* expression in WT (black lines) and *phyB* mutants (red lines) during dark incubation. (e,f) By reverse transcription and quantitative real-time PCR, relative transcript levels of *PIL1* (e) and *PIF4/PIF5* (f) were normalized to those of *GAPDH* (a control). (g) Stabilization of PIF4/PIF5 proteins in *phyB_{off}* conditions. The 35S-driven PIF4/PIF5-Myc and tubulin (TUB; a control) were detected by anti-Myc and anti-tubulin antibodies, respectively. PIF4/PIF5 levels were normalized to TUB protein levels.

To gauge if PIF4/PIF5 transcription factors become active during dark incubation, we examined the expression of the PIF target gene *PHYTOCHROME-INTERACTING FACTOR 3-LIKE 1 (PIL1)* during dark incubation, as PIFs directly regulate *PIL1* expression³⁴. *PIL1* mRNA levels increased and plateaued at 2 DDI in WT (Fig. 2e). However, *PIL1* mRNA levels did not rise in *pifQ* mutants; in contrast, *phyB* mutants showed much higher *PIL1* mRNA levels than WT. The altered expression of *PIL1* in *pifQ* and *phyB* mutants provides indirect evidence that phyB negatively regulates the activity of PIF4/PIF5 in the light; the increased activity of PIF4/PIF5 in the prolonged darkness could be caused by the increase of *PIF4/PIF5* transcription, PIF4/PIF5 protein stability or both. To investigate this, we first examined the expression levels of *PIF4/PIF5* during dark incubation and found that their expression in *phyB* mutants was almost the same as in WT (Fig. 2f). This suggests that phyB does not regulate PIF4/PIF5 at the transcriptional level, but may possibly act at the post-translational level. Indeed, we found that in *PIF4-OX* and *PIF5-*

OX plants, the levels of PIF4/PIF5 proteins increased in darkness (Fig. 2g). Intermittent pulses of red light (*PhyB_{on}*) inhibited the increase of PIF4/PIF5 proteins, but pulses of red light followed by far-red light (*PhyB_{off}*) did not, indicating that the PIF4/PIF5 proteins are stabilized when phyB converts to the inactive Pr form.

ELF3 inhibits leaf senescence by repressing PIF4/PIF5. A previous study showed that an ELF3-ELF4-LUX complex represses the expression of *PIF4/PIF5* in the evening phase³⁰. Thus, we examined if ELF3 inhibits leaf senescence by repressing *PIF4/PIF5* transcription during dark incubation. We found that *elf3* mutants, similar to *PIF4-OX* and *PIF5-OX* plants, senesced much faster than WT in both detached and attached leaves of adult plants (Fig. 3a; Supplementary Fig. 8a), while *ELF3-OX* plants, similar to *pif4* and *pif5* mutants, exhibited delayed senescence (Fig. 3b), as shown by Chl levels (Fig. 3c), ion




Figure 3 | ELF3 inhibits senescence. (a–d) Inhibition of senescence by ELF3 in darkness. Three-week-old detached leaves (upper) or whole plants (lower) grown in long days were transferred to complete darkness and photographed at 4 DDI (a) or at 7 DDI (b). Total Chl levels (c) and ion leakage rates (d) were measured at indicated DDIs. *elf3* is *elf3-8* (Supplementary Table 3). *PIF4-OX*, *PIF5-OX* and *ELF3-OX* indicate the plants overexpressing *PIF4*, *PIF5* and *ELF3* under the control of 35S promoter, respectively. (e) Independent regulation of senescence by ELF3 and phyB. The 7-day-old seedlings grown in continuous light conditions were transferred to darkness for 16 days with intermittent red (PhyB_{on}) or red/far-red light pulses (PhyB_{off}) as shown in Fig. 1a. phyB_{off}/phyB_{on} indicates that total Chl levels of WT, *phyB* and *elf3* mutants in PhyB_{off} conditions are normalized to those in PhyB_{on} conditions. (f) Repression of *PIF4*/*PIF5* transcription by ELF3. The 7-day-old WT (black bars), *elf3* mutants (red bars) and *ELF3-OX* plants (blue bars) were transferred to darkness for 2 days. *PIF4*/*PIF5* expression was normalized to *GAPDH* expression. (g,h) *PIF4*/*PIF5* are epistatic to *ELF3*. Three-week-old plants grown in long days were transferred to darkness for 7 days (g) and total Chl levels were measured (h). DDI indicates day(s) of dark incubation. Data are means ± s.d. of at least four biological replicates. **P* < 0.05, ***P* < 0.01 (Student’s *t*-test).

leakage rates (Fig. 3d), the expression levels of *SEN4* and *Lhcb1* (Supplementary Fig. 5) and chloroplast structure and photosystem protein levels (Supplementary Fig. 6). *ELF3* functions as one of the critical factors in light input into the circadian

clock and in the regulation of flowering time^{35,36}; therefore, we further examined whether senescence is also affected by mutations of other regulators of the circadian clock, flowering time and photomorphogenesis, such as *CIRCADIAN-CLOCK*

ASSOCIATED 1 (*CCA1*), LATE ELONGATED HYPOCOTYL (*LHY*), *ELF4*, *LUX*, FLAVIN-BINDING, KELCH REPEAT, F-BOX 1 (*FKF1*), CRYPTOCHROME 2 (*CRY2*), *GIGANTEA* (*GI*), *CONSTANS* (*CO*), CONSTITUTIVE PHOTOMORPHOGENIC 1 (*COP1*) and *DE-ETIOLATED 1* (*DET1*) (refs 36,37). We found that none of the mutants for these regulators showed altered senescence phenotypes during dark incubation (Supplementary Fig. 8b). Although *CRY1* was suggested to regulate PIF4/PIF5 for shade avoidance responses³⁸, we found that detached leaves of *cry1*, *cry2* and *cry1 cry2* double mutants did not show an early or delayed senescence phenotype during dark incubation (Supplementary Fig. 8c). These results indicate that among the components we tested, *ELF3* functions as a unique negative regulator of senescence induction.

ELF3 acts as a phyB signalling component in seedling development³⁵; we thus tested whether *ELF3* inhibits senescence through red light-activated phyB signalling. Although both *phyB* and *elf3* mutants senesced faster than WT, *elf3* mutants still responded to red light pulses comparable to WT, whereas *phyB* mutants did not (Fig. 3e), indicating that *ELF3* suppresses DIS independent of phyB. Since the expression of *PIF4/PIF5* increased during dark incubation independent of phyB activity (Fig. 2f), we measured the mRNA levels of *PIF4/PIF5* in *elf3* mutants and *ELF3-OX* plants at 2 DDI (Fig. 3f). We found that the *PIF4* expression was higher in *elf3* mutants than in WT, and lower in *ELF3-OX* plants. *PIF5* expression also increased more in *elf3* mutants. Furthermore, the *pif4 elf3* and *pif5 elf3* double mutants senesced much later than WT, but similar to *pif4* and *pif5* single mutants at 7 DDI (Fig. 3g,h), demonstrating that *pif4* and *pif5* are epistatic to *elf3*. Consistent with this, *ELF3-OX PIF4-OX* plants senesced faster than WT, but similar to *PIF4-OX* plants (Supplementary Fig. 9). These results indicate that *ELF3* represses *PIF4/PIF5* transcription independent of phyB, while the Pfr form of phyB destabilizes *PIF4/PIF5* proteins.

PIF4/PIF5 promote ethylene and ABA signalling. To investigate how *PIF4/PIF5* promote DIS at the molecular level, we performed microarray analysis and identified 677 differentially expressed genes (DEGs) between WT and *pifQ* mutants at 2 DDI (WT/*pifQ*, twofold, false discovery rate (FDR) < 0.05; Fig. 4a, left circle). We compared the DEGs of WT/*pifQ* with those of senescing WT at 2 DDI (DIS (2 days/0 days)) (Fig. 2a, right circle) from previously reported microarray data³⁹. We found that the two sets share 132 DEGs ($P < 10^{-72}$, hypergeometric test), showing a high correlation coefficient ($R = 0.85$; Supplementary Fig. 10a). The common DEGs have enriched functional categories of ageing, leaf senescence, photosynthesis, Chl metabolism and light response (Supplementary Fig. 10b). Moreover, although the changes were less than twofold, Gene Set Enrichment Analysis (GSEA, see Methods) showed that the non-shared DEGs of senescing WT also significantly correlated with the changes in *pifQ* mutants (Supplementary Fig. 10c). The results indicate that *PIF4/PIF5* function as the major regulators of SAG expression in senescing leaves during dark incubation.

To further explore how *PIF4/PIF5* regulate the expression of SAGs, we tested whether the 132 shared DEGs include direct targets of *PIF4/PIF5* transcription factors. Thirty two of the shared DEGs are PIF-direct targets ($P = 0.0062$, hypergeometric test), according to previously reported *PIF4/PIF5* chromatin immunoprecipitation (ChIP)-seq data^{34,40}. Their promoters are enriched with the G-box element (CACGTG), a typical binding motif for PIF transcription factors^{34,40–43} (Fig. 4b, upper panel). In contrast, the promoters of the indirectly regulated DEGs are enriched with ABRE-like elements (YACGT), the known binding

sites of bZIP transcription factors, such as the positive ABA signalling component *ABI5*⁴⁴ (Fig. 4b, lower panel). These results suggest that *PIF4/PIF5* directly activate the expression of many SAGs and also indirectly function through ABA signalling.

Ethylene signalling also promotes senescence^{15–16}; we thus examined whether *PIF4/PIF5* activate both ABA and ethylene signalling for inducing senescence. The GSEA indicated that both ABA- and ethylene-induced genes are downregulated in *pifQ* mutants at 2 DDI and are upregulated in senescing WT (Supplementary Table 1). The suppression of ABA and ethylene signalling in *pifQ* mutants might be partly due to the decreased expression of *EIN3* (a positive ethylene signalling component) and the Group A bZIP transcription factors *ABI5* and *EEL* (Fig. 4c,d,h–j), which are direct targets of *PIF4/PIF5*, as shown by ChIP assays (Fig. 4d). Consistent with the decreased expression of *ABI5*, *EEL* and *EIN3* in *pifQ* mutants, the bZIP- and *EIN3*-induced genes, which were identified from previously reported microarray data (see Methods), were significantly downregulated in *pifQ* mutants and upregulated in senescing WT (Supplementary Table 1). We found that *abi5*, *eel* and *ein3* mutations delayed senescence, and *abi5* and *eel* mutations showed additive effects (Fig. 4e–g), further supporting the hypothesis that decreased expression of *ABI5*, *EEL* and *EIN3* is closely associated with delayed senescence in *pifQ* mutants. Since *ELF3* delays senescence by repressing the transcription of *PIF4/PIF5* (Fig. 3f), *ELF3* should repress the expression of *ABI5*, *EEL* and *EIN3* during dark incubation. Indeed, their mRNA expression was severely downregulated in *ELF3-OX* plants and upregulated in *elf3* mutants (Fig. 4h–j). These results indicate that *PIF4/PIF5* regulate senescence partly through the two bZIPs and *EIN3*.

PIF4/PIF5 activate *ORE1* through multiple feed-forward loops.

We next investigated if *PIF4*, *PIF5* and their direct target transcription factors (*ABI5*, *EEL* and *EIN3*) jointly affect the expression of known senescence-promoting regulators. To date, 3,744 senescence-associated loci in *Arabidopsis* have been curated³, and 89 loci are known to promote senescence. Among the 89 senescence-promoting genes³ (Supplementary Data 1), we found that *ORE1*, encoding a key senescence-promoting NAC transcription factor, was commonly induced by all three groups of transcription factors (Fig. 5a). *ORE1* expression was downregulated in *ELF3-OX* plants and *pifQ*, *abi5*, *eel* and *ein3* mutants, and upregulated in *elf3* mutants and *PIF4-OX* plants (Fig. 5b), indicating that *ORE1* is a target senTF acting downstream of *ELF3*, *PIF4*, *PIF5*, *ABI5*, *EEL* and *EIN3*. Consistent with this, *ore1* mutants, similar to *phyB* and *pif4* mutants, did not respond to red light pulse treatment during dark incubation (Fig. 5c,d). These results indicate that *ORE1* integrates phyB-mediated light signalling to promote senescence in light-deprived conditions.

We found that the *ORE1* promoter harbours three potential PIF binding sites (G-box motifs), several bZIP binding sites (ABRE motifs) and an *EIN3* binding site (Fig. 5e), suggesting that *ORE1* is a direct target of not only *PIF4/PIF5* but also of bZIPs and *EIN3*. To examine whether *PIF4*, *PIF5*, *ABI5*, *EEL* and *EIN3* directly activate *ORE1* transcription, we next performed ChIP assays and found that all of the tested TFs bind to the *ORE1* promoter (Fig. 5f). Collectively, the direct activation of *ORE1* transcription by *PIF4/PIF5* and by *PIF4/PIF5*-targeted transcription factors (*ABI5*, *EEL* and *EIN3*) indicates that these transcription factors establish multiple coherent feed-forward regulatory loops to induce leaf senescence; *PIF4/PIF5* directly activate the expression of *ABI5*, *EEL* and *EIN3*, and in turn, *PIF4*, *PIF5*, *ABI5*, *EEL* and *EIN3* directly activate *ORE1* transcription.




Figure 4 | PIF4/PIF5 directly induce bZIPs and EIN3. (a) Significant overlap between DEGs (twofold, FDR < 0.05) of PIF-induced genes (WT/*pifQ*, this study) and DIS genes (previous microarray data³⁹). Light-grown 7-day-old WT and *pifQ* seedlings were transferred to darkness for 2 days (WT versus *pifQ*), and 4-week-old adult WT plants were transferred to darkness for 0 and 2 days (2 DDI versus 0 DDI). The number in parentheses indicates the PIF4/PIF5-direct target genes retrieved from the previous ChIP-seq database^{34,40}. (b) Enrichment of the G-box motifs in the PIF-direct target promoters and the ABRE-like element in the PIF-indirect target promoters, identified by the MEME software using 500 bp promoter sequences. (c) Heat map showing the expression of *EIN3*, *EIL1* and group A bZIPs in the microarray analysis from this study (WT/*pifQ*) and two independent sets of DIS microarray data from previous studies^{39,69}. Black circles indicate the PIF4/PIF5-direct target genes according to previous ChIP-seq analyses^{34,40}. The scale bar (bottom) indicates fold changes as log₂ values. The *ABF2* probe is absent in the DIS microarray (Affymetrix). (d) Direct binding of PIF4/PIF5 to the promoters of *ABI5*, *EEL* and *EIN3* by ChIP assays. Values were normalized to the values for *PP2A* as a non-binding control. Data are means ± s.d. of two biological replicates. (e–g) Delayed senescence of *abi5*, *eel*, *abi5 eel*, *ein3* and *pifQ* mutants. Light-grown 7-day-old seedlings were transferred to darkness for 10 days (e). Total Chl levels were measured at 10 DDI (f). Data are means ± s.d. of at least four biological replicates. Detached leaves of 3-week-old plants were transferred to darkness and photographed at 7 DDI (g). (h–j) Regulation of *ABI5*, *EEL* and *EIN3* expression by ELF3 and PIF4/5. The mRNA levels were determined by quantitative real-time PCR using light-grown 7-day-old seedlings transferred to darkness for 0, 2 and 4 DDI. *GAPDH* was used as a control. DDI indicates day(s) of dark incubation. Data are means ± s.d. of four biological replicates. **P* < 0.05, ***P* < 0.01 (Student's *t*-test).




Figure 5 | PIF4/PIF5 induce *ORE1* through multiple feed-forward loops. (a) Identification of *ORE1* as a common target of PIFs, bZIPs and EIN3. Among the literature-curated senescence-promoting genes³, DEGs (FDR < 0.05) were identified from the microarray data for PIF-induced genes (this study), ABF2/3/4-induced genes⁶⁷ and EIN3-induced genes (GSE21762). (b) Regulation of *ORE1* expression by *ELF3*, *PIF4/PIF5*, *ABI5*, *EEL* and *EIN3*. *ORE1* mRNA levels were determined by quantitative real-time PCR at 0 (black bars), 2 (grey bars) and 4 (white bars) DDIs. *GAPDH* was used as a control. Data are means \pm s.d. of at least four independent biological replicates. (c,d) Senescence induction in *phyB_{off}* conditions requires *ORE1*. Light-grown 7-day-old seedlings were transferred to darkness for 16 days with intermittent red light pulses (*phyB_{on}*) or red/far-red light pulses (*phyB_{off}*) as shown in Fig. 1a and total Chl levels (d) were measured to determine the degree of senescence. Data are means \pm s.d. of four biological replicates. (e) A diagram of the *ORE1* promoter showing the positions of G-box elements (red boxes), ABRE-like elements (green boxes), EIN3 binding sites (EBS; blue box) and two ChIP amplicons (*ORE1-1* and *ORE1-2*; black horizontal lines). (f) Direct binding of *PIF4*, *PIF5*, *ABI5*, *EEL* and *EIN3* to the *ORE1* promoter *in vivo*. Enrichment of *ORE1* promoter by *PIF4* (black), *PIF5* (red), *ABI5* (blue), *EEL* (green) and *EIN3* (yellow) were examined by ChIP-qPCR. Values were normalized to the values for *PP2A*, a non-binding control. Data are means \pm s.d. of two biological replicates. * $P < 0.05$, ** $P < 0.01$ (Student's *t*-test).

ABI5 and EEL directly promote Chl degradation. *PIF4*, *PIF5* and their direct target transcription factors (*EIN3* and bZIPs) may have other senescence-related targets in addition to *ORE1*. Microarray data analysis indicated that among the SAGs, *PIF4*, *PIF5* and bZIPs induce four Chl catabolism-associated genes, including *STAYGREEN 1* (*SGR1*, also known as *NYE1*) and *NON-YELLOW COLORING 1* (*NYC1*) (refs 45,46) (Supplementary Table 2). *SGR1* and *NYC1* expression was downregulated in *ELF3-OX* plants and *pifQ* mutants, and upregulated in *elf3* mutants and *PIF4-OX* plants (Fig. 6a). We also found the bZIP binding motif (ABRE motif) in the promoters of *SGR1* and *NYC1* (Fig. 6b). Consistent with the presence of ABREs, ChIP assays revealed that *ABI5* and *EEL* directly bind to the promoters of *SGR1* and *NYC1* (Fig. 6c) and the expression levels of *SGR1* and *NYC1* were significantly downregulated in *abi5 eel* double mutants during DIS (Fig. 6d). Although the promoters of *SGR1* and *NYC1* also contain a putative PIF binding motif (G-box motif; Supplementary Fig. 11a), *PIF4/PIF5* did not bind to the G-box motifs of *SGR1* and *NYC1* (Supplementary Fig. 11b). The results indicate that *PIF4/PIF5* indirectly promote Chl degradation through their target bZIPs during leaf senescence.

Discussion

Light deprivation initiates leaf senescence in plants. In this study, we reveal the molecular genetic pathway by which light deprivation accelerates senescence, via increased levels of *PIF4/PIF5* mRNAs and *PIF4/PIF5* proteins, which are negatively regulated by *ELF3* and *phyB*, respectively (Fig. 7). The increased activity of *PIF4/PIF5* promotes the expression of *ORE1* not only by directly binding to its promoter but also indirectly through *EIN3* and bZIPs (*ABI5* and *EEL*). These TFs, in turn, activate the expression of hundreds of SAGs that mediate the highly ordered decomposition of cellular components, such as Chls and photosynthetic proteins in chloroplasts, finally leading to leaf senescence^{1,3,4}.

We used plate-grown young seedlings rather than adult plants for the majority of the senescence assays, because it is easier to synchronize the growth of seedlings and to reduce unknown variation caused by environmental fluctuations. However, as we have shown, *phyB* and *elf3* mutants senesce early, while *pif4*, *pif5* and *pifQ* mutants senesce later, not only in seedlings but also in detached, attached or individually covered leaves of adult plants, suggesting that the senescence pathways identified in seedlings are likely important for senescence at other stages of




Figure 6 | bZIPs directly induce *SGR1* and *NYC1* for Chl breakdown. (a) Gene expression analysis of *SGR1* and *NYC1* in light-grown 7-day-old WT (Col), *elf3* (*elf3-8*), *ELF3-OX*, *pifQ* and *PIF4-OX* plants at 0, 2 and 4 DDI. (b) The promoter regions of *SGR1* and *NYC1* showing the position of ABRE motifs (green boxes) and ChIP amplicons (black thick underlines). (c) ChIP assays showing direct binding of ABI5 and EEL to the promoters of *SGR1* and *NYC1*. The *PP2A* locus was used as a non-binding control. Data are means \pm s.d. of two independent biological replicates. (d) Decreased expression of *SGR1* and *NYC1* in *abis5 eel* double mutants. Gene expression was measured by quantitative real-time PCR using light-grown 7-day-old seedlings transferred to darkness for 0 and 4 days. The relative mRNA levels of *SGR1* and *NYC1* were normalized to *GAPDH* expression as a control. Data are means \pm s.d. of four independent biological replicates. ** $P < 0.01$ (Student's *t*-test).

development. Overall, our data show that PIF4/PIF5 promote senescence throughout development, while phyB and ELF3 repress PIF4/PIF5 to prevent precocious senescence in the presence of light. Our study revealed the role of PIF4/PIF5 in senescence in relation to light signalling, and other works showed that PIF4 also mediates high temperature signalling to promote hypocotyl growth⁴⁷. Thus, additional work is required to investigate whether other signals from abiotic stresses, such as high temperature, also regulate senescence through PIF4/PIF5.

The identified pathway links light signalling to senescence and consists of multiple coherent feed-forward loops, which likely function to make the pathway robust. In conditions with sufficient light, ELF3 and phyB convey light information to PIF4/PIF5, resulting in the reduction of PIF4/PIF5 mRNA and protein levels (Figs 2g and 3f). Under prolonged darkness, however, the process reverses and PIF4/PIF5 mRNA and protein levels increase. The increased PIF4/PIF5 activate the expression of *EIN3* and two bZIPs (*ABI5* and *EEL*) by directly associating with their promoters (Fig. 4c,d,h-j), thus affecting regulators of ethylene and ABA signalling, respectively. In turn, the increased *EIN3* and bZIPs directly activate the expression of *ORE1* (Fig. 5f), a major senescence-promoting NAC transcription factor that regulates hundreds of SAGs. Notably, not only *EIN3* and bZIPs but also PIF4/PIF5 directly bind to the *ORE1* promoter to activate its expression (Fig. 5f), indicating that PIF4/PIF5 form coherent feed-forward loops to activate the expression of *ORE1* (Fig. 7). Such coherent feed-forward loops make a pathway less prone to being disturbed by ephemeral environmental fluctuations⁴⁸.

Similar coherent feed-forward loops between *EIN2* and *ORE1* were previously proposed⁵; *EIN2* activates the expression of *ORE1* both by repressing the expression of *miR164*, which cleaves the *ORE1* mRNA, and by a *miR164*-independent pathway. *EIN2* also inhibits the expression of *miR164* through *EIN3*, which directly binds to the *miR164* promoter¹⁷. Since we showed that *EIN3* directly binds to the *ORE1* promoter to activate its expression (Fig. 5b,f), these results suggest that a previously proposed *miR164*-independent pathway likely functions through *EIN3* and thus *EIN3*, *miR164*, and *ORE1* form a coherent feed-forward loop. Such a loop implies that the cascade of coherent feed-forward loops regulates the expression of *ORE1* during senescence. We note that *EIN3* also directly binds to the promoter of *NAP*, a major senescence-promoting NAC transcription factor and one of the 89 senescence-promoting genes³ (Supplementary Data 1), and induces *NAP* transcription⁴⁹, indicating that *EIN3* activates the expression of both *ORE1* and *NAP*.

Our study showed that ELF3 represses senescence independently of its previously reported roles in light signalling. A previous study showed that ELF3 interacts with phyB to regulate the phyB signal transduction pathway³⁵. Although ELF3 represses senescence as phyB does, the *elf3* mutants still respond to red light pulse treatment similar to WT, indicating that ELF3 regulates senescence independent of phyB activity^{35,50}. The fact that ELF3 represses the expression of *PIF4/PIF5* mRNA and phyB promotes the degradation of PIF4/PIF5 proteins further supports the independent roles of ELF3 and phyB in senescence (Figs 2g and 3f). Our study also suggests that ELF3 represses




Figure 7 | A proposed model for PIF4/PIF5-induced leaf senescence.

PIF4/PIF5 promote DIS by forming multiple coherent feed-forward loops with bZIPs (*ABI5* and *EEL*), *EIN3* and *ORE1*, all of which are inhibited by *phyB* and *ELF3*.

senescence independently of its roles in circadian clock, flowering time and diurnal control of hypocotyl elongation. First, *ELF3* is required to maintain the circadian clock under continuous light⁵¹ and its mRNA expression is directly repressed by *CCA1* for the regulation of the diurnal control of hypocotyl length and flowering time⁵². Besides *elf3* mutants, however, the circadian-clock mutants including *cca1*, *lhy1* and *det1* do not exhibit an altered senescence phenotype in darkness (Supplementary Fig. 8b), indicating that the dysfunction of the circadian clock *per se* does not cause the early senescence symptoms of *elf3* mutants. Second, *ELF3* interacts with *COP1*, which allows *COP1* to destabilize *GI* downstream of *CRY2* to control flowering time³⁶. However, *cop1*, *gi*, *fkf1*, *co* and *cry2* mutants do not show any senescence-associated phenotype in darkness, suggesting that the flowering-time defect in *elf3* mutants is irrelevant to the early senescence in darkness. Third, *ELF3* forms an evening complex with *ELF4* and *LUX* and represses the expression of *PIF4/PIF5* mRNAs for diurnal control of hypocotyl elongation³⁰. Unlike the *elf3* mutants, *elf4* and *lux* mutants senesce at the same rate as WT in darkness, indicating that *ELF3* acts as the transcription repressor of *PIF4/PIF5* in the absence of *ELF4* or *LUX*. Taken together, our data suggest that the inhibition of senescence represents a novel function of *ELF3*.

The increase of *PIF4/PIF5* expression independent of *phyB* (Fig. 2f) raises the possibility that other light signalling components also suppress senescence by repressing the expression of *PIF4/PIF5* in connection with *ELF3*. A previous study showed that *ELF3* is destabilized when light-grown seedlings were transferred to darkness³⁵. It is not known, however, which light signalling components regulate *ELF3* stability. A few possibilities exist. First, the inactivation of other phytochromes might be responsible for the destabilization of *ELF3*, which leads to the upregulation of *PIF4/PIF5* expression in the dark, but the role of *phyA* in senescence remains inconclusive. We showed that the *Arabidopsis phyA* mutants do not show any senescence phenotype in our conditions (Supplementary Fig. 2). Unlike our analysis, however, other studies have reported two contradictory roles of *phyA* in senescence: the overexpression of oat *phyA* delayed senescence in tobacco⁵³, whereas the pea *phyA* mutants

exhibited delayed senescence⁵⁴. These contrasting results could be due to the different senescence conditions (dark-induced versus natural senescence) or different plant species (*Arabidopsis* versus tobacco versus pea). Alternatively, *phyA* proteins from different plant species might have different functions. For example, *Arabidopsis phyA* does not mediate the red light-low fluence response (LFR)⁵⁵, whereas monocot *phyA* mediates the red light-LFR even when it is expressed in *Arabidopsis*⁵⁶. Thus, the suppression of senescence by oat *phyA* could be due to its unique ability to mediate the LFR, a function usually reserved for *phyB*⁵⁷. The delayed, rather than early, senescence of pea *phyA* mutants also excludes pea *phyA* as the repressor of *PIF4/PIF5* mRNA expression in the dark. Currently, the role of three minor phytochromes (*phyC*, *phyD* and *phyE*) in senescence remains to be examined. Second, the activation of the expression of *PIF4/PIF5* during senescence might require the absence of blue light, rather than presence of red light. Previous studies showed that blue light pulses inhibit DIS in papaya⁵⁸. In soybean, *CRY2a* interacts with and inhibits a *CRY*-interacting CRYPTOCHROME-INTERACTING BASIC-HELIX-LOOP-HELIX 1 (*CIB1*) transcription factor that promotes leaf senescence by directly upregulating the expression of *WRKY53* (ref. 59). Since low red to far-red ratio also promotes leaf senescence in soybean⁶⁰, both phytochrome and cryptochrome signals can delay leaf senescence in soybean. However, we found that cryptochrome signalling does not regulate senescence in *Arabidopsis* (Supplementary Fig. 8c). Since blue light is perceived not only by *CRY* but also by phototropins and Light-Oxygen-Voltage-sensing-domain containing F-box proteins²³, our data do not exclude the possibility that other blue light receptors repress senescence through *ELF3*. Further investigation is needed to determine which light signals regulate the expression of *PIF4/PIF5* mRNA through *ELF3* during senescence.

The inclusion of *ABI5* and *EIN3* in the *PIF4/PIF5*-initiated feed-forward loops suggests that the identified regulatory loops integrate signals from light, abiotic stress and ageing, to regulate senescence (Fig. 7). Previous studies showed that ABA signalling increases the activity of *ABI5* and its homologues, Group A bZIPs, in response to abiotic stresses, both at the transcriptional and post-transcriptional levels^{44,61} (AtGenExpress; abiotic stress). Activated *ABI5* and its homologues trigger a subset of ABA responses, including senescence (Fig. 4e–g), seedling growth arrest and abiotic stress tolerance^{61,62}. Our data indicate that *PIF4/PIF5* also activate the expression of *ABI5* and its homologues at the transcriptional level, by directly binding to their promoters (Fig. 4c,d,h–j). Light decreases the activity of *PIF4/PIF5*; this finding indicates that light and abiotic stresses antagonistically regulate the expression of *ABI5* and its homologues. This finding also indicates that the integration of light and abiotic stress signalling determines the output of the *PIF4/PIF5*-initiated coherent feed-forward loops. Similarly, the *PIF4/PIF5*-initiated coherent feed-forward loops likely integrate developmental stage signals through *EIN3*, as shown by a gradual increase of *EIN3* expression during ageing, and the consequent promotion of age-dependent senescence¹⁶. In conclusion, identification of *EIN3* and *ABI5* as constituents of the loops in our model suggests that light signalling (represented by *PIF4/PIF5*) interacts with abiotic stress signalling (*ABI5* and *EEL*)⁶² and developmental age signalling (*EIN3*)¹⁶ to regulate senescence (Fig. 7). The interplay among these signals provides the mechanism by which plants decide when to senesce in response to ageing and dynamic environmental conditions.

Methods

Plant materials and growth conditions. *Arabidopsis thaliana* plants were grown at 22 to 24 °C under long days (16-h light/8-h dark) in a growth room equipped

with cool-white fluorescent light (90 to $100 \mu\text{mol m}^{-2} \text{s}^{-1}$) for general growth and seed harvest. For phenotypic analysis, surface-sterilized seeds were plated on Murashige and Skoog (MS) agar plates (half-strength MS, 0.8% phytoagar and 0.05% MES, pH 5.7) and stratified for 3 days at 4°C in darkness before being transferred to the growth room. For DIS, 7-day-old seedlings, detached (4th or 5th) rosette leaves of 3-week-old plants with 3 mM MES buffer (pH 5.8) or whole plants were incubated in complete darkness for the indicated periods. For quantitative reverse transcriptase PCR and dark treatment with or without light pulse irradiance, 7-day-old seedlings grown on MS agar plates grown under continuous light ($100 \mu\text{mol m}^{-2} \text{s}^{-1}$) were incubated in darkness with or without 5 min of red light ($13 \mu\text{mol m}^{-2} \text{s}^{-1}$) followed by darkness or 5 min of far-red light ($3.2 \mu\text{mol m}^{-2} \text{s}^{-1}$) for the indicated periods.

Plasmid construction and plant transformation. For *Arabidopsis* transgenic plants expressing genes under the control of the 35S promoter, the full-length *PHYB*, *PIF4*, *PIF5*, *ABI5*, *EEL* and *EIN3* cDNAs were amplified with specific primers (Supplementary Table 4), cloned into the binary vectors, pBI121 (*PIF4* and *PIF5*) and pCAMBIA1300, bearing either GFP (*PHYB*), Myc (*PIF4* and *PIF5*) or FLAG (*ABI5*, *EEL* and *EIN3*) tag, and transformed into WT (Col-0) by *Agrobacterium*-mediated transformation. 35S:FLAG-*ABI5* and 35S:FLAG-*EEL* are hypersensitive to ABA and 35S:*EIN3*-FLAG is hypersensitive to ethylene (Supplementary Fig. 12).

Measurement of Chl pigments. For measurement of total Chl concentration, Chl pigments were extracted with 80% ice-cold acetone from leaf tissues of plants grown under long days. The Chl concentration was determined using an ultraviolet/visible spectrophotometer according to the previous method⁶⁵.

Reverse transcription and quantitative real-time PCR. Total RNA from plant tissues was isolated using the Total RNA Extraction Kit (iNtRON Biotechnology, Korea). For RT, first-strand cDNAs were prepared with 5 μg of total RNA using M-MLV reverse transcriptase and oligo(dT)₁₅ primer (Promega) and diluted to 100 μl with water. Gene expression levels were determined by qPCR analysis. Twenty microlitres of qPCR mixture contained 2 μl of first-strand cDNAs, 10 μl of 2X QuantiTect SYBR Green I Master (Roche) and 0.25 μM of the forward and reverse primers for each gene. The qPCR analysis was performed using the Light Cycler 2.0 (Roche Diagnostics). Relative expression levels of each gene were normalized to mRNA levels of glyceraldehyde phosphate dehydrogenase (*GADPH*) as a loading control. The gene-specific primers for qPCR are listed in Supplementary Table 4.

Measurement of ion leakage rate. To measure ion leakage rate caused by DIS, approximately 10 whole seedlings or 10 rosette leaves in each condition were placed in a tube with 6 ml of 0.4 M mannitol solution. The tubes were placed at room temperature for 3 h with gentle shaking. Conductivity of incubated solution was measured before and after boiling for 10 min, using an electroconductivity metre (CON6 METRE, LaMOTTE Co., USA). The ion leakage rate was calculated by the ratio of initial to total conductivity.

SDS-PAGE and immunoblot analysis. *PIF4*-Myc and *PIF5*-Myc seedlings were collected and immediately frozen in liquid nitrogen under a dim green light. The seedlings were then ground in liquid nitrogen and homogenized in denaturing buffer (100 mM NaH₂PO₄, 10 mM Tris-HCl, 8 M urea, pH 8.0) by vigorous vortexing. The debris was removed by centrifugation at 20,000 g for 10 min at 4°C . For immunoblot analysis, the supernatants were separated on an 8% SDS-polyacrylamide gel, and the proteins were transferred to a nitrocellulose membrane (Hybond ECL, Amersham) using transfer buffer (5.8 g l⁻¹ Tris base, 29 g l⁻¹ glycine, 20% methanol and 0.01% SDS). For protein detection, rabbit polyclonal anti-Myc antibody (Santa Cruz, USA) with 1/1,000 dilution for *PIF4*-Myc and *PIF5*-Myc and mouse monoclonal anti-tubulin antibody (Sigma, USA) with 1/10,000 dilution for a loading control in PBS buffer containing 0.05% Tween 20 were used. Blots were washed three times with the same buffer and then incubated with appropriate secondary antibodies. After washing three times, the horseradish peroxidase activity of secondary antibodies was detected using an ECL detection kit (AbFRONTIER, Korea). Uncropped blot images of Fig. 2g are shown in Supplementary Fig. 13.

Microarray analysis. For microarray analysis, 7-day-old light-grown seedlings of WT and *pijQ* mutants were incubated in the dark for 2 days before RNA extraction. Total RNA was extracted using the Spectrum Plant Total RNA kit (Sigma-Aldrich, USA). Three biological replicates of WT and *pijQ* mutants were grown and sampled for RNA purification. Two μg of total RNA was used for probe synthesis. The Agilent Arabidopsis Genome 44 K chip (version 4) was used. The analysis of microarray data was done in R version 2.15.0. The limma package was used for background correction and for normalization within arrays and between arrays⁶⁴. Then, a linear model was fitted using *lmFit* followed by statistical calculation using *eBayes*. FDR less than 0.05 was applied to identify DEGs.

Microarray data were deposited in the NCBI Gene Expression Omnibus (GEO) with accession number GSE52646. For the identification of DEGs of WT DIS and hormonal transcription factor-activated genes, raw data deposited in NCBI GEO and EMBL-EBI were obtained and processed as described above.

ChIP assay. The plants overexpressing *PIF4*-Myc, *PIF5*-Myc, *ABI5*-FLAG, *EEL*-FLAG and *EIN3*-FLAG under the cauliflower mosaic virus 35S promoter (see Plasmid construction and plant transformation) were grown in continuous white light for 7 days and then transferred to darkness for 3 days before cross-linking for 20 min with 1% formaldehyde under vacuum. Chromatin complexes were isolated and sonicated with slight modification⁴³. Anti-Myc monoclonal antibody (mouse, Cell Signaling) or anti-FLAG polyclonal antibody (rabbit, Sigma), and Protein A agarose/salmon sperm DNA (Millipore) were used for immunoprecipitation. After reverse cross-linking and protein digestion, DNA was purified by QIAquick PCR Purification Kit (Qiagen).

GSEA. GSEA⁶⁵ was conducted with Software GSEA v2.08 (Broad Institute, MIT) according to the online user guide (<http://www.broadinstitute.org/gsea>). ABA and ACC-responsive gene sets were generated based on previously processed microarray data⁶⁶. Sets of genes activated by *ABI5*, *ABF2/3/4* and *EIN3* were identified from public microarray data from previous studies^{67,68} and from NCBI GEO deposition (GSE21762) with twofold criteria (FDR < 0.05) as described in Microarray Analysis. The set of genes activated by *ABI5* was further selected for genes whose expression is induced by ABA treatment (AtGenExpress). Parameters were set as gene_set for permutation type, 1,500 for maximal size of gene set and default for other parameters.

References

- Lim, P. O., Kim, H. J. & Nam, H. G. Leaf senescence. *Annu. Rev. Plant Biol.* **58**, 115–136 (2007).
- Rousseaux, M. C., Hall, A. J. & Sánchez, R. A. Far-red enrichment and photosynthetically active radiation level influence leaf senescence in field-grown sunflower. *Physiol. Plant.* **96**, 217–224 (1996).
- Liu, X. *et al.* LSD: a leaf senescence database. *Nucleic Acids Res.* **39**, D1103–D1107 (2011).
- Kim, J. H. *et al.* Trifurcate feed-forward regulation of age-dependent cell death involving miR164 in *Arabidopsis*. *Science* **323**, 1053–1057 (2009).
- Balazadeh, S., Riaño-Pachón, D. M. & Mueller-Roeber, B. Transcription factors regulating leaf senescence in *Arabidopsis thaliana*. *Plant Biol.* **10**, 63–75 (2008).
- Balazadeh, S. *et al.* A gene regulatory network controlled by the NAC transcription factor ANAC092/AtNAC2/ORE1 during salt-promoted senescence. *Plant J.* **62**, 250–264 (2010).
- Matallana-Ramirez, L. P. *et al.* NAC transcription factor ORE1 and senescence induced BIFUNCTIONAL NUCLEASE1 (BFN1) constitute a regulatory cascade in *Arabidopsis*. *Mol. Plant* **6**, 14321452 (2013).
- He, X.-J. *et al.* AtNAC2, a transcription factor downstream of ethylene and auxin signaling pathways, is involved in salt stress response and lateral root development. *Plant J.* **44**, 903–916 (2005).
- Guo, Y. & Gan, S. AtNAP, a NAC family transcription factor, has an important role in leaf senescence. *Plant J.* **46**, 601–612 (2006).
- Kim, Y.-S., Sakuraba, Y., Han, S.-H., Yoo, S.-C. & Paek, N.-C. Mutation of the *Arabidopsis* NAC016 transcription factor delays leaf senescence. *Plant Cell Physiol.* **54**, 1660–1672 (2013).
- Yang, S.-D., Seo, P. J., Yoon, H.-K. & Park, C.-M. The *Arabidopsis* NAC transcription factor VN12 integrates abscisic acid signals into leaf senescence via the COR/RD genes. *Plant Cell* **23**, 2155–2168 (2011).
- Wu, A. *et al.* JUNGBRUNNEN1, a reactive oxygen species-responsive NAC transcription factor, regulates longevity in *Arabidopsis*. *Plant Cell* **24**, 482–506 (2012).
- Gepstein, S. & Thimann, K. V. The role of ethylene in the senescence of Oat leaves. *Plant Physiol.* **68**, 349–354 (1981).
- Grbić, V. & Bleeker, A. B. Ethylene regulates the timing of leaf senescence in *Arabidopsis*. *Plant J.* **8**, 595–602 (1995).
- Oh, S. A. *et al.* Identification of three genetic loci controlling leaf senescence in *Arabidopsis thaliana*. *Plant J.* **12**, 527–535 (1997).
- Li, Z., Peng, J., Wen, X. & Guo, H. ETHYLENE-INSENSITIVE3 is a senescence-associated gene that accelerates age-dependent leaf senescence by directly repressing miR164 transcription in *Arabidopsis*. *Plant Cell* **25**, 3311–3328 (2013).
- Nooden, L. D. & Leopold, A. C. *Senescence and Aging in Plants* (Academic Press, 1988).
- Lee, I. C. *et al.* Age-dependent action of an ABA-inducible receptor kinase, RPK1, as a positive regulator of senescence in *Arabidopsis* leaves. *Plant Cell Physiol.* **52**, 651–662 (2011).
- Zhang, K., Xia, X., Zhang, Y. & Gan, S.-S. An ABA-regulated and Golgi-localized protein phosphatase controls water loss during leaf senescence in *Arabidopsis*. *Plant J.* **69**, 667–678 (2012).

20. Ueda, J., Kato, J., Yamane, H. & Takahashi, N. Inhibitory effect of methyl jasmonate and its related compounds on kinetin-induced retardation of oat leaf senescence. *Physiol. Plant.* **52**, 305–309 (1981).
21. Morris, K. *et al.* Salicylic acid has a role in regulating gene expression during leaf senescence. *Plant J.* **23**, 677–685 (2000).
22. Gan, S. & Amasino, R. M. Inhibition of leaf senescence by autoregulated production of cytokinin. *Science* **270**, 1986–1988 (1995).
23. Whitelam, G. C. & Halliday, K. J. *Light and Plant Development* (Blackwell, 2007).
24. Biswal, U. C. & Biswal, B. Photocontrol of leaf senescence. *Photochem. Photobiol.* **39**, 875–879 (1984).
25. Rousseaux, M. C., Ballaré, C. L., Jordan, E. T. & Vierstra, R. D. Directed overexpression of PHYA locally suppresses stem elongation and leaf senescence responses to far-red radiation. *Plant Cell Environ.* **20**, 1551–1558 (1997).
26. Jeong, J. & Choi, G. Phytochrome-interacting factors have both shared and distinct biological roles. *Mol. Cells* **35**, 371–380 (2013).
27. Park, E. *et al.* Phytochrome B inhibits binding of phytochrome-interacting factors to their target promoters. *Plant J.* **72**, 537–546 (2012).
28. Bauer, D. *et al.* Constitutive photomorphogenesis 1 and multiple photoreceptors control degradation of phytochrome interacting factor 3, a transcription factor required for light signaling in *Arabidopsis*. *Plant Cell* **16**, 1433–1445 (2004).
29. Park, E. *et al.* Degradation of phytochrome interacting factor 3 in phytochrome-mediated light signaling. *Plant Cell Physiol.* **45**, 968–975 (2004).
30. Nusinow, D. A. *et al.* The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* **475**, 398–402 (2011).
31. Borthwick, H. A., Hendricks, S. B., Toole, E. H. & Toole, V. K. Action of light on lettuce-seed germination. *Bot. Gaz.* **115**, 205–225 (1954).
32. Leivar, P. *et al.* Multiple phytochrome-interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness. *Curr. Biol.* **18**, 1815–1823 (2008).
33. Shin, J. *et al.* Phytochromes promote seedling light responses by inhibiting four negatively-acting phytochrome-interacting factors. *Proc. Natl Acad. Sci. USA* **106**, 7660–7665 (2009).
34. Hornitschek, P. *et al.* Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling. *Plant J.* **71**, 699–711 (2012).
35. Liu, X. L., Covington, M. F., Fankhauser, C., Chory, J. & Wagner, D. R. ELF3 encodes a circadian clock-regulated nuclear protein that functions in an *Arabidopsis* PHYB signal transduction pathway. *Plant Cell* **13**, 1293–1304 (2001).
36. Yu, J.-W. *et al.* COP1 and ELF3 control circadian function and photoperiodic flowering by regulating GI stability. *Mol. Cell* **32**, 617–630 (2008).
37. Imaizumi, T. *Arabidopsis* circadian clock and photoperiodism: time to think about location. *Curr. Opin. Plant Biol.* **13**, 83–89 (2010).
38. Keller, M. M. *et al.* Cryptochrome 1 and phytochrome B control shade-avoidance responses in *Arabidopsis* via partially-independent hormonal cascades. *Plant J.* **67**, 195–207 (2011).
39. van der Graaff, E. *et al.* Transcription analysis of *Arabidopsis* membrane transporters and hormone pathways during developmental and induced leaf senescence. *Plant Physiol.* **141**, 776–792 (2006).
40. Oh, E., Zhu, J. Y. & Wang, Z. Y. Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nat. Cell Biol.* **14**, 802–809 (2012).
41. Martinez-Garcia, J. F., Huq, E. & Quail, P. H. Direct targeting of light signals to a promoter element-bound transcription factor. *Science* **288**, 859–863 (2000).
42. Zhang, Y. *et al.* A quartet of PIF bHLH factors provides a transcriptionally centered signaling hub that regulates seedling morphogenesis through differential expression-patterning of shared target genes in *Arabidopsis*. *PLoS Genet.* **9**, e1003244 (2013).
43. Oh, E. *et al.* Genome-wide analysis of genes targeted by PHYTOCHROME INTERACTING FACTOR 3-LIKE5 during seed germination in *Arabidopsis*. *Plant Cell* **21**, 403–419 (2009).
44. Choi, H., Hong, J., Ha, J., Kang, J. & Kim, S. Y. ABFs, a family of ABA-responsive element binding factors. *J. Biol. Chem.* **275**, 1723–1730 (2000).
45. Park, S.-Y. *et al.* The senescence-induced staygreen protein regulates chlorophyll degradation. *Plant Cell* **19**, 1649–1664 (2007).
46. Kusaba, M. *et al.* Rice NON-YELLOW COLORING1 is involved in light-harvesting complex II and grana degradation during leaf senescence. *Plant Cell* **19**, 1362–1375 (2007).
47. Koini, M. A. *et al.* High temperature-mediated adaptations in plant architecture require the bHLH transcription factor PIF4. *Curr. Biol.* **19**, 408–413 (2009).
48. Le, D.-H. & Kwon, Y.-K. A coherent feedforward loop design principle to sustain robustness of biological networks. *Bioinformatics* **29**, 630–637 (2013).
49. Kim, H. J. *et al.* Gene regulatory cascade of senescence-associated NAC transcription factors activated by ETHYLENE-INSENSITIVE2-mediated leaf senescence signalling in *Arabidopsis*. *J. Exp. Bot.* **65**, 4023–4036 (2014).
50. Reed, J. W. *et al.* Independent action of ELF3 and phyB to control hypocotyl elongation and flowering time. *Plant Physiol.* **122**, 1149–1160 (2000).
51. Hicks, K. A. *et al.* Conditional circadian dysfunction of the *Arabidopsis* early-flowering 3 mutant. *Science* **274**, 790–792 (1996).
52. Lu, S. X. *et al.* CCA1 and ELF3 interact in the control of hypocotyl length and flowering time in *Arabidopsis*. *Plant Physiol.* **158**, 1079–1088 (2012).
53. Cherry, J. R., Hershey, H. P. & Vierstra, R. D. Characterization of tobacco expressing functional Oat phytochrome: domains responsible for the rapid degradation of Pfr are conserved between monocots and dicots. *Plant Physiol.* **96**, 775–785 (1991).
54. Weller, J. L., Murfet, I. C. & Reid, J. B. Pea mutants with reduced sensitivity to far-red light define an important role for phytochrome A in day-length detection. *Plant Physiol.* **114**, 1225–1236 (1997).
55. Whitelam, G. C. *et al.* Phytochrome A null mutants of *Arabidopsis* display a wild-type phenotype in white light. *Plant Cell* **5**, 757–768 (1993).
56. Boylan, M. T. & Quail, P. H. Phytochrome A overexpression inhibits hypocotyl elongation in transgenic *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **88**, 10806–10810 (1991).
57. Reed, J. W., Nagatani, A., Elich, T. D., Fagan, M. & Chory, J. Phytochrome A and phytochrome B have overlapping but distinct functions in *Arabidopsis* development. *Plant Physiol.* **104**, 1139–1149 (1994).
58. Biswal, B. & Choudhury, N. K. Photocontrol of chlorophyll loss in papaya leaf discs. *Plant Cell Physiol.* **27**, 1439–1444 (1986).
59. Meng, Y., Li, H., Wang, Q., Liu, B. & Lin, C. Blue light-dependent interaction between cryptochrome2 and CIB1 regulates transcription and leaf senescence in Soybean. *Plant Cell* **25**, 4405–4420 (2013).
60. Guaiamet, J. J., Willems, J. G. & Montaldi, E. R. Modulation of progressive leaf senescence by the red:far-red ratio of incident light. *Bot. Gaz.* **150**, 148–151 (1989).
61. Cutler, S. R., Rodriguez, P. L., Finkelstein, R. R. & Abrams, S. R. Abscisic acid: emergence of a core signaling network. *Annu. Rev. Plant Biol.* **61**, 651–679 (2010).
62. Lopez-Molina, L. & Chua, N. H. A null mutation in a bZIP factor confers ABA-insensitivity in *Arabidopsis thaliana*. *Plant Cell Physiol.* **41**, 541–547 (2000).
63. Lichtenthaler, H. K. in *Methods in Enzymology* Vol. 148 (eds Douce, R. & Packer, L.) 350–382 (Academic Press, 1987).
64. Smyth, G. K. in *Bioinformatics and Computational Biology Solutions Using R and Bioconductor Statistics for Biology and Health* Ch. 23 (eds Gentleman, R. *et al.*) 397–420 (Springer, 2005).
65. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl Acad. Sci. USA* **102**, 15545–15550 (2005).
66. Nemhauser, J. L., Hong, F. & Chory, J. Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* **126**, 467–475 (2006).
67. Fujita, Y. *et al.* Three SnRK2 protein kinases are the main positive regulators of abscisic acid signaling in response to water stress in *Arabidopsis*. *Plant Cell Physiol.* **50**, 2123–2132 (2009).
68. Reeves, W. M., Lynch, T. J., Mobin, R. & Finkelstein, R. R. Direct targets of the transcription factors ABA-insensitive(ABI)4 and ABI5 reveal synergistic action by ABI4 and several bZIP ABA response factors. *Plant Mol. Biol.* **75**, 347–363 (2011).
69. Usadel, B. *et al.* Global transcript levels respond to small changes of the carbon status during progressive exhaustion of carbohydrates in *Arabidopsis* rosettes. *Plant Physiol.* **146**, 1834–1861 (2008).

Acknowledgements

We thank Drs D.R. Wagner, M. Ishiura and P.H. Quail for providing the *ELF3-OX*, *pcl1-1* and *elf4-101* seeds, respectively. This work is supported by the National Research Foundation of Korea (NRF grant no. 2011-0017308 to N.-C.P.) and (NRF-2012R1A2A1A01003133 to G.C.).

Author contributions

Y.S., J.J., M.-Y.K. and J.K. performed the experiments. N.-C.P. and G.C. designed the study and supervised the work. All authors discussed the results and made substantial contributions to the manuscript.

Additional information

Accession codes: Gene information from this article can be found in the Arabidopsis Genome Initiative or GenBank/EMBL databases as the following accession numbers: *ABI5*, At1g16540; *CO*, At5g15840; *CRY1*, At4g08920; *CRY2*, At1g04400; *DETI*; At4g10180; *EEL*, At1g30230; *ELN3*, At3g20770; *ELF3*, At2g25930; *ELF4*, At2g40080; *GAPDH*, At3g26650; *Lhcb1*, At1g29910; *LUX*, At3g46640; *NYC1*, At4g13250; *ORE1*,

At5g39610; *PHYA*, At1g09570; *PHYB*, At2g18790; *PIF1*, At2g20180; *PIF3*, At1g09530; *PIF4*, At2g43010; *PIF5*, At3g59060; *PILI*, At2g46970; *PP2A*, At1g13320; *RGA*, At2g01570; *SAG12*, At5g45890; *SEN4*, At4g30270; *SGR1*, At4g22920. Microarray data were deposited in the NCBI Gene Expression Omnibus (GEO) with accession number GSE52646.

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Sakuraba, Y. *et al.* Phytochrome-interacting transcription factors PIF4 and PIF5 induce leaf senescence in *Arabidopsis*. *Nat. Commun.* 5:4636 doi: 10.1038/ncomms5636 (2014).