

## PHYTOMEDICINAL VALUE OF *Moringa oleifera* WITH SPECIAL REFERENCE TO ANTIPARASITICS

**Tabassam Fatima, Muhammad Sohail Sajid\*, Muhammad Jawad-ul-Hassan, Rao Muhammad Siddique and Zafar Iqbal**

**Department of Parasitology, Faculty of Veterinary Science, University of Agriculture, Faisalabad-38040, Pakistan**  
\*Corresponding author's e-mail: drsohailuaf@hotmail.com

Plants are claimed as folk medicine for their therapeutic activity. *Moringa (M.) oleifera*, known as the "miracle tree", is greatly esteemed for its unique nutritional and medicinal value. Nutritively, it contains essential, disease-preventing nutrients. The methanolic and ethanolic extracts of plants have anthelmintic activity through paralysis of helminths. The compounds found responsible so far for their anthelmintic activities include: niazirinin, glycoside, niazirin and three glycosides from mustard oil, niaziminin A, B and 4-[4'-O-acetyl- $\alpha$ -L-rhamnosyloxy] benzyl] isothiocyanate isolated from leaves; D-glucose, D-mannose, ascorbic acid, protein and polysaccharide isolated from mature flowers; 0-[2'-hydroxy-3'-(2"-heptenyloxy)] propylundecanoate, methyl-p-hydroxybenzoate, thiocarbonates, isothiocyanate, nitriles, 0-ethyl-4-[( $\alpha$ -1-rhamnosyloxy)-benzyl] carbamate, and  $\beta$ -sitosterol isolated from whole pods of *M. oleifera*. Parasites are one of the major causes of diseases in human and animals. Due to parasitic infections livestock industry suffers from huge economic losses. The parasites which have been treated using different parts of *M. oleifera* include: Dracunculiasis (guinea worm), schistosomes and trypanosomes. Plant extracts may serve as potential candidates in future to exterminate helminthiasis in human and livestock populations. Hence, isolation, characterization and *in vivo* and *in vitro* efficacy trials of *M. oleifera* derivatives on scientific grounds are direly needed to elucidate it as a noteworthy candidate as neutraceutical and anthelmintic.

**Keywords:** Phytomedicine, *Moringa oleifera*, antiparasitic, neutraceutical, ethnopharmacology

### INTRODUCTION

Plants are being used all over the world for the cure of countless disease conditions. Treatment of diseases using plants or their extracts is called as "phytotherapy". It is sometimes also called as herbalism, plant medicine and herblore. Phytotherapy is considered under the category of Pharmacognosy, which studies medicines acquired from substances occurring in nature.

Since ancient times, traditions and folklore declare plants and their parts to be used against ailments. In northern Iraq, about 60,000 years old at the burial site of Shanidar IV, huge amount of pollens have been recovered belonging to 8 species of plants, of which 7 are used as phytomedicine nowadays (Solecki and Ralph, 1975). Records testify the use and study of herbs even old as 5000 years. At that time; the Sumerians used laurel, caraway and thyme for medicinal purpose (Sumner, 2000). Egyptians, considered expert in the use of herbal remedies, used castor oil, coriander, mint, indigo, garlic and opium etc. in 1000 BC (Sumner, 2000). Usage and art of cultivation is mentioned in the first part of Bible about mandrake, caraway, wheat, barley and rye. Herbs e.g. turmeric were used in Ayurveda medicine since 1900 BC in India (Aggarwal *et al.*, 2007). Sushruta in 6<sup>th</sup> century BC explained 700 herbal preparations, 64 mineral preparations and 57 preparations of animal origin. Ma-

Huang shrub is considered as the reason for the introduction of ephedrine to the list of modern medicines (Sumner, 2000). Greeks and Romans were expert in the usage of herbal therapy against various ailments. Scripts from Hippocrates paved the way for Western medicine, so as the scripts of Galen (Loudon, 2002). Hippocrates used to prescribe empirical herbal mixtures along with the prescription of rest and balanced diet; whereas, Galen used plants, animals and mineral ingredient sources for drugs formulation and prescribed larger dosage regimen (Loudon, 2002). Arab traders also had access to the markets of China and India, from where they purchased raw plant materials. In 13<sup>th</sup> century, Ibn-al-Baitar depicted 1400 plants, foods and drugs, and he also discovered 300 of his own (Boulanger, 2002). In the same century, Abu-al-Abbas, Al-Nabati and Andalusian-Arab botanist and educator of Ibn-al-Baitar, initiated experimentation in the field of Materia Medica (Huff, 1996). He verified transcripts that were supported by pillars of observation and testing. This advancement of the "Materia Medica" leads to the development of pharmacology.

Among all other diseases, parasites are major havoc for livestock in all the tropical countries (Githiori *et al.*, 2004). Parasites either external or internal, when attack their host, lead to stunted growth, drop down productivity, increase in death rate and serious economic disturbance (Iqbal *et al.*, 1993; Sykes, 1994; Kochapakdee *et al.*, 1995). This review

comprehensively describes few selected herbs having antiparasitic activity and potential of *Moringa oleifera* as an emerging candidate for phytochemical control of parasites.

**Plants having anti-parasitic activity:** The plant kingdom is the oldest best known opulent pool of herbal antibacterials, anthelmintics and insecticides (Satyavati *et al.*, 1976; Zaman *et al.*, 2012; Babar *et al.*, 2012; Hamad *et al.*, 2013). Uncountable figures of medicinal/herbal plants have been used against parasites. Up till now, no complete record of plants being used as antiparasitics is there. Usually the knowledge about antiparasitics passes from one generation to the next (Cox, 2000). Round the globe, the major plants having antiparasitic activity are: *Allium sativum* (Iqbal *et al.*, 2001), *Annona senegalensis* (Ibrahim *et al.*, 1984), *Acacia albida* (Nwude and Ibrahim, 1980), *Adhatoda vesica* (Lateef *et al.*, 2003), *Agati gratifolia* (Kalesaraj, 1975), *Ageratum conyzoides* (Sharma *et al.*, 1979), *Aglaiod odoratissima* (Nanda *et al.*, 1987), *Agrimonia (Ag.) eupatori*, *Ag. pilosa* (Xiao and Lin, 1986), *Alangium lamarckii* (Dubey and Gupta, 1969), *Albizia (Al.) anthelmintica* (Minja, 1989), *Al. coriavera* and *Al. lebbeck*, *Aloe barteri*, *Alpinia calcarata* and *Cucuruma aramatica* (Kalesaraj, 1975), *Amomum aromaticum*, *Amomora wallichii*, *Calamintha umberosa*, *Picus religiosa*, *Sentia myrtina* and *Sumplocos crataegoides* (Kaushik *et al.*, 1981), *Anacardium occidentale* (Garg and Kasera, 1982a,b), *Ananas (An.) comosus*, *An. sativus* (Neogi *et al.*, 1964), *An. sativus* (Chakraborty *et al.*, 1976), *Annona(Ann.) cherimolia*, *Ann. muricata*, *Ann. brasiliensis* and *Molinema dessetae* (Bories *et al.*, 1991), *Anogeissus (Ano.) leiocarpus* (Ibrahim *et al.*, 1984), *Ano. leiocarpus*, *Securinega virosa*, *Khaya (K.) senegalensis* and *Nauuclea latifolia*, *Anthocephalus indicus* (Kaushik *et al.*, 1981), *Areca (Ar.) catechu*, *Artabotrys odoratissimus* (Siddiqui and Garg, 1990), *Artemisia (Art.) abrotanum* (Krause, 1993), *Art. absinthium* (Guarrera, 1999), *Art. annua* (Shuhua *et al.*, 2000), *Art. brevifolia* (Iqbal *et al.*, 2004), *Art. herbaalba*, *Art. inflorescence* (Hammond *et al.*, 1997), *Art. maritime* (Sherif *et al.*, 1987), *Art. mesatlantica*, *Art. monosperma* (Abu-Niaaj *et al.*, 1996), *Art. pallens* (Nakhare and Garg, 1991), *Art. scoparia* (Naqvi *et al.*, 1991), *Azadirachta (Az.) indica*, *Melia azedarach*, *An. comosus*, *Vernonia anthelmintica*, *Embelia ribes*, *Fumaria parviflora* and *Caesalpinia crista*, *Bixa orellana*, *Boswellia (Bo.) dalzelii* (Nwude and Ibrahim, 1980), *Bo. serrata* (Girgune *et al.*, 1978), *Buddleia asiatica*, *Butea (Bu.) frondosa* (Shilaskar and Parashar, 1989), *Bu. frondosa* (Mehta and Parashar, 1966), *Bu. frondosa* (Lal *et al.*, 1976), *Bu. monosperma* (Iqbal *et al.*, 2006), *Bu. superba*, *Caesalpinia crista* (Akhtar *et al.*, 1985), *Calliandra (Ca.) calothrysus* (Parker and Palmer, 1991), *Ca. portoricensis* and *Calotropis (Cal.) procera* (Iqbal *et al.*, 2005), *Carica papaya* (Lal *et al.*, 1976), *Carum copticum* (Kalesaraj, 1975), *Cassia (Cs.) alata*, *Cs. accidentalis* (Ibrahim *et al.*, 1984), *Emblea myrobalans* (Gaind *et al.*, 1964), *Chenopodium album*

(Akhtar *et al.*, 1999), *Chloroylon swientenia*, *Chrysanthemum spp.* (Rebrassier, 1934), *Chrysophyllum cainito*, *Cinnamomum tamala* (Girgune *et al.*, 1978), *Cissampelos mucromata* (Minja, 1989), *Citrus (Ci.) acida*, *Ci. aromatic* and *Ci. medico* (Kalesaraj, 1975), *Combretum mucronatum* (Sofowora, 1982), *Commiphora mukul* (Kakrani and Kalyani, 1984), *Croton macrostachys* (Minja, 1989), *Cucurbita (Cu.) rhexicana* (Shrivastava and Singh, 1967), *Cu. moschata* (Xiao and Lin, 1986), *Cu. pepo* and *Momordica charantia* (Sharma *et al.*, 1971), *Cyathocline lyrata* (Shrivastava, 1979), *Cymbopogon (Cy.) nardus* and *Cy. citratus* (Kokate and Varma, 1971), *Cyperus rotundus* (Girgune *et al.*, 1979), *Datura (Da.) quercifolia* and *Da. metal* (Kaushik *et al.*, 1981), *Diospyros mollis* (Sen *et al.*, 1974), *Dodonea viscosa* (Sharma and Singh, 1989), *Embelia (Em.) kilimandschiraca* (Minja, 1989), *Em. schimperi* (Bogh *et al.*, 1996), *Em. ribes* (Akhtar *et al.*, 1990; Qureshi and Sabir, 1979), *Erythrina senegalensis* (Nwude and Ibrahim, 1980), *Eupatorium triplinerve* (Garg and Nakhare, 1993), *Evodia rutaecarpa* (Perrett and Whitfield, 1995), *Feruia foetidissima*, *Ficus religiosa* (Iqbal *et al.*, 2001), *Flemingia (Fl.) vestita* (Pal and Tandon, 1998), *Fl. vestita* (Kar *et al.*, 2002), *Fumaria parviflora* (Kailani *et al.*, 1995), *Gardenia lucida* (Girgune *et al.*, 1979), *Hedychium (He.) coronarium* and *He. spicatum* (Dixit and Varma, 1975), *Helleborus niger* (Kalesaraj, 1975), *Heracleum sosnowskyi*, *Hyoscyamus niger* (Akhtar and Ahmad, 1990), *Inula racemosa* (Mishra *et al.*, 1979), *Juglans regia*, *Musa paradisaca* and *Scindapsus officinalis* (Sharma *et al.*, 1971), *K. senegalensis*, *Lagenaria siceraria* (Akhtar and Riffat, 1987), *Lantana camavarav*, *Lawsonia inermis* (Nwude and Ibrahim, 1980), *Leucaena leucocephala*, *Limnophila conferta* (Reddy *et al.*, 1991), *Litsea chinensis* (Mishra *et al.*, 1979), *Macuna pruriita* (Neogi *et al.*, 1964), *Mangifera indica* (Kalesaraj, 1975), *Matteuccia orientalis* (Shiramizu *et al.*, 1993), *Mitragyna stipulosa* (Sofowora, 1982), *M. oleifera* (Akhtar and Ahmad, 1990), *Nicotiana (N.) tabacum* and *Nigella sativa* (Kailani *et al.*, 1995), *Peganum harmala* (Akhtar and Ahmed, 1991), *Piper (Pi.) betle* (Ali and Mehta, 1970), *Psitacia integrifolia* (Mishra *et al.*, 1979), *Psoralea coylifolia* (Akhtar, 1985), *Punica granatum* (Prakash *et al.*, 1980), *Quisqualis indica* (Xiao and Lin, 1986), *Randia dumetorum* (Mishra *et al.*, 1979), *Rapanea melanoploeo*s and *Sapindus trifoliatum* (Lal *et al.*, 1976), *Saussurea lappa* (Akhtar and Hassan, 1985), *Senecio lyratiparus* (Minja, 1989), *Solanum nodiflorum* (Nwude and Ibrahim, 1980), *Spigelia anthelmia* (Assis *et al.*, 2003), *Swertia chirata* (Shilaskar and Parashar, 1989), *Tiinospora rumphii*, *Tribulus terrestris* (Chakraborty *et al.*, 1979), *Uvaria (U.) hookeri*, *U. narum* (Padmaja *et al.*, 1993), *Vernonia (V.) amygdalina* (Alawa *et al.*, 2003), *V. anthelmintica* (Mehta and Parashar, 1966), *Withania (W.) coagulans* (Singh *et al.*, 1982), *Zanthoxylum alatum* (Kalyani *et al.*, 1989) and *Zingiber officinale* (Adewunmi *et al.*, 1990).

Predominant reports of medicinal plants in Pakistan include those of Akhtar *et al.*, 1984, 1985, 1987, 1990, 1991 and 1999; Abbas *et al.*, 2010; Hussain *et al.*, 2010; Iqbal *et al.*, 1993, 2001, 2004, 2005 and 2006; Sindhu *et al.*, 2010; Awais *et al.*, 2011; Badar *et al.*, 2011; Hussain *et al.*, 2011; Ashfaq and Ashfaq, 2012. According to their findings the most important medicinal plants present in Pakistan are *Acacia nilotica*, *Allium (Al.) cepa*, *Al. sativum*, *Aloe vera*, *Areca catechu*, *Az. indica*, *Brassica campestris*, *Cal. procera*, *Capsicum annum*, *Citrullus colocynthis*, *Ci. medica*, *Coriandrum sativum*, *Curcuma longa*, *Eruca sativa*, *Ferula asafetida*, *Foeniculum vulgare*, *Linum usitatissimum*, *Mallotus philippensis*, *Me. azedarach*, *Mentha arvensis*, *M. oleifera*, *N. tabacum*, *Phoenix dactylifera*, *Pi. nigrum*, *Terminalia chebula*, *Trachyspermum ammi*, *Triticum aestivum*, *V. anthelmintica*, *W. coagulans* and *Ziziphus jujube*.

**Introduction to *Moringa*:** *Moringa* belongs to family Moringaceae which contains thirteen species (Olson, 2002) from tropics to subtropics of the world in the form of tiny herbs to massive trees (Ramachandran *et al.*, 1980). It is being cultivated in Ethiopia, Kenya, Afghanistan, Namibia, Bangladesh, Angola, Pakistan, Madagascar, Africa and India (Stephenson and Fahey, 2004; Trapti *et al.*, 2009). *Moringa* is also widely distributed on tropics of the world including Ghana, Uganda, Egypt, Sierra Leone, Philippines, Nicaragua, Kenya, Haiti and many more. It can thrive well with minimal moisture contents as its roots store moisture contents for long period. It also thrives well in Pakistan, Bangladesh, Sri Lanka, Tropical Africa, Arabia, Cambodia and Central North and South America (Qaiser *et al.*, 1973; Sofowora *et al.*, 1982; Somali *et al.*, 1984; Morton *et al.*, 1991; Mughal *et al.*, 1999; Shahzad *et al.*, 2013).

Most extensively cultivated species under this family is *M. oleifera*. Vernacular names of *Moringa* in different languages of the world include: Rawag (Arabic), Daintha, Dandalonbin (Burmese), La Ken (Chinese), Maissang, Moring, Moxing (Konkani), Moringa, Moringueiro (Portuguese), Murunga (Sinhalese), Angela, Ben, Moringa (Spanish), Mrongo, Mzunze (Swahili), Sahjan (Hindi), Horseradish tree, Drumstick tree, Moringa (English), *Moringa oleifera* (Latin), Surajana (Sanskrit), Amukira (Tamil), Keramaddinagaddi and Nugge (Kannada), Mulakkaya (Telgu), Muringa, Murinna and Sigru (Malayalam), Achajhada and Shevga (Marathi), Saragvo (Gujarati), Sajiwani or Swejan (Nepali), Sojne danta (Bengali), Sajana or Sujuna (Oriya), Suhanjana (Punjabi), Sojina (Assamese) and Murunga, Sojina (Sinhalese). Specifically, *M. oleifera* is also known as: the Ben oil tree, Benzoline tree, Kelor, Marango, Mlonge, Moonga, Mulangay, Nebeday, Saijhan, Sajna, Miracle tree and "Mothers best friend" (Coppin, 2008). It is being cultivated in Himalayas in north western India and all over the tropics.

A diverse range of species belonging to genus *Moringa* have been reported from Africa (*M. arborea*, *M. borziana*, *M. longituba*, *M. oleifera*, *M. peregrine*, *M. pygmaea*, *M. rivae*, *M. ruspoliana*, *M. stenopetala*, *M. drouhardii*, *M. hildebrandtii*, *M. ovalifolia*) and from India (*M. concanensis*, *M. oleifera*) is cultivated.

**History of the miracle plant:** *M. oleifera* is highly valued member of genus *Moringa*, which comprises of thirteen species (Mabberley, 1997). Plantation of *Moringa* started from the West Indies in the 19<sup>th</sup> century. Later on, people from Asia and West Africa started using it as food supplement. Mostly, the leaves and pods were used as food while seed oil was used for lubrication. *Moringa* is the tree of tropical and sub-tropical countries with medium size, no or very rare branching, flowers are creamy white and sweetly scented (Sastri, 1962). Silvi-culture of *M. oleifera* is being introduced to elicit poverty and malnutrition therefore improving health status (Parrotta *et al.*, 2001).

**Medicinal activity of *Moringa*:** The identified species of *Moringa* have been reported for their specified geographical uses like; traditional use to combat anemia, anxiety, asthma, blackheads, blood impurities, bronchitis, catarrh, chest congestion, cholera, conjunctivitis, cough, diarrhea, eye and ear infections, fever, glandular swelling, headaches, abnormal blood pressure, hysteria, pain in joints, pimples, psoriasis, respiratory disorders, scurvy, semen deficiency, sore throat, sprain, tuberculosis (India), intestinal worms (Malaysia and Puerto Rico), skin infections and sores (Guatemala), for anemia and glandular swelling (Philippines).

As *M. oleifera* is a medicinal plant, so all of its parts are utilized against various diseases worldwide. Roots being used against rheumatism, as anti-inflammatory, cardiac/circulatory tonic, carminative, joint pains, kidney pain, as laxative, as abortion inducer, rubefacient, stimulant in paralytic afflictions and vesicant (Sastri, 1962; Dahot, 1988; Ruckmani *et al.*, 1998). Leaves are being utilized against catarrh, scurvy, purgation, applied as poultice, rubbed for headaches, fevers, otitis, sore throat, bronchitis, eye infections; leaf extract regulates the level of glucose in the body and lowers down the swelling in glands (Sastri, 1962; Dahot, 1988; Morton, 1991). Stem and bark are used as rubefacient, vesicant, to cure eye diseases, prevent the spleenomegaly and tubercle formation in the neck, eradicates tumors and resolves the ulceration. Root bark extract is dropped in ear to relieve aches and also relieves tooth aches (Bhatnagar *et al.*, 1961). Gum acts as rubefacient and astringent. Moreover, gum when emulsified with sesame oil, acts as preventive therapy against asthma, dysentery, fever, headache, intestinal complaints and acts as abortion inducer. Flowers are of great value in the field of medicine so used as abortion inducer, anti inflammatory, relieves the muscular diseases, antitumerous, against spleenomegaly, lowers down the level of fats and lipids in the serum,

decrease lipids levels of liver and heart (Bhattacharya *et al.*, 1982; Dahot, 1998; Mehta *et al.*, 2003). Seed extract decreases liver lipid peroxides (Faizi *et al.*, 1998; Lalas and Tsaknis, 2002).

The research also supports the traditional medicinal uses of *M. oleifera* as antipyretic (Sutar *et al.*, 2009) and antiasthmatic (Mahajan *et al.*, 2007 and 2009; Agarwal *et al.*, 2008) due to its ethanolic extracts present in leaf and seed. Its wound healing ability is attributable to seeds containing ethanol and ethyl acetate (Hukkeri *et al.*, 2006; Rathi *et al.*, 2006). In addition, it is antithyroid (Tahiliani and Kar, 2000), anti-inflammatory, antiarthritic, analgesic (Rao, 1998; Mahajan *et al.*, 2007; Sutar *et al.*, 2008; Sashidhara *et al.*, 2009), hypocholesterolemic (Zlatkis *et al.*, 1953; Ghazi *et al.*, 2000; Mehta *et al.*, 2003; Chumark *et al.*, 2008), antimicrobial (Kurup and Rao, 1954; Das *et al.*, 1957; Bhatnagar *et al.*, 1961; Eilert *et al.*, 1981; Caceres *et al.*, 1991; Dahot *et al.*, 1998; Nikkon *et al.*, 2003; Chuang *et al.*, 2007; Ayanbimpe *et al.*, 2009; Lürling and Beekman, 2010), anaphylactic, hepatoprotective (Ruckmani *et al.*, 1998; Pari and Kumar, 2002; Hamza, 2007; Fakurazi *et al.*, 2008; Selvakumar and Natarajan, 2008; Hamza, 2010), radioprotective (Rao *et al.*, 2001), antiulcer (Akhtar and Ahmad, 1995; Pal *et al.*, 1995; Ruckmani *et al.*, 1998; Debnath and Guha, 2007; Devaraj *et al.*, 2007), antispasmodic (Caceres *et al.*, 1992; Gilani *et al.*, 2007), antihyperglycemic (Ndong *et al.*, 2007; Jaiswal *et al.*, 2009). Major compounds in *M. oleifera* to fight against diseases are: 4-(4'-*O*-acetyl- $\alpha$ -L-rhamnopyranosyloxy) benzyl isothiocyanate, 4-( $\alpha$ -L-rhamnopyranosyloxy) benzyl isothiocyanate, niazimicin, pterygospermin, benzyl isothiocyanate and 4-( $\alpha$ -L-rhamnopyranosyloxy) benzyl glucosinolate (Kerharo, 1969; Sashidhara *et al.*, 2009). Table 1 provides the list of important chemical constituents present in *M. oleifera* which have extensively been reported by Fahey (2005). Isoleucine builds proteins and enzymes, leucine works with isoleucine to enhance energy level. Lysine helps in accurate absorption of calcium and aids in the antibody production. Methionine predominantly supports the sulfur provision to the body therefore decreases the irritation in the bladder. Phenylalanine yields the chemical transmitters between nerve cells and brain. Threonine is an integral portion of fibrous proteins. Tryptophan supports immune system and reduces cholesterol level while valine helps in muscle coordination. Body manufactures non-essential amino acids, when nutrient supply is proper. These non-essential amino acids are also found abundantly in Moringa having diverse types of functions e.g. alanine helps in energy build up, arginine is helpful in the release of the growth hormones and improves immune status, aspartic acid helps rid from cellular waste, cysteine functions as an antioxidant, glutamic acid is food for the brain, glycine

promotes the oxygen supply which is integral part in the cell-making phenomenon, histidine is used in the treatment of rheumatoid arthritis, allergies, ulcers, and anemia while serine helps in glucose preservation in the liver as well as in muscles whereas proline is tremendously significant for the appropriate functioning of tendons and joints. It also helps to sustain the integrity and strength of heart muscles. Tryrosine helps in nerve impulse transmission process so in this way it helps coverup depression, improves memory, increases mental alertness, promote healthy functioning of the thyroid, adrenal and pituitary glands.

**Nutritive value of Moringa:** The key components of Moringa are omega-3 and omega-6 fatty acids, deic, saponins, palmitic acid, glycoside, stearic acid, gum, vitamins (A, B1, B2, B3 and C), protein, minerals (calcium, iron, phosphorus, magnesium) (Kasolo, 2010). The leaves, flowers and pods are used as substantial sources of ascorbic acid, alpha-tocopherol beta-carotene, calcium, folic acid, iron, nicotinic acid, pyridoxine, vitamins, and riboflavin (Dahot, 1988). A compound, pterygospermin, has been stated to possess potent antibiotic and antifungal activity (Das *et al.*, 1957). The root bark contains alkaloids namely moringinine and moringine (Shukla *et al.*, 1988). Upon the nutritional analysis of moringa pods, fresh raw leaves and dried leaf powder have been explored (<http://www.edlagman.com/moringa/moringa-fresh-leaf-vs-dried-leaf.pdf>) containing moisture (86.9%, 75% and 7.5%), calories (26.0, 92.0 and 205.0), proteins (2.5g, 6.7g, 27.1g), fat (0.1g, 1.7g and 2.3g), carbohydrates (3.7g, 13.4g and 38.2g), fiber contents (4.8g, 0.9g and 19.2g) respectively. High levels of calcium, phosphorous, magnesium, potassium, copper, iron, oxalic acid and sulphur and other trace minerals are also observed. Vitamins profile is fabulous like it contains carotene, thiamin, riboflavin, nicotinic acid, ascorbic acid and tocopherol acetate. Earlier studies on the phytochemical analysis has shown the presence of alkaloids, amino acids, carbohydrates, flavonoids, glycosides, proteins, saponins, steroids and tannins in ethanolic extracts of plant (Rastogi *et al.*, 2009;). The leaves are rich source of vitamin A, B and C, protein and essential amino acids. Cattles which are kept on feeding of Moringa plant, have shown an increase of 32% in weight gain and 43% to 65% in milk production (Mathur, 2006). Moringa leaves contains a plenty full of vitamin A, calcium, iron, vitamin C, potassium and high levels of protein (Singh, 2007; Silja, 2008). The seeds have higher contents of oil i.e. about 35% while left over meal is having about 65% of protein content (Anwar *et al.*, 2005). The dry seed powder and defatted meal is a flocculent and can be used in water purification to settle down sediments and undesirable organisms (Muyibi *et al.*, 2009). Traditionally its roots are applied as plaster to reduce the swelling and rheumatism.

**Table 1. A review of chemical constituents of various body parts of *Moringa oleifera***

Parts	Phytochemical constituents	References
Leaves	Glycoside niazirin, niazirinin and three mustard oil glycosides, 4-[4'-O-acetyl- α-L-rhamnosyloxy) benzyl] isothiocyanate, niaziminin A and B	Faizi <i>et al.</i> , 1994; Verma <i>et al.</i> , 2009; Anwar <i>et al.</i> , 2005; Makkar and Becker, 1996; Geervani and Devi, 1981; Das, 1965; Lakshminarayana <i>et al.</i> , 2005; Seshadri and Nambiar, 2003
Stem	4-hydroxymellein, vanillin, β-sitosterone, octacosanic acid and β-sitosterol	Saluja <i>et al.</i> , 1978; Faizi <i>et al.</i> , 1994
Bark	4-(α-L-rhamnopyranosyloxy)-benzylglucosinolate	Kerharo, 1969
Whole gum exudates	L-arabinose, D-galactose, D-glucuronic acid, L-rhamnose, Dmannose, D-xylose and leucoanthocyanin	Bhattacharya <i>et al.</i> , 1982
Roots	4-(α-L-rhamnopyranosyloxy) benzylglucosinolate and Benzylglucosinolate	Sashidhara <i>et al.</i> , 2009
Mature flowers	D-mannose, D-glucose, protein, ascorbic acid, polysaccharide	Faizi <i>et al.</i> , 1994; Ruckmani <i>et al.</i> , 1998
Whole pods	Nitriles, isothiocyanate, thiocarbonates, 0-[2'-hydroxy-3'-(2''-heptenoxy)] propylundecanoate, 0-ethyl-4-[( α -1-rhamnosyloxy)-benzyl] carbamate, methyl-p-hydroxybenzoate and β-sitosterol	Faizi <i>et al.</i> , 1998; Nagar <i>et al.</i> , 1981, 1982; Roy <i>et al.</i> , 2007
Leaves and Pods	Glycosides-carbamate,thiocarbamate, and isothiocyanate Ascorbic acid, oestrogenic substances, β-sitosterol, iron, calcium, phosphorus, copper, Vitamin A, B, C, α-tocopherol, riboflavin, nicotinic acid, folic acid, pyridoxine, β-carotene, proteins, essential amino acids – methionine, cystine, tryptophan and lysine Novel bioactive nitrile glycosides-Niaziridin and niazirin α and gamma tocopherols	Faizi <i>et al.</i> , 1994; Makkar and Becker, 1996; Tsaknis <i>et al.</i> , 1999; Lako <i>et al.</i> , 2007; Verma <i>et al.</i> , 1976; Shanker <i>et al.</i> , 2007; Jadhav <i>et al.</i> , 2000
Leaves, flowers and fresh beans		Guevara <i>et al.</i> , 1999
Mature seeds	Crude protein, Crude fat, carbohydrate, methionine, cysteine, 4-(α- L-rhamnopyranosyloxy)-benzylglucosinolate, benzylglucosinolate, moringyne, mono-palmitic and di-oleic triglyceride, benzyl isothiocynate	Eilert and Wolters, 1981; Lalas and Tsaknis, 2002; Bennett <i>et al.</i> , 2003; Katre <i>et al.</i> , 2008; Fahey <i>et al.</i> , 2001
Seed oil	Vitamin A, beta carotene, precursor of Vitamin A	Anwar <i>et al.</i> , 2005; Tsaknis <i>et al.</i> , 1999; Anwar and Bhanger, 2003; Ogunbinu <i>et al.</i> , 2009; Banerji <i>et al.</i> , 2009

**Antiparasitic activity of *M. oleifera*:** It is observed that *M. oleifera* is having potent anthelmintic activity and its gum is being used as an anti-filarial agent (Kushwaha *et al.*, 2011). The most targeted parasite species with the use of *M. oleifera* are helminths including Dracunculiasis, schistosomes and trypanosomes (Fahey, 2005). *In vitro* activity of *M. oleifera* has shown some antiprotozoan activity (Kohler *et al.*, 2002) and is larvical because of soluble lectin present in its seed extract. It hinders the process of larval development and due to its haemagglutinating activity therefore it causes mortality in *Aedes aegypti* (Coelho *et al.*, 2009; Ferreira *et al.*, 2009). Water extract of *M. oleifera* is known to have larvicidal, pupicidal as well as adult mosquito killer properties against the *Culex quinquefasciatus* (Ashfaq and Ashfaq, 2012). It is being used as antimarial (Eilert *et al.*, 1980; Gbeassor *et al.*,

1990). Major anthelmintic activity is shown by the ethanolic extracts of the plant which when used at various concentrations can cause paralysis and death of the worms. Various concentrations of Moringa extract take varied period of time for paralysis and death of the worms as 25 mg/ml takes  $14.32 \pm 2.2$  min and  $63.57 \pm 12.6$  min, 50 mg/ml takes  $8 \pm 1.6$  min and  $52.33 \pm 3.1$  min while 100 mg/ml takes  $6 \pm 1.2$  min and  $45 \pm 11.4$  min respectively, as experimentally noted by Gosh *et al.*, (2005).

**Miscellaneous uses of *Moringa oleifera*:** Some of the uses for Moringa encompass: alley cropping for biomass, leaves and treated seed-cake as animal forage (Mahatab *et al.*, 1987), leaves for biogas production, crushed leaves as domestic cleaning agent, wood for blue dye preparation, living trees for fencing the houses, seed-cake as natural fertilizer, juices extracted from the leaves as foliar nutrient,

leaves as green manure, gum from tree trunks, powdered seeds used to clarify honey and sugar cane juice, flower nectar as honey (Rajan, 1986), whole plant is used as medicine, ornamental plantings, dumped off leaves as biopesticide, wood to prepare pulp (Guha, *et al.*, 1968), bark for roping, bark and gum used in tanneries for tanning hides due to presence of tannin, dry seed powder for water purification purpose (Muyibi *et al.*, 2009).

**Conclusions:** Chemotherapy remains the corner stone in the treatment of majority of infectious diseases in the under developed countries. However, the efficacy is reduced over time against pathogens (including parasites) due to development of resistance against these drugs. In addition, the drug toxicity in the host leads to depressed immune status making it more prone to other infectious diseases (Chang, 2011). Under the given circumstances, use of plants and/or plant-derived medicines for the cure of infectious diseases, in general, and parasitic diseases, in specific, is justified. Lot of work has already been carried out during the past decade by various scientists of our country regarding antiparasitic efficacy of plants and their scientific validation (Abbas *et al.*, 2010; Hussain *et al.*, 2010; Sindhu *et al.*, 2010; Awais *et al.*, 2011; Badar *et al.*, 2011; Hussain *et al.*, 2011). Keeping in view the expected benefits and detailed nutritive and medicinal profile of *M. oleifera*, it is suggested that effective candidates from this miraculous plant should be extracted for potent activity against prevalent parasitic agents in livestock and human population. As discussed above, *M. oleifera*, an edible tree, being cultivated worldwide in almost all tropical countries, and is progressively being utilized to meet nutritional deficiency. Leaves are rich in nutrients and have high protein content therefore widely used in medicine by physicians, healers, nutritionists and community leaders, for the treatment of nutrition deficiency and other ailments related to it, but yet no clinical trials are applied for the testification of its efficacy to add it up as nutrition supplement. The development and implementation of dietary *M. oleifera* leaves has been described in "Diffusion of innovations theory" and it highlights the need for a scientific consensus on the nutritional benefits. It is suitable and economically friendly nutrient-rich food stuff for nutrient deficient populations. *M. oleifera* is capable to meet its high rate of overall utilization, macro and micronutrients availability, swift growth capability, high leaf yield and can survive in harsh climatic conditions. Therefore it is strongly recommended that *M. oleifera* should be adopted as a unique pan-tropical dietary plant not only for humans because of its good nutritional profile, but also for livestock rations as it contains high level of vitamins and minerals and its seeds are rich in protein. At present, cultivation of this plant is done in selected southern zone of Punjab; Researchers should try to explore the details of this plant on logical

scientific grounds to provide guidelines for its use in rationing and human supplementation. There is a dire need to start the cultivation of Moringa on national scale. In this connection, the agriculture section of the Government should facilitate farmers for technical and financial assistance whatever possible to boost up its production making it easily available for humans as well as livestock.

## REFERENCES

- Abbas, R.Z., Z. Iqbal and M.N. Khan. 2010. Anticoccidial activity of *Curcuma longa* L. in Broilers. *Braz. Arch. Biol. Technol.* 53:63-67.
- Abu-Niaaj, L., M. Abu-Zarga and S. Abdalla. 1996. Isolation and inhibitory effects of eupatilin, a flavone isolated from *Artemisia monosperma* Del., on rat isolated smooth muscle. *Int. J. Pharmacol.* 34:134-140.
- Adewunmi, C.O., O.B.O. Gtuitimein and P. Furu. 1990. Molluscicidal and antischistosomal activities of *Zingiber officinale*. *Planta Med.* 56:374-376.
- Agarwal, R., S.K. Gupta, S.S. Agrawal, S. Srivastava and R. Saxena. 2008. Oculohypotensive effects of *Foeniculum vulgare* in experimental models of glaucoma. *Indian J. Physiol. Pharmacol.* 52 : 77-83.
- Aggarwal, B.B., C. Sundaram, N. Malani and H. Ichikawa. 2007. Curcumin: the Indian solid gold. *Adv. Exp. Med. Biol.* 595:1-75
- Akhtar, A.H. and K.U. Ahmad. 1995. Anti-ulcerogenic evaluation of the methanolic extracts of some indigenous medicinal plants of Pakistan in aspirin-ulcerated rats. *J. Ethnopharmacol.* 46:1-6.
- Akhtar, M.S. and I. Ahmad. 1990. Anthelmintic and phytochemistry studies on *Hyoscyamus niger* Linn. (Ajwain Khurasani) seeds and *Morrinda oleifera*, Lam (Sohanjna) roots. *J. Pharmacol.* 3:75-81.
- Akhtar, M.S. and I. Ahmad. 1991. Evaluation of antinematodal efficacy of tetrahydrohaunine in goats. *Vet. Arhiv.* 6:307-311.
- Akhtar, M.S. and I.J. Hassan. 1985. Evaluation of *Saussurea lappa* roots (Qust) against natural infection of gastrointestinal nematodes in sheep. *Pak. J. Agri. Sci.* 22:1-7.
- Akhtar, M.S. and S. Riffat. 1984. Efficacy of *Melia azedarach* Linn. (Bakain) and morantel against naturally acquired gastrointestinal nematodes in goats. *Pak. Vet. J.* 4:176-179.
- Akhtar, M.S. and S. Riffat. 1987. Evaluation of antcestodal activity of *Lagenaria siceraria* (Kaddoo) seeds in sheep. *Pak. Vet. J.* 7:139-141.
- Akhtar, M.S., I. Javed, C.S. Hayat and B.H. Shah. 1985. Efficacy and safety of *Caesalpinia crista* Linn. seeds and its extracts in water and methanol against natural *Neoascaris vitulorum* infection in buffalo calves. *Pak. Vet. J.* 5:192-196.
- Akhtar, M.S., Z. Iqbal and M.N. Khan. 1999. Evaluation of anthelmintic activity *Chenopodium album* (Bathu)

- against nematodes in sheep. Int. J. Agric. Biol. 1:121-124.
- Alawa, C.B.I., A.M. Adamu, J.O. Gefu, O.J. Ajanusi, P.A. Abdu, N.P. Chiezey, J.N. Alawa and D.D. Bowman. 2003. *In vitro* screening of two Nigerian medicinal plants (*Vernonia amygdalina* and *Annona senegalensis*) for anthelmintic activity. Vet. Parasitol. 113:73-81.
- Ali, S.M. and R.K. Mehta. 1970. Preliminary pharmacological and anthelmintic studies of the essential oil of *Piper betle* L. Indian J. Pharmacol. 32:132-133.
- Anwar, F. and M.I. Bhanger. 2003. Analytical characterization of *Moringa oleifera* seed oil grown in temperate regions of Pakistan. J. Agric. Food Chem. 51:6558-6563.
- Anwar, F., M. Ashraf and M.I. Bhanger. 2005. Interprovenance variation in the composition of *Moringa oleifera* oil seeds from Pakistan. J. Am. Oil Chem. Soc. 82:45-51.
- Ashfaq, M. and U. Ashfaq. 2012. Evaluation of mosquitocidal activity of water extract of *Moringa oleifera* seeds against *Culex quinquefasciatus* (Diptera: culicidae) in Pakistan. Pak. Entomol. 34:21-26.
- Aslam, M., M. Ismat, R.H. Qureshi, S. Nawaz and I.A. Mehmood. 1994. Paddy yield affected by planting techniques in a salt-affected soil. Pak. J. Agri. Sci. 31:40-405.
- Assis, L.M., C.M.L. Bevilaqua, S.M. Morais, L.S. Vieira, C.T.C. Costa and J.A.L. Souza. 2003. Ovicidal and larvicidal activity *in vitro* of *Spigelia anthelmia* Linn. extracts on *Haemonchus contortus*. Vet. Parasitol. 117:43-49.
- Awais, M.M., M. Akhtar, F. Muhammad, A.U. Haq and M.I. Anwar. 2011. Immunotherapeutic effects of some sugar cane (*Saccharum officinarum* L.) extracts against coccidiosis in industrial broiler chickens. Exp. Parasitol. 128:104-110.
- Ayanbimpe, G.M., T.K. Ojo, E. Afolabi, F. Opara, S. Orsaah and O.S. Ojerinde. 2009. Evaluation of extracts of *Jatropha curcas* and *Moringa oleifera* in culture media for selective inhibition of saprophytic fungal contaminants. J. Clin. Lab. Anal. 23:161-164.
- Badar, N., Z. Iqbal, M.N. Khan and M.S. Akhtar. 2011. *In vitro* and *in vivo* anthelmintic activity of *Acacia nilotica* (L.) Willd. Ex Delile Bark and Leaves. Pak. Vet. J. 31:185-191.
- Babar, W., Z. Iqbal, M.N. Khan and G. Muhammad. 2012. An inventory of the plants used for parasitic ailments of animals. Pak. Vet. J. 32: 183-187.
- Banerji, R., A. Bajpai and S.C. Verma. 2009. Oil and fatty acid diversity in genetically variable clones of *Moringa oleifera* from India. J. Oleo. Sci. 58:9-16.
- Bennett, R.N., F.A. Mellon, N. Foidl, J.H. Pratt, M.S. Dupont, L. Perkins and P.A. Kroon. 2003. Profiling glucosinolates and phenolics in vegetative and reproductive tissues of the multipurpose trees *Moringa oleifera* L. (horseradish tree) and *Moringa stenopetala* L. J. Agric. Food Chem. 51:3546-3553.
- Bhatnagar, S.S., H. Santapau, J.D. Desa, A.C. Maniar, N.C. Ghadially and S. Yellore. 1961. Biological activity of Indian medicinal plants. Part 1. Antibacterial, anti-tubercular and antifungal action. Indian J. Med. Res. 49:799-805.
- Bhattacharya, S.B., A.K. Das and N. Banerji. 1982. Chemical investigations on the gum exudate from sajna (*Moringa oleifera*). Carbohyd. Res. 102:253-262.
- Bogh, H.O., J. Andreassen and J. Lemmich. 1996. Anthelmintic usage of extracts of *Embelia schimperi* from Tanzania. J. Ethnopharmacol. 50:35-42.
- Bories, C., L. Loiseau, D. Cortes, S.H. Myint, R. Hoquenuller, P. Gayral, A. Cave and A. Laurens. 1991. Antiparasitic activity of *Annona muricata* and *Annona cheromolia* seeds. Planta Med. 54:434-436.
- Boulanger, D. 2002. The Islamic contribution to science, mathematics and technology", OISE Papers, in STSE Education.
- Caceres, A., A. Saravia, S. Rizzo, L. Zabala, E. De Leon and F. Nave. 1992. Pharmacologic properties of *Moringa oleifera* 2: Screening for antispasmodic, anti-inflammatory and diuretic activity. J. Ethnopharmacol. 36:233-237.
- Caceres, A., O. Cabrera, O. Morales, P. Mollinedo and P. Mendia. 1991. Pharmacological properties of *Moringa oleifera* 1: Preliminary screening of antimicrobial activity. J. Ethnopharmacol. 33:213-216.
- Chakraborty, B., N.M. Ray and S. Sikdar. 1976. Anthelmintic and pharmacological study with an alcoholic extract of *Ananas sativus*. Indian J. Anim. Health. 15:117-119.
- Chakraborty, B., N.M. Ray and S. Sikdar. 1979. Study of anthelmintic property of *Tribulus terrestris* Linn. Indian J. Anim. Health. 1:23-25.
- Chang, A. 2011. Chemotherapy, chemoresistance and the changing treatment landscape for NSCLC. Lung Cancer. 71:3-10.
- Chuang, P.H., C.W. Lee, J.Y. Chou, M. Murugan, B.J. Shieh and H.M. Chen. 2007. Antifungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. Bioresource Technol. 98:232-236.
- Chumark, P., P. Khunawat, Y. Sanvarinda, S. Phornchirasilp, N.P. Morales, L. Phivthong-ngam, P. Ratanachamnong, S. Srisawat and K.S. Pongrapeeporn. 2008. The *in vitro* and *ex vivo* antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves. J. Ethnopharmacol. 116:439-446.
- Coelho, J.S., N.D. Santos, T.H. Napoleao, F.S. Gomes, R.S. Ferreira and R.B. Zingali. 2009. Effect of *Moringa oleifera* lectin on development and mortality of *Aedes aegypti* larvae. Chemosphere. 77:934-938.
- Coppin, J. 2008. A study of nutritional and medicinal values of *Moringa oleifera* leaves from susaharan Africa: Ghana, Rwanda Senegal and Zambia. M.Sc. Thesis.

- Cox, P.A. 2000. Will tribal knowledge survive the millennium. *Sci.* 287:44-45.
- Dahot, M.U. 1998. Antimicrobial activity of small portion of *Moringa oleifera* leaves. *J. Islamic Acad. Sci.* 11:27-32.
- Das, B.R., P.A. Kurup, R.P.L. Narasimha and A.S. Ramaswamy. 1957. Antibiotic principle from *Moringa pterygosperma* VII. Antibacterial activity and chemical structure of compounds related to pterygospermin. *Indian J. Med. Res.* 45:191-196.
- Debnath, S. and D. Guha. 2007. Role of *Moringa oleifera* on enterochromaffin cell count and serotonin content of experimental ulcer model. *Indian J. Exp. Biol.* 45:726-731.
- Devaraj, V.C., M. Asad and S. Prasad. 2007. Effect of leaves and fruits of *Moringa oleifera* on gastric and duodenal ulcers. *Pharmaceutical Biotechnol.* 45:332-338.
- Dixit, V.K. and K.C. Varma. 1975. Anthelmintic properties of essential oils from rhizomes of *Hedychium coronarium* Koenig and *Hedychium spicatum* Koenig. *Indian J. Pharm.* 37:143-144.
- Dubey, M.P. and I. Gupta. 1969. Anthelmintic activity of *Alangium lamarckii*. *Indian J. Physiol. Pharmacol.* 12:35-36.
- Eilert, U., B. Wolters and A. Nahrstedt. 1980. Antibiotic principle of seeds of *Moringa oleifera*. *Planta Med.* 39(3):235.
- Eilert, U., B. Wolters and A. Nahrstedt. 1981. The antibiotic principle of seeds of *Moringa oleifera* and *Moringa stenopetala*. *Planta Med.* 42:55-61.
- Fahey J.W. 2005. *Moringa oleifera*: a review of the medical evidence for its nutritional, therapeutic and prophylactic properties. Part 1. In: *Trees for Life Journal*. <http://www.tfljournal.org/article.php/20051201124931586>.
- Fahey, J.W., A.T. Zalcmann and P. Talalay. 2001. The chemical diversity and distribution of glucosinolates and isothiocyanates among plants. *Phytochemistry*. 56:5-51.
- Faizi, S., B.S. Siddiqui, R. Saleem, K. Aftab, F. Shaheen and A.H. Gilani. 1998. Hypotensive constituents from the pods of *Moringa oleifera*. *Planta Med.* 64:225-228.
- Faizi, S., B.S. Siddiqui, R. Saleem, K. Aftab, F. Shaheen and A.H. Gilani. 1998. Bioactive compounds from the leaves and pods of *Moringa oleifera*. *New Trends in Natural Products Chem.* 175-183.
- Faizi, S., B.S. Siddiqui, R. Saleem, S. Siddiqui, K. Aftab, F. Shaheen and A.H. Gilani. 1994. Isolation and structure elucidation of new nitrile and mustard oil glycosides from *Moringa oleifera* and their effect on blood pressure. *J. Nat. Prod.* 57:1256-1261.
- Fakurazi, S., I. Hairuszah and U. Nanthini. 2008. *Moringa oleifera* Lam. prevents acetaminophen induced liver injury through restoration of glutathione level. *Food Chem. Toxicol.* 46:2611-2615.
- Ferreira, P.M.P., A.F.F.U. Carvalho, D.F. Farias, N.G. Cariolano, V.M.M. Melo, M.G.R. Queiroz, A.M.C. Martins and J.G. Machado-Neto. 2009. Larvicidal activity of the water extract of *Moringa oleifera* seeds against *Aedes aegypti* and its toxicity upon laboratory animals. *An. Acad. Bras. Cienc.* 81:207-216.
- Gaind, K.N., H.C. Mital and S.R. Khanna. 1964. Anthelmintic activity of Triphala. *Indian J. Pharmacol.* 26:106-107.
- Garg, S.C. and H.L. Kasera. 1982a. Anthelmintic activity of *Callistemon viminalis*. *Fitoterapia*. 53:179-181.
- Garg, S.C. and H.L. Kasera. 1982b. *In vitro* anthelmintic activity of the essential oil of *Anacardium occidentale*. *Indian Perfumer*. 26:239-240.
- Garg, S.C. and S. Nakhare. 1993. Studies on the essential oils from the flowers of *Eupatorium triplinerve*. *Indian Perfumer*. 37:318-323.
- Gbeassor, M., A.Y. Kedjagni, K. Koumaglo, C. de Souza, K. Agbo, K. Aklikokou and K. A. Amegbo. 1990. *In vitro* antimalarial activity of six medicinal plants. *Phytotherapy Res.* 4:(3)115-117
- Geervani, P. and A. Devi. 1981. Influence of protein and fat on the utilization of carotene from drumstick (*Moringa oleifera*) leaves. *Indian J. Med. Res.* 74: 548-553.
- Ghasi, S., E. Nwobodo and J.O. Ofili. 2000. Hypocholesterolemic effects of crude extract of leaf of *Moringa oleifera* Lam. in high-fat diet fed wistar rats. *J. Ethnopharmacol.* 69:21-25.
- Gilani, A.H., S. Bashir and A. Khan. 2007. Pharmacological basis for the use of *Borago officinalis* in gastrointestinal, respiratory and cardiovascular disorders. *J. Ethnopharmacol.* 114(3):393-399.
- Girgune, J.B., N.K. Jain and B.D. Garg. 1978. Anthelmintic activity of some essential oils. *Indian Perfumer*. 22:296-297.
- Girgune, J.B., N.K. Jain and B.D. Garg. 1979. Antimicrobial and anthelmintic activity of essential oil from *Gardenia lucida* Roxb. *Indian Perfumer*. 23:213-215.
- Githiori, J.B., J. Hogland, P.J. Waller and R.L. Baker. 2004. Evaluation of anthelmintic properties of some plants used as livestock dewormers against *Haemonchus contortus* infection in sheep. *Parasitol.* 129:245-253.
- Guarrera, P.M. 1999. Traditional antihelminthic, antiparasitic and repellent uses of plants in Central Italy. *J. Ethnopharmacol.* 68:183-192.
- Guevara, A.P., C. Vargas, H. Sakurai, Y. Fujiwara, K. Hashimoto, T. Maoka, M. Kozuka, Y. Ito, H. Tokuda and H. Nishino. 1999. An antitumor promoter from *Moringa oleifera* Lam. *Mutat Res.* 440:181-188.
- Guha, S.R.D., S.N. Dhoundiyal and G.M. Mathur. 1968. Mechanical pulps for newsprint grade papers from *Moringa pterygosperma*. *Indian Forester*. 94(8):634-638.
- Hamad, K.K., Z. Iqbal, Z.U.D. Sindhu and G. Muhammad. 2013. Antinematicidal activity of *Nicotiana tabacum* L. leaf extracts to control benzimidazole-resistant *Haemonchus contortus* in sheep. *Pak. Vet. J.* 33: 85-90.
- Hammond, J.A., D. Fielding and S.C. Bishop. 1997. Prospects for plant anthelmintics in tropical veterinary medicine. *Vet. Res. Commun.* 21:213-228.

- Hamza, A.A. 2007. *Curcuma longa*, *Glycyrrhiza glabra* and *Moringa oleifera* ameliorate diclofenac-induced hepatotoxicity in rats. Am. J. Pharmacol. Toxicol. 2:80-88.
- Hamza, A.A. 2010. Ameliorative effects of *Moringa oleifera* Lam. seed extract on liver fibrosis in rats. Food Chem. Toxicol. 48:345-355.
- Huff, T. 1996. The rise of early modern science: Islam, China, and the West. Cambridge University Press. p. 218.
- Hukkeri, V.I., C.V. Nagathan, R.V. Karadi and B.S.M. Patil. 2006. Antipyretic and wound healing activities of *Moringa oleifera* Lam. in rats. Indian J. Pharm. Sci. 68:124-126.
- Hussain, A., M.N. Khan, M.S. Sajid, Z. Iqbal, M.K. Khan, R.Z. Abbas, M.A. Raza and G.R. Needham. 2010. *In Vitro* Screening of the Leaves of *Musca paradisiaca* for anthelmintic activity. J. Anim. Plant Sci. 20:5-8.
- Hussain, A., M.N. Khan, Z. Iqbal, M.S. Sajid and M.K. Khan. 2011. Anthelmintic activity of *Trianthema portulacastrum* L. and *Musa paradisiaca* L. against gastrointestinal nematodes of sheep. Vet. Parasitol. 179:92-99.
- Ibrahim, M.A., N. Nwude, R.A. Ogunsusi and Y.O. Aliu. 1984. Screening of the West African plants for antihelmintic activities. ILCA Bulletin. 17:19-23.
- <http://www.edlagman.com/moringa/moringa-fresh-leaf-vs-dried-leaf.pdf> (accessed on January 28, 2014)
- Iqbal, Z., M. Akhtar, M.N. Khan and M. Riaz. 1993. Prevalence and economic significance of haemonchosis in sheep and goats slaughtered at Faisalabad abattoir. Pak. J. Agri. Sci. 30:51-53.
- Iqbal, Z., M. Lateef, A. Jabbar, G. Muhammad and M.N. Khan. 2005. Anthelmintic activity of *Calotropis procera* (Ait.) Ait. F. flowers in sheep. J. Ethnopharmacol. 102:256-261.
- Iqbal, Z., M. Lateef, A. Jabbar, M.N. Ghayur and A.H. Gilani. 2006. *In vivo* anthelmintic activity of *Butea monosperma* against Trichostrongylid nematodes in sheep. Fitoterapia. 77:137-140.
- Iqbal, Z., M. Lateef, M. Ashraf and A. Jabbar. 2004. Anthelmintic activity of *Artemisia brevifolia* in sheep. J. Ethnopharmacol. 93:265-268.
- Iqbal, Z., Q.K. Nadeem, M.N. Khan, M.S. Akhtar and F.N. Waraich. 2001b. *In vitro* anthelmintic activity of *Allium sativum*, *Zingiber officinale*, *Curcurbita mexicana* and *Ficus religiosa*. Int. J. Agr. Biol. 3:454-457.
- Iqbal, Z., W. Babar, Z.U.D. Sindhu, R.Z. Abbas and M.S. Sajid. 2012. Evaluation of anthelmintic activity of different fractions of *Azadirachta indica* A. Juss seed extract. Pak. Vet. J. 32:579-583.
- Jadhav, S.L., S.R. Sharma, S.C. Pal, S.B. Kasture and V.S. Kasture. 2000. Chemistry and pharmacology of *Moringa oleifera* and *Moringa concanescens* Niro. Indian Drugs. 37(3):139-144.
- Jaiswal, D., R.P. Kumar, A. Kumar, S. Mehta and G. Watal. 2009. Effect of *Moringa oleifera* Lam. leaves aqueous extract therapy on hyperglycemic rats. J. Ethnopharmacol. 123:392-396.
- Kailani, S.U.R., M.S. Akhtar and M. Ashraf. 1995. Antifasciolic efficacy of indigenous plant drugs: Kalonji, Shahterah and Karanjwa in buffaloes. Pak. J. Pharm. Sci. 8:17-27.
- Kakrani, H.K. and G.A. Kalyani. 1984. Anthelmintic activity of the essential oil of *Commiphora mukul*. Fitoterapia. 55:232-234.
- Kalesaraj, R. 1975. Screening of some indigenous plants for anthelmintic action against human *Ascaris lumbricoides*. Indian J. Physiol. Pharmacol. 19:47-49.
- Kalyani, G.A., K.S. Aithal and K.K. Srivastava. 1989. *In vitro* anthelmintic activity of essential oil from the fruits of *Zanthoxylum limonella*. Fitoterapia. 2:160-162.
- Kar, P.K., V. Tandon and N. Saha. 2002. Anthelmintic efficacy of *Flemingia vestita*: genistein-induced effect on the activity of nitric oxide synthase and nitric oxide in the trematode parasite, *Fasciolopsis buski*. Parasitol. Int. 51:249-257.
- Kasolo, J. 2010. Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities. J. Med. Plant Res. 4(9):753-775.
- Katre, U.V., C.G. Suresh, M.I. Khan and S.M. Gaikwad. 2008. Structure-activity relationship of a hemagglutinin from *Moringa oleifera* seeds. Int. J. Biol. Macromol. 42:203-207.
- Kaushik, R.K., J.C. Katiyar and A.B. Sen. 1981. A new *in vitro* screening technique for anthelmintic activity using *Ascaridia galli* as a test parasite. Indian J. Anim. Sci. 51:869-872.
- Kerharo, P.J. 1969. Un remede populaire Sengalais: Le 'Nebreday' (*Moringa oleifera* Lam.) emploie therapeutiques en milieu Africain chimie et pharmacologie. Plantes Med. Phytother. 3:14-21.
- Kochapakdee, S., W. Pralomkarn, S. Choldumrongku and S. Saithanoo. 1995. Change in live-weight gain, blood constituents and worm egg counts in Thai native and crossbred goats raised in village environments in southern Thailand. Asian-Aust. J. Anim. Sci. 8:241-247.
- Kohler, I., K. Jenett-Siems, K. Siems, M.A. Hernandez, R.A. Ibarra, W.G. Berendsohn, U. Bienzle and E. Eich. 2002. *In vitro* antiplasmodial investigation of medicinal plants from El Salvador. Z Naturforsch C. 57(3-4):277-281.
- Kokane, C.K. and K.C. Varma. 1971. Anthelmintic activity of some essential oils. Indian J. Hosp. Pharm. 8:150-151.
- Krause, H.D. 1993. Treatment of helminth infections with Abrotanum. Biologische Tiermedizin. 10:13-19.
- Kurup, P.A. and P.I. Rao. 1954. Antibiotic principle from *Moringa pterygosperma* IV. The effect of addition of vitamins and amino acids on the anti-bacterial activity. Indian J. Med. Res. 42:101-107.
- Kushwaha, V., K. Saxena, S.K. Verma, V. Lakshmi, R.K. Sharma and P.K. Murthy, 2011. Antifilarial activity of gum from *Moringa oleifera* Lam. On human lymphatic

- filaria Brugia malayi. Chron. Young Scientists 2(4):201-206.
- Lako, J., V.C. Trencerry, M. Wahlqvist, N. Wattanapenpaiboon, S. Sotheeswaran and R. Premier. 2007. Phytochemical flavonols, carotenoids and the antioxidant properties of a wide selection of Fijian fruit, vegetables and other readily available foods. Food Chem. 101:1727-1741.
- Lakshminarayana, R., M. Raju, T.P. Krishnakantha and V. Baskaran. 2005. Determination of major carotenoids in a few Indian leafy vegetables by highperformance liquid chromatography. J. Agric. Food Chem. 53:2838-2842.
- Lal, J., S. Chandra, V.R. Prakash and M. Sabir. 1976. *In vitro* anthelmintic action of some indigenous medicinal plants on *Ascaridia galli* worms. Indian J. Physiol. Pharmacol. 20:64-68.
- Lalas, S. and J. Tsaknis. 2002. Characterization of *Moringa oleifera* seed oil variety "Periyakulum 1". J. Food Compost. Anal. 15:65-77.
- Lateef, M., Z. Iqbal, M.N. Khan, M.S. Akhtar and A. Jabbar. 2003. Anthelmintic activity of *Adhatoda vesica* roots. Int. J. Agri. Biol. 5:86-90.
- Loudon, I., 2002. Western medicine: An illustrated history. Oxford University Press. p.54.
- Lürling, M. and W. Beekman. 2010. Anti-cyanobacterial activity of *Moringa oleifera* seeds. J. Appl. Phycol. 22(4):503-510.
- Mabberley, D.J. 1997. The plant book. 2nd ed., Cambridge Univ. Press, Cambridge, UK. p.467.
- Mahajan, S.G. and A.A. Mehta. 2007. Inhibitory action of ethanolic extract of seeds of *Moringa oleifera* Lam. on systemic and local anaphylaxis. J. Immunotoxicol. 4:287-294.
- Mahajan, S.G., A. Banerjee, B.F. Chauhan, H. Padh, M. Nivsarkar and A.A. Mehta. 2009. Inhibitory effect of n-butanol fraction of *Moringa oleifera* Lam. seeds on ovalbumin-induced airway inflammation in a guinea pig model of asthma. Int. J. Toxicol. 28:519-527.
- Mahajan, S.G., R.G. Mali and A.A. Mehta. 2007. Effect of *Moringa oleifera* Lam. Seed extract on toluene di isocyanate-induced immune-mediated inflammatory responses in rats. J. Immunotoxicol. 4:85-96.
- Mahajan, S.G., R.G. Mali and A.A. Mehta. 2007. Protective effect of ethanolic extract of seeds of *Moringa oleifera* Lam. against inflammation associated with development of arthritis in rats. J. Immunotoxicol. 4:39-47.
- Mahatab, S.N., A. Ali and A.H.M. Asaduzzaman. 1987. Nutritional potential of sajna leaves in goats. Livestock Advisor. 12(12):9-12.
- Makkar, H.P.S. and K. Becker. 1996. Nutritional value and antinutritional components of whole and ethanol extracted *Moringa oleifera* leaves. Anim. Feed Sci. Technol. 63:211-228.
- Mathur, B. 2006. Moringa for cattle fodder and plant growth. President, Trees for life. www.treesforlife.org.
- Mehta, K., R. Balaraman, A.H. Amin, P.A. Bafna and O.D. Gulati. 2003. Effect of fruits of *Moringa oleifera* on the lipid profile of normal and hypercholesterolaemic rabbits. J. Ethnopharmacol. 86:191-195.
- Mehta, R.K. and G.C. Parashar. 1966. Effect of *Butea frondosa*, *Vernonia anthelmintica*, and *Carica Papaya* against Oxyurids in mice. Indian Vet. J. 43:73-78.
- Minja, M.M.J. 1989. Collection of Tanzanian medicinal plants for biological activity studies. In: Proc. 7th Tanzania Vet. Assoc. Sci. Conf., Arusha. 7:67-78.
- Mishra, S.H., R.S. Gaud, R.A. Sharma and S.C. Chaturvedi. 1979. Anthelmintic activity of some essential oils. Indian Perfumer. 23:208-209.
- Morton, J.F. 1991. The horseradish tree, *Moringa pterygosperma* (Moringaceae). A boon to Arid lands? Econ. Bot. 45:318-333.
- Mughal, M.H., G. Ali, P.S. Srivastava and M. Iqbal. 1999. Improvement of drumstick (*Moringa pterygosperma* Gaertn.) - A unique source of food and medicine through tissue culture. Hamdard Med. 42:37-42.
- Muyibi, S.A., M.J.M.M. Noor, D.T. Ong and K.W. Kai. 2001. *Moringa oleifera* seeds as a flocculant in waste sludge treatment. Int. J. Envir. Stud. 58:185-189.
- Nagar, P.K., R.I. Iyer and P.K. Sircar. 1981. Auxins in *Moringa pterygosperma* Gaertn fruits. Indian J. Exp. Biol. 19:487.
- Nagar, P.K., R.I. Iyer and P.K. Sircar. 1982. Cytokinins in developing fruits of *Moringa pterygosperma* Gaertn. Physiol. Plant. 55:45-50.
- Nakhare, S. and S.C. Garg. 1991. Anthelmintic activity of some essential of *Artemisia pallens* Wall. Ancient Sci. Life. 10:185-186.
- Nanda, A., M.A. Iyengar, C.S. Narayan and D.R. Kalkarni. 1987. Investigations on the root bark of *Aglai odoratissima*. Fitoterapia. 58:189-191.
- Naqvi, S.A.H., S.B. Vohora and M.S.Y. Khan. 1991. Antibacterial, antifungal and anthelmintic studies of *Artemisia scoparia*. Herba Hungarica. 30:54-60.
- Ndong, M., M. Uehara, S. Katsumata and K. Suzuki. 2007. Effects of oral administration of *Moringa oleifera* Lam. on glucose tolerance in goto-kakizaki and wistar rats. J. Clin. Biochem. Nutr. 40:229-233.
- Neogi, N.C., P.A.C. Baliga and R.K. Srivastava. 1964. Anthelmintic activity of some indigenous drugs. Indian J. Pharmacol. 26:37.
- Nikkon, F., Z.A. Saud, M.H. Rehman and M.E. Haque. 2003. *In vitro* antimicrobial activity of the compound isolated from chloroform extract of *Moringa oleifera* Lam. Pak. J. Biol. Sci. 22:1888-1890.
- Nwude, N. and M.A. Ibrahim. 1980. Plants used in traditional veterinary medical practice in Nigeria. J. Vet. Pharmacol. Therapeut. 3:261-273.
- Ogunbinu, A.O., G. Flamini, P.L. Cioni, M.O. Adebayo and I.A. Ogunwande. 2009. Constituents of *Cajanus cajan* (L.) Millsp, *Moringa oleifera* Lam., *Heliotropium indicum* L. and *Bidens pilosa* L. from Nigeria. Nat. Prod. Commun. 4:573-578.

- Olson, M.E. 2002. Combining data from DNA sequence and morphology for a phylogeny of Moringaceae (Brassicales). *Syst. Bot.* 27:55-73.
- Padmaja, V., V. Thankamany and A. Hisham. 1993. Antibacterial, antifungal and anthelmintic activities of root barks of *Uvaria hookeri* and *Uvaria narum*. *J. Ethnopharmacol.* 40:181-186.
- Pal, P. and V. Tandon. 1998. Anthelmintic efficacy of *Flemingia vestita* Leguminosae: Genisteininduced alterations in the activity of tegumental enzymes in the cestode, *Raillietina echinobothrida*. *Parasitol. Int.* 47:233-243.
- Pal, S.K., P.K. Mukherjee and B.P. Saha. 1995. Studies on the antiulcer activity of *Moringa oleifera* leaf extract on gastric ulcer models in rats. *Phytotherapy Research.* 9:463-465.
- Pari, L. and N.A. Kumar. 2002. Hepatoprotective activity of *Moringa oleifera* on antitubercular drug-induced liver damage in rats. *J. Med. Food.* 5:171-177.
- Parker, R.J. and B. Palmer. 1991. Lack of anthelmintic effect of *Calliandra calothrysus* in sheep. *Aust. Vet. J.* 68:309.
- Parrotta, J.A. 2001. In: Healing plants of peninsular India 2001; CABI publication, New York, p. 528-530.
- Perrett, S. and P.J. Whitfield. 1995. Atanine (3-Dimethylallyl-4-methoxy-2- quinolone), an alkaloid with anthelmintic activity from the Chinese medicinal plant, *Evodia rutaecarpa*. *Planta Med.* 61:276-278.
- Prakash, V., K.C. Singhal and R.R. Gupta. 1980. Anthelmintic activity of *Punica granatum* and *Artemisia silversiana*. *Indian J. Pharmacol.* 12:62.
- Qaiser, M. 1973. Moringaceae. In flora of West Pakistan, Nasir, E. and S.I. Ali (Editors), University of Karachi Press, Karachi. p. 1-4.
- Qureshi, M.A. and M. Sabir. 1979. Preliminary study on anthelmintic efficacy of Embellia seeds (Babarung) against tapeworms of poultry. *Pak. J. Sci.* 31:218-220.
- Rajan, B.K.C. 1986. Agriculture and farm forestry in semi-arid tracts of Karnataka. *My Forest.* 22(1):41-49.
- Ramachandran, C., K.V. Peter and P.K. Gopalakrishnan. 1980. Drumstick (*Moringa oleifera*): a multipurpose Indian vegetable. *Econ. Bot.* 34:276-283.
- Rao, A.V., P.U. Devi and R. Kamath. 2001. *In vivo* radio protective effect of *Moringa oleifera* leaves. *Indian J. Exp. Biol.* 39:858-863.
- Rao, K.S. and S.H. Mishra. 1998. Anti-inflammatory and anti-hepatotoxic activities of the roots of *Moringa pterygosperma* Gaertn. *Indian J. Pharmaceut. Sci.* 60:12-16.
- Rastogi, T., V. Bhutda, K. Moon, P.B. Aswar and S.S. Khadabadi. 2009. Comparative Studies on Anthelmintic Activity of *Moringa Oleifera* and *Vitex Negundo*. *Asian J. Res. Chem.* 2:181-182.
- Rathi, B.S., S.L. Bodhankar and A.M. Baheti. 2006. Evaluation of aqueous leaves extract of *Moringa oleifera* Linn. for wound healing in albino rats. *Indian J. Exp. Biol.* 44:898-901.
- Rebrassier, R.E. 1934. Pyrethrum as an anthelmintic for *Ascaridia lineata*. *J. Am. Vet. Med. Assoc.* 84:645-648.
- Reddy, G.B.S., A.B. Melkhani, G.A. Kalyani, R.J. Venkata, A. Shirwaikar, M. Kotian, R. Ramani, K.S. Aithal, A.L. Udupa, G. Bhat and K.K. Srinivasan. 1991. Chemical and pharmacological investigations of *Limnophila conferta* and *L. heterophylla*. *Int. J. Pharmacogn.* 29:145-153.
- Roy, S.K., K. Chandra, K. Ghosh, S. Mondal, D. Maiti, A.K. Ojha, D. Das, S. Mondal, I. Chakraborty and S.S. Islam. 2007. Structural investigation of a heteropolysaccharide isolated from the pods (fruits) of *Moringa oleifera* (Sajina). *Carbohydr. Res.* 342: 2380-2389.
- Ruckmani, K., S. Davimani, B. Jayakar and R. Anandan. 1998. Anti-ulcer activity of the alkali preparation of the root and fresh leaf juice of *Moringa oleifera* Lam. *Ancient Sci. Life.* 17:220-223.
- Ruckmani, K., S. Kavimani, R. Anandan and B. Jaykar. 1998. Effect of *Moringa oleifera* Lam. on paracetamol-Induced Hepatotoxicity. *Indian J. Pharm. Sci.* 60:33-35.
- Saluja, M.P., R.S. Kapil and S.P. Popli. 1978. Studies on medicinal plants: part VI. Chemical constituents of *Moringa oleifera* Lamk. (hybrid variety) and isolation of 4-hydroxymellein. *Indian J. Chem. B.* 16:1044-1045.
- Sashidhara, K.V., J.N. Rosaiah, E. Tyagi, R. Shukla, R. Raghbir and S.M. Rajendran. 2009. Rare dipeptide and urea derivatives from roots of *M. oleifera* as potential anti-inflammatory and anti-nociceptive agents. *Eur. J. Med. Chem.* 44:432-436.
- Sastri, B.N. 1962. The Wealth of India: Council of Scientific and Industrial Research, New Delhi. p.425.
- Satyavati, G.V., M.K. Raina and M. Sharma. 1976. Medicinal Plants of India. Indian Council of Medical Research, New Delhi, 1:201-206.
- Selvakumar, D. and P. Natarajan. 2008. Hepatoprotective activity of *Moringa oleifera* Lam. leaves in carbon tetrachloride induced hepato-toxicity in albino rats. *Pharmacog. Mag.* 4:97-98.
- Sen, H.G., B.S. Joshi, P.C. Parthasarathy and V.N. Kamat. 1974. Anthelmintic efficacy of *Diosptrol* and its derivatives. *Arzneimittelforschung.* 24:2000-2003.
- Seshadri, S. and V.S. Nambiar. 2003. Kanjero (*Digera arvensis*) and drumstick leaves (*Moringa oleifera*): nutrient profile and potential for human consumption. *World Rev. Nutr. Diet.* 91:41-59.
- Shahzad, U., M.A. Khan, M.J. Jaskani, I.A. Khan and S.S. Korban. 2013. Genetic diversity and population structure of *Moringa oleifera*. *Conserv. Genet.* 14: 1161-1172.
- Shanker, K., M.M. Gupta, S.K. Srivastava, D.U. Bawankule, A. Pal, and S.P.S. Khanuja. 2007. Determination of bioactive nitrile glycoside(s) in drumstick (*Moringa oleifera*) by reverse phase HPLC. *Food Chem.* 105:376-382.
- Sharma, G.P., N.K. Jain and B.D. Garg. 1979. Anthelmintic activity of some essential oils. *Indian Perfumer.* 23:210-212.

- Sharma, L.D., H.S. Bagha and P.S. Srivastava. 1971. *In vitro* anthelmintic screening of indigenous medicinal plants against *Haemonchus contortus* (Rudolphi, 1803) Cobbold, 1898 of sheep and goats. Indian J. Anim. Res. 5:33-38.
- Sharma, P.K. and V. Singh. 1989. Ethnobotanical studies in northwest and Trans-Himalaya. Ethno veterinary medicinal plants used in Jammu and Kashmir. Indian J. Ethnopharmacol. 27:63-70.
- Sherif, A., R.G. Hall and M. el-Amamy. 1987. Drugs, insecticides and other agents from *Artemisia*. Med. Hypotheses. 23:187-193.
- Shilaskar, D.V. and G.C. Parashar. 1989. Evaluation of indigenous anthelmintics: *In vitro* screening of some indigenous plants for their anthelmintic activity against *Ascaridia galli*. Indian J. Indg. Med. 6:49-53.
- Shiramizu, k., T. Tuchida and M. Aru. 1993. Anthelmintic effects of a crude preparation containing the root of *Matteuccia orientalis* on bovine Fasciolosis. J. Japan Vet. Med. Assoc. 46:561-564.
- Shrivastava, M.C. and S.W. Singh. 1967. Anthelmintic activity of *Cucurbita maxima* seeds. Indian J. Med. Res. 55:629-632, 746-748.
- Shrivastava, R. 1979. Anthelmintic properties of essential oil of *Cyathocline lyrata* Cass. Indian J. Pharm. Sci. 41:228-229.
- Shuhua, X., J. Chollet, N.A. Weiss, R.N. Bergquist and M. Tanner. 2000. Preventive effect of artemether in experimental animals infected with *Schistosoma mansoni*. Parasitol. Int. 49:19-24.
- Shukla S., R. Mathur and A.O. Prakash. 1988. Antifertility profile of the aqueous extract of *Moringa oleifera* roots. J Ethnopharmacol. 22:51-62.
- Siddiqui, N. and S.C. Garg. 1990. *In vitro* anthelmintic activity of some essential oils. Pak. J. Sci. Indust. Res. 33:536-537.
- Silja, V.P., K.S. Varma and K.V. Mohanan. 2008. Ethnomedicinal plant knowledge of the Mullu kuruma tribe of Wayanad district, Kerala. Indian J. Trad. Know. 7:604-612.
- Sindhu, Z.D., Z. Iqbal, M.N. Khan, N.N. Jonsson and M. Siddique. 2010. Documentation of ethnoveterinary practices used for treatment of different ailments in a selected hilly area of Pakistan. Int. J. Agri. Biol. 12:353-358.
- Singh, R.K., A. Singh and A.K. Sureja. 2007. Sustainable use of ethnobotanical resources. Indian J. Trad. Know. 6:521-530.
- Singh, S., M.C. Srivastava and J.P. Tewari. 1982. Anthelmintic activity of *Zanthoxylum alatum* (Tejbal). Indian Drugs. 19:348-353.
- Sofowora, A. 1982. Medicinal plants and traditional medicine in Africa. John Wiley and Sons Ltd, New York, p. 214-218.
- Solecki and S. Ralph. 1975. Shanidar IV, a Neanderthal Flower Burial in Northern Iraq. Sci. 190:880-881.
- Somali, M.A., M.A. Bajnedi and S.S. Al-Faimani. 1984. Chemical composition and characteristics of *Moringa peregrina* seeds and seed oil. J. Amer. Oil Chem. Soc. 61: 85-86.
- Stephenson, K.K. and J.W. Fahey. 2004. Development of tissue culture methods for the rescue and propagation of endangered *Moringa* spp. germplasm. Econ. Bot. 58:16-124
- Sumner, J. 2000. The natural history of medicinal plants. Timber Press, p. 17-18.
- Sutar, N., C.G. Bonde, V.V. Patil, S.B. Narkhede, A.P. Patil and R.T. Kakade. 2008. Analgesic activity of seeds of *Moringa oleifera* Lam. Int. J. Green Pharm. 2:108-110.
- Sutar, N.G., V.V. Patil, T.A. Deshmukh, N.M. Jawle, V.R. Patil and S.C. Bhangale. 2009. Evaluation of anti-pyretic potential of seeds of *Moringa oleifera* Lam. Int. J. Green Pharm. 3:148-150.
- Sykes, A.R. 1994. Parasitism and production in farm ruminants. Anim. Prod. 59:155-172.
- Tahiliani, P. and A. Kar. 2000. Role of *Moringa oleifera* leaf extract in the regulation of thyroid hormone status in adult male and female rats. Pharmacol. Res. 41:319-323.
- Trapti, R., V. Bhutda, K. Moon, P.B. Aswar, and S.S. Khadabadi. 2009. Comparative Studies on Anthelmintic Activity of *Moringa Oleifera* and *Vitex Negundo*. Asian J. Res. Chem. 2(2):181-182.
- Tsaknis, J., S. Lalas, V. Gergis, V. Dourtoglou and V. Spiliotis. 1999. Characterization of *Moringa oleifera* variety Mbolo seed oil of Kenya. J. Agric. Food Chem. 47:4495-4499.
- Verma, A.R., M. Vijaykumar, C.S. Mathela and C.V. Rao. 2009. *In vitro* and *in vivo* antioxidant properties of different fractions of *Moringa oleifera* leaves. Food Chem. Toxicol. 47:2196-2201.
- Verma, S.C., R. Banerji, G. Misra and S.K. Nigam. 1976. Nutritional value of Moringa. Current Sci. 45:769-770.
- Xiao, P.G. and F.S. Lin. 1986. Traditional anti-parasitic drugs in China. Parasitol. Today. 2:353-355.
- Zaman, M.A., Z. Iqbal, M.N. Khan and G. Muhammad. 2012. Anthelmintic activity of a herbal formulation against gastrointestinal nematodes of sheep. Pak. Vet. J. 32: 117-121.
- Zlatkis, A., B. Zak, G.J. Boyle. 1953. A new method for direct determination of serum cholesterol. J. Lab. Clin. Med. 41:486.