

DTIC  
ELECTE  
MAR 4 1986  
S A D

2

# Pineal-Induced Depression of Free Thyroxine in Syrian Hamsters

George M. Vaughan and Basil A. Pruitt, Jr.

U.S. Army Institute of Surgical Research, Brooke Army Medical Center,  
Fort Sam Houston, Texas

AD-A165 036

Blind hamsters had no alteration of the dialyzable fraction of serum thyroxine (T4) but had depressed total and free T4 concentrations compared to controls. Prevention of the effects of blinding by pinealectomy indicates pineal influence on circulating free T4 concentration. Parallel changes in free T4 and the free T4 index indicate adequacy of the index in representing pineal-induced changes in free T4.

Key words: pineal, free thyroxine, thyroxine, hamsters, blinding

DTIC FILE COPY

## INTRODUCTION

When Syrian hamsters are exposed to a short photoperiod (less than 12 hr light per day) for 2-3 months or are subjected to bilateral orbital enucleation, their circulating total thyroxine (T4) concentration falls. This effect is brought about by the pineal gland; pinealectomy prevents it [Vriend et al., 1979; Vriend, 1983]. In the above and other studies [G.M. Vaughan et al., 1982; M.K. Vaughan et al., 1982] of light restriction, the pineal, and T4 in this species the status of circulating free T4 has been approached by

Received March 20, 1985; accepted May 20, 1985.

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. In conducting the research described in this report the investigators adhered to the Guide for Laboratory Animal Facilities and Care, as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences, National Research Council.

Address reprint requests to Library Branch, U.S. Army Institute of Surgical Research, Fort Sam Houston, TX 78234-6200.

Reproduced from best available copy.

© 1985 ALAN R. LISS, INC. I 86 3 3 139

simultaneous observations of the free T<sub>4</sub> index (FT<sub>4</sub>I), which depends on the in vitro uptake (T<sub>3</sub>U) of added radiotracer triiodothyronine (T<sub>3</sub>) from the serum onto a solid matrix. The FT<sub>4</sub>I ( $T_4 \times T_3U$ ) thus contained a correction for possible alterations of protein binding of T<sub>4</sub> and provided an indirect assessment of its free concentration. Because the FT<sub>4</sub>I changed with the same pattern exhibited by T<sub>4</sub> in the above studies, it was tentatively concluded that the pineal gland, in the condition of light deprivation, is capable of influencing circulating free T<sub>4</sub> concentration (FT<sub>4</sub>).

However, the general use of the FT<sub>4</sub>I to represent changes of FT<sub>4</sub> has recently been called into question. For example, in a number of conditions and species, including various nonthyroidal illnesses in humans [Chopra et al., 1979, 1984; Woeber and Maddux, 1981; Oppenheimer et al., 1982] and burn injury in rats [Shirani et al., 1985], human [Vaughan et al., 1985a], and hamsters [Vaughan et al., 1985b], the total T<sub>4</sub> and FT<sub>4</sub>I were often depressed but the free or dialyzable fraction of T<sub>4</sub> (T<sub>4</sub>DF) was elevated out of proportion to any change in T<sub>3</sub>U. This often resulted in free T<sub>4</sub> concentrations ( $FT_4 = T_4 \times T_4DF$ ) that were not depressed or were less depressed than predicted by the FT<sub>4</sub>I in these conditions. The T<sub>4</sub>DF is that portion of radiotracer T<sub>4</sub> that crosses a protein-impermeable membrane from the serum into buffer without involvement of a solid matrix. Thus such discrepancy between the T<sub>3</sub>U and the T<sub>4</sub>DF (and hence between the FT<sub>4</sub>I and FT<sub>4</sub>) might be due to a circulating factor in some conditions that not only inhibits binding of T<sub>4</sub> to serum proteins but also to the solid matrix of the T<sub>3</sub>U test [Oppenheimer et al., 1982]. It is thus necessary to show that, in the experimental condition under study, the changes in FT<sub>4</sub>I do reflect those of FT<sub>4</sub> directly assessed by observations of the T<sub>4</sub>DF.

Because of these considerations, in the present study we have assessed whether or not the pineal gland, in the condition of its activation by blindness in Syrian hamsters, can influence free T<sub>4</sub> concentration.

## MATERIALS AND METHODS

Male golden hamsters, *Mesocricetus auratus*, were purchased from Charles River Breeding Laboratories, Inc. (Wilmington, Massachusetts), at about 90 gm body weight (young adults), maintained in our animal quarters in a light/dark (L/D) cycle of 14/10 hr at 21°C with lights on at 0700 hr and given standard laboratory chow and tap water ad libitum. After 2 weeks, they were divided into three groups, according to surgical procedures performed under pentobarbital anesthesia: sham-pinealectomy (SHPX), blinding by bilateral orbital enucleation together with sham-pinealectomy (BL + SHPX), and blinding plus pinealectomy (BL + PX). Pinealectomy was performed according to the method of Hoffman and Reiter [1965], and the sham procedure involved replacing the disc of skull bone without performing pinealectomy. The animals were maintained in the L/D 14/10 cycle until sacrifice by guillotine 11 weeks after surgery, when trunk serum was obtained and both testes and the dorsal prostate were cleaned of fat and connective tissue and weighed. Successful pinealectomy without disruption of the brain

structures in the vicinity was verified grossly at autopsy. T<sub>4</sub> was measured by radioimmunoassay and T<sub>3</sub>U by radioassay (charcoal matrix) with kits from Diagnostic Products (Los Angeles, California). The dialyzable fraction of T<sub>4</sub> (T<sub>4</sub>DF) was determined in duplicate serum aliquots at the Nichols Institute, San Juan Capistrano, California, by equilibrium dialysis at 37°C [Sterling and Brenner, 1966]. The FT<sub>4</sub>I was calculated as the product of T<sub>4</sub> and T<sub>3</sub>U, and the FT<sub>4</sub> as the product of T<sub>4</sub> and T<sub>4</sub>DF (Sterling and Brenner, 1966). Data were analysed by the Student-Newman-Keuls test [Bruning and Kintz, 1977] and analysis of covariance [Dixon, 1983].

## RESULTS


Reduction of testicular and prostatic mass was observed in the blind animals, an effect obliterated by additional pinealectomy (Fig. 1). Blinding produced no detectable effect on T<sub>3</sub>U or T<sub>4</sub>DF, though a small reduction of mean T<sub>3</sub>U was seen in the BL+PX group (Table 1). T<sub>4</sub>, FT<sub>4</sub>I, and FT<sub>4</sub> were all depressed after blinding, unless the animals were additionally pinealectomized (Fig. 1). The correlation of FT<sub>4</sub>I with FT<sub>4</sub> was significant (Fig. 1), and there were no significant differences among slopes for individual groups (not shown).

## DISCUSSION

Compared to values in controls, lower mean serum T<sub>4</sub> without major alteration of T<sub>3</sub>U has been reported in blind humans [Dieckhues, 1974], an effect mediated by the pineal gland in Syrian hamsters [Vriend et al., 1977; Vriend et al., 1979; G.M. Vaughan, et al. 1982; M.K. Vaughan, et al., 1982; Vriend, 1983]. We now report that free T<sub>4</sub> concentration is depressed by blinding and that this effect appears to be mediated by the pineal gland. The small reduction of mean T<sub>3</sub>U in BL+PX is not explained, but blinding alone appeared to have no effect on T<sub>3</sub>U, and FT<sub>4</sub>I appeared to be a relatively reliable index of FT<sub>4</sub> (Fig. 1).

The observed blinding-induced suppression of the reproductive system and the prevention of this by pinealectomy serve to indicate that the animals in this study exhibited the expected pineal-mediated reproductive responses [Reiter, 1980] and thus adequately represent the usual hamster model for demonstrating pineal function. The parallel changes between the reproductive system and circulating T<sub>4</sub> have been noted previously, and the possibility has been considered that the changes in T<sub>4</sub> passively result from pineal-induced hypogonadism and the possible resultant effects on T<sub>4</sub> serum binding proteins [Vriend et al., 1977, 1979; G.M. Vaughan et al., 1982]. It was interpreted that the effects of the pineal on T<sub>4</sub> and free T<sub>4</sub> were likely independent of such a potential effect of hypogonadism, because the FT<sub>4</sub>I varied with the same pattern as did total T<sub>4</sub>. This interpretation is confirmed in the present study, which shows that the free fraction (T<sub>4</sub>DF) of T<sub>4</sub> and hence its serum transport binding in the hamster was not significantly altered in the presence of pineal-induced hypogonadism and that, like T<sub>4</sub> and FT<sub>4</sub>I,

Rx \_\_\_\_\_  
 Distribution/ \_\_\_\_\_  
 Availability Codes \_\_\_\_\_  
 Dist \_\_\_\_\_  
 Special \_\_\_\_\_  
 AY 21



**Fig. 1.** Testicular and prostate weights as absolute (g) or relative to body weight (mg/g), T<sub>4</sub>, FT<sub>4I</sub>, and FT<sub>4</sub> after sham pinealectomy (SHPX), blinding (BL + SHPX), or combined blinding and pinealectomy (BL + PX). Means ± SE (top and middle panels) and rectilinear regression (bottom panel).

**TABLE 1. T3U and T4DF After Sham Pinealectomy (SHPX), Blinding (BL), and/or Pinealectomy (PX)**

	SHPX	BL + SHPX	BL + PX
n in group	11	12	8
T3U (%)	38.04 ±.319	38.66 ±.293	36.63** ±.404
T4DF (%)	.0638 ±.004	.0581 ±.004	.0504 ±.006

\*\*P < 0.01 vs. SHPX and BL + SHPX.

Means ± SE.

FT4 was actually depressed in this condition. Furthermore, it was previously shown that the depressed T4 in blind hamsters was not a result of their gonadal regression, because castration alone did not alter serum T4 [Vriend et al., 1979]. Of course, the possibility remains that the pineal-dependency of the blinding-induced suppressions of both the reproductive and thyroidal systems results from some pineal-dependent common central neural control mechanism [Vriend, 1983].

The magnitude of the dialyzable fraction (T4DF) in hamsters appears to be similar to that in humans [Sterling and Brenner, 1966]. Because humans with altered levels of serum thyroxine-binding protein (and consequently abnormal total serum T4 concentration) often have normal free (dialyzable) concentrations but are euthyroid, it is thought that the free concentration is the bioavailable portion important for hormone action [Ingbar and Woeber, 1981]. Although somewhat more than the dialyzable fraction can be transported into some rat tissues [Pardridge, 1981], the precise fraction of T4 that is active in hamsters particularly at the level of the pituitary has not been elucidated. In the rat and human with primary reduction of thyroid function, it appears that T4 rather than T3 in the circulation determines feedback in the pituitary for thyrotrophin (TSH) secretion, via local T4 to T3 conversion [Connors and Hedge, 1981; Larsen, 1982]. The observation that both total and free T4 in serum are lower in blind than in control hamsters is important, because it indicates that the pineal gland can induce a condition in which bioavailable T4 in the circulation is reduced, a finding previously only suspected on the basis of indirect FT4I results [Vriend et al., 1977]. Whether or not the reduced FT4 results in other physiologic changes in these animals remains to be determined.

The site of the suppressive action of the pineal on serum T4 might be some element of the hypothalamopituitary unit (central), the thyroid, the organs responsible for T4 conversion or degradation (peripheral), or a combination of the above. TSH was not determined in the present animals because of an insufficient quantity of serum. As reviewed previously for the Syrian hamster [G.M. Vaughan, et al., 1982], in animals blind for 2-3 months, mean TSH is not consistently altered at a time when serum T4 is always low. Thus it is not unreasonable to suggest that the effect of the pineal is at least partly at the level of the pituitary or hypothalamus [Vriend, 1983], causing an inadequacy in the compensatory elevation of TSH, which normally should be a response to a fall in free T4 concentration [Larsen, 1982].

## ACKNOWLEDGMENTS

We thank James Lasko, Len Seraiffe, and Sandy Coggins for their technical work.

## LITERATURE CITED

1. Bruning, J.L., B.L. Kintz (1977) Computational Handbook of Statistics. Scott, Foresman and Company, Glenview, Illinois.
2. Chopra, I.J., T-S. Huang, R.E. Hurd, A. Beredo, D.H. Solomon (1984) A competitive ligand binding assay for measurement of thyroid hormone-binding in serum and tissues. *J. Clin. Endocrinol. Metab.* 58:619-628.
3. Chopra, I.J., D.H. Solomon, G.W. Hepner, A.A. Morgenstein (1979) Misleadingly low free thyroxine index and usefulness of reverse triiodothyronine measurement in nonthyroidal illness. *Ann. Intern. Med.* 90:905-912.
4. Connors, J.M., and G.A. Hedge (1981) Feedback regulation of thyrotropin by thyroxine under physiological conditions. *Am. J. Physiol.* 240:E308-E313.
5. Dieckhues, B. (1974) Die Bedeutung der Lichtperzeption durch das Auge auf den Hormonhaushalt des Menschen. *Klin. Monatsbl. Augenheilkd.* 165:291-296.
6. Dixon, W.J. (ed) (1983) BMDP Statistical Software. University of California Press, Berkeley.
7. Hoffman, RA, and R.J. Reiter (1965) Rapid pinealectomy in hamsters and other small rodents. *Anat. Rec.* 153:19-21.
8. Ingbar, S.H., and K.A. Woebber (1981) The thyroid gland. In: *Textbook of Endocrinology*. R.H. Williams, ed. W.B. Saunders, Philadelphia, p. 117.
9. Larsen, P.R. (1982) Thyroid-pituitary interaction: Feedback regulation of thyrotropin secretion by thyroid hormones. *N. Engl. J. Med.* 306:23-32.
10. Oppenheimer, J.H., H.L. Schwartz, C.N. Mariash, F.E. Kaiser (1982) Evidence for a factor in the sera of patients with nonthyroidal disease which inhibits iodothyronine binding by solid matrices, serum proteins, and rat hepatocytes. *J. Clin. Endocrinol. Metab.* 54:757-766.
11. Pardridge, W.M. (1981) Transport of protein-bound hormones into tissues in vivo. *Endocrinol. Rev.* 2:103-123.
12. Reiter, R.J. (1980) The pineal and its hormones in the control of reproduction in mammals. *Endocrinol. Rev.* 1:109-131.
13. Shirani, K.Z., G.M. Vaughan, B.A. Pruitt, Jr., A.D. Mason, Jr. (1985) Reduced serum T<sub>4</sub> and T<sub>3</sub> and their altered transport binding after burn injury in rats. *J. Trauma* (in press).
14. Sterling, K., and M.A. Brenner (1966) Free thyroxine in human serum: Simplified measurement with the aid of magnesium precipitation. *J. Clin. Invest.* 45:155-163.
15. Vaughan, G.M., A.D. Mason, Jr., W.F. McManus, B.A. Pruitt, Jr. (1985a) Alterations of mental status and thyroid hormones after thermal injury. *J. Clin. Endocrinol. Metab.* 60:1221-1225.
16. Vaughan, G.M., K.Z. Shirani, M.K. Vaughan, B.A. Pruitt, Jr., A.D. Mason, Jr. (1985b) Hormonal changes in a hamster burn model. *Endocrinol.* (in press).
17. Vaughan, G.M., M.K. Vaughan, L.G. Seraiffe, R.J. Reiter (1982) Thyroid hormones in male hamsters with activated pineals or melatonin treatment. *Prog. Clin. Biol. Res.* 92:187-196.
18. Vaughan, M.K., M.C. Powanda, G.C. Brainard, L.Y. Johnson, R.J. Reiter (1982) Effects of blinding or afternoon melatonin injections on plasma cholesterol, triglycerides, glucose, TSH and thyroid hormone levels in male and female Syrian hamsters. *Prog. Clin. Biol. Res.* 92:177-186.
19. Vriend, J. (1983) Evidence for pineal gland modulation of the neuroendocrine-thyroid axis. *Neuroendocrinology* 36:68-78.
20. Vriend, J., R.J. Reiter, G.R. Anderson (1979) Effects of the pineal and melatonin on thyroid activity of male golden hamsters. *Gen. Comp. Endocrinol.* 38:189-195.
21. Vriend, J., J.W. Sackman, R.J. Reiter (1977) Effects of blinding, pinealectomy and superior cervical ganglionectomy on free thyroxine index of male golden hamsters. *Acta Endocrinol* 86:758-762.
22. Woebber, K.A., and B.A. Maddux (1981) Thyroid hormone binding in non-thyroidal illness. *Metabolism* 30:412-416.