

# Plant Cell Cultures - An Alternative and Efficient Source for the Production of Biologically Important Secondary Metabolites

Vanisree Mulabagal and Hsin-Sheng Tsay\*

*Institute of Biotechnology,  
Chaoyang University of Technology,  
Wufeng, Taichung county 413, Taiwan, R.O.C.*

**Abstract:** Plant cell culture systems represent a potential renewable source of valuable medicinal compounds, flavors, fragrances, and colorants, which cannot be produced by microbial cells or chemical synthesis. Biotechnological applications of plant cell cultures presents the most updated reviews on current techniques in plant culture in the field. The evolving commercial importance of the secondary metabolites has in recent years resulted in a great interest, in secondary metabolism, and particularly in the possibility to alter the production of bioactive plant metabolites by means of cell culture technology. The principle advantage of this technology is that it may provide continuous, reliable source of plant pharmaceuticals and could be used for the large-scale culture of plant cells from which these metabolites can be extracted. In addition to its importance in the discovery of new medicines, plant cell culture technology plays an even more significant role in solving world hunger by developing agricultural crops that provide both higher yield and more resistance to pathogens and adverse environmental and climatic conditions. This paper describes the callus and suspension culture methods that we have established in our laboratory for the production of bioactive secondary metabolites from medicinal plants.

**Keywords:** bioreactors; callus cultures; secondary metabolites; suspension cultures; pharmaceuticals; precursor feeding.

## 1. Introduction

Medicinal plants are the most exclusive source of life saving drugs for the majority of the world's population. Bioactive compounds currently extracted from plants are used as food additives, pigments, dyes, insecticides, cosmetics and perfumes and fine chemicals [Balandrin and Klocke, 1988]. These compounds belong to a group collectively known as secondary metabolites. Studies on plant

secondary metabolites have been increasing over the last 50 years. The molecules are known to play a major role in the adaptation of plants to their environment, but also represent an important source of pharmaceuticals [Ramachandra Rao and Ravishankar, 2002].

In recent years, traditional system of medicine has become a topic of global importance. Although modern medicine may be available in developed countries, herbal medicines [phytopharmaceuticals] have often maintained popularity for historical and cultural reasons.

---

\* Corresponding author: e-mail: [hstsay@mail.cyut.edu.tw](mailto:hstsay@mail.cyut.edu.tw)

Many of the plant species that provide medicinal herbs have been scientifically evaluated for their possible medical applications. It has been mentioned that natural habitats for medicinal plants are disappearing fast and together with environmental and geopolitical instabilities; it is increasingly difficult to acquire plant-derived compounds. This has prompted industries, as well as scientists to consider the possibilities of investigation into cell cultures as an alternative supply for the production of plant pharmaceuticals.

Advances in biotechnology particularly methods for culturing plant cell cultures, should provide new means for the commercial processing of even rare plants and the chemicals they provide. These new technologies will extend and enhance the usefulness of plants as renewable resources of valuable chemicals. There has been considerable interest in plant cell cultures as a potential alternative to traditional agriculture for the industrial production of secondary metabolites [Dicosmo and Misawa, 1995]. Plant cell culture technologies were introduced at the end of 1960s as a possible tool for both studying and producing plant secondary metabolites. Different strategies using cell cultures systems have been extensively studied with the objective of improving the production of bioactive secondary metabolites. Cell culture systems could be used for the large scale culturing of plant cells from which secondary metabolites can be extracted. The advantage of this method is that it can ultimately provide a continuous, reliable source of natural products. The major advantages of cell cultures includes (i) synthesis of bioactive secondary metabolites is running in controlled environment, independently from climatic and soil conditions; (ii) negative biological influences that affect secondary metabolites production in the nature are eliminated (microorganisms and insects); (iii) it is possible to select cultivars with higher production of secondary metabolites; (iv) with automatization of cell growth control and metabolic processes regu-

lation, cost price can decrease and production increase. The scheme of production of some important plant pharmaceuticals produced in cell cultures has been presented in Table 1. The objectives of many industries are to develop plant cell culture techniques to the stage where they yield secondary products more cheaply than extracting either the whole plant grown under natural conditions or synthesizing the product. Although the production of pharmaceuticals using plant cell cultures have been highlighted, other uses have also been suggested as new route for synthesis, for products from plants difficult to grow, or in short supply, as a source of novel chemicals and as biotransformation systems. It is expected that the use, production of market price and structure would bring some of the other compounds to a commercial scale more rapidly and *in vitro* culture products may see further commercialization. Recent research results indicate that plant cell suspension cells can be used for recombinant protein production under controlled conditions [Fischer et al., 1999]. The aim of the present review is to focus the successful research accomplishments obtained on callus and suspension cultures for production of bioactive secondary metabolites in our research laboratory.

## **2. Accumulation of secondary metabolites in plant cell cultures**

For plant cell culture techniques to become economically viable, it is important to develop methods that would allow for consistent generation of high yields of products from cultured cells [Berlin and Sasse, 1985]. Careful selection of productive cells and cultural conditions resulted in accumulation of several products in higher levels in cultured cells. In order to obtain yields in high concentrations for commercial exploitation, efforts have focused on the stimulation of biosynthetic activities of cultured cells using various methods (Ramachandra Rao, 2000; Dixon, 1999; Ravishankar and Venkataraman, 1993; Buite-

laar and Tramper, 1992).

Table 1. Bioactive secondary metabolites from plant cell cultures

Plant name	Active ingredient	Culture type	Reference
<i>Agave amaniensis</i>	Saponins	Callus	Andrijany et al., 1999.
<i>Ailanthus altissima</i>	Alkaloids	Suspension	Anderson et al., 1987.
<i>Ailanthus altissima</i>	Canthinone alkaloids	Suspension	Anderson et al., 1986.
<i>Allium sativum</i> L.	Alliin	Callus	Malpathak and David , 1986.
<i>Aloe saponaria</i>	Tetrahydroanthracene gluco- sides	Suspension	Yagi et al., 1983.
<i>Ambrosia tenuifolia</i>	Altamisine	Callus	Goleniowski and Trippi , 1999.
<i>Anchusa officinalis</i>	Rosmarinic acid	Suspension	De-Eknamkul and Ellis, 1985.
<i>Brucea javanica</i> (L.) Merr.	Canthinone alkaloids	Suspension	Liu et al., 1990.
<i>Bupleurum falcatum</i>	Saikosaponins	Callus	Wang and Huang, 1982.
<i>Camellia Sinensis</i>	Theamine, $\gamma$ -glutamyl derivatives	Suspension	Orihara and Furuya, 1990.
<i>Canavalia ensiformis</i>	L-Canavanine	Callus	Ramirez et al., 1992.
<i>Capsicum annuum</i> L.	Capsaicin	Suspension	Johnson et al., 1990.
<i>Cassia acutifolia</i>	Anthraquinones	Suspension	Nazif et al., 2000.
<i>Catharanthus roseus</i>	Indole alkaloids	Suspension	Moreno et al., 1993.
<i>Catharanthus roseus</i>	Catharanthine	Suspension	Zhao et al., 2001.
<i>Choisya ternata</i>	Furoquinoline alkaloids	Suspension	Sejourne et al., 1981.
<i>Chrysanthemum cinerariaefolium</i>	Pyrethrins	Callus	Rajasekaran et al., 1991.
<i>Chrysanthemum cinerariaefolium</i>	Chrysanthemic acid and pyre- thrins	Suspension	Kueh et al., 1985.
<i>Cinchona</i> L.	Alkaloids	Suspension	Koblitz et al., 1983.
<i>Cinchona robusta</i>	Robustaquinones	Suspension	Schripsema et al., 1999.
<i>Cinchona</i> spe.	Anthraquinones	Suspension	Wijnsma et al., 1985.
<i>Cinchona succirubra</i>	Anthraquinones	Suspension	Khouri et al., 1986.
<i>Citrus</i> sp.	Naringin, Limonin	Callus	Barthe et al., 1987.
<i>Coffea arabica</i> L.	Caffeine	Callus	Waller et al., 1983.
<i>Cornus kousa</i>	Polyphenols	Suspension	Ishimaru et al., 1993.
<i>Corydalis ophiocarpa</i>	Isoquinoline alkaloids	Callus	Iwasa and Takao, 1982.
<i>Croton sublyratus</i> Kurz	Plaunotol	Callus	Morimoto and Murai, 1989.
<i>Cruciata glabra</i>	Anthraquinones	Suspension	Dornenburg and Knorr, 1996.
<i>Cryptolepis buchanani</i> Roem. & Schult	Cryptosin	Callus	Venkateswara et al., 1987.
<i>Digitalis purpurea</i> L.	Cardenolides	Suspension	Hagimori et al., 1982
<i>Dioscorea deltoidea</i>	Diosgenin	Suspension	Heble and Staba, 1980.
<i>Dioscorea doryophora</i> Hance	Diosgenin	Suspension	Huang et al., 1993.
<i>Duboisia leichhardtii</i>	Tropane alkaloids	Callus	Yamada and Endo, 1984.
<i>Ephedra</i> spp.	L- Ephedrine D-pseudoephedrine	Suspension	O'Dowd et al., 1993.

continued ...

<i>Eriobotrya japonica</i>	Triterpenes	Callus	Taniguchi et al., 2002.
<i>Eucalyptus tereticornis</i> SM.	Sterols and Phenolic compounds	Callus	Venkateswara et al., 1986.
<i>Eucommia ulmoides</i>	Chlorogenic acid	Suspension	Wang et al., 2003.
<i>Fumaria capreolata</i>	Isoquinoline alkaloids	Suspension	Tanahashi and Zenk, 1985.
<i>Gentiana</i> sp.	Secoiridoid glucosides	Callus	Skrzypczak et al., 1993.
<i>Ginkgo biloba</i>	Ginkgolide A	Suspension	Carrier et al., 1991.
<i>Glehnia littoralis</i>	Furanocoumarin	Suspension	Kitamura et al., 1998.
<i>Glycyrrhiza echinata</i>	Flavonoids	Callus	Ayabe et al., 1986.
<i>Glycyrrhiza glabra</i> var. <i>glandulifera</i>	Triterpenes	Callus	Ayabe et al., 1990.
<i>Hyoscyamus niger</i>	Tropane alkaloids	Callus	Yamada and Hashimoto, 1982.
<i>Isoplexis isabellina</i>	Anthraquinones	Suspension	Arrebola et al., 1999.
<i>Linum flavum</i> L.	5-Methoxypodophyllo-toxin	Suspension	Uden et al., 1990.
<i>Lithospermum erythrorhizon</i>	Shikonin derivatives	Suspension	Fujita et al., 1981.
<i>Lithospermum erythrorhizon</i>	Shikonin derivatives	Suspension	Fukui et al., 1990.
<i>Lycium chinense</i>	Cerebroside	Suspension	Jang et al., 1998.
<i>Morinda citrifolia</i>	Anthraquinones	Suspension	Zenk et al., 1975.
<i>Morinda citrifolia</i>	Anthraquinones	Suspension	Bassetti et al., 1995.
<i>Mucuna pruriens</i>	L-DOPA	Suspension	Wichers et al., 1993.
<i>Mucuna pruriens</i>	L-DOPA	Callus	Brain K. R., 1976.
<i>Nandina domestica</i>	Alkaloids	Callus	Ikuta and Itokawa, 1988.
<i>Nicotiana rustica</i>	Alkaloids	Callus	Tabata and Hiraoka, 1976.
<i>Nicotiana tabacum</i> L.	Nicotine	Suspension	Mantell et al., 1983.
<i>Nothapodytes foetida</i>	Camptothecin	Callus	Thengane et al., 2003.
<i>Ophiorrhiza pumila</i>	Camptothecin related alkaloids	Callus	Kitajima et al., 1998.
<i>Panax ginseng</i>	Saponins and Sapogenins	Callus	Furuya et al., 1973.
<i>Panax notoginseng</i>	Ginsenosides	Suspension	Zhong and Zhu, 1995.
<i>Papaver bracteatum</i>	Thebaine	Callus	Day et al., 1986.
<i>Papaver somniferum</i> L.	Alkaloids	Callus	Furuya et al., 1972.
<i>Papaver somniferum</i>	Morphine , Codeine	Suspension	Siah and Doran, 1991.
<i>Peganum harmala</i> L.	$\beta$ -Carboline alkaloids	Suspension	Sasse et al., 1982.
<i>Phytolacca americana</i>	Betacyanin	Suspension	Sakuta et al., 1987.
<i>Picrasma quassioides</i> Bennett	Quassin	Suspension	Scragg and Allan, 1986.
<i>Podophyllum hexandrum</i> royle	Podophyllotoxin	Suspension	Uden et al., 1989. Chattopadhyay et al., 2002.
<i>Polygala amarella</i>	Saponins	Callus	Desbene et al., 1999.
<i>Polygonum hydropiper</i>	Flavonoids	Suspension	Nakao et al., 1999.
<i>Portulaca grandiflora</i>	Betacyanin	Callus	Schroder and Bohm, 1984.
<i>Ptelea trifoliata</i> L.	Dihydrofuro [2,3-b] quinolinium alkaloids	Callus	Petit-Paly et al., 1987.
<i>Rauwolfia sellowii</i>	Alkaloids	Suspension	Rech et al., 1998.
<i>Rauwolfia serpentina</i> Benth.	Reserpine	Suspension	Yamamoto and Yamada, 1986.

continued ...

<i>Rauvolfia serpentina</i> x <i>Rhazya stricta</i> Hybrid plant	3-Oxo-rhazinilam	Callus	Gerasimenko et al., 2001.
<i>Ruta</i> sp.	Acridone and Furoquinoline alkaloids and coumarins	Callus	Baumert et al., 1992.
<i>Salvia fruticosa</i>	Rosmarinic acid	Callus & suspension	Karam et al., 2003
<i>Salvia miltiorrhiza</i>	Lithospermic acid B and Rosmarinic acid	Callus	Morimoto et al., 1994.
<i>Salvia miltiorrhiza</i>	Cryptotanshinone	Suspension	Miyasaka et al., 1989.
<i>Sapium sebiferum</i>	Tannin	Callus & suspension	Neera and Ishimaru, 1992.
<i>Scopolia parviflora</i>	Alkaloids	Callus	Tabata et al., 1972.
<i>Scutellaria columnae</i>	Phenolics	Callus	Stojakowska and Kisiel, 1999.
<i>Solanum chrysotrichum</i> (Schldl.)	Spirostanol saponin	Suspension	Villarreal et al., 1997.
<i>Solanum laciniatum</i> Ait	Solasodine	Suspension	Chandler and Dodds, 1983.
<i>Solanum paludosum</i>	Solamargine	Suspension	Badaoui et al., 1996.
<i>Stizolobium hassjoo</i>	L-DOPA	Suspension	Huang et al., 2002.
<i>Tabernaemontana divaricata</i>	Alkaloids	Suspension	Sierra et al., 1992.
<i>Taxus</i> spp.	Taxol	Suspension	Wu et al., 2001.
<i>Taxus baccata</i>	Taxol, baccatin III	Suspension	Cusido et al., 1999.
<i>Taxus cuspidata</i>	Taxoids	Suspension	Ketchum et al., 2003.
<i>Tecoma sambucifolium</i>	Phenylpropanoid glycosides	Callus	Pletsch et al., 1993
<i>Thalictrum minus</i>	Berberin	Suspension	Kobayashi et al., 1987.
<i>Thalictrum minus</i>	Berberin	Suspension	Nakagawa et al., 1986.
<i>Torreya nucifera</i> var. <i>radicans</i>	Diterpenoids	Suspension	Orihara et al., 2002.
<i>Trigonella foenumgraecum</i>	Saponins	Suspension	Brain and Williams, 1983.

Culture productivity is critical to the practical application of plant cell culture technology to production of plant-specific bioactive metabolites. Until now, various strategies have been developed to improve the production of secondary metabolites using plant cell cultures. The tissue culture cells typically accumulate large amounts of secondary compounds only under specific conditions. That means maximization of the production and accumulation of secondary metabolites by plant tissue cultured cells requires (i) manipulating the parameters of the environment and medium, (ii) selecting high yielding cell clones, (iii) precursor feeding, and (iv) elicitation.

**Optimization of cultural conditions:** number of chemical and physical factors like media components, phytohormones, pH, temperature, aeration, agitation, light affecting production of secondary metabolites has been extensively studied (Lee and Shuler, 2000; Wang et al., 1999; Fett-Neto et al., 1995. Goleniowski and Trippi, 1999). Several products were found to be accumulating in cultured cells at a higher level than those in native plants through optimization of cultural conditions. Manipulation of physical aspects and nutritional elements in a culture is perhaps the most fundamental approach for optimization of culture productivity. For example, ginsenosides by *Panax ginseng* [Choi et al., 1994; Furuya et al., 1984; Franklin and

Dixon, 1994; Furuya, 1988;], rosmarinic acid by *Coleus blumei* [Ulbrich et al., 1985], shikonin by *Lithospermum erythrorhizon* [Takahashi and Fujita, 1991], ubiquinone-10 by *Nicotiana tabacum* [Fontanel and Tabata, 1987], berberin by *Coptis japonica* [Matsubara et al., 1989], were accumulated in much higher levels in cultured cells than in the intact plants.

**Selection of high-producing strains:** plant cell cultures represent a heterogeneous population in which physiological characteristics of individual plant cells are different. Synthesis of several products in high amounts using selection and screening of plant cell cultures have been described by Berlin and Sasse [1985]. Cell cloning methods provide a promising way of selecting cell lines yielding increased levels of product. A strain of *Euphorbia milli* accumulated about 7-fold the level of anthocyanins produced by the parent culture after 24 selections [Yamamoto et al., 1982]. Selection can be easily achieved if the product of interest is a pigment (Fujita et al., 1984). Cell cloning using cell aggregates of *Coptis japonica* [Yamada and Sato, 1981], and obtained strain, which grew faster and produced a higher amount of berberin and cultivated the strain in a 14 L bioreactor. Selected cell line increased growth about 6-fold in 3 weeks and the highest amount of alkaloid was produced 1.2 g/L of the medium and the strain was very stable, producing a high level of berberin even after 27 generations. Increased capsaicin and rosmarinic acid in PEP cell lines of *Capsicum annuum* were reported (Salgado-Garciglia and Ochoa-Alejo, 1990). Selective agents such as 5-methyltryptophan, glyphosate and biotin have also been studied to select high-yielding cell lines (Amrhein et al., 1985; Watanabe et al., 1982; Widholm, 1974).

**Precursor feeding:** exogenous supply of a biosynthetic precursor to culture medium may also increase the yield of the desired product.

This approach is useful when the precursors are inexpensive. The concept is based on the idea that any compound, which is an intermediate, in or at the beginning of a secondary metabolite biosynthetic route, stands a good chance of increasing the yield of the final product. Attempts to induce or increase the production of plant secondary metabolites, by supplying precursor or intermediate compounds, have been effective in many cases (Silvestrini et al., 2002; Moreno et al., 1993; Whitmer et al., 1998). For example, amino acids have been added to cell suspension culture media for production of tropane alkaloids, indole alkaloids etc. Addition of phenylalanine to *Salvia officinalis* cell suspension cultures stimulated the production of rosmarinic acid [Ellis and Towers, 1970]. Addition of the same precursor resulted stimulation of taxol production in *Taxus* cultures [Fett-Neto et al., 1993 and 1994]. Feeding ferulic acid to cultures of *Vanilla planifolia* resulted in increase in vanillin accumulation (Romagnoli and Knorr, 1988). Furthermore, addition of leucine, led to enhancement of volatile monoterpenes in cultures of *Perilla frutiscens*, where as addition of geraniol to rose cell cultures led to accumulation of nerol and citronellol (Mulder-Krieger et al., 1988).

**Elicitation:** plants produce secondary metabolites in nature as a defense mechanism against attack by pathogens. Elicitors are signals triggering the formation of secondary metabolites. Use of elicitors of plant defense mechanisms, i.e. elicitation, has been one of the most effective strategies for improving the productivity of bioactive secondary metabolites (Roberts and Shuler, 1997). Biotic and abiotic elicitors which are classified on their origin are used to stimulate secondary metabolite formation in plant cell cultures, thereby reducing the process time to attain high product concentrations (Barz et al., 1988; Eilert, 1987; DiCosmo and Tallevi, 1985). Production of many valuable secondary metabolites using various elicitors were reported

(Wang and Zhong, 2002a, 2002b; Dong and Zhong, 2001; Hu et al., 2001; Lee and Shuler, 2000).

### 3. Cell culture protocols for production of important secondary metabolites established in our laboratory

Most applications of plant cell cultures in biotechnology are aimed at the production of bioactive secondary metabolites. This has included production of taxol, morphine and codeine, ginsenosides, L-DOPA, berberine, diosgenin, capsaicin, podophyllotoxin, shikonin derivatives, ajmalicine, vincristine and vinblastine [Dicosmo and Misawa, 1995]. Since cell suspension cultures are preferred for large-scale production due to its rapid growth cycles they have been used for generating large amounts of cells for quantitative or qualitative analysis of growth responses and metabolism of novel chemicals. In our laboratory, we focus on the production of some important pharmaceuticals in plant cell cultures. We have successfully established cell cultures for production of taxol from *Taxus mairei*, imperatorin from *Angelica dahurica*, diosgenin from *Dioscorea doryophora*, gentipicroside and swertiamarin from *Gentiana*, cryptotanshinone from *Salvia miltiorrhiza* and the results were described in the present review.

#### 3.1. Case study 1: Production of taxol from *Taxus mairei* by cell suspension cultures

As early as 1969, Wani and his colleagues discovered a novel anticancer diterpene amide, "taxol" from the Pacific yew (*Taxus brevifolia*) extract [Wani et al., 1971]. In 1983, taxol was approved to enter phase I clinical trials for ovarian cancer by the Food & Drug Administration (FDA) in the USA. It has been approved for clinical treatment of ovarian and breast cancer by the FDA, and it also had significant activity in the treatment of patients

with malignant melanoma, lung cancer, and other solid tumors [Wickremesinhe and Arteca, 1993 and 1994]. Taxol is considered as the prototype of a new class of cancer chemotherapeutic agents [Cragg et al., 1993].

However the supply of taxol for clinical use is limited. It depends on extraction from yew trees, and the bark is the only commercial source. The thin bark of yew tree contains 0.001% of taxol by dry weight basis. A century old tree yields an average of 3 kg of bark, corresponding to 300 mg of taxol, which is approximately a single dose in the course of a cancer treatment. Because of the scarcity of the slow growing trees and the relatively low content of taxol [Cragg et al., 1993], alternative sources are needed to meet the increasing demand for the drug. The total synthesis of taxol on an industrial-scale seems economically unrealistic due to the complexity of the chemical structure of this molecule [Holton et al., 1994; Nicolaou et al., 1994]. The plant cell culture of *Taxus* spp. is considered as one of the approach available to provide a stable supply of taxol and related taxane derivatives [Slichenmyer and Von Horf, 1991].

For exploiting the source of taxol, we collected different tissues of *Taxus mairei*, a species found in Taiwan at an altitude of about 2000m above the sea level. The extracts of bark and leaf tissues were analyzed by using HPLC for the content of taxol and taxol related compounds. The HPLC analysis revealed that the amounts of taxol and taxol related compounds varies in individual plants, and the principle components such as docetaxel, baccatin III, and 10-deacetylbaaccatin in leaf extract were higher than those in bark extracts [Lee et al., 1995]. *Taxus mairei* calli were induced from needle and stem explants on B5 medium [Gamborg's et al., 1968] supplemented with 2 mg/l 2,4-D or NAA. Different cell lines were established using stem and needle-derived callus. One of the cell lines, after precursor feeding and 6 weeks of incubation, produced 200 mg taxol per liter cell suspension cultures.

### 3.2. Case study 2: Formation of imperatorin from *Angelica dahurica* var. *formosana* by cell suspension cultures

*Angelica dahurica* var. *formosana* commonly known as "Bai-Zhi" in Chinese is a valuable medicinal herb used in the treatment of headache and psoriasis in China [Zhou, 1980]. The constituent imperatorin has been suggested as the major active ingredient for curing the skin disease [Zhou et al., 1988]. *Angelica dahurica* var. *formosana* is a perennial and indigenous plant in Taiwan [Chen et al., 1994].

As the production of Bai-Zhi by conventional cultural method is far short to meet the demand, cell suspension culture methods for production of imperatorin was therefore emphasized. For establishing a rapidly growing and finely dispersed cell suspension culture, the best medium composition was 1/2 strength of MS medium [Murashige and Skoog's, 1962] supplemented with 1 mg/L 2,4-D, 0.1 mg/L kinetin, and 3% sucrose. All the cultures were routinely sub cultured at an interval of 14 days.

In a growth cycle of the suspension cells, the production of imperatorin was maximum between the 10<sup>th</sup> and 14<sup>th</sup> day. Growth conditions, especially the nutritional factors optimal for imperatorin production were also investigated. It was found that MS basal media was the best among the media tested on imperatorin production. Influence of auxins on imperatorin production were investigated. Deletion of auxins from the medium was beneficial to imperatorin production and addition of BA (0.5-1 mg/L) promoted synthesis of imperatorin in suspension cells.

The effects of the ratio of ammonium nitrate to nitrate in the medium had also been studied. A moderate ammonium nitrate to nitrate ratio (2:1) increased the amount of imperatorin production. Increasing phosphate concentration (1 mM to 2 mM) promoted imperatorin production. Glucose was found to be better carbon source than sucrose and

fructose in terms of their effects on imperatorin production. The possible stimulating effect of elicitors on imperatorin synthesis were studied. Addition of vanadyl sulphate to cell suspension cultures enhance the accumulation of imperatorin, depending on its concentration and the growth stage of the cells. The most significant effect was recorded when vanadyl sulphate at the concentration of 30 mg/L was added to the medium on the 10<sup>th</sup> day of the culture. A rapid and drastic increase in imperatorin synthesis was observed by adding 20 g/L of adsorbent Amberlite XAD-7 to suspension cells on the 10<sup>th</sup> day of culture. The quantity of imperatorin produced by this treatment (460  $\mu$ g/gd. w.) was 140 folds higher than that produced by the check treatment [Tsay, 1999; Tsay et al., 1994]. These results showed that the cultured cells possessed the biosynthetic potential of the intact plants from which they were derived. The biosynthetic potential could be further facilitated by the addition of proper stimulants to the culture medium.

### 3.3. Case study 3: Diosgenin production from *Dioscorea doryophora* by cell suspension cultures

*Dioscorea* spp. (Dioscoreaceae) is frequently used as tonic in Chinese traditional medicine. *Dioscorea doryophora* Hance, tubers are in high demand as it is used not only as crude drug but also as food. The most active ingredient discovered in the tuber is diosgenin, which can be used as a precursor for many important medicinal steroids such as prednisolone, dexamethasone, norethisterone and metenolone etc. [Tsukamoto et al., 1936].

The purpose of the present study was to establish the cell suspension cultures of *Dioscorea doryophora* Hance for the production of diosgenin, which can be used as an alternative source for the synthesis of steroids [Yeh et al., 1994]. Callus was successfully induced on MS basal medium supplemented with 2.0 mg/L 2, 4-D and 0.2 mg/L BA. Cell suspen-



sion cultures was successfully established by subculturing callus into a liquid medium containing MS basal medium supplemented with 0.2 mg/L 2, 4-D. Sucrose concentration optimal for diosgenin production was investigated. Although 6% sucrose was an optimum concentration for the growth and increasing dry weight of the cell suspension culture, 3% sucrose gave better results of the diosgenin production. Differences in the quality and diosgenin content of the induced callus were observed among the six source organs. Analysis by HPLC revealed that both stem-node and microtuber derived suspension cells contained diosgenin. The content of microtuber derived cell suspension culture contains 3.2% diosgenin per gram dry weight, where as, the stem-node derived cultures contains only 0.3%. Callus from microtuber showed yellowish color and granular in appearance and contained the highest diosgenin content (3.3-3.5%). As the amount of diosgenin obtained from tuber derived cell suspension is high and comparable with that found in the intact tuber [Chen, 1985], cell suspension culture could be used for production of diosgenin.

### 3.4. Case study 4: Establishment of *Gentiana davidii* var. *formosana* (Hayata) T. N. Ho cell suspension cultures

*Gentiana davidii* var. *formosana* (Hayata) T. N. Ho (Gentianaceae), commonly known as Long-dan in Chinese is a perennial herb indigenous to Taiwan. Regular consumption of Long-dan has been believed to enhance memory, prevent obesity and aging, and protect the liver by removing toxic metabolites [Lou and Chin, 1996 and Zheng et al., 1997]. The entire dried herb, collected from the wild habitat, is used as a crude drug in traditional Chinese medicine in Taiwan. Bitter principles of Gentianaceae constitute many pharmacologically important compounds, which justify the use of most species of this family in traditional medicine or for the preparation of bitter

tonic [Rodriguez et al., 1996]. Secoiridoid glycosides are the main compounds with medicinal properties in roots of *Gentiana* species [Skrzypczak et al., 1993].

We have optimized the conditions for establishment of cell suspension cultures of *G. davidii* for the production of gentipicroside and swertiamarin, the two pharmacologically important compounds. Callus was initiated by culturing stem explants of *G. davidii* var. *formosa* on MS basal medium supplemented with 0.2 mg/L kinetin and 1.0 mg/L  $\alpha$ -naphthaleneacetic acid (NAA). Fast growing cell suspension cultures were established by subculturing the callus in MS basal medium supplemented with 0.2 mg/L kinetin and 3% sucrose [Chueh et al., 2000].

Optimal cell growth was obtained when callus cultured in 25 mL liquid MS basal medium supplemented with 0.2 mg/L kinetin and 3% sucrose, pH between 4.2-5.2, incubated in light  $2.33 \mu E m^{-2} s^{-1}$  at  $25 \pm 1^\circ C$ , under 80-100 rev/min shaker speed. The maximum content of the two principles, swertiamarin and gentipicroside, in cell suspension were obtained after 12 and 24 days of culture respectively. Using this standard protocol, it may be possible to study the effect of precursor feeding on the content of active principles.

### 3.5. Case study 5: Production of cryptotanshinone from callus cultures of *Salvia miltiorrhiza* Bunge.

*Salvia* is an important genus consisting of ca. 900 species in the family Lamiaceae and some species of *Salvia* have been cultivated worldwide for use in folk medicine. Dan-shen, the dried roots of *Salvia miltiorrhiza* Bunge, is one of the most popular Chinese medicines and widely used for promoting blood circulation to remove blood stasis, clearing away heat, relieving vexation, nourishing blood and tranquilizing the mind and cooling blood to relieve carbuncles [Verpoorte, 1998]. The major active principles of the organic extract of Dan-shen are tanshinones, the quinoid

diterpenes [Bruneton, 1995]. Since Dan-shen preparations constitute a basis for considerable commercial activity, there is a continued interest in development of biotechnology-based approaches to the production of tanshinones [Hu and Alfermann, 1993; Miyasaka et al., 1989 and Shimomura et al., 1991]. In our continuing study of the application of tissue culture of medicinal plants to the production of bioactive secondary metabolites, we have adopted an approach to the production of cryptotanshinone from *S. miltiorrhiza* through callus cultures.

In this experiment effect of  $N^6$ -benzyladenine (BA) on cryptotanshinone formation in callus cultures of *Salvia miltiorrhiza* was examined. Primary callus was induced by culturing leaf explants on Murashige and Skoog's (MS) basal medium supplemented with 1.0 mg/L of 2,4-dichlorophenoxyacetic acid (2,4-D) in darkness. The callus proliferated further on MS basal medium containing 1.0 mg/L 2,4-D and 0.5 mg/L BA and was analyzed for cryptotanshinone by high performance liquid chromatography (HPLC). The HPLC results indicated that it contained small amounts of cryptotanshinone ( $0.26 \pm 0.05$  mg/g dry wt.). Omission of 2,4-D from the medium resulted in a marked increase in the content of cryptotanshinone in callus. The HPLC analysis revealed that the content of cryptotanshinone in the callus cultured on the MS basal medium supplemented with 0.1, 0.2, 0.5, 1.0, and 2.0 mg/L of BA was significantly higher than the marketed crude drug (processed underground parts of *S. miltiorrhiza*). Maximum yield of cryptotanshinone ( $4.59 \pm 0.09$  mg/g dry wt.) was observed in the callus cultured on MS basal medium supplemented with 0.2 mg/L BA for sixty days [Wu et al., 2003].

#### 4. Conclusions

The use of plant cell culture for the production of chemicals and pharmaceuticals has made great strides building on advances in

plant science. The increased use of genetic tools and an emerging picture of the structure and regulation of pathways for secondary metabolism will provide the basis for the production of commercially acceptable levels of product. The increased appeal of natural products for medicinal purposes coupled with the low product yields and supply concerns of plant harvestation has renewed interest in large-scale plant cell culture technology. Knowledge of biosynthetic pathways of desired compounds in plants as well as in cultures is often still in its infancy, and consequently, strategies are needed to develop an information based on a cellular and molecular level. Because of the complex and incompletely understood nature of plant cells in *in vitro* cultures, case-by-case studies have been used to explain the problems occurring in the production of secondary metabolites from cultured plant cells. These results shows that plant cell culture systems have potential for commercial exploitation of secondary metabolites. The introduction of the techniques of molecular biology, so as to produce transgenic cultures and to effect the expression and regulation of biosynthetic pathways, is also likely to be a significant step towards making cell cultures more generally applicable to the commercial production of secondary metabolites.

#### References

- [ 1 ] Amrhein, N., Johanning, D., and Smart, C. C. 1985. A glyphosate-tolerant plant tissue cultures. In: Neumann, K. H., Barz, W., Reinhard, E., editors. "*Primary and Secondary Metabolism of Plant Cell Cultures*". Berlin: Springer-Verlag: 355-361.
- [ 2 ] Anderson, L. A., Hay, C. A., Roberts, M. F., and Phillipson, J. D. 1986. Studies on *Ailanthus altissima* cell suspension cultures. *Plant Cell Reports*, 5: 387-390.
- [ 3 ] Anderson, L. A., Roberts, M. F., and Phillipson, J. D. 1987. Studies on *Ailan-*

- thus altissima cell suspension cultures. The effect of basal media on growth and alkaloid production. *Plant Cell Reports*, 6: 239-241.
- [ 4] Andrijany, V. S., Indrayanto, G., and Soehono, L. D. 1999. Simultaneous effect of calcium, magnesium, copper and cobalt on sapogenin steroids content in callus cultures of *Agave amaniensis*. *Plant Cell Tissue and Organ Culture*, 55: 103-108.
- [ 5] Arrebola, M. L., Ringbom, T., and Verpoorte, R. 1999. Anthraquinones from *Isoplexis isabelliana* cell suspension cultures. *Phytochemistry*, 52: 1283-1286.
- [ 6] Ayabe, S., Iida, K., and Furuya, T. 1986. Induction of stress metabolites in immobilized *Glycyrrhiza echinata* cultured cells. *Plant Cell Reports*, 3: 186-189.
- [ 7] Ayabe, S., Takano, H., Fujita, T., Hirota, H., and Takahashi, T. 1990. Triterpenoid biosynthesis in tissue cultures of *Glycyrrhiza glabra* var. *glandulifera*. *Plant Cell Reports*, 9: 181-184.
- [ 8] Badaoui, H. E1., Muguet, B., and Henry, M. 1996. Production of solamargine by *in vitro* cultures of *Solanum paludosum*. *Plant Cell Tissue and Organ Culture*, 45: 123-127.
- [ 9] Balandrin, M. J. and Klocke, J. A. 1988. Medicinal, aromatic and industrial materials from plants. In: Bajaj, Y. P. S., editor. "Biotechnology in Agriculture and Forestry. Medicinal and Aromatic plant". 4, Springer-Verlag, Berlin, Heidelberg: 1-36.
- [10] Barthe, G. A., Jourdan, P. S., McIntosh, C. A., and Mansell, R.L. 1987. Naringin and limonin production in callus cultures and regenerated shoots from *Citrus* sp. *Journal of Plant Physiology*, 127: 55-65.
- [11] Barz, W., Daniel, S., Hinderer, W., Jaques, U., Kessmann, H., Koster, J., and Tiemann, K. 1988. Elicitation and metabolism of phytoalexins in plant cell cultures. In: Pais, M., Mavituna, F., Novais, J, editors. "Plant Cell Biotechnology". NATO ASI Series. Berlin: Springer-Verlag: 211-230.
- [12] Bassetti, L., Hagendoorn, M., and Johannes, T. 1995. Surfactant-induced non-lethal release of anthraquinones from suspension cultures of *Morinda citrifolia*. *Journal of Biotechnology*, 39: 149-155.
- [13] Baumert, A., Groger, D., Kuzovkina, I. N., and Reisch, J. 1992. Secondary metabolites produced by callus cultures of various *Ruta* species. *Plant Cell Tissue and Organ Culture*, 28: 159-162.
- [14] Berlin, J. and Sasse, F. 1985. Selection and screening techniques for plant cell cultures. *Advanced Biochemistry and Engineering*, 31: 99-132.
- [15] Brain, K. R. 1976. Accumulation of L-DOPA in cultures from *Mucuna pruriens*. *Plant Science Letters*, 7: 157-161.
- [16] Brain, K. R. and Williams, M. H. 1983. Evidence for an alternative route from sterol to sapogenin in suspension cultures from *Trigonella foenumgraecum*. *Plant Cell Reports*, 2: 7-10.
- [17] Bruneton, J. 1995. Pharmacognosy, Phytochemistry, Medicinal plants. *Intercept UK*, 522.
- [18] Buitelaar, R. M. and Tramper, J. 1992. Strategies to improve the production of secondary metabolites with plant cell cultures: a literature review. *Journal of Biotechnology*, 23: 111-143.
- [19] Carrier, D., Chauret, N., Mancini, M., Coulombe, P., Neufeld, R., Weber, M., and Archambault, J. 1991. Detection of ginkgolide A in *Ginkgo biloba* cell cultures. *Plant Cell Reports*, 10: 256-259.
- [20] Chandler, S. and Dodds, J. 1983. Solasodine production in rapidly proliferating tissue cultures of *Solanum laciniatum* Ait. *Plant Cell Reports*, 2: 69-72.
- [21] Chattopadhyay, S., Srivastava, A. K., Bhojwani, S. S., and Bisaria, V. S. 2002. Production of podophyllotoxin by plant cell cultures of *Podophyllum hexandrum* in bioreactor. *Journal of Bioscience and*

- Bioengineering*, 93 ,2: 215-220.
- [22] Chen, A. H. 1985. Study on application of diosgenin-I. Analysis of diosgenin constituent of plants from Taiwan. *Science Monthly*, 43: 79-85.
- [23] Chen, C. C., Chang, W. T., Chang, Y. S., and Tsay, H. S. 1994. Studies on the tissue culture of *Angelica dahurica* var. *formosana* II. Establishment of cell suspension culture and evaluation of cultural conditions. *Journal of Chinese Medicine*, 5: 123-134.
- [24] Choi, K. T., Ahn, I. O., and Park, J. C. 1994. Production of ginseng saponin in tissue culture of ginseng (*Panax ginseng* C.A. Mayer). *Russian Journal of Plant Physiology*, 41: 784-788.
- [25] Chueh, F. S., Chen, C. C., Sagare, A. P., and Tsay, H. S. 2000. Quantitative determination of secoiridoid glucoside in in vitro propagated plants of *Gentiana davidii* var. *formosana* by high performance liquid chromatography. *Planta Medica*, 67: 70-73.
- [26] Cragg, G. M., Schepartz, S. A., Suffness, M., and Grever, M. R. 1993. The taxol supply crisis. New NCI policies for handling the large-scale production of novel natural product anticancer and anti-HIV agents. *Journal of Natural Products*, 56: 1657-1668.
- [27] Cusido R. M., Palazon, J., Navia-Osorio, A., Mallol, A., Bonfill, M., Morales, C., and Pinol, M.T. 1999. Production of taxol and baccatin III by a selected *Taxus baccata* callus line and its derived cell suspension culture. *Plant Science*, 146: 101-107.
- [28] Day, K. B., Draper, J., and Smith, H. 1986. Plant regeneration and thebaine content of plants derived from callus culture of *Papaver bracteatum*. *Plant Cell Reports*, 5: 471-474.
- [29] De-Eknamkul, D. and Ellis, B. E. 1985. Effects of macronutrients on growth and rosmarinic acid formation in cell suspension cultures of *Anchusa officinalis*. *Plant Cell Reports*, 4: 46-49.
- [30] Desbene, S., Hanquet, B., Shoyama, Y., Wagner, H., and Lacaille-Dubois, M. 1999. Biologically active triterpene saponins from callus tissue of *Polygala amarella*. *Journal of Natural Products*, 62: 923-926.
- [31] Dicosmo, F. and Misawa, M. 1995. Plant cell and tissue culture: Alternatives for metabolite production. *Biotechnology Advances*, 13 , 3: 425-453.
- [32] Dicosmo, F. and Tallevi, S. G. 1985. Plant cell cultures and microbial insult: interactions with biotechnological potential. *Trends in Biotechnology*, 3: 110-111.
- [33] Dixon, R. A. 1999. Plant natural products: the molecular genetic basis of biosynthetic diversity. *Current Opinion in Biotechnology*, 10: 192-197.
- [34] Dong, H. D. and Zhong, J. J. Significant improvement of taxane production in suspension cultures of *Taxus chinensis* by combining elicitation with sucrose feed. *Biochemical engineering Journal*, 8: 145-150.
- [35] Dornenburg, H. and Knorr, D. 1996. Semicontinuous processes for anthraquinone production with immobilized *Cruciata glabra* cell cultures in a three-phase system. *Journal of Biotechnology*, 50: 55-62.
- [36] Eilert, U. 1987. Elicitation: methodology and aspects of application. In: Constabel, F., Vasil, I., editors. "Cell Culture and Somatic Cell Genetics of Plants". 4, San Diego: Academic Press: 153-196.
- [37] Ellis, B. E. and Towers, G. H. N. 1970. Biogenesis of rosmarinic acid in *Mentha*. *Journal of Biochemistry*, 118: 291-297.
- [38] Fett-Neto, A. G., Melanson, S. J., Sakata, K., and DiCosmo, F. 1993. Improved growth and taxol yield in developing calli of *Taxus cuspidata* by medium composition modification. *Biotechnology*, 11: 731-734.
- [39] Fett-Neto A. G., Pennington, J. J., and DiCosmo, F. 1995. Effect of white light

- on taxol and baccatin III. Accumulation in cell cultures of *Taxus cuspidata* Sieb and Zucc. *Journal of Plant Physiology*, 146: 584-590.
- [40] Fett-Neto, A. G., Stewart, J. M., Nicholson, S. A., Pennington, J. J., and Di-Cosmo, F. 1994. Improved taxol yield by aromatic carboxylic acid and amino acid feeding to cell cultures of *T. cuspidata*. *Biotechnology Bioengineering*, 44: 967-971.
- [41] Fischer, R., Liao, Y. C., Hoffmann, K., Schillberg, S., and Emans, N. 1999. Molecular farming of recombinant antibodies in plants. *Biological Chemistry*, 380: 825-839.
- [42] Fontanel, A. and Tabata, M. 1987. Production of secondary metabolites from plant tissue and cell cultures. *Nestle Research News*: 92-103.
- [43] Franklin, C. I. and Dixon, R. A. 1994. Initiation and maintenance of callus and cell suspension cultures. In: Dixon, R. A., Gonzales, R. A. (eds.), "*Plant Cell Culture-A Practical Approach*". 2<sup>nd</sup> edition. IRL Press, Oxford: 1-25.
- [44] Fujita, Y., Hara, Y., Ogino, T., and Suga, C. 1981. Production of shikonin derivatives by cell suspension cultures of *Lithospermum erythrorhizon*. *Plant Cell Reports*, 1: 59-60.
- [45] Fujita, y., Takahashi, S., and Yamada, Y. 1984. Selection of cell lines with high productivity of shikonin derivatives through protoplasts of *Lithospermum erithrorhizon*. *Proceedings of European congress Biotechnology*, 1: 161-166.
- [46] Fukui, H., Tani, M., and Tabata, M. 1990. Induction of shikonin biosynthesis by endogenous polysaccharides in *Lithospermum erythrorhizon* cell suspension cultures. *Plant Cell Reports*, 9: 73-76.
- [47] Furuya, T. 1988. Saponins (ginseng saponins). In: Vasil, I.K. (Ed.), "*Cell Cultures and Somatic Cell Genetics of Plants*". 5, Academic Press, San Diego, CA: 213-234.
- [48] Furuya, T., Ikuta, A., and Syono, K. 1972. Alkaloids from callus cultures of *Papaver somniferum*. *Phytochemistry*, 11: 3041-3044.
- [49] Furuya, T., Kojima, H., Syono, K., Ishi, T., Uotani, K., and Nishio, M. 1973. Isolation of saponin and sapogenins from callus tissue of *Panax ginseng*. *Chemical & Pharmaceutical Bulletin*, 21: 98-101.
- [50] Furuya, T., Yoshikawa, T., Orihara, Y., and Oda, H. 1984. Studies of the culture conditions for *Panax ginseng* cells in jar fermentors. *Journal of Natural. Products*, 47: 70-75.
- [51] Gamborg, O. L., Miller, R. A., and Ojima, K. 1968. Nutrient requirements of suspension cultures of soyabean root cells. *Experimental Cell Ressearch*. 50: 151-158.
- [52] Gerasimenko, I., Sheludko, Y., and Stockigt, J. 2001. 3-Oxo-rhazinilam: A new indole alkaloid from *Rauwolfia serpentina* x *Rhazya stricta* hybrid plant cell cultures. *Journal of Natural Products*, 64: 114-116.
- [53] Goleniowski, M. and Trippi, V. S. 1999. Effect of growth medium composition on psilostachyinolides and altamisine production. *Plant Cell Tissue and Organ Culture*, 56: 215-218
- [54] Hagimori, M., Matsumoto, T., and Obi, Y. 1982. Studies on the production of *Digitalis* cardenolides by plant tissue culture. III. Effects of nutrients on digitoxin formation by shoot-forming cultures of *Digitalis purpurea* L. grown in liquid media. *Plant Cell Physiology*, 23, 7: 1205-1211.
- [55] Heble, M. R. and Staba, E. 1980. Steroid metabolism in stationary phase cell suspensions of *Dioscorea deltoidea*. *Planta Medica Supplement*: 124-128.
- [56] Holton, R. A., Somoza, C., Kim, H. B., Liang, F., Biediger, R J., Boatman, P. D., Shindo, M., Smith, C. C., Kim, S., Nadizadeh, H., Suzuki, Y., Tao, C., Vu, P., Tang, S., Zhang, P., Murthi, K. K., Gen-

- tile, L. N., and Liu, J. H. 1994. First total synthesis of taxol. 1. Functionalization of the B ring. *Journal of American Chemical Society*, 116: 1597-1600.
- [57] Hu, W. W., Yao, H. U., and Zhong, J. J. 2001. Improvement of *Panax notoginseng* cell cultures for production of ginseng saponin and polysaccharide by high-density cultivation of pneumatically agitated bioreactors. *Biotechnology Progress*, 17: 838-846.
- [58] Hu, Z. B. and Alfermann, A.W. 1993. Diterpenoid production in hairy root cultures of *Salvia miltiorrhiza*. *Phytochemistry*, 32: 699-703.
- [59] Huang, W. W., Cheng, C. C., Yeh, F. T., and Tsay, H. S. 1993. Tissue culture of *Dioscorea doryophora* HANCE 1. Callus induction from different source organs and the measurement of diosgenin content. *Chinese Medicinal College Journal*, 2, 2: 151-160.
- [60] Huang, S. H., Shen, Y. W., and Chan, H. S. 2002. Development of bioreactor operation strategy for L-DOPA production using *Stizolobium hassjoo* suspension culture. *Enzyme and Microbial Technology*, 30, 6: 779-791.
- [61] Ikuta, A. and Itokawa, H. 1988. Alkaloids of tissue cultures of *Nandina domestica*. *Phytochemistry*, 27, 7: 2143-2145.
- [62] Ishimaru, K., Arakawa, H., and Neera, S. 1993. Polyphenol production in cell cultures of *Cornus kousa*. *Phytochemistry*, 32, 5: 1193-1197.
- [63] Iwasa, K. and Takao, N. 1982. Formation of alkaloids in *Corydalis ophiocarpa* callus cultures. *Phytochemistry*, 21, 3: 611-614.
- [64] Jang, Y. P., Lee, Y. J., Kim, Y. C., and Huh, H. 1998. Production of a hepatoprotective cerebroside from suspension cultures of *Lycium chinense*. *Plant Cell Reports*, 18: 252-254.
- [65] Johnson, T., Ravishankar, G. A., and Venkataraman, L. V. 1990. In vitro capsaicin production by immobilized cells and placental tissues of *Capsicum annum* L. grown in liquid medium. *Plant Science*, 70: 223-229.
- [66] Karam, N. S., Jawad, F. M., Arikat, N. S., and Shibli, R. A. 2003. growth and rosmarinic acid accumulation in callus, cell suspension, and root cultures of wild *Salvia fruticosa*. *Plant Cell Tissue and Organ Culture*, 73: 117-121.
- [67] Ketchum, R. E. B., Rithner, C. D., Qiu, D., Kim, Y. S., Williams, R. M, and Croteau, R. B. 2003. *Taxus* metabolites: methyl jasmonates preferentially induces production of taxoids oxygenated at C-13 in *Taxus media* cell cultures. *Phytochemistry*, 62 (6): 901-909.
- [68] Khouri, H. E., Ibrahim, R. K., and Rideau, M. 1986. Effects of nutritional factors on growth and production of anthraquinone glucosides in cell suspension cultures of *Cinchona succirubra*. *Plant Cell Reports*, 5: 423-426.
- [69] Kitajima, M., Fischer, U., Nakamura, M., Ohsawa, M., Ueno, M., Takayama, H., Unger, M., Stockigt, J., and Aimi, N. 1998. Anthraquinone from *Ophiorrhiza pumila* tissue and cell cultures. *Phytochemistry*, 48, 1: 107-111.
- [70] Kitamura, Y., Ikenaga, T., Ooe, Y., Hirakawa, N., and Mizukami, H. 1998. Induction of furanocoumarin biosynthesis in *Glehnia littoralis* cell suspension cultures by elicitor treatment. *Phytochemistry*, 48, 1: 113-117.
- [71] Kobayashi, Y., Fukui, H., and Tabata, M. 1987. An immobilized cell culture system for berberine production by *Thalictrum minus* cells. *Plant Cell Reports*, 6: 185-186.
- [72] Koblitz, H., Koblitz, D., Schmauder, H. P., and Groger, D. 1983. Studies on tissue cultures of the genus *Cinchona* L. alkaloid production in cell suspension cultures. *Plant Cell Reports*, 2: 122-125.
- [73] Kueh, J. S. H., MacKenzie, I. A., and Pattenden, G. 1985. Production of chry-

- santhemiac acid and pyrethrins by tissue cultures of *Chrysanthemum cinerariaefolium*. *Plant Cell Reports*, 4: 118-119.
- [74] Lee, C. Y., Lin, F. L., Yang, C. T., Wang, L. H., Wei, H. L., and Tsay, H. S. 1995. Taxol production by cell cultures of *Taxus mairei*. Proceeding of Symposium. on development and utilization of resources of medicinal plants in Taiwan, Taiwan Agricultural Research Institute, Taiwan, April 21, 1995, TARI Special Publication 48: 137-148.
- [75] Lee, C. W. T. and Shuler, M. L. 2000. The effect of inoculum density and conditioned medium on the production of ajmalicine and catharanthine from immobilized *Catharanthus roseus* cells. *Biotechnology Bioengineering*, 67: 61-71.
- [76] Liu, K. C. S., Yang, S. H., Roberts, M. F., and Phillipson, J. D. 1990. Production of canthin-6-one alkaloids by cell suspension cultures of *Brucea javanica* (L.) Merr. *Plant Cell Reports*, 9: 261-263.
- [77] Lou, Z. C. and Chin, P. 1996. Longdan. In "*Species Systematization and Quality Evaluation of Commonly Used Chinese Traditional Drugs*". North-China edition, 3: 437-504. Beijing Medical University and Peking Union Medical College Press., Beijing, China.
- [78] Malpathak, N. P. and David, S. B. 1986. Flavor formation in tissue cultures of garlic (*Allium sativum* L.). *Plant Cell Reports*, 5: 446-447.
- [79] Mantell, S. H., Pearson, D. W., Hazell, L. P., and Smith, H. 1983. The effect of initial phosphate and sucrose levels on nicotine accumulation in batch suspension cultures of *Nicotiana tabacum* L. *Plant Cell Reports*, 2: 73-33.
- [80] Matsubara, K., Shigekazu, K., Yoshioka, T., Fujita, Y., and Yamada, Y. 1989. High density culture of *Coptis japonica* cells increases berberine production. *Journal of Chemical Technology and Biotechnology*, 46: 61-69.
- [81] Miyasaka, H., Nasu, M., and Yoneda, K. 1989. *Salvia miltiorrhiza*: In vitro production of cryptotanshinone and feruginol. In "*Biotechnology in agriculture and forestry*". 7, Medicinal and Aromatic Plants II ed. By Bajaj, Y.P.S. Springer-Verlag Berlin, Heidelberg: 417- 430.
- [82] Moreno, P. R. H., van der Heijden, R., and Verpoorte, R. 1993. Effect of terpenoid precursor feeding and elicitation on formation of indole alkaloids in cell suspension cultures of *Catharanthus roseus*. *Plant Cell Reports*, 12: 702-705.
- [83] Morimoto, S., Goto, Y., and Shoyama, Y. 1994. Production of lithospermic acid B and rosmarinic acid in callus tissue and regenerated plantlets of *Salvia miltiorrhiza*. *Journal of Natural Products*, 57, 6: 817-823.
- [84] Morimoto, H. and Murai, F. 1989. The effect of gelling agents on plaunotol accumulation in callus cultures of *Croton sublyratus* Kurz. *Plant Cell Reports*, 8: 210-213.
- [85] Mulder-Krieger, T., Verpoorte, R., Svendse, A., and Scheffer, J. 1988. Production of essential oils and flavours in plant cell and tissue cultures. A review. *Plant Cell Tissue and Organ Culture*, 13: 85-114.
- [86] Murashige, T. and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- [87] Nakagawa, K., Fukui, H., and Tabata, M. 1986. Hormonal regulation of berberine production in cell suspension cultures of *Thalictrum minus*. *Plant Cell Reports*, 5: 69-71.
- [88] Nakao, K., Ono, K., and Takio, S. 1999. The effect of calcium on flavanol production in cell suspension cultures of *Polygonum hydropiper*. *Plant Cell Reports*, 18: 759-763.
- [89] Neera, S. and Ishimaru, K. 1992. Tannin production in cell cultures of *Sapium sebiferum*. *Phytochemistry*, 31, 3: 833-836.
- [90] Nicolaou, K.C., Yang, Z., Liu, J. J., Ueno,

- H., Nantermet, P. G., Guy, R. K., Claiborne, C.F. 1994. Renaud, J., Couladouros, E.A., Paulvannan, K., and Sorensen, E. J. Total synthesis of taxol. *Nature*, 367: 630-634.
- [91] Nazif, N. M., Rady, M. R., and Seif E1-Nasr, M. M. 2000. Stimulation of anthraquinone production in suspension cultures of *Cassia acutifolia* by salt stress. *Fitoterapia*, 71: 34-40.
- [92] O' Dowd, N., McCauley, P. G., Richardson, D. H. S., and Wilson, G. 1993. Callus production, suspension culture and in vitro alkaloid yields of *Ephedra*. *Plant Cell Tissue and Organ Culture*, 34: 149-155.
- [93] Orihara, Y. and Furuya, T. 1990. Production of theanine and other  $\gamma$ -glutamyl derivatives by *Camellia sinensis* cultured cells. *Plant Cell Reports*, 9: 65-68.
- [94] Orihara, Y., Yang, J. W., Komiya, N., Koge, K., and Yoshikawa, T. 2002. Abietane diterpenoid from suspension cultured cells of *Torreya nucifera* var. *radicans*. *Phytochemistry*, 59: 385-389.
- [95] Petit-Paly, G., Montagu, M., Viel, C., Rideau, M., and Chenieux, J. 1987. Dihydrofuro [2,3-b] quinolinium alkaloids in cultured cells of *Ptelea trifolia* L. *Plant Cell Reports*, 6: 309-312.
- [96] Pletsch, M., Piacente, S., Pizza, C., and Charlwood, B. V. 1993. The accumulation of phenylpropanoid glycosides in tissue cultures of *Tecoma sambucifolium*. *Phytochemistry*, 34 (1): 161-165.
- [97] Rajasekaran, T., Rajendran, L., Ravishankar, G. A., and Venkataraman, L. V. 1991. Influence of nutrient stress on pyrethrin production by cultured cells of pyrethrum (*Chrysanthemum cinerariaefolium*). *Current Science*, 60, 12: 705-707.
- [98] Ramachandra Rao, S. and Ravishankar, G. A. 2002. Plant cell cultures: Chemical factories of secondary metabolites. *Biotechnology Advances*, 20: 101-153.
- [99] Ramirez, M., Alpizar, L., Quiroz, J., and Oropeza, C. 1992. Formation of L-canavanine in *in vitro* cultures of *Canavalia ensiformis* (L) DC. *Plant Cell Tissue and Organ Culture*, 30: 231-235.
- [100] Ramachandra Rao, S. 2000. Biotechnological production of phyto-pharmaceuticals. *Journal of Biochemistry Molecular Biology Biophysics*, 4: 73-102.
- [101] Ravishankar, G. A. and Venkataraman, L. V. 1993. Role of plant cell cultures in food biotechnology: commercial prospectus and problems. New Delhi: Oxford IBH Press: 255-274.
- [102] Rech, S. B., Batista, C. V. F., Schripsema, J., Verpoorte, R., and Henriques, A. T. 1998. Cell cultures of *Rauwolfia sellowii*: growth and alkaloid production. *Plant Cell Tissue and Organ Culture*, 54: 61-63.
- [103] Roberts, S. C. and Shuler, M. L. 1997. Large scale plant cell culture. *Current opinion in Biotechnology*, 8: 154-159.
- [104] Rodriguez, S., Wolfender, J. L., Hostettmann, K., Odontuya, G., and Purev, O. 1996. Corniculoside, a new biosidic ester secoiridoid from *Halemia corniculata*. *Helvetica Chimica Acta*, 79: 363-370.
- [105] Romagnoli, L. G. and Knorr, D. 1988. Effects of ferulic acid treatment on growth and flavour development of cultured *Vanilla planifolia* cells. *Food Biotechnology*, 2: 93-104.
- [106] Sakuta, M., Takagi, T., and Komamine, A. 1987. Effects of nitrogen source on betacyanin accumulation and growth in suspension cultures of *Phytolacca americana*. *Physiologia Plantarum*, 71: 459-463.
- [107] Salgado, Garciglia, R., and Ochoa-Alejo, N. 1990. Increased capsaicin content in PFP resistant cells of chilli pepper (*Capsicum annum*). *Plant Cell Reports*, 8: 617-620.
- [108] Sasse, F., Heckenberg, U., and Berlin, J. 1982. Accumulation of  $\beta$ -carboline alkaloids and serotonin by cell cultures of



- Peganum harmala L. *Plant Physiology*, 69: 400-404.
- [109] Schripsema, J., Ramos Valdivia, A., and Verpoorte, R. 1999. Robustaquinones, novel anthraquinones from an elicited *Cinchona robusta* suspension culture. *Phytochemistry*, 51: 55-60.
- [110] Schroder, W. and Bohm, H. 1984. Betacyanin concentrations in young cell cultures from *Portulaca grandiflora* an analysis of variation. *Plant Cell Reports*, 3: 14-17.
- [111] Scragg, A. H. and Allan, E. J. 1986. Production of triterpenoid quassin and cell suspension cultures of *Picrasana quassioides* Bennett. *Plant Cell Reports*, 5: 356-359.
- [112] Sejourne, M., Viel, C., Bruneton, J., Rideau, M., and Chenieux, J. C. 1981. Growth and furoquinoline alkaloid production in cultured cells of *Choisya ternata*. *Phytochemistry*, 20, 2: 353-355.
- [113] Shimomura, K., Kitazawa, T., Okamura, N., and Yagi, A. 1991. Tanshinone production in adventitious roots and regenerates of *Salvia miltiorrhiza*. *Journal of Natural Products*, 54: 1583.
- [114] Siah, C. L. and Doran, P. M. 1991. Enhanced codeine and morphine production in suspended *Papaver somniferum* cultures after removal of exogenous hormones. *Plant Cell Reports*, 10: 349-353.
- [115] Sierra, M. I., Heijden, R., Leer, T., and Verpoorte, R. 1992. Stability of alkaloid production in cell suspension cultures of *Tabernaemontana divaricata* during long-term subculture. *Plant Cell Tissue and Organ Culture*, 28: 59-68.
- [116] Silvestrini, A., Pasqua, G., Botta, B., Monacelli, B., van der Heijden, R., and Verpoorte, R. 2002. Effect of alkaloid precursor feeding on a *Camptotheca acuminata* cell line. *Plant Physiology and Biochemistry*, 40: 749-753.
- [117] Skrzypczak, L., Wesolowska, M., and Skrzypczak, E. 1993. *Gentiana* species XII: In vitro culture, regeneration, and production of secoirridoid glucosides. In: Bajaj, Y.P.S, editor. "Biotechnology in Agriculture and Forestry". 21, Medicinal and aromatic plants IV. Berlin, Heidelberg: Springer-Verlag: 172-186.
- [118] Slichenmyer, W. J. and Von Horf, D. D. 1991. Taxol: a new and effective anti-cancer drug. *Anti-Cancer Drugs*, 2: 519-530.
- [119] Stojakowska, A. and Kisiel, W. 1999. Secondary metabolites from a callus culture of *Scutellaria columnae*. *Fittoterapia*, 70: 324-325.
- [120] Tabata, M. and Hiraoka, N. 1976. Variation of alkaloid production in *Nicotiana rustica* callus cultures. *Physiology Plantarum* 38: 19-23.
- [121] Tabata, M., Yamamoto, H., Hiraoka, N., and Konoshima, M. 1972. Organization and alkaloid production in tissue cultures of *Scopolia parviflora*. *Phytochemistry*. 11: 949-955.
- [122] Takahashi, D. and Fjita, Y. 1991. In: "Plant Cell Culture in Japan". Komamine, A., Misawa, M., and Dicosmo, F., editors. Cosmetic materials: 72-78.
- [123] Tanahashi, T. and Zenk, M.H. 1985. Isoquinoline alkaloids from cell suspension cultures of *Fumaria capreolata*. *Plant Cell Reports*, 4: 96-99.
- [124] Taniguchi, S., Imayoshi, Y., Kobayashi, E., Takamatsu, Y., Ito, H., Hatano, T., Sakagami, H., Tokuda, H., Nishino, H., Sugita, D., Shimura, S., and Yoshida, T. 2002. Production of bioactive triterpenes by *Eriobotrya japonica* calli. *Phytochemistry*, 59: 315-323.
- [125] Thengane, S. R., Kulkarni, D. K., Shrikhande, V. A., Joshi, S. P., Sonawane, K. B., and Krishnamurthy, K. V. 2003. Influence of medium composition on callus induction and camptothecin(s) accumulation in *Nothapodytes foetida*. *Plant Cell Tissue and Organ Culture*, 72: 247-251.
- [126] Tsay, H. S. 1999. Tissue culture technology of medicinal herbs and its appli-

- cation of medicinal herbs and its application in Taiwan. In: Chou, C.H., G.R.Waller, C. Reinhardt (eds), Biodiversity and allelopathy: from organisms to ecosystems in The Pacific. Academia Sinica, Taipei, Taiwan: 137-144.
- [127] Tsay, H. S., Chang, W. D., Chen, C. C., and Chang, Y. S. 1994. The production of imperatorin from *Angelica dahurica* var. *formosana* by cell suspension culture. *Journal of Agricultural Association China*, 168: 32-48.
- [128] Tsukamoto, T., Yeno, Y., and Ohta, Z. 1936. Glycosides of *Dioscorea tokoro* I. Diocin, dioscoreasapooxin and diosgenin. *Journal of Pharmaceutical Society Japan*, 56: 135.
- [129] Uden, W., Pras, N. N., Visser, J. F., and Malingre, T. M. 1989. Detection and identification of Podophyllotoxin produced by cell cultures derived from *Podophyllum hexandrum* royle. *Plant Cell Reports*, 8: 165-168.
- [130] Uden, W., Pras, N., Vosseveld, E. M., Mol, J. N. M., and Malingre, T. M. 1990. Production of 5 methoxypodophyllotoxin in cell suspension cultures of *Linum flavum* L. *Plant Cell Tissue and Organ Culture*, 20: 81-87.
- [131] Ulbrich, B., Weisner, W., and Arens, H. 1985. In primary and secondary metabolism of plant cell cultures, (Neumann, K. H. and Reinhard, E., editors), Springer-Verlag (Berlin): 293-303.
- [132] Venkateswara, R., Sankara Rao, S., and Vaidyanathan, C. S. 1986. Phytochemical constituents of cultured cells of *Eucalyptus tereticornis* SM. *Plant Cell Reports*, 3: 231-233.
- [133] Venkateswara, R., Sankara Rao, K., and Vaidyanathan, C. S. 1987. Cryptosin-a new cardenolide in tissue culture and intact plants of *Cryptolepis buchanani* Roem. & Schult. *Plant Cell Reports*, 6: 291-293.
- [134] Verpoorte, R. 1998. Exploration of nature's chemodiversity: the role of secondary metabolites as leads in drug development. *Drug Discovery Today*, 3: 232-238.
- [135] Villarreal, M. L., Arias, C., Feria-Velasco, A., Ramirez, O. T., and Quintero, R. 1997. Cell suspension culture of *Solanum chrysotrichum* (Schldl.) - A plant producing an antifungal spirostanol saponin. *Plant Cell Tissue and Organ Culture*, 50: 39-44.
- [136] Waller, G. R., MacVean, C. D., and Suzuki, T. 1983. High production of caffeine and related enzyme activities in callus cultures of *Coffea arabica* L. *Plant Cell Reports*, 2: 109-112
- [137] Wang, P. J. and Huang, C. I. 1982. Production of saikosaponins by callus and redifferentiated organs of *Bupleurum falatum* L. In: Fujiwara A. (ed.) *Plant Tissue Culture*. Maruzen, Tokyo: 71-72.
- [138] Wang, J., Liao, X., Zhang, H., Du, J, and Chen, P. 2003. Accumulation of chlorogenic acid in cell suspension cultures of *Eucommia ulmoides*. *Plant Cell Tissue and Organ Culture*, 74: 193-195.
- [139] Wang, H. Q., Yu, J. T. and Zhong, J. J. 1999. Significant improvement of taxane production in suspension cultures of *Taxus chinensis* by sucrose feeding strategy. *Process Biochemistry*, 35: 479-483.
- [140] Wang, Z. Y. and Zhong, J. J. 2002a. Combination of conditioned medium and elicitation enhances taxoid production in bioreactor cultures of *Taxus chinensis* cells. *Biochemical Engineering Journal*, 12: 93-97.
- [141] Wang, W. and Zhong, J. J. 2002b. Manipulation of ginsenoside heterogeneity in cell cultures of *Panax notoginseng* by addition of jasmonates. *Journal of Biosciences Bioengineering*, 93: 154-159.
- [142] Wani, M. C., Taylor, H. L., Wall, M. E., Coggon, P., and McPhail, A. T. 1971. Plant antitumor agents VI. The isolation and structure of taxol, a novel antileukemic and antitumor agent from *Taxus*

- brevifolia. *Journal of American Chemical Society*, 93: 2325-2327.
- [143] Wataneba, K., Yano, S. Y., and Yamada, Y. 1982. Selection of cultured plant cell lines producing high levels of biotin. *Phytochemicals*, 21: 513-516.
- [144] Whitmer, S., Canel, C., Hallard, D., Goncalves, C. and Verpoorte, R. 1998. Influence of precursor availability on alkaloid accumulation by transgenic cell lines of *Catharanthus roseus*. *Plant Physiology*, 116: 853-857.
- [145] Wichers, H J., Visser, J. F., Huizing, H. J., and Pras, N. 1993. Occurrence of L-DOPA and dopamine in plants and cell cultures of *Mucuna pruriens* and effects of 2, 4-D and NaCl on these compounds. *Plant Cell Tissue and Organ Culture*, 33: 259-264.
- [146] Wickremesinha, E. R. M. and Arteca, R. N. 1993. *Taxus* callus cultures: Initiation, growth optimization, characterization and taxol production. *Plant Cell Tissue and Organ Culture*, 35: 181-193.
- [147] Wickremesinha, E. R. M. and Arteca, R. N. 1994. *Taxus* cell suspension cultures: optimizing growth and production of taxol. *Journal of Plant Physiology*, 144: 183-188.
- [148] Widholm, J. M. 1974. Evidence for compartmentation of tryptophan in cultured plant tissues. Free tryptophan levels and inhibition of anthranilate synthetase. *Physiology Plantarum*, 30: 323-326.
- [149] Wijnsma, R., Go, J. T. K. A., Weerden, van I. N., Harkes, P. A. A., Verpoorte, R., and Svendsen, A. B. 1985. Anthraquinones as phytoalexins in cell and tissue cultures of *Cinchona* sp. *Plant Cell Reports*, 4: 241-244.
- [150] Wu, J., Wang, C., and Mei, X. 2001. Stimulation of taxol production and excretion in *Taxus* spp cell cultures by rare earth chemical lanthanum. *Journal of Biotechnology*, 85: 67-73.
- [151] Wu, C. T., Vanisree M., Satish M. N., Chen, C. L., Yang, T. F, and Tsay, H.S. 2003. Isolation and quantitative analysis of cryptotanshinone, an active quinoid diterpene formed in the callus of *Salvia miltiorrhiza* Bunge. *Biological and Pharmaceutical Bulletin*, 26, 6: 845-848.
- [152] Yagi, A., Shoyama, Y., and Nishioka, I. 1983. Formation of tetrahydroanthracene glucosides by callus tissue of *Aloe saponaria*. *Phytochemistry*, 22, 6: 1483-1484.
- [153] Yamada, Y. and Endo, T. 1984. Tropane alkaloids in cultured cells of *Duboisia leichhardtii*. *Plant Cell Reports*, 3: 186-188.
- [154] Yamada, Y. and Hashimoto, T. 1982. Production of tropane alkaloids in cultured cells of *Hyoscyamus niger*. *Plant Cell Reports*, 1: 101-103.
- [155] Yamada, Y, and Sato, F. 1981. Production of berberine in cultured cells of *Coptis japonica*. *Phytochemistry*, 20: 545-547.
- [156] Yamamoto, Y., Mizuguchi, R., and Yamada, Y. 1982. Selection of a high and stable pigment-producing strain in cultured *Euforbia millii* cells. *Theoretical and Applied Genetics*, 61: 113-116.
- [157] Yamamoto, O. and Yamada, Y. 1986. Production of reserpine and its optimization in cultured *Rauwolfia serpentina* Benth. cells. *Plant Cell Reports*, 5: 50-53.
- [158] Yeh, F. T., Huang, W. W., Cheng, C. C., Na, C., and Tsay, H. S. 1994. Tissue culture of *Dioscorea doryophora* Hance. II. Establishment of suspension culture and the measurement of diosgenin content. *Chinese Agronomy Journal*, 4: 257-268.
- [159] Zenk, M. H., El-Shagi, H., and Schulte, U. 1975. Anthraquinone production by cell suspension cultures of *Morinda citrifolia*. *Planta Medica Supplement*: 79-101.
- [160] Zhao, J., Zhu, W., and Hu, Q. 2001. Enhanced catharanthine production in *Catharanthus roseus* cell cultures by

- combined elicitor treatment in shake flasks and bioreactors. *Enzyme and Microbial Technology*, 28: 673-681.
- [161] Zheng, H. Z., Dong, Z. H, and She, J. 1997. Longdan. In *Modern Study of Traditional Chinese Medicine*, 2: 1398-1408. Beijing Xue Yuan Press., Beijing, China.
- [162] Zhong, J. J. and Zhu, Q. X. 1995. Effect of initial phosphate concentration on cell growth and ginsenoside saponin production by suspended cultures of *Panax notoginseng*. *Applied Biochemistry Biotechnology*, 55: 241-246.
- [163] Zhou, G. M. 1980. Studies on useful compounds of Bai-Zhi for healing Yin-Hsieh Ping. *Chung-Chen-Yau Research*, 4: 33. (in Chinese)
- [164] Zhou, G. M., Yu, C. G., Han, Y. C., and Mun, C. T. 1988. Studies on Bai-Zhi. IV. The toxicity test of useful compounds. *Medical Journal of China Hospital*, 8: 220-221. (in Chinese)