

Plant nutrient acquisition and utilisation in a high carbon dioxide world

T. R. Cavagnaro^{A,C}, R. M. Gleadow^A and R. E. Miller^{A,B}

^ASchool of Biological Sciences, Monash University, Clayton, Vic. 3800, Australia.

^BThe Australian Centre for Biodiversity, Monash University, Clayton, Vic. 3800, Australia.

^CCorresponding author. Email: timothy.cavagnaro@monash.edu

Abstract. Producing enough food to meet the needs of an increasing global population is one of the greatest challenges we currently face. The issue of food security is further complicated by impacts of elevated CO₂ and climate change. In this viewpoint article, we begin to explore the impacts of elevated CO₂ on two specific aspects of plant nutrition and resource allocation that have traditionally been considered separately. First, we focus on arbuscular mycorrhizas, which play a major role in plant nutrient acquisition. We then turn our attention to the allocation of resources (specifically N and C) *in planta*, with an emphasis on the secondary metabolites involved in plant defence against herbivores. In doing so, we seek to encourage a more integrated approach to investigation of all aspects of plant responses to eCO₂.

Additional keywords: arbuscular mycorrhizas, cyanogenesis, elevated CO₂, food security, nitrogen, secondary metabolism.

Introduction

Plant nutrient demand, acquisition and utilisation are all altered by changes in atmospheric CO₂ concentrations. In the context of looming fertiliser shortages (Cordell *et al.* 2009) and increasing global demand for food (Rosegrant and Cline 2003), we need to understand how all facets of plant nutrient dynamics will respond to a changing climate. In this viewpoint article, we focus on the impacts of elevated atmospheric CO₂ (eCO₂) on two specific aspects of plant nutrients that have traditionally been considered separately: acquisition and utilisation.

First, we consider the impacts of eCO₂ on the formation and functioning of arbuscular mycorrhizas (AM), which play a major role in plant nutrient acquisition. Second, we discuss the impacts of eCO₂ on plant allocation of resources (specifically N and C) to secondary metabolites involved in defence against herbivores. Finally, we begin to explore how these two quite different aspects of plant biology may be linked, and conclude with several readily testable hypotheses with a view to stimulating further work in this area. In so doing, we seek to better understand where our own areas of research can inform one another and to encourage a more integrated approach to investigating all aspects of plant responses to eCO₂.

Impacts of eCO₂ on arbuscular mycorrhizas

In this first section, we outline the benefits to plants of forming AM, in terms of nutrient acquisition, and discuss the impact of eCO₂ on acquisition of nutrients by AM and on C supply to the fungi. We also discuss how AM may be relatively more important with rising atmospheric CO₂. The simple model in Fig. 1 summarises the potential role of AM in nutrient supply and as

a C sink in a high CO₂ world, and forms a reference point for this section.

Nutrient acquisition by AM under eCO₂

Arbuscular mycorrhizal associations are formed by the majority of terrestrial plants, including most crop species, with a specialised group of soil fungi. They play important roles in plant nutrition by providing access to soil-derived nutrients not necessarily otherwise accessible to roots (Smith and Read 2008). To date, the majority of research, both under ambient and eCO₂ conditions, has focussed on the role of AM in plant P nutrition. This is not surprising, given that in some crops, up to 100% of plant P can be delivered via the AM fungal pathway (Smith *et al.* 2004). However, AM also play an important, albeit less well studied, role in the acquisition of other essential nutrients, including N, Zn and others (Cavagnaro 2008). With global P supply expected to peak in coming decades (Cordell *et al.* 2009), we predict that the AM contribution to plant P nutrition in agricultural systems will become increasingly important. Furthermore, given the importance of both N (e.g. Daepf *et al.* 2001) and P supply (e.g. Campbell and Sage 2006) in determining the response of plants to eCO₂, it is essential that we understand how the acquisition of nutrients by AM is likely to be affected by eCO₂.

It is well established that growth of C₃ plants usually increases under eCO₂, primarily because the current daytime atmospheric CO₂ concentration (approximately 380 ppm; IPCC 2007) is still suboptimal for Rubisco, the enzyme that catalyses carbon fixation (Stitt and Krapp 1999; Ainsworth and Rogers 2007). Critically, this increase in C₃ plant growth under eCO₂ is dependent on access to sufficient nutrients. In a meta-analysis of the growth




Fig. 1. Potential links between $e\text{CO}_2$, agricultural management, AM and plant nutrient acquisition. NB: Emphasis is on nutrient uptake and supply to (C_3) plants, not C supply to the fungi. (1) Environmental factors aside from $e\text{CO}_2$, e.g. edaphic, temperature, etc. (2) Tillage, nutrient management, crop rotations (including fallow). (3) AM function in this example refers to increased nutrient uptake and delivery to the plant. (4) Looming P shortage, limited access due to increased cost. (5) Impact on crop yields, nutrient content or both.

responses of herbaceous and woody species to $e\text{CO}_2$, for example, the $e\text{CO}_2$ stimulation of aboveground biomass by +20.1% was more than halved to +8.8% when nitrogen supply was limited (de Graaff *et al.* 2006). A range of other studies across grasses, crops and other functional groups report similar nutrient (N) constraints on the magnitude of the $e\text{CO}_2$ response (e.g. Wand *et al.* 1999; Nowak *et al.* 2004; Reich *et al.* 2006; Jackson *et al.* 2008). Phosphorus supply can also limit the stimulation of photosynthesis and growth by $e\text{CO}_2$; indeed, P limitation may be more critical in some systems (Edwards *et al.* 2005). Phosphorus constraints on the response to $e\text{CO}_2$ have been reported in trees and grasses (Conroy *et al.* 1992; Stöcklin *et al.* 1998), and have been widely reported among legumes (Grünzweig and Körner 2003; Edwards *et al.* 2005; Rogers *et al.* 2009), many of which form AM.

Due to greater plant biomass and carbohydrate accumulation, plants grown at $e\text{CO}_2$ almost always have lower concentrations of nutrients such as P and N (Treseder and Allen 2000; Jifon *et al.*

2002). Lower concentrations of P in the tissues of plants when grown under $e\text{CO}_2$ have been shown to be alleviated, to varying extents, by the formation of AM in some, but not all, cases (Gavito *et al.* 2002; Gavito *et al.* 2003; Cavagnaro *et al.* 2007). For example, plant P concentrations of non-mycorrhizal black locust seedlings (*Robinia pseudoacacia* L.) declined by 50% under $e\text{CO}_2$, whereas in mycorrhizal seedlings, the decline in P concentration was 22% (Olesniewicz and Thomas 1999). In a study of two *Citrus* spp., AM colonisation under $e\text{CO}_2$ stimulated plant growth and P acquisition in the more mycorrhizal-dependent of the species but not in the other (Syvertsen and Graham 1999). The effect of $e\text{CO}_2$ on AM colonisation and plant P nutrition may also vary over time (Rouhier and Read 1998; Staddon *et al.* 1999). Interestingly, in a study in *Pisum sativum* L. under $e\text{CO}_2$, the greater shoot P content and concentration of plants with AM compared with non-mycorrhizal plants could not be attributed to hyphal P uptake (Gavito *et al.* 2002). We suggest that important insights into AM and P nutrition under $e\text{CO}_2$ are

likely to come where the relative contributions of both root and fungal uptake are considered, as has been done under ambient CO₂ conditions (Smith *et al.* 2004).

As indicated previously, equally important although much less well understood, are the impacts of eCO₂ on plant acquisition of other nutrients such as N via the AM pathway. There have been fewer studies on AM-mediated N uptake both under ambient CO₂ and eCO₂. Arbuscular mycorrhizal fungi (AMF) can acquire N both as NO₃⁻ and NH₄⁺, and transfer it to plants (Ames *et al.* 1983; Johansen *et al.* 1993). Uptake of N in its more mobile form NO₃⁻ by AMF is likely to be important in agricultural soils, where NO₃⁻ is often the dominant form of N in the soil (Bloom *et al.* 2010), but will also be important in dry soils, as limited water can interfere with NO₃⁻ supply to the roots (via mass flow) (Smith *et al.* 1985; Tobar *et al.* 1994; Allen *et al.* 2003). While AMF can take up both NH₄⁺ and NO₃⁻, it has been shown that they can take up and deliver to the plant up to 10 times as much N when supplied as NH₄⁺ (Tanaka and Yano 2005). Provision of N as NH₄⁺ could become increasingly important in the future, as recent studies indicate that eCO₂ inhibits NO₃⁻ assimilation in leaves (Bloom *et al.* 2010).

Any improvements in plant N nutrition resulting from the formation of AM at eCO₂ are likely to be important, but to date only a handful of studies have investigated this directly. In one study, AMF were found to improve plant N acquisition, after a period of selection of the fungal community over several generations under eCO₂ concentrations (Gamper *et al.* 2005). Similarly, in a study on tomatoes (*Solanum lycopersicum* L.), positive mycorrhizal effects on plant N content were greater under eCO₂ than under ambient CO₂ conditions, although differences were small (Cavagnaro *et al.* 2007). Chen *et al.* (2007) found that the effects of mycorrhizal colonisation on N uptake under eCO₂ were species-specific, with N uptake enhanced by colonisation under eCO₂ in *Plantago lanceolata* L., but not in *Festuca arundinacea* Schreb. These three examples serve to highlight the potential for AMF to improve plant N acquisition under eCO₂; there is a need for further dedicated investigations in this area. The assessment of plant nutritional changes due to AM under eCO₂ may be complicated to some extent by the different ways in which plant nutrient status is reported, e.g. total nutrient content per plant or total nutrient uptake versus nutrient concentration. While total nutrient uptake may reflect AM function and be more commonly reported in the AM literature, tissue nutrient concentrations may be more important for plant nutritional value. In addition, most studies of the role of AM in plant nutrition focus on changes in nutrient concentrations (or contents); they rarely consider the fate of those nutrients *in planta*. In particular, the fate of acquired N has not been investigated in studies of AM, despite the implications for food security (see below).

Demand for C resources by AM

Producing and supporting AM comes at a cost to plants because C is required to both establish and maintain the association with the fungi. It has been hypothesised that the relative cost of forming AM is likely to decrease at eCO₂ (Treseder and Allen 2000; Jifon *et al.* 2002). This is essentially because the fungi rely solely upon the plant for their C supply; thus any increase in C assimilation and photosynthate supply to roots under eCO₂ is predicted to reduce

the relative cost of the association to the plant (Lovelock *et al.* 1997). Consistent with this, the majority of studies in which AMF biomass is measured (e.g. as extra-radical hyphal length per g soil) report either significant increases in AMF biomass or a neutral response, with no reported decreases in AM biomass under eCO₂ to date (Treseder 2004; Hu *et al.* 2006; Lukac *et al.* 2010). By contrast, while most studies under eCO₂ similarly report either no change or increases in the most common measure of AM formation, the percent root colonised by AMF (Hartwig *et al.* 2002; Lukac *et al.* 2003; Gamper *et al.* 2004; Treseder 2004; Hu *et al.* 2006), decreases in percent root colonisation have also been reported (Rillig and Allen 1999; Olsrud *et al.* 2010). It is important to recognise that considering AM responses to eCO₂ in terms of AM colonisation of roots alone is likely to be an oversimplification, since the extent of colonisation is a function of both fungal and root growth (Alberton *et al.* 2005; Smith and Read 2008). Consequently, when the well-established stimulatory effects of eCO₂ on root growth are taken into account, differences in colonisation are often eliminated (Alberton *et al.* 2005). Further, changes in percent root colonisation may not be the best measure of mycorrhizal response to eCO₂ as percent root colonisation (1) does not necessarily correlate with nutrient transfer to the plant (Smith and Read 2008), (2) may underestimate the stimulatory effect of eCO₂ on the AMF as the fungal biomass response to eCO₂ is typically greater than any increase in root colonisation (Staddon *et al.* 2004; Treseder 2004; Hu *et al.* 2006), and (3) may be found not to change, despite significant eCO₂ effects on a range of other measures of AM such as the intensity of colonisation (i.e. hyphal number per root length) or the frequency and abundance of arbuscules and vesicles (Rillig *et al.* 1998; Gamper *et al.* 2004).

AM have both intra- and extra-radical phases of colonisation. The intra-radical phase involves fungal penetration of the root epidermis, and growth between and within the cortical cells, where arbuscules and hyphal coils, across which nutrients are transferred, are formed (Smith and Read 2008). Since nutrients are transferred from the fungi to the plant across the arbuscular interface, impacts on arbuscule production may be more informative than impacts on total colonisation. Arbuscules themselves exhibit variable responses to eCO₂ (Klironomos *et al.* 1998); however, given their rapid rates of turnover, this is perhaps not unexpected. The impacts of eCO₂ on the extra-radical phase of colonisation, which involves the growth of the fungi into the soil beyond the roots, will also be important (Rillig 2004; Collins-Johnson *et al.* 2005), especially where function is also taken into account. For example, in one study, an increase in hyphal production under eCO₂ did not result in an increase in P uptake (Sanders *et al.* 1998).

Further studies done at finer spatial scales (Cavagnaro *et al.* 2001; Dickson 2004) and using multiple time points (Garcia *et al.* 2008) will be important in determining the impact of eCO₂ on both the intra- and extra-radical phases of colonisation. Ideally, we should move beyond simple counts of percent root colonisation (Alberton *et al.* 2005) and embrace the information afforded by (1) advanced microscopic techniques to give measurements of changes in the area of symbiotic interface (Bago *et al.* 1998; Dickson and Kolesik 1999), (2) visualisation of metabolically active and inactive fungal biomass (Dickson and Smith 2001; van Aarle *et al.* 2005), and (3) genomic and

proteomic analyses of nutrient transport processes (Glassop *et al.* 2005; González-Guerrero *et al.* 2005, 2007). All these methods, which are more costly and time-consuming than quantification of the percentage of the root length colonised by the fungi, would also need to be verified against both the rate of nutrient acquisition and impact on plant growth and development. Given that AM exhibit a considerable functional diversity in terms of their impacts on plant growth and nutrition (Smith *et al.* 2004; Cavagnaro *et al.* 2005), and the diversity in the eCO₂ response of plant species from different functional groups (e.g. C₃, C₄, legumes, and woody or herbaceous) there is a need to study the responses of a wide variety of plant–fungal combinations under eCO₂ and differing nutrient supply (Alberton *et al.* 2005; Collins-Johnson *et al.* 2005), using this wider spectrum of techniques. This, however, is only half of the story; once nutrients have been acquired, it is equally as important to consider their fate *in planta*.

Impacts of eCO₂ on the nutritive value of plants and chemical defence

In this second section, we move our attention to the fate of nutrients (especially N) once they have been taken up or delivered to plants. Once plants acquire nutrients, irrespective of the mechanisms involved, they are incorporated into a wide array of primary and secondary metabolites, including nucleic acids, amino acids and secondary metabolites. Many secondary metabolites play an important role in defence, acting to protect plants from losing valuable biomass and nutrients to herbivores, including humans (Agrawal and Fishbein 2006). Any change in allocation in resources between the different metabolites within the plant is important to us because the nutritional value of a plant as food depends on their relative proportions (Gleadow and Woodrow 2002; Lindroth 2010). When plants are grown under eCO₂, the abundance of carbohydrates and the reduced demand for leaf N by the photosynthetic apparatus (Stitt and Krapp 1999) allows plants to invest in both resource acquisition (i.e. roots) and defence (Lambers 1993; Coley *et al.* 2002). Changes in not only N content and allocation under eCO₂ but also in P, P : C ratios and C : N : P ratios may also affect plant yield and nutritional status (Loladze 2002). Thus, the impacts of eCO₂ on plant stoichiometry are expected to change the economics of the plant defence equation and, therefore, their nutritional value (Lambers 1993; Coley *et al.* 2002; Bidart-Bouzat and Imeh-Nathaniel 2008). Consistent with the view of Bennett *et al.* (2006), we propose that the formation of AM may further alter the economics of this equation through their provisioning of mineral nutrients to plants.

The economic trade-off between investment in primary production and defence depends on the availability of input resources (including C : N ratios), the efficiency of the metabolic apparatus, the developmental stage of the plant and the value of the particular organ for growth and reproduction (see Fig. 2; Gleadow and Woodrow 2000; Boege and Marquis 2005; Cornelissen *et al.* 2008; Haugen *et al.* 2008). Chemical defences are often classified as C- and N-containing secondary metabolites (CCSMs and NCSMs) (Rhoades 1979). CCSMs (e.g. phenolics and tannins) act by reducing the digestibility of plant tissue (Agrawal and Fishbein 2006). While few studies have examined the effect of AM on foliar secondary metabolites,

one study under ambient CO₂ found that AM colonisation of *Plantago lanceolata* increased concentrations of C-based defences (Gange and West 1994), suggesting that AM can affect changes in CCSMs. Despite species specificity in the eCO₂ effects on CCSMs (Lindroth 2010), leaves of plants grown at eCO₂ tend to contain higher concentrations of some CCSMs (particularly tannins and some phenolics, but not terpenoids) in proportion to the higher C : N ratio (Lincoln *et al.* 1993; Lawler *et al.* 1997; Coley *et al.* 2002; Agrawal and Fishbein 2006). The increased synthesis of CCSMs may be particularly pronounced in plants that are also N-limited, consistent with the higher C : N ratio (Peñuelas and Estiarte 1998). The formation of AM may further alter this balance (C : N) under eCO₂ through demand for C and supply of N.

NCSMs (e.g. cyanogenic glycosides, glucosinolates, alkaloids) are usually toxic and act in lower concentrations than CCSMs (Gleadow and Woodrow 2002). In general, the effects of eCO₂ on N-based defences have received little research attention (Gleadow *et al.* 2009c; Lindroth 2010). Given the lower concentration of tissue N in C₃ plants grown at eCO₂, it was originally hypothesised that the concentration of NCSMs would be lower under eCO₂ (Lambers 1993). Experimentally, however, the concentration of NCSMs has been found to be higher in many plants grown under eCO₂ either on a mass basis (Ziska *et al.* 2005; Matros *et al.* 2006; Bidart-Bouzat and Imeh-Nathaniel 2008; Gleadow *et al.* 2009a, 2009b) or relative to plant protein (e.g. Gleadow *et al.* 1998, 2009a, 2009c), although there are exceptions (see Bazin *et al.* 2002; Bidart-Bouzat and Imeh-Nathaniel 2008). This response appears to be linked to the higher nitrogen use efficiency (NUE) of C₃ plants grown at eCO₂ (Drake *et al.* 1997; Ainsworth and Long 2005), which effectively 'frees' N from the primary metabolism to be incorporated into secondary metabolites (see Fig. 2; Gleadow *et al.* 2009a, 2009b). Thus, many plants, especially those investing in N-based defences, may become more toxic to herbivores under eCO₂ (see Fig. 2). The impact of AM on this N allocation *in planta* has, to our knowledge, not been reported in the scientific literature. Changes in N-based defences in C₄ plants under eCO₂ have, to date, not been investigated, although C₄ plants are not expected to show such a reallocation of resources, as NUE is not affected to the same degree (Ghannoum *et al.* 1997; Gleadow *et al.* 2009c). However, changes in plant water use, soil water and nutrient availability (Dijkstra *et al.* 2010), and AM associations under eCO₂ may affect nutrient content and allocation in C₄ plants.

Any increase in NCSMs in staple crops is an important but, until recently, little considered response of plants to eCO₂ (Gleadow *et al.* 2009b). The group of NCSMs with which we are most familiar is the cyanogenic glycosides, which break down when cyanogenic plant tissue is macerated to release hydrogen cyanide and either an aldehyde or a ketone (Zagrobelny Bak *et al.* 2008). Because over half of all crop plants have at least some parts that are cyanogenic (Jones 1998), we consider N allocation to cyanogenesis especially important in the context of food security. The capacity of herbivores to convert toxic cyanide (HCN) to the less toxic thiocyanate (SCN) is known to be dependent on the overall dose and rate of ingestion, and the availability of sulfur-rich proteins (Westley 1988; Cardoso *et al.* 1998; Gleadow and Woodrow 2002).




Fig. 2. Resource allocation within plants. Plant morphology, biochemistry and the efficiency of the physiological processes are integrated with nutrient supply and partitioning. Any change in resource supply or resource-use efficiency affects the partitioning of resources and the defence–growth equilibrium. (1) Assimilation rate is controlled by the size and effectiveness of the photosynthetic machinery, especially (2) Rubisco, as it accounts for ~25% of leaf N. A smaller but variable proportion of resources are allocated to the (3) synthesis of (C- or N-based) defensive secondary compounds, here classified on whether they contain (4) nitrogen (NCSM) or (5) not (CCSM). (6) Defended plants are assumed to have greater longevity, but at the cost of (7) high growth rates. (8) Plants grown at elevated atmospheric [CO₂] accumulate more total non-structural carbohydrates, which inhibit Rubisco synthesis. Low nutrient supply and elevated atmospheric [CO₂] is associated with (9) greater root growth (root : shoot ratio), and (10) decreased leaf area ratio and specific leaf area. (11) These, in turn, affect the ability of plants to acquire water and transpiration rates (WUE).

The ability of animals, including humans, to tolerate toxins and other anti-feedants in food is dependent on the toxin concentration, the availability of other food and importantly, total protein intake. Leaves of plants grown at eCO₂ almost always have less leaf protein on a per mass basis (Cotrufo *et al.* 1998). Since almost all of the N in the leaves in cereal crops is translocated to grain during senescence (Martre *et al.* 2003), low leaf protein is expected to impact grain protein as well. Indeed, studies in wheat (*Triticum aestivum* L.) and rapeseed (*Brassica napus* L.) report significant declines in total protein and essential amino acids, as well as a range of macro- and micro-nutrients (Högy and Fangmeier 2008; Högy *et al.* 2009, 2010). The protein concentration in wheat, barley (*Hordeum vulgare* L.) and rice (*Oryza sativa* L.) is forecast to be 10–15% lower in plants by 2030, reducing the nutritional value (Taub *et al.* 2008; Wieser *et al.* 2008; Gregory *et al.* 2009; Erbs *et al.* 2010), and affecting properties such as the amino acid balance and bread-making quality (Kimball *et al.* 2001; Wieser *et al.* 2008). Herbivores generally compensate for the lower plant protein concentration under eCO₂ by eating more (Lindroth *et al.* 1993; Agrell *et al.* 2006; Schädler *et al.* 2007; Stiling and Cornelissen 2007). Significantly, some studies indicate that this compensatory feeding may not be possible in species containing NCSMs (Schädler *et al.* 2007).

In the future, herbivores, including humans, may well face the twofold problem of decreased protein and increased defence

chemicals in their diets. The defence–protein balance is closely linked to plant N status, which, in turn, is dependent on nutrient (be it via the root or AMF uptake pathways) and CO₂ supply (Gleadow *et al.* 2009a; Gleadow and Woodrow 2000). This highlights the need for an integrated approach that not only considers the impacts of eCO₂ on plant mineral nutrient status and the mechanisms by which plants acquire those nutrients, but also the flow-on effects on plant primary and secondary metabolism.

Integration, hypotheses and conclusions

In this viewpoint article, we have deliberately turned our attention to two quite different aspects of plant nutrition. First, we considered the impacts of eCO₂ on AM. We predict that these associations will play an increasingly important role in a high CO₂ world, especially when considered in light of increasing scarcity of synthetic fertilisers and increased demand for nutrients as (C₃) plants become less C-limited. Second, we turned our attention to the impacts of eCO₂ on the fate of nutrients *in planta*, with an emphasis on secondary metabolites with a role in plant defence. These compounds highlight the importance of considering not only the concentration of nutrients in plant tissues (as is typically measured in studies of AM, with some exceptions), but also the forms in which they are present. It is for these reasons that we have turned our attention to the nexus between AM and plant chemical




Fig. 3. Hypothesised links between eCO₂, plant (C₃) N nutrition, AM and plant defences. (1) AM function in this example refers to increased nutrient uptake and delivery to the plant. (2) Decreased tissue N leads to increased rates of herbivory as herbivores consume more plant tissue to satisfy their dietary N requirements.

defence in the context of a high CO₂ world. While the potential overlap between the two distinct aspects of plant nutrition we consider here are clear, there has been essentially no investigation of such links. Thus, in this final section, we present several hypotheses related to these issues. In doing so, our aim is to stimulate further research in this area.

Hypothesis 1

The formation of AM will lead to improved plant N nutrition under eCO₂ and thence to increased supply of N to N-based defence compounds (see Fig. 3).

Rationale

Mycorrhizal uptake of N (Tanaka and Yano 2005) can help alleviate the N deficiency (under low soil N conditions) typically observed under eCO₂ (Cotrufo *et al.* 1998). Since the rate of herbivory can be increased under eCO₂ as herbivores need to increase their consumption of plant tissue to meet their dietary requirements (Veteli *et al.* 2002), plants which allocate more N to defence compounds will be at a competitive advantage. Consequently, the capacity of AM to increase plant supply of nutrients (including N) will be of increased benefit to plants (see Fig. 3). To test this hypothesis, there is a need for studies of AM impacts on plant N nutrition and the production of N-based defence compounds under eCO₂ conditions.

Hypothesis 2

Plants (C₃) grown under eCO₂ are less C-limited and thus, the capacity of plants to meet the demand for C by AMF and C-based defence compounds is increased (see Fig. 4).




Fig. 4. Hypothesised links between eCO₂, C assimilation, AM and (C₃) plant defences. (1) AM function in this example refers to increased nutrient uptake and delivery to the plant.

Rationale

As atmospheric CO₂ concentrations increase, C₃ photosynthesis will be less C-limited. As a consequence, more C will be available to support both the synthesis of C-based defence compounds, and also the establishment and maintenance of AM (Bennett *et al.* 2006). However, with the increased rates of herbivory under eCO₂ due to compensatory feeding (Schädler *et al.* 2007), there may be no net benefit. Again, there is a need for dedicated studies in which the competing demands for C by AMF and C-based defence compounds are assessed under eCO₂.

Conclusion

In this viewpoint article, we have explored the impacts of eCO₂ on aspects of plant biology that are typically considered in isolation from one another. Many different but equally important complex interactions between different aspects of plant biology exist, and are also in need of consideration in unison rather than isolation. The growth of C₃ plants is typically increased under eCO₂, but it is likely that this will only be realised where N and P supply (and other nutrients) to plants is adequate. With much of the world's population having limited access to costly fertilisers, the potential yield increases that eCO₂ make possible may not be fully realised. Moreover, changes in the nutritional status of plants in the future could negatively impact on subsistence farmers who have fewer options available to them. We contend that if we are to discover how plants (especially food plants) will respond to environmental change, the plant sciences need to take an integrated approach to the study of plant responses to the environment, and one that is not constrained by discipline boundaries.

Acknowledgements

The authors apologise to those of our colleagues whose work it was not possible to cite in this short viewpoint article. We also thank our friends and colleagues for valuable discussions over the years. We also thank two anonymous reviewers for their valuable comments which helped us to focus and refine our hypotheses. Our work in this area is funded by AusAID, the Australian Research Council and Monash University.

References

- Agrawal AA, Fishbein M (2006) Plant defense syndromes. *Ecology* **87**, S132–S149. doi:10.1890/0012-9658(2006)87[132:PDS]2.0.CO;2
- Agrell J, Anderson P, Oleszek W, Stochmal A, Agrell C (2006) Elevated CO₂ levels and herbivore damage alter host plant preferences. *Oikos* **112**, 63–72. doi:10.1111/j.0030-1299.2006.13614.x
- Ainsworth EA, Long SP (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist* **165**, 351–372. doi:10.1111/j.1469-8137.2004.01224.x
- Ainsworth EA, Rogers A (2007) The response of photosynthesis and stomatal conductance to rising [CO₂]: mechanisms and environmental interactions. *Plant Cell and Environment* **30**, 258–270. doi:10.1111/j.1365-3040.2007.01641.x
- Alberton O, Kuyper TW, Gorissen A (2005) Taking mycoecentrism seriously: mycorrhizal fungal and plant responses to elevated CO₂. *New Phytologist* **167**, 859–868. doi:10.1111/j.1469-8137.2005.01458.x
- Allen MF, Swenson W, Querejeta JI, Egerton-Warburton LM, Treseder KK (2003) Ecology of mycorrhizae: a conceptual framework for complex interactions among plants and fungi. *Annual Review of Phytopathology* **41**, 271–303. doi:10.1146/annurev.phyto.41.052002.095518
- Ames RN, Reid CPP, Porter LK, Cambardella C (1983) Hyphal uptake and transport of nitrogen from two ¹⁵N-labelled sources by *Glomus mosseae*, a vesicular arbuscular mycorrhizal fungus. *New Phytologist* **95**, 381–396. doi:10.1111/j.1469-8137.1983.tb03506.x
- Bago B, Azcón-Aguilar C, Goulet A, Piché Y (1998) Branched adsorbing structures (BAS): a feature of the extraradical mycelium of symbiotic arbuscular mycorrhizal fungi. *New Phytologist* **139**, 375–388. doi:10.1046/j.1469-8137.1998.00199.x
- Bazin A, Goverde M, Erhardt A, Shykoff JA (2002) Influence of atmospheric carbon dioxide enrichment on induced response and growth compensation after herbivore damage in *Lotus corniculatus*. *Ecological Entomology* **27**, 271–278. doi:10.1046/j.1365-2311.2002.00409.x
- Bennett AE, Alers-Garcia J, Bever JD (2006) Three-way interactions among mutualistic mycorrhizal fungi, plants, and plant enemies: hypotheses and synthesis. *American Naturalist* **167**, 141–152. doi:10.1086/499379
- Bidart-Bouzat MG, Imeh-Nathaniel A (2008) Global change effects on plant chemical defenses against insect herbivores. *Journal of Integrative Plant Biology* **50**, 1339–1354. doi:10.1111/j.1744-7909.2008.00751.x
- Bloom AJ, Burger M, Asensio JSR, Cousins AB (2010) Carbon dioxide enrichment inhibits nitrate assimilation in wheat and *Arabidopsis*. *Science* **328**, 899–903. doi:10.1126/science.1186440
- Boege K, Marquis RJ (2005) Facing herbivory as you grow up: the ontogeny of resistance in plants. *Trends in Ecology & Evolution* **20**, 441–448. doi:10.1016/j.tree.2005.05.001
- Campbell CD, Sage RF (2006) Interactions between the effects of atmospheric CO₂ content and P nutrition on photosynthesis in white lupin (*Lupinus alba* L.). *Plant, Cell & Environment* **29**, 844–853. doi:10.1111/j.1365-3040.2005.01464.x
- Cardoso A, Ernesto M, Cliff J, Egan SV, Bradbury JH (1998) Cyanogenic potential of cassava flour: field trial in Mozambique of a simple kit. *International Journal of Food Sciences and Nutrition* **49**, 93–99. doi:10.3109/09637489809089388
- Cavagnaro TR (2008) The role of arbuscular mycorrhizas in improving plant zinc nutrition under low soil zinc concentrations: a review. *Plant and Soil* **304**, 315–325. doi:10.1007/s11104-008-9559-7
- Cavagnaro TR, Smith FA, Lorimer MF, Haskard KA, Ayling SM, Smith SE (2001) Quantitative development of Paris-type arbuscular mycorrhizas formed between *Asphodelus fistulosus* and *Glomus coronatum*. *New Phytologist* **149**, 105–113. doi:10.1046/j.1469-8137.2001.00001.x
- Cavagnaro TR, Smith FA, Smith SE, Jakobsen I (2005) Functional diversity in arbuscular mycorrhizas: exploitation of soil patches with different phosphate enrichment differs among fungal species. *Plant, Cell & Environment* **164**, 485–491.
- Cavagnaro TR, Sokolow SK, Jackson LE (2007) Mycorrhizal effects on growth and nutrition of tomato under elevated atmospheric carbon dioxide. *Functional Plant Biology* **34**, 730–736. doi:10.1071/FP06340
- Chen X, Tu C, Burton MG, Watson DM, Burkey KO, Hu S (2007) Plant nitrogen acquisition and interactions under elevated carbon dioxide: impact of endophytes and mycorrhizae. *Global Change Biology* **13**, 1238–1249. doi:10.1111/j.1365-2486.2007.01347.x
- Coley PD, Massa M, Lovelock CE, Winter K (2002) Effects of elevated CO₂ on foliar chemistry of saplings of nine species of tropical tree. *Oecologia* **133**, 62–69. doi:10.1007/s00442-002-1005-6
- Collins-Johnson N, Wolf J, Reyes MA, Panter A, Koch GW, Redman A (2005) Species of plants and associated arbuscular mycorrhizal fungi mediate mycorrhizal responses to CO₂ enrichment. *Global Change Biology* **11**, 1156–1166. doi:10.1111/j.1365-2486.2005.00967.x

- Conroy JP, Milham PJ, Barlow EWR (1992) Effect of nitrogen and phosphorus availability on the growth-response of *Eucalyptus grandis* to high CO₂. *Plant, Cell & Environment* **15**, 843–847. doi:10.1111/j.1365-3040.1992.tb02152.x
- Cordell D, Drangert J-O, White S (2009) The story of phosphorus: global food security and food for thought. *Global Environmental Change* **19**, 292–305. doi:10.1016/j.gloenvcha.2008.10.009
- Cornelissen T, Fernandes GW, Vasconcelos-Neto J (2008) Size does matter: variation in herbivory between and within plants and the plant vigor hypothesis. *Oikos* **117**, 1121–1130. doi:10.1111/j.0030-1299.2008.16588.x
- Cotrufo MF, Ineson P, Scott A (1998) Elevated CO₂ reduces the nitrogen concentration of plant tissues. *Global Change Biology* **4**, 43–54. doi:10.1046/j.1365-2486.1998.00101.x
- Daepf M, Nösberger J, Lüscher A (2001) Nitrogen fertilization and developmental stage alter the response of *Lolium perenne* to elevated CO₂. *New Phytologist* **150**, 347–358. doi:10.1046/j.1469-8137.2001.00109.x
- de Graaff MA, van Groeningen KJ, Six J, Hungate BA, van Kessel C (2006) Interactions between plant growth and nutrient dynamics under elevated CO₂: a meta analysis. *Global Change Biology* **12**, 2077–2091. doi:10.1111/j.1365-2486.2006.01240.x
- Dickson S (2004) The Arum–Paris continuum of mycorrhizal symbioses. *New Phytologist* **163**, 187–200. doi:10.1111/j.1469-8137.2004.01095.x
- Dickson S, Kolesik P (1999) Visualisation of mycorrhizal fungal structures and quantification of their surface area and volume using laser scanning confocal microscopy. *Mycorrhiza* **9**, 205–213. doi:10.1007/s005720050268
- Dickson S, Smith SE (2001) Cross walls in arbuscular trunk hyphae form after loss of metabolic activity. *New Phytologist* **151**, 735–742. doi:10.1046/j.0028-646x.2001.00225.x
- Dijkstra FA, Blumenthal D, Morgan JA, LeCain DR, Follett RF (2010) Elevated CO₂ effects on semi-arid grassland plants in relation to water availability and competition. *Functional Ecology* **24**, 1152–1161. doi:10.1111/j.1365-2435.2010.01717.x
- Drake BG, González-Meler MA, Long SP (1997) More efficient plants: a consequence of rising atmospheric CO₂? *Annual Review of Plant Physiology and Plant Molecular Biology* **48**, 609–639. doi:10.1146/annurev.arplant.48.1.609
- Edwards EJ, McCaffrey S, Evans JR (2005) Phosphorus status determines biomass response to elevated CO₂ in a legume: C₄ grass community. *Global Change Biology* **11**, 1968–1981.
- Erbs M, Manderscheid R, Jansen G, Seddig S, Pacholski A, Weigel H-J (2010) Effects of free-air CO₂ enrichment and nitrogen supply on grain quality parameters and elemental composition of wheat and barley grown in a crop rotation. *Agriculture Ecosystems & Environment* **136**, 59–68. doi:10.1016/j.agee.2009.11.009
- Gamper H, Peter M, Jansa J, Lüscher A, Hartwig UA, Leuchtman A (2004) Arbuscular mycorrhizal fungi benefit from 7 years of free air CO₂ enrichment in well-fertilized grass and legume monocultures. *Global Change Biology* **10**, 189–199. doi:10.1111/j.1529-8817.2003.00734.x
- Gamper H, Hartwig UA, Leuchtman A (2005) Mycorrhizas improve nitrogen nutrition of *Trifolium repens* after 8 yr of selection under elevated atmospheric CO₂ partial pressure. *New Phytologist* **167**, 531–542. doi:10.1111/j.1469-8137.2005.01440.x
- Gange AC, West HM (1994) Interactions between arbuscular mycorrhizal fungi and foliar-feeding insects in *Plantago lanceolata* L. *New Phytologist* **128**, 79–87. doi:10.1111/j.1469-8137.1994.tb03989.x
- García MO, Ovasapyan T, Greas M, Treseder KK (2008) Mycorrhizal dynamics under elevated CO₂ and nitrogen fertilization in a warm temperate forest. *Plant and Soil* **303**, 301–310. doi:10.1007/s11104-007-9509-9
- Gavito ME, Bruhn D, Jakobsen I (2002) P uptake by arbuscular mycorrhizal hyphae does not increase when the host plant grows under atmospheric CO₂ enrichment. *New Phytologist* **154**, 751–760. doi:10.1046/j.1469-8137.2002.00404.x
- Gavito ME, Schweiger P, Jakobsen I (2003) P uptake by arbuscular mycorrhizal hyphae: effect of soil temperature and atmospheric CO₂ enrichment. *Global Change Biology* **9**, 106–116. doi:10.1046/j.1365-2486.2003.00560.x
- Ghannoum O, von Caemmerer S, Barlow EWR, Conroy JP (1997) The effect of CO₂ enrichment and irradiance on the growth, morphology and gas exchange of a C₃ (*Panicum laxum*) and a C₄ (*Panicum antidotale*) grass. *Australian Journal of Plant Physiology* **24**, 227–237. doi:10.1071/PP96077
- Glassop D, Smith SE, Smith FW (2005) Cereal phosphate transporters associated with the mycorrhizal pathway of phosphate uptake in roots. *Planta* **222**, 688–698. doi:10.1007/s00425-005-0015-0
- Gleadow RM, Woodrow IE (2000) Temporal and spatial variation in cyanogenic glycosides in *Eucalyptus cladocalyx*. *Tree Physiology* **20**, 591–598.
- Gleadow RM, Woodrow IE (2002) Constraints on effectiveness of cyanogenic glycosides in herbivore defense. *Journal of Chemical Ecology* **28**, 1301–1313. doi:10.1023/A:1016298100201
- Gleadow RM, Foley WJ, Woodrow IE (1998) Enhanced CO₂ alters the relationship between photosynthesis and defence in cyanogenic *Eucalyptus cladocalyx* F. Muell. *Plant, Cell & Environment* **21**, 12–22. doi:10.1046/j.1365-3040.1998.00258.x
- Gleadow RM, Edwards E, Evans J (2009a) Changes in nutritional value of cyanogenic *Trifolium repens* at elevated CO₂. *Journal of Chemical Ecology* **35**, 476–478. doi:10.1007/s10886-009-9617-5
- Gleadow RM, Evans J, McCaffrey S, Cavagnaro TR (2009b) Growth and nutritive value of cassava (*Manihot esculenta* Cranz.) are reduced when grown at elevated CO₂. *Plant Biology* **11**, 76–82. doi:10.1111/j.1438-8677.2009.00238.x
- Gleadow R, Cavagnaro T, O'Donnell N, Evans J, Neale A, Blomstedt C, Hamill J (2009c) Unbalancing global resources: will plants be edible in a high CO₂ world? *Comparative Biochemistry and Physiology. Part A, Molecular & Integrative Physiology* **153**(Suppl. 1), s225. doi:10.1016/j.cbpa.2009.04.559
- González-Guerrero M, Azcón-Aguilar C, Mooney M, Valderas A, MacDiarmid CW, Eide DJ, Ferrol N (2005) Characterization of a *Glomus intraradices* gene encoding a putative Zn transporter of the cation diffusion facilitator family. *Fungal Genetics and Biology* **42**, 130–140. doi:10.1016/j.fgb.2004.10.007
- González-Guerrero M, Cano C, Azcón-Aguilar C, Ferrol N (2007) GintMT1 encodes a functional metallothionein in *Glomus intraradices* that responds to oxidative stress. *Mycorrhiza* **17**, 327–335. doi:10.1007/s00572-007-0108-7
- Gregory PJ, Johnson SN, Newton AC, Ingram JSI (2009) Integrating pests and pathogens into the climate change/food security debate. *Journal of Experimental Botany* **60**, 2827–2838. doi:10.1093/jxb/erp080
- Grünzweig JM, Körner C (2003) Differential phosphorus and nitrogen effects drive species and community responses to elevated CO₂ in semi-arid grassland. *Functional Ecology* **17**, 766–777. doi:10.1111/j.1365-2435.2003.00797.x
- Hartwig UA, Wittmann P, Braun R, Hartwig-Räz B, Jansa J, Mozafar A, Lüscher A, Leuchtman A, Frossard E, Nösberger J (2002) Arbuscular mycorrhiza infection enhances the growth response of *Lolium perenne* to elevated atmospheric pCO₂. *Journal of Experimental Botany* **53**, 1207–1213. doi:10.1093/jexbot/53.371.1207
- Haugen R, Steffes L, Wolf J, Brown P, Matzner S, Siemens DH (2008) Evolution of drought tolerance and defense: dependence of tradeoffs on mechanism, environment and defense switching. *Oikos* **117**, 231–244. doi:10.1111/j.2007.0030-1299.16111.x

- Högy P, Fangmeier A (2008) Effects of elevated atmospheric CO₂ on grain quality of wheat. *Journal of Cereal Science* **48**, 580–591. doi:10.1016/j.jcs.2008.01.006
- Högy P, Wieser H, Köhler P, Schwadorf K, Breuer J, Franzaring J, Muntiferung R, Fangmeier A (2009) Effects of elevated CO₂ on grain yield and quality of wheat: results from a 3-year free-air CO₂ enrichment experiment. *Plant Biology* **11**(Suppl. 1), 60–69. doi:10.1111/j.1438-8677.2009.00230.x
- Högy P, Franzaring J, Schwadorf K, Breuer J, Schultze W, Fangmeier A (2010) Effects of free-air CO₂ enrichment on energy traits and seed quality of oilseed rape. *Agriculture Ecosystems & Environment* **139**, 239–244. doi:10.1016/j.agee.2010.08.009
- Hu S, Tu C, Chen X, Gruver JB (2006) Progressive N limitation of plant response to elevated CO₂: a microbiological perspective. *Plant and Soil* **289**, 47–58. doi:10.1007/s11104-006-9093-4
- IPCC (2007) Technical summary. In 'Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change'. pp. 19–91. (Cambridge University Press: Cambridge)
- Jackson LE, Burger M, Cavagnaro TR (2008) Roots, nitrogen transformations, and ecosystem services. *Annual Review of Plant Biology* **59**, 341–363. doi:10.1146/annurev.arplant.59.032607.092932
- Jifon JL, Graham JH, Drouillard DL, Syvertsen JP (2002) Growth depression of mycorrhizal *Citrus* seedlings grown at high phosphorus supply is mitigated by elevated CO₂. *New Phytologist* **153**, 133–142. doi:10.1046/j.0028-646X.2001.00294.x
- Johansen A, Jakobsen I, Jensen ES (1993) External hyphae of vesicular-arbuscular mycorrhizal fungi associated with *Trifolium subterraneum*. 3. Hyphal transport of ³²P and ¹⁵N. *New Phytologist* **124**, 61–68. doi:10.1111/j.1469-8137.1993.tb03797.x
- Jones DA (1998) Why are so many food plants cyanogenic? *Phytochemistry* **47**, 155–162. doi:10.1016/S0031-9422(97)00425-1
- Kimball BA, Morris CE, Pinter PJ Jr, Wall GW, Hunsaker DJ, Adamsen FJ, LaMorte RL, Leavitt SW, Thompson TL, Matthias AD, Brooks TJ (2001) Elevated CO₂, drought and soil nitrogen effects on wheat grain quality. *New Phytologist* **150**, 295–303. doi:10.1046/j.1469-8137.2001.00107.x
- Klironomos JN, Ursic M, Rillig M, Allen MF (1998) Interspecific differences in the response of arbuscular mycorrhizal fungi to *Artemisia tridentata* grown under elevated atmospheric CO₂. *New Phytologist* **138**, 599–605. doi:10.1046/j.1469-8137.1998.00141.x
- Lambers H (1993) Rising CO₂, secondary plant metabolism, plant–herbivore interactions and litter decomposition: theoretical considerations. *Vegetation* **104–105**, 263–271. doi:10.1007/BF00048157
- Lawler IR, Foley WJ, Woodrow IE, Cork SJ (1997) The effects of elevated CO₂ atmospheres on the nutritional quality of *Eucalyptus* foliage and its interaction with soil nutrient and light availability. *Oecologia* **109**, 59–68. doi:10.1007/s004420050058
- Lincoln DE, Fajer ED, Johnson RH (1993) Plant insect herbivore interactions in elevated CO₂ environments. *Trends in Ecology & Evolution* **8**, 64–68. doi:10.1016/0169-5347(93)90161-H
- Lindroth RL (2010) Impacts of elevated atmospheric CO₂ and O₃ on forests: phytochemistry, trophic interactions and ecosystem dynamics. *Journal of Chemical Ecology* **36**, 2–21. doi:10.1007/s10886-009-9731-4
- Lindroth RL, Kinney KK, Platz CL (1993) Responses of deciduous trees to elevated atmospheric CO₂: productivity, phytochemistry and insect performance. *Ecology* **74**, 763–777. doi:10.2307/1940804
- Loladze I (2002) Rising atmospheric CO₂ and human nutrition: toward globally imbalanced plant stoichiometry. *Trends in Ecology & Evolution* **17**, 457–461. doi:10.1016/S0169-5347(02)02587-9
- Lovelock C, Kyllö D, Popp M, Isopp H, Virgo A, Winter K (1997) Symbiotic vesicular arbuscular mycorrhizae influence maximum rates of photosynthesis in tropical tree seedlings grown under elevated CO₂. *Australian Journal of Plant Physiology* **24**, 185–194. doi:10.1071/PP96089
- Lukac M, Calfapietra C, Godbold DL (2003) Production, turnover and mycorrhizal colonization of root systems of three *Populus* species grown under elevated CO₂ (POPFACE). *Global Change Biology* **9**, 838–848. doi:10.1046/j.1365-2486.2003.00582.x
- Lukac M, Calfapietra C, Lagomarsino A, Loreto F (2010) Global climate change and tree nutrition: effects of elevated CO₂ and temperature. *Tree Physiology* **30**, 1209–1220. doi:10.1093/treephys/tpq040
- Martre P, Porter JR, Jamieson PD, Triboni E (2003) Modeling grain nitrogen accumulation and protein composition to understand the sink/source regulations of nitrogen remobilization for wheat. *Plant Physiology* **133**, 1959–1967. doi:10.1104/pp.103.030585
- Matros A, Amme S, Kettig B, Buck-Sorlin GH, Sonnwald U, Mock HP (2006) Growth at elevated CO₂ concentrations leads to modified profiles of secondary metabolites in tobacco cv. SamsunNN and to increased resistance against infection with potato virus Y. *Plant, Cell & Environment* **29**, 126–137. doi:10.1111/j.1365-3040.2005.01406.x
- Nowak RS, Ellsworth DS, Smith SD (2004) Functional responses of plants to elevated atmospheric CO₂ – do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytologist* **162**, 253–280. doi:10.1111/j.1469-8137.2004.01033.x
- Olesniewicz KS, Thomas RB (1999) Effects of mycorrhizal colonization on biomass production and nitrogen fixation of black locust (*Robinia pseudoacacia*) seedlings grown under elevated atmospheric carbon dioxide. *New Phytologist* **142**, 133–140. doi:10.1046/j.1469-8137.1999.00372.x
- Olsrud M, Carlsson BA, Svensson BM, Michelsen A, Melillo JM (2010) Responses of fungal root colonization, plant cover and leaf nutrients to long-term exposure to elevated atmospheric CO₂ and warming in a subarctic birch forest understorey. *Global Change Biology* **16**, 1820–1829. doi:10.1111/j.1365-2486.2009.02079.x
- Peñuelas J, Estiarte M (1998) Can elevated CO₂ affect secondary metabolism and ecosystem function? *Trends in Ecology & Evolution* **13**, 20–24. doi:10.1016/S0169-5347(97)01235-4
- Reich PB, Hobbie SE, Lee TD, Ellsworth DS, West JB, Timan D, Knops JMH, Naeem S, Trost J (2006) Nitrogen limitation constraints sustainability of ecosystem response to CO₂. *Nature* **440**, 922–925. doi:10.1038/nature04486
- Rhoades DF (1979) 'Evolution of chemical defence against herbivores.' (Academic Press: New York)
- Rillig MC (2004) Arbuscular mycorrhizae and terrestrial ecosystem processes. *Ecology Letters* **7**, 740–754. doi:10.1111/j.1461-0248.2004.00620.x
- Rillig MC, Allen MF (1999) What is the role of arbuscular mycorrhizal fungi in plant-to-ecosystem responses to elevated atmospheric CO₂? *Mycorrhiza* **9**, 1–8. doi:10.1007/s005720050257
- Rillig MC, Allen MF, Klironomos JN, Field CB (1998) Arbuscular mycorrhizal percent infection and infection intensity of *Bromus hordeaceus* grown in elevated atmospheric CO₂. *Mycologia* **90**, 199–205. doi:10.2307/3761295
- Rogers A, Ainsworth EA, Leakey ADB (2009) Will elevated carbon dioxide concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiology* **151**, 1009–1016. doi:10.1104/pp.109.144113
- Rosegrant MW, Cline SA (2003) Global food security: challenges and policies. *Science* **302**, 1917–1919. doi:10.1126/science.1092958
- Rouhier H, Read DJ (1998) The role of mycorrhiza in determining the response of *Plantago lanceolata* to CO₂ enrichment. *New Phytologist* **139**, 367–373. doi:10.1046/j.1469-8137.1998.00200.x
- Sanders IR, Streitwolf-Engel R, van der Heijden MGA, Boller T, Wiemken A (1998) Increased allocation to external hyphae of arbuscular mycorrhizal fungi under CO₂ enrichment. *Oecologia* **117**, 496–503. doi:10.1007/s004420050685
- Schädler M, Roeder M, Brandl R, Matthies D (2007) Interacting effects of elevated CO₂, nutrient availability and plant species on a generalist invertebrate herbivore. *Global Change Biology* **13**, 1005–1015. doi:10.1111/j.1365-2486.2007.01319.x

- Smith SE, Read DJ (2008) 'Mycorrhizal symbiosis.' (Academic Press Ltd: Cambridge, UK)
- Smith SE, St John BJ, Smith FA, Nicholas DJD (1985) Activity of glutamine synthetase and glutamate dehydrogenase in *Trifolium subterraneum* L. and *Allium cepa* L.: effects of mycorrhizal infection and phosphate nutrition. *New Phytologist* **99**, 211–227. doi:10.1111/j.1469-8137.1985.tb03651.x
- Smith SE, Smith FA, Jakobsen I (2004) Functional diversity in arbuscular mycorrhizal (AM) symbioses: the contribution of the mycorrhizal P uptake pathway is not correlated with mycorrhizal responses in growth or total P uptake. *New Phytologist* **162**, 511–524. doi:10.1111/j.1469-8137.2004.01039.x
- Staddon PL, Fitter AH, Graves JD (1999) Effect of elevated atmospheric CO₂ on mycorrhizal colonisation, external hyphal production and phosphorus inflow in *Plantago lanceolata* and *Trifolium repens* in association with the arbuscular mycorrhizal fungus *Glomus mosseae*. *Global Change Biology* **5**, 347–358. doi:10.1046/j.1365-2486.1999.00230.x
- Staddon PL, Gregersen R, Jakobsen I (2004) The response of two *Glomus* mycorrhizal fungi and a fine endophyte to elevated atmospheric CO₂, soil warming and drought. *Global Change Biology* **10**, 1909–1921. doi:10.1111/j.1365-2486.2004.00861.x
- Stiling P, Cornelissen T (2007) How does elevated carbon dioxide (CO₂) affect plant–herbivore interactions? A field experiment and meta-analysis of CO₂-mediated changes on plant chemistry and herbivore performance. *Global Change Biology* **13**, 1823–1842. doi:10.1111/j.1365-2486.2007.01392.x
- Stitt M, Krapp A (1999) The molecular physiological basis for the interaction between elevated carbon dioxide and nutrients. *Plant, Cell & Environment* **22**, 583–621. doi:10.1046/j.1365-3040.1999.00386.x
- Stöcklin J, Schweizer K, Körner C (1998) Effects of elevated CO₂ and phosphorus addition on productivity and community composition of intact monoliths from calcareous grassland. *Oecologia* **116**, 50–56. doi:10.1007/s004420050562
- Syvrtsen JP, Graham JH (1999) Phosphorus supply and arbuscular mycorrhizas increase growth and net gas exchange response of two *Citrus* spp. grown at elevated [CO₂]. *Plant and Soil* **208**, 209–219. doi:10.1023/A:1004553315041
- Tanaka Y, Yano K (2005) Nitrogen delivery to maize via mycorrhizal hyphae depends on the form of N supplied. *Plant, Cell & Environment* **28**, 1247–1254. doi:10.1111/j.1365-3040.2005.01360.x
- Taub DR, Miller B, Allen H (2008) Effects of elevated CO₂ on the protein concentration of food crops: a meta-analysis. *Global Change Biology* **14**, 565–575. doi:10.1111/j.1365-2486.2007.01511.x
- Tobar R, Azcón R, Barea JM (1994) Improved nitrogen uptake and transport from ¹⁵N-labelled nitrate by external hyphae of arbuscular mycorrhizae under water-stressed conditions. *New Phytologist* **126**, 119–122. doi:10.1111/j.1469-8137.1994.tb07536.x
- Treseder KK (2004) A meta-analysis of mycorrhizal responses to nitrogen, phosphorus, and atmospheric CO₂ in field studies. *New Phytologist* **164**, 347–355. doi:10.1111/j.1469-8137.2004.01159.x
- Treseder KK, Allen MF (2000) Mycorrhizal fungi have a potential role in soil carbon storage under elevated CO₂ and nitrogen deposition. *New Phytologist* **147**, 189–200. doi:10.1046/j.1469-8137.2000.00690.x
- van Aarle IM, Cavagnaro TR, Smith SE, Smith FA, Dickson S (2005) Metabolic activity of *Glomus intraradices* in *Arum*- and *Paris*-type arbuscular mycorrhiza colonization. *New Phytologist* **166**, 611–618. doi:10.1111/j.1469-8137.2005.01340.x
- Veteli TO, Kuokkanen K, Julkenen-Tiitto R, Roininen H, Tahvanainen J (2002) Effects of elevated CO₂ and temperature on plant growth and defensive chemistry. *Global Change Biology* **8**, 1240–1252. doi:10.1046/j.1365-2486.2002.00553.x
- Ward SJE, Midgley GF, Jones MH, Curtis PS (1999) Responses of wild C₄ and C₃ grass (Poaceae) species to elevated atmospheric CO₂ concentration: a meta-analytic test of current theories and perceptions. *Global Change Biology* **5**, 723–741. doi:10.1046/j.1365-2486.1999.00265.x
- Westley J (1988) Mammalian cyanide detoxification with sulfane sulfur. *Ciba Foundation Symposium* **140**, 201–218.
- Wieser H, Manderscheid R, Erbs M, Weigel H-J (2008) Effects of elevated atmospheric CO₂ concentrations on the quantitative protein composition of wheat grain. *Journal of Agricultural and Food Chemistry* **56**, 6531–6535. doi:10.1021/jf8008603
- Zagrobelyny M, Bak S, Møller BL (2008) Cyanogenesis in plants and arthropods. *Phytochemistry* **69**, 1457–1468. doi:10.1016/j.phytochem.2008.02.019
- Ziska LH, Emche SD, Johnson EL, George K, Reed DR, Sicher RC (2005) Alterations in the production and concentration of selected alkaloids as a function of rising atmospheric carbon dioxide and air temperature: implications for ethno-pharmacology. *Global Change Biology* **11**, 1798–1807. doi:10.1111/j.1365-2486.2005.001029.x

Manuscript received 4 June 2010, accepted 18 November 2010