

<https://helda.helsinki.fi>

Plant-plant interactions and N fertilization shape soil bacterial and fungal communities

Guo, Qingxue

2019-01

Guo , Q , Yan , L , Korpelainen , H , Niinemets , Ü & Li , C 2019 , ' Plant-plant interactions and N fertilization shape soil bacterial and fungal communities ' , Soil Biology & Biochemistry , vol. 128 , pp. 127-138 . <https://doi.org/10.1016/j.soilbio.2018.10.018>

<http://hdl.handle.net/10138/320598>

<https://doi.org/10.1016/j.soilbio.2018.10.018>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 *The Revision for SBB13116.R1*

2

3 **Plant-plant interactions and N fertilization shape soil**
4 **bacterial and fungal communities**

5

6 Qingxue Guo ^{1, †}, Lijuan Yan ^{2, †},

7 Helena Korpelainen ³, Ülo Niinemets ⁴, Chunyang Li ^{1, *}

8

9 ¹ College of Life and Environmental Sciences, Hangzhou Normal University,
10 Hangzhou 310036, China

11 ² Aquatic Geomicrobiology, Institute of Biodiversity, Friedrich Schiller University
12 Jena, Dornburger Str. 159, D-07743 Jena, Germany

13 ³ Department of Agricultural Sciences, Viikki Plant Science Centre, P.O. Box 27,
14 FI-00014 University of Helsinki, Finland

15 ⁴ Institute of Agricultural and Environmental Sciences, Estonian University of Life
16 Sciences, Kreutzwaldi 1, 51014 Tartu, Estonia

17 [†] Authors contributed equally to this study

18 ^{*} Corresponding author: Chunyang Li, E-mail: licy@hznu.edu.cn

19

20

21

22

23 **Abstract** The impact of conspecific and heterospecific neighboring plants on soil
24 bacterial and fungal communities has never been explored in a forest ecosystem. In
25 the present study, we first investigated soil microbial communities in three plantations:
26 *Larix kaempferi* monoculture, *L. olgensis* monoculture and their mixture. Then, a
27 two-year growth experiment was conducted to investigate the effects of intra- and
28 inter-specific plant interactions of *L. kaempferi* and *L. olgensis* on rhizosphere
29 microbial communities in different nitrogen conditions. The results demonstrated
30 clear differences in the beta-diversity and composition of bacteria and fungi among
31 the three plantations, which implied different effects of plant-plant interactions on soil
32 microbial communities. The results of the pot experiment showed that *L. kaempferi*
33 suffered from greater negative effects from its conspecific neighbor regardless of the
34 N fertilization, whereas the negative effect declined when *L. kaempferi* was grown
35 with *L. olgensis* under N fertilization. Changes in intra- and inter-specific plant
36 interactions significantly impacted the chemical and biological properties of soil
37 under N fertilization, with lower concentrations of NH_4^+ , and lower soil microbial
38 biomass (C_{Mic}) and soil carbon nitrogen biomass (N_{Mic}) under intra-specific plant
39 interactions of *L. kaempferi* (KK) compared to inter-specific interactions of *L.*
40 *kaempferi* and *L. olgensis* (KO). N fertilization increased bacterial and fungal alpha
41 diversities in the rhizosphere soil of KO. For the beta diversity, the PERMANOVA
42 results demonstrated that there was a significant impact of intra- and inter-specific
43 plant interactions on soil microbial communities, with KK significantly differing from
44 intra-specific plant interactions of *L. olgensis* (OO) and KO. The two plant species

45 and N fertilization showed specific effects on the soil microbial composition,
46 particularly on the fungal community. Both *L. olgensis* and N fertilization increased
47 the abundance of *Ascomycota* but reduced that of *Basidiomycota*, and even shifted the
48 dominance of *Basidiomycota* to *Ascomycota* in KO under N fertilization. Based on
49 our results, we suggest that *L. kaempferi* planted with *L. olgensis* under N fertilization
50 may be an efficient way to promote the productivity of plantations.

51

52 **Keywords** Forest plantations; Plant neighbors; Plant-soil feedback; Fungal
53 communities; Carbon investment

54

55

56

57

58

59

60

61

62

63

64

65 **1. Introduction**

66

67 Many types of forest plantations (mostly monocultures) have been established around
68 the world to provide wood products or restore degraded lands (Paul et al., 2010;
69 Richards et al., 2010). However, the productivity of monocultures has commonly
70 declined due to reasons, such as decreasing nitrogen (N) availability or autotoxicity
71 (O’Hehir and Nambiar, 2010; Chen et al., 2014; Deng et al., 2014). For example, the
72 declined productivity of Chinese fir (*Cunninghamia lanceolata*) monocultures is
73 mainly caused by a novel allelochemical, cyclic dipeptide (Chen et al., 2014), which
74 alters fungal communities in soil by inhibiting the spore germination of *Glomus*
75 *cunninghamia* and *Gigaspora alboaurantiaca* (Xia et al., 2016). Mixed-forest
76 plantations always have higher productivity than monocultures (Lovelock and Ewel,
77 2005; Richards et al., 2010). Lovelock and Ewel (2005) have found differences in
78 arbuscular mycorrhizal compositions between different tropical tree mixtures and
79 respective monocultures, and the fungal diversity was correlated with the net primary
80 productivity of plantations. Similarly, Mortimer et al. (2015) have observed that the
81 biomass of soil fungi and bacteria as well as the productivity of tea were higher in a
82 mixture of *Alnus nepalensis* and tea species (*Camellia sinensis* var., *assamica*)
83 compared with a tea monoculture.

84

85 Plants interact intensively and reciprocally with soil microbial communities
86 (Kuzyakov and Xu, 2013). Previous studies have confirmed that plants show specific
87 effects on the structure of their own microbial communities, for example, by secreting
88 organic compounds and through input on litter (reviewed in Bakker et al., 2013a;

89 Cline and Zak, 2015; Kumar et al., 2016; Cheeke et al., 2017). However, the majority
90 of these studies have focused on isolated plants or have overlooked the impacts of
91 plant-plant interactions. Conspecific or heterospecific plant neighbors always cause
92 different effects on processes, such as nitrogen uptake, the secretion of root exudates
93 or growth (McKane et al., 2002; Kozovits et al., 2005; Broz et al., 2010; Kunstler et
94 al., 2015; Guo et al., 2017). If plants suffer from their conspecific or heterospecific
95 neighbors, for example as a result of competition, there may be consequent effects on
96 the microbial communities of soil. As research increasingly demonstrates the
97 connection between plants and microbes, there remains a need to understand how
98 intra- and inter-specific plant-plant interactions impact belowground microbial
99 communities.

100

101 Plant-plant interactions are affected by nitrogen availability (Wilson and Tilman, 1991;
102 Guo et al., 2017; Broadbent et al., 2018) and plant-soil feedback processes (Harrison
103 and Bardgett, 2010; Hendriks et al., 2015). A negative effect means that a particular
104 species changes its biological community or abiotic soil properties in a way that its
105 growth is reduced, while the other species shows less harmful or even beneficial
106 effects (Harrison and Bardgett, 2010). Reduced plant growth can result from changes
107 in the composition of beneficial arbuscular mycorrhizal fungi (Castelli and Casper,
108 2003). It has been discovered that nitrogen fertilization can alleviate plant-soil
109 negative effects and plant-plant interactions (Guo et al., 2017), probably not only by
110 increasing the N level of soil but also by improving microbial communities (Xia et al.,

111 2016). Many studies have reported that nitrogen availability largely impacts bacterial
112 and fungal communities (e.g. Fierer et al., 2012; Dietrich et al., 2017; Carrara et al.,
113 2018; Treseder et al., 2018). Based on a nitrogen fertilization experiment, Yuan et al.
114 (2016) found that *Chloroflexi* and *Bacteroidetes* show positive responses to N
115 fertilization, while *Acidobacteria* and *Verrucomicrobia* have negative responses.
116 Treseder and Allen (2002) discovered that the fungal community of soil tends to shift
117 from *Gigasporaceae* under low N availability to *Glomeraceae* under high N. A few
118 studies have investigated the effects of intra- and inter-specific plant-plant interactions
119 on soil bacterial communities at different N levels in grass lands or agroecosystems
120 (Bakker et al., 2013b, 2014; Pivato et al., 2017). For instance, Pivato et al. (2017)
121 showed that the total bacterial abundance in the rhizosphere of some inter-specific
122 plant-plant mixtures was significantly higher than that of either plant species in a
123 monoculture at a low nitrogen level in an agroecosystem, whereas the difference
124 disappeared at a high nitrogen level. Thus, their results implied that the impacts of
125 plant-plant interactions on soil bacterial communities largely depend on N levels.
126 However, there is less knowledge of the effects of N availability and plant-plant
127 interactions on soil bacterial and fungal communities in forest ecosystems.

128

129 Larch plantations are widely distributed and have a high economic and ecological
130 value across the Northern Hemisphere in areas such as Siberia, north-eastern China
131 and Japan (Agathokleous et al., 2017). The larches *Larix kaempferi* and *L. olgensis*
132 are two important plantation species. *L. kaempferi* grown in isolation has a higher

133 growth rate, biomass production and N accumulation than does *L. olgensis* (Li et al.,
134 2016; Guo et al., 2017). However, our previous results have suggested that the growth
135 of *L. kaempferi* is suppressed when grown in a mixed culture with *L. olgensis* in soil
136 from a *L. kaempferi* plantation (Guo et al., 2017), which may result from the presence
137 of different soil microbial communities. The present study consists of two parts.
138 Firstly, bulk soil samples were collected from three types of plantations (*L. kaempferi*
139 monoculture, *L. olgensis* monoculture and a mixed plantation of these two species).
140 The plantations are characterized by even-aged individuals and there are some
141 herbaceous species in low abundance. This setting enables studies on intra- and
142 inter-specific interactions in natural conditions. Secondly, a two-year pot experiment
143 was established, including isolated plants and combinations of two plants under intra-
144 and inter-specific interactions with and without N fertilization. In the pot experiment,
145 we used soil collected from a mature *L. kaempferi* monoculture to investigate the
146 effects of plant-plant interactions and N fertilization on soil microbial communities.
147 We aimed to answer the following questions: 1) do different plantations contain
148 specific types of soil bacterial and fungal communities? Based on the pot experiment,
149 2) if the intra- or inter-specific plant interactions of the two studied species were
150 different, would the rhizosphere soil bacterial and fungal communities also differ from
151 each other? 3) If N fertilization differently affects intra- and inter-specific plant
152 interactions, how does the interplay of N fertilization and plant interactions affect
153 rhizosphere soil communities?

154 **2. Materials and methods**

155

156 In the first part of this study, bulk soil samples were collected from three types of
157 plantations: *L. kaempferi* monoculture plantation, *L. olgensis* monoculture plantation
158 and a mixed-plantation of these two species. In the second part of the study,
159 rhizosphere soil samples were collected from the pot experiment. The pot experiment
160 was conducted outside on an open space at the Qingyuan Experimental Station of
161 Forest Ecology, Institute of Applied Ecology, Chinese Academy of Sciences, Liaoning
162 Province, Northeast China (41°51'N, 124°54'E, 560 m above sea level). The mean
163 annual rainfall is 811 mm and the mean annual air temperature varies between 3.9 °C
164 and 5.4 °C. *L. kaempferi* and *L. olgensis* plantations are widely distributed in the study
165 area.

166

167 *2.1. Plantation selection*

168

169 The following principles were followed when selecting the three types of plantations:
170 1) similar age, 2) close location, and 3) similar exposure and slope. Eventually, we
171 found only one pure *L. kaempferi* plantation (about 20 years old, 3.6 ha and 2050
172 stems per ha), one pure *L. olgensis* plantation (about 20 years old, 4.0 ha and 1750
173 stems per ha) and one mixed plantation (about 24 years old, 6.0 ha and 600 *L.*
174 *kaempferi* stems per ha, and 640 *L. olgensis* stems per ha), about 30 km away from the
175 research station. The two monoculture plantations were adjacent, and the *L. olgensis*
176 plantation experienced thinning forest management in 2010 according to the

177 description of a local farmer. The mixed plantation was about 400 m away from the
178 two monoculture plantations but on another hillside.

179

180 In the center of each plantation, ten plots (10 m × 10 m) were established. From each
181 plot, ten bulk soil samples (0-10 cm in depth) were collected from ten subplots (1 m ×
182 1 m) and then mixed to form one composite sample. Finally, ten samples from each
183 plantation were collected. Each bulk soil sample was divided into two equal
184 subsamples: one subsample was used for the analysis of soil properties, while the
185 other subsample was stored at -20 °C for the subsequent DNA extraction. Five
186 subsamples from each plantation were randomly selected for later DNA extraction.

187

188 *2.2. Pot experiment and sampling*

189

190 For the pot experiment, soil was collected from the *L. kaempferi* plantation near the
191 research station and then homogenized (pH 5.65, C 18.61 g kg⁻¹ and total N 1.82 g
192 kg⁻¹) early October, 2013. One-year old seedlings of *L. kaempferi* and *L. olgensis* with
193 approximately the same crown size and height were chosen from a local nursery
194 garden to establish intra- and inter-specific plant interactions. Before planting, all
195 roots were carefully washed with sterile water to remove adhering soil. Two seedlings
196 were planted 10 cm apart from each other into each plastic pot (external diameter and
197 height 56 cm and 33 cm, respectively) late October, 2013. KK refers to two *L.*
198 *kaempferi* seedlings in a pot, KO refers to one *L. kaempferi* and one *L. olgensis* in a

199 pot, and OO refers to two *L. olgensis* seedlings in a pot. Each seedling combination
200 included thirty-two pots. In May 2014, sixteen pots were randomly selected from each
201 treatment to carry out N fertilization by applying 5.1 g urea (46.3% N). N application
202 was repeated three more times (each time 5.1 g urea), in June 2014, and in May and
203 June 2015. In addition, single *L. kaempferi* seedlings (K) and single *L. olgensis*
204 seedlings (O) were planted into pots in late October, 2013, six pots for each species.
205 Similarly, three pots from both K and O were randomly selected to apply urea as
206 above. The combination of results from the plantations and pot experiment will
207 provide a better understanding of the effects of intra- and inter-specific plant
208 interactions on soil bacterial and fungal communities.

209

210 Double-plant pots were harvested twice, late August, 2014 and early September, 2015.

211 The harvested plants were dried at 70°C for 72h to determine their dry masses.

212 Neighbor effect index (NEI), estimated to measure plant-plant interactions, used the
213 following formulas (Manea and Leishman, 2011):

$$214 \text{NEI}_{k/kk} = (Y_{k/kk} - Y_k)/(Y_{k/kk} + Y_k), \text{NEI}_{o/oo} = (Y_{o/oo} - Y_o)/(Y_{o/oo} + Y_o),$$

$$215 \text{NEI}_{k/ko} = (Y_{k/ko} - Y_k)/(Y_{k/ko} + Y_k), \text{NEI}_{o/ko} = (Y_{o/ko} - Y_o)/(Y_{o/ko} + Y_o),$$

216 where $\text{NEI}_{k/kk}$ and $\text{NEI}_{o/oo}$ indicate intra-specific plant interactions for *L. kaempferi*
217 and *L. olgensis*, respectively; $\text{NEI}_{k/ko}$ and $\text{NEI}_{o/ko}$ indicate inter-specific plant
218 interactions for *L. kaempferi* and *L. olgensis*, respectively. Y_k and Y_o are the biomass
219 of *L. kaempferi* and *L. olgensis* grown in isolation, respectively. $Y_{k/kk}$ and $Y_{o/oo}$ are the
220 biomass of *L. kaempferi* and *L. olgensis* with intra-specific plant interactions,

221 respectively. $Y_{k/ko}$ and $Y_{o/ko}$ are the biomass of *L. kaempferi* and *L. olgensis* with
222 inter-specific plant interactions, respectively. Because the biomass of isolated plants is
223 higher than that of plants exposed to plant-plant interactions (Kozovits et al., 2005;
224 Guo et al., 2017), the NEI values are negative. A more negative NEI value indicates
225 that a plant suffers a more negative impact from its neighbor.

226

227 Soils were sampled following the last harvest. Using a soil corer (4 cm in diameter),
228 soil samples were collected from four pots of KK, KO and OO, as well as from
229 corresponding pots with N fertilization. The sampling position was at the center
230 between the two plants in a pot. Then loose soil was removed by careful shaking and
231 the tightly-adhering soil was sampled from root surfaces as rhizosphere soil.
232 Rhizosphere soil samples were collected also from the pots of K and O as well as
233 from corresponding pots with N fertilization. The sampling center was 5 cm away
234 from the basal stem. Each soil sample was divided into two equal subsamples: one
235 subsample was used for the analysis of soil properties, while the other subsample was
236 stored at -20 °C for the subsequent DNA extraction. Three subsamples from each
237 treatment were chosen for later DNA extraction.

238

239 *2.3. Determination of soil properties*

240

241 Soil pH was determined from soil-water suspensions (1:2.5 v/v). Available
242 phosphorus (AP) was extracted with sodium bicarbonate and then determined using

243 the molybdenum blue method (Xiong et al., 2015) Soil organic matter (SOM) was
244 determined using the potassium dichromate external heating method (Ciavatta et al.,
245 1991). Total N (TN) was determined by the Kjeldahl method (Buchi K370,
246 Switzerland). Soil NH_4^+ and NO_3^- were extracted with 1M KCl and the
247 indophenol-blue colorimetric and double wavelength (220 nm and 275 nm) methods
248 (Nie et al., 2018), respectively. Soil microbial biomass carbon (C_{Mic}), nitrogen (N_{Mic})
249 and phosphorus (P_{Mic}) were determined by the chloroform fumigation extraction
250 method (Yao et al., 2017). For the pot experiment, two-way ANOVAs were applied to
251 check the impact of N application and plant-plant interactions on the soil properties.

252

253 *2.4. DNA extraction and Illumina sequencing*

254

255 Genomic DNA was extracted using the Power Soil kit (MO BIO Laboratories,
256 Carlsbad, CA, USA) following the manufacturer's instructions. The primers used for
257 the amplification of the V4-V5 hypervariable regions of the bacterial 16S rRNA gene
258 were 515F (5'-GTGYCAGCMGCCGCGGTA-3') and 909R
259 (5'-CCCGYCAATTCMTTTRAGT-3') (Yao et al., 2017). The primers used for the
260 amplification of the partial ITS region of fungi were ITS4:
261 5'TCCTCCGCTTATTGATATGC-3' and ITS3_KYO2:
262 5'GATGAAGAACGYAGYRAA-3' (Bokulich and Mills, 2013). The PCR reactions,
263 quality control and purification processes followed the instructions of Yao et al.
264 (2014). A library was constructed and all sequences were generated with the

265 Illumina's MiSeq platform using paired-end reads. All above-mentioned steps were
266 completed at the Environmental Genome Platform of Chengdu Institute of Biology,
267 Chinese Academy of Sciences, China.

268

269 *2.5. Bioinformatics*

270

271 The paired-end reads of the bacterial 16S rRNA gene and fungal ITS region
272 amplicons were processed using the mothur pipeline (V.1.35.1) (Schloss et al., 2009),
273 based on the MiSeq standard operating procedure (Kozich et al., 2013). For quality
274 control, the 16S and ITS sequences that contained ambiguous (N) bases and
275 homopolymers longer than 8 nucleotides were screened out. The remaining sequences
276 were pre-clustered to allow for up to 1 bp difference per 100 bp bases to remove
277 potential sequencing errors before the identification of the chimeric sequences using
278 the UCHIME algorithm. The SILVA full length reference sequences (V.128) were
279 used for the alignments of the bacterial 16S rRNA sequences. For the fungal ITS
280 sequence analysis, due to a lack of reference templates for sequence alignment, we
281 trimmed the raw sequences to the same size (300 bp) after the removal of the forward
282 primer sequence. After removing the chimeric sequences, the unique bacterial 16S
283 rRNA gene sequences were classified using the SILVA reference database (V.128),
284 and fungal ITS sequences were classified using the mothur-formatted UNITE ITS
285 reference database (UNITE v6_sh_99) with the default bootstrapping algorithm
286 (cutoff value 80%). All sequences were assigned to operational taxonomic units

287 (OTUs) using the OptiClust clustering algorithm at 97% similarity. Bacterial 16S
288 sequences were assigned to OTUs using classify as the split method, whereas fungal
289 ITS sequences were assigned into OTUs using fasta as the split method based on
290 nearest neighbor clustering. Singletons were removed from both bacterial and fungal
291 ITS datasets. For the bacterial OTU dataset, OTUs that were classified as
292 non-bacterial or chloroplast were removed. For the fungal OTU dataset, all non-fungal
293 OTUs were removed. The raw sequencing data were deposited in the NCBI Sequence
294 Read Archive (SRA) database under accession no. SRP125300.

295

296 *2.6. Alpha diversity estimation*

297

298 The alpha diversity of bacteria and fungi in each sample was rarefied and estimated at
299 the sampling depths of 2873 and 5558, respectively, using the R package phyloseq
300 (McMurdie and Holmes, 2013). The normalization of the alpha diversity data was
301 visualized using Q-Q plots. The homogeneity of variances was examined using the
302 Bartlett test. The differences of the means between treatments and their interactions
303 were tested using ANOVA, assuming that the alpha diversity data were normally
304 distributed and the variances were equal among treatment groups. The non-parametric
305 Kruskal-Wallis rank sum test of each single treatment factor was used when the data
306 were not normally distributed or the group variances were heterogeneous. Tukey's
307 multiple comparisons of means were used when the differences between groups were
308 significant. The significance of the differences was concluded at the 95% confidence

309 level ($P < 0.05$).

310

311 2.7. Beta diversity estimation

312

313 The relative abundance of each OTU was calculated by dividing its read count by the

314 total read count of the corresponding sample, prior to the beta diversity analysis.

315 Principal coordinate analyses (PCoA) were applied to the relative abundance data to

316 visualize the broad pattern of bacterial and fungal communities between treatment

317 groups based on the Bray-Curtis distance, using the R package phyloseq (McMurdie

318 and Holmes, 2013). PERMANOVA was used to assess, whether the treatment groups

319 of Species and N fertilization and their interaction resulted in a different bacterial

320 community composition with the default 999 permutations. Betadisper was used to

321 test, whether the dispersions of observations between the treatment groups were equal

322 with 999 permutations. PERMANOVA and Betadisper were performed based on the

323 Bray-Curtis distance using the R package Vegan (Oksanen et al., 2007). Pairwise

324 PERMANOVA tests were used when the differences were significant between a priori

325 groups following PERMANOVA tests. Constrained analysis of principle coordinates

326 (CAP) was performed to visualize the significant differences in the community

327 composition between the treatment groups based on the Bray-Curtis distance using the

328 R package phyloseq (McMurdie and Holmes, 2013) with 999 permutations. The

329 significance of the differences was concluded at the 95% confidence level ($P < 0.05$).

330

331 2.8. Biomarker discovery

332

333 The linear discriminant analysis effect size LEfSe (Segata et al., 2011) was used to
334 determine the differentially abundant OTUs 1) among treatments without interaction
335 (isolated *L. kaempferi* and *L. olgensis*) and with intra-specific and inter-specific
336 interactions, irrespective of the N treatment (Class: Interaction; Subclass: N treatment),
337 and 2) between the N-treated and control pots (Class: N treatment; Subclass:
338 Interaction) in the pot experiment. For the soil sampled from the three plantations,
339 LEfSe was used to explore the OTUs with differential abundance between the
340 different interaction patterns (Class: Interaction).

341

342

343

344

345

346

347

348

349

350

351 **3. Results**

352

353 *3.1. Neighbor effect index*

354

355 *L. kaempferi* suffered stronger negative effects from its conspecific neighbor
356 compared to *L. olgensis*, regardless of N fertilization. However, the negative effect
357 declined when *L. kaempferi* was grown with *L. olgensis* under N fertilization. *L.*
358 *olgensis* suffered less negative effect from *L. kaempferi* when they were grown
359 together (Fig. 1).

360

361 *3.2. Changes in soil properties*

362

363 The mixed plantation soil had higher NH_4^+ , C_{Mic} and N_{Mic} contents compared to soil
364 from the two monoculture plantations (Table S1). In the pot experiment, soil pH,
365 SOM and the NO_3^- concentration were significantly impacted by N fertilization and
366 plant-plant interactions (Table 1). N fertilization decreased rhizosphere soil pH in
367 each treatment (Table S2). The intra-specific plant interactions of KK showed lower
368 NH_4^+ , C_{Mic} and N_{Mic} compared to KO under N fertilization (Table S2).

369

370 *3.3. Taxonomic composition and alpha diversity*

371

372 The bacterial communities of the bulk soil samples from the three plantations and
373 from the pot experiment were dominated by *Proteobacteria*, *Acidobacteria* and
374 *Actinobacteria*, whereas the fungal communities were dominated by *Basidiomycota*

375 and *Ascomycota* (Figs. 2 and 3, Supplementary Fig. S1). The abundance of
376 *Basidiomycota* was lower, whereas the abundance of *Ascomycota* became higher in
377 KO and OO relative to KK, especially under N fertilization in the pot experiment (Fig.
378 3).

379

380 In the bulk soil samples from the three plantations, the average Shannon values of the
381 bacterial and fungal communities were 5.98 and 2.52, respectively, and neither intra-
382 nor inter-specific plant interactions had any effects on the bacterial and fungal
383 diversity ($P > 0.05$, Supplementary Fig. S2). In the rhizosphere soil of single-plant
384 pots, N treatment showed no significant effect on bacterial and fungal diversity
385 ($P > 0.05$, Figs. 3a and 4a). However, N treatment increased bacterial diversity ($P <$
386 0.001 , Fig. 4b) and also fungal diversity in the rhizosphere soil of KO ($P < 0.05$, Fig.
387 5d). In addition, plant-plant interactions had a significant impact on bacterial diversity
388 ($P < 0.001$, Fig. 4b). For instance, the rhizosphere soil of KK exhibited a lower
389 bacterial diversity compared to KO ($P < 0.001$) and OO ($P < 0.001$).

390

391 3.4. Beta diversity

392

393 The PCoA showed that the plantation soils had distinct compositions of both bacterial
394 and fungal communities (Supplementary Fig. S3), unlike the rhizosphere soils from
395 the pot experiment (Supplementary Fig. S4). Intra- and inter-specific interactions had
396 a significant effect on bacterial and fungal community structures in soil from the three

397 plantations (PERMANOVA, $P < 0.01$, Supplementary Fig. S5). In the rhizosphere soil
398 of double-plant pots, plant-plant interactions and N treatment significantly affected
399 both the bacterial and fungal community composition (Supplementary Figs. S6e-h,
400 PERMANOVA, $P < 0.01$). Pairwise PERMANOVA demonstrated that the structure of
401 both bacterial and fungal communities in the rhizosphere soil of KK was different
402 from those of KO and OO (Supplementary Figs. S6e and f, PERMANOVA, $P < 0.05$).

403

404 3.5. Biomarker discovery

405

406 We used the LEfSe analysis to discover biomarkers (different abundances of bacterial
407 and fungal taxa) in the plantation and pot experiment soil. From phyla to genera, the
408 three plantations showed their specific influence on bacterial and fungal compositions
409 (Supplementary Fig. S7).

410

411 In the rhizosphere soil of single plants, the abundance of the bacterial orders like
412 *Ktedonobacterales* and *Sphingomonadales* and the fungal order *Chaetothyriales* and
413 the genus *Ciliophora* decreased, whereas the fungal family *Corynesporascaceae*
414 became more prevalent under N fertilization (Fig. 6a and b). In the rhizosphere soil of
415 double-plant pots, N fertilization significantly increased the abundance of the bacterial
416 orders *Gemmatimonadales*, *Xanthomonadales*, *Propionibacteriales*, *Methylophilales*
417 and *JG37-AG-4*, whereas the abundance of the phylum *Acidobacteria*, including the
418 orders *Solibacterales*, *Blastocatellales* and *Subgroup 6* and the orders *Rhodocyclales*

419 and *Rhodospirillales*, reduced under N fertilization (Fig. 7a). The abundance of
420 *Basidiomycota*, such as class *Tritirachiomycetes* and order *Boletales*, reduced but that
421 of *Ascomycota*, such as the order *Hypocreales*, increased under N fertilization in the
422 double-plant pots (Fig. 7b).

423

424 In the pot experiment, we also identified the bacterial and fungal taxa responding to
425 species and plant interactions, irrespective of N fertilization (Figs. 6c and d, Fig. 8).

426 The orders *Rhodospirillales*, *Rhizobiales*, *Acidimicrobiales* and *Acidobacteriales* were
427 more abundant in the rhizosphere soils of K than O (Fig. 6c). Among fungal taxa,
428 *Ascomycota* preferred the soil of O, whereas *Basidiomycota* and the order
429 *Rhizophydiales* preferred the soil of K (Fig. 6d).

430

431 In the double-plant pots, the bacterial orders *Acidobacteriales* and *Micrococcales*
432 were generally more abundant in the soil of KK, whereas *Actinobacteria* became
433 more prevalent in the soil of KO (Fig. 8a). The taxa from the phyla *Firmicutes*,
434 *Chloroflexi* and *Planctomycetes* were more frequent in the soil of OO (Fig. 8a). The
435 fungal taxa *Basidiomycota* were generally more abundant in the rhizosphere soil of
436 KK, whereas fungi from the other dominant phylum *Ascomycota* were more abundant
437 in the rhizosphere soil of OO (Fig. 8b). In addition, the orders *Microascales* and
438 *Hymenochaetales* became more abundant in the rhizosphere soil of KO (Fig. 8b).

439 **4. Discussion**

440

441 Many plant attributes, such as physiological processes, root exudates and leaf N
442 content largely depend on the identity of a plant's neighbors and on the soil nitrogen
443 content (Broz et al., 2010; Pierik et al., 2013; Chen et al., 2017). It follows that soil
444 microbial communities can be differently impacted depending on, whether plants
445 grow in the absence of neighbors or in the presence of conspecific or heterospecific
446 neighbors in different environments.

447

448 *4.1. Different plantations have specific bacterial and fungal communities*

449

450 Different plantations have specific effects on soil chemical properties (Nazaries et al.,
451 2015; Suz et al., 2017). For example, a pine plantation was found to decrease soil pH
452 and increase the soil nitrate (NO_3^-) content, whereas a birch plantation declined the
453 total carbon and NO_3^- contents (Nazaries et al., 2015). Our results also showed that
454 the chemical traits of soil were different among the three studied plantations,
455 particularly in the KK plantation. Differences in the chemical properties of soil largely
456 depend on plantation characteristics, such as litter production and decomposition
457 (Hättenschwiler et al., 2005; Helfrich et al., 2015), nutrient mineralization (Richards
458 et al., 2010) and root exudates (Chen et al., 2014). Guo et al. (2016) have revealed
459 that inter-specific interactions of *L. kaempferi* and *L. olgensis* decrease leaf C/N
460 compared to intra-specific plant interactions without N fertilization, which may lead
461 to biochemically heterogeneous plant litter in different types of plantations.

462

463 It has been shown that functions, compositions or diversities of soil microbe
464 communities closely correlate with soil properties, such as pH, and soil carbon and
465 nitrogen contents (Bakker et al., 2013b, 2014; Nazaries et al., 2015; Gunina et al.,
466 2017; Suz et al., 2017). Suz et al. (2017) found that there were no significant
467 differences in root colonization by generalist ectomycorrhizal fungi between mixed
468 and pure plantations but, on average, there were more generalist ectomycorrhizal
469 fungi in mixed plantations, where plants connect with their neighbors through
470 common mycorrhizal networks. In this study, the phylum level abundance and the
471 Shannon index of bacterial and fungal communities showed little difference; however,
472 the beta diversity and biomarkers of bacterial and fungal taxa displayed greater
473 dissimilarities, possibly different nutrient translocation or litter decomposition. For
474 example, the abundance of *Basidiomycota* was higher but the abundance of
475 *Ascomycota* was lower in KO and OO plantations compared with KK. The ability to
476 degrade lignin is mainly conserved in *Basidiomycota* (Baldrian, 2006). The
477 accumulation of soil organic matter exerts a positive and significant effect on the
478 fungal beta-diversity (Cline and Zak, 2015), thus implying an important role of soil
479 organic matter in shaping fungal communities and in regulating ecosystem carbon
480 dynamics (Cheeke et al., 2017).

481

482 *4.2. Direct and indirect effects of N fertilization on microbial communities*

483

484 Nitrogen addition directly drives changes in bacterial and fungal communities in

485 different terrestrial ecosystems by increasing N availability (Cox et al., 2010; Fierer et
486 al., 2012; Yuan et al., 2016; Treseder et al., 2018; Zhu et al., 2018). Our results found
487 a decrease in the abundance of *Acidobacteria* after N fertilization in the pot
488 experiment. An enhanced soil N availability increased the abundance of copiotrophic
489 bacterial taxa, including *Proteobacteria* and *Bacteroidetes*, but lowered the proportion
490 of oligotrophic *Acidobacteria* (Fierer et al., 2012). Similarly, Yuan et al., (2016) have
491 found that *Acidobacteria* show a negative response to N fertilization. For fungal
492 communities, shifts in the fungal composition have been considered to be the main
493 driver of the decomposition response to enhanced N addition (Baldrian, 2006; Cox et
494 al., 2010; Morrison et al., 2016). Cox et al. (2010) have reported that *Russula*
495 *ochroleuca* and *Thelephora terrestris* respond positively to increasing N, whereas
496 *Pseudotomentella tristis* and *Piloderma* respond negatively to increasing N. In our pot
497 experiment, we discovered a shift from *Basidiomycota* to *Ascomycota* under the
498 combined effects of *L. oligensis* and N fertilization. Changes in bacterial and fungal
499 communities reflect corresponding alterations in functional consequences (Cox et al.,
500 2010; Dietrich et al., 2017).

501

502 Nitrogen fertilization affects soil microbial communities also indirectly through
503 changing chemical properties of soil. We found that N fertilization significantly
504 impacted soil pH and soil organic matter, which indirectly control microbial
505 communities (Fierer and Jackson, 2006; Cline and Zak, 2015; Yuan et al., 2016;
506 Carrara et al., 2018). Soil pH is a major driver in shaping soil microbial communities

507 (Constancias et al., 2015; Ochoa-Hueso et al., 2018). For instance, the relative
508 abundances of *Bacteroidetes*, *Planctomycetes* and *Thaumarchaeota* show a positive
509 correlation with soil pH across an agricultural landscape (Constancias et al., 2015).
510 Fierer and Jackson (2006) have suggested that any significant deviation from
511 extracellular pH should impose stress on single-celled organisms and restrict the
512 survival of taxa exposed to pH beyond their optimum.

513

514 *4.3. Effects of plant-plant interactions*

515

516 Mortimer et al. (2015) have reported fewer significant differences in the chemical
517 properties of soil, whereas the ectomycorrhizal biomass, as well as the biomass of
518 Gram-positive, Gram-negative, and actinomycetes bacteria were significantly higher
519 in a mixed plantation than in a monoculture. Our results also demonstrated greater
520 differences in the composition and beta-diversity of soil microbes between
521 monocultures and a mixed-plantation. These results indicated that intra- and
522 inter-specific plant interactions possibly have great impacts on soil bacterial and
523 fungal communities. Our PERMANOVA and LEfse results indicated significant
524 differences between KK and KO as well as KK and OO, with a higher bacterial and
525 fungal diversity in KO compared to KK. At first, we observed a stronger neighbor
526 effect from conspecific species in both species. Available P, total N (under fertilization)
527 and NH_4^+ in the rhizosphere soil of a double-plant *L. kaempferi* (KK) pot were the
528 lowest among all double-plant pots, implying a higher demand for nutrients and a

529 stronger intra-specific competition than previously suggested by Kunstler et al. (2015).
530 Changes in soil nutrients induced by plant-plant competition contribute to alterations
531 in soil microbes (Fierer et al., 2012; Yuan et al., 2016; Treseder et al., 2018; Zhu et al.,
532 2018). Secondly, we observed a weakened neighbor effect caused by *L. kaempferi* on
533 *L. olgensis* in the inter-specific plant interactions. There is a general acceptance that
534 soil microbes are important in determining plant-plant interactions in different
535 environments (Hodge and Fitter, 2013; Keymer and Lankau, 2017). *L. olgensis* has
536 been found to enhance its ability to absorb NO_3^- under inter-specific plant interactions
537 (Guo et al., 2018), which may result from the increase in the fungal phylum
538 *Ascomycota* (Leroy et al., 2017). Finally, we found that after N fertilization, the
539 neighbor effects on both species were less negative in KO, which implied declined
540 inter-specific plant competition compared to KK and OO. In the present study, strong
541 combined effects between plant-plant interactions and N fertilization on the
542 rhizosphere soil were observed, particularly on the fungal communities. Both *L.*
543 *olgensis* and N fertilization increased the abundance of fungal species belonging to
544 *Ascomycota* but reduced abundance of *Basidiomycota* species, and their combined
545 effect even led to a dominance shift from *Basidiomycota* to *Ascomycota* and to a
546 higher fungal diversity. The changes in rhizosphere soil microbes were possibly an
547 important reason to drive changes in plant-plant interactions.

548

549 *4.4. Negative feedback from L. kaempferi plantation conditioned soil*

550

551 In the pot experiment, the soil was selected from a *L. kaempferi* plantation. Lower
552 bacterial and fungal diversities of KK were probably caused by *L. kaempferi*
553 continuing to prefer its own soil microbial communities, whereas *L. olgensis* selected
554 different microbes and increased bacterial and fungal diversity particularly under N
555 fertilization. Guo et al. (2017) have reported that the *L. kaempferi* conditioned soil
556 showed negative effects on the growth of *L. kaempferi*. The more negative neighbor
557 effect of *L. kaempferi* on its conspecific neighbors also confirmed that. A given plant
558 species alters its biological soil communities and abiotic soil properties that may
559 decrease its own growth rate, resulting in a negative feedback (Harrison and Bardegett,
560 2010; Hendriks et al., 2015). The introduction of *L. olgensis* to *L. kaempferi*
561 conditioned soil greatly changed the composition, abundance and diversity of
562 bacterial and fungal communities, particularly under N fertilization. Previously, Van
563 der Putten et al. (2016) have emphasized the role of bacterial and fungal communities
564 in the development of plant-soil feedback under environmental changes through plant
565 species loss or nitrogen enrichment. For example, the negative effects of Chinese fir
566 conditioned soil on Chinese fir is alleviated by an introduced foreign plant species
567 through increasing arbuscular mycorrhizal fungi and improving chemical properties of
568 soil (Xia et al., 2016). The above results suggested that an introduced foreign plant
569 species could improve soil conditions by changing microbial communities to alleviate
570 the negative soil feedback.

571

572 **5. Conclusions**

573

574 Our work demonstrated that intra- and inter-specific plant interactions were
575 differently affected by plant neighbors and N fertilization, and there were distinct
576 changes in soil microbial communities. In turn, the changing soil bacterial and fungal
577 communities probably influence plant-plant interactions. Based on the present study,
578 we suggest that *L. olgensis* and *L. kaempferi* growing together with N fertilization
579 may be an efficient way to promote the productivity of plantations. The introduced
580 foreign plant species and N fertilization could improve the chemical and biological
581 conditions of soil and, consequently, plantation productivity. However, further studies
582 are needed to explore, how soil microbes mediate plant-plant interactions in different
583 ecosystems all around the world. Such knowledge would be crucial for revealing and
584 understanding plant-soil-microbe relationships.

585

586

587 **Acknowledgments** This work was supported by the Talent Program of the Hangzhou
588 Normal University (2016QDL020). We are grateful to the Qingyuan Experimental
589 Station, Institute of Applied Ecology, Chinese Academy of Sciences for providing
590 experimental sites and support.

591

592 **References**

593 Agathokleous, E., Vanderstock, A., Kita, K., Koike, T., 2017. Stem and crown growth
594 of Japanese larch and its hybrid F₁ grown in two soils and exposed to two

595 free-air O₃ regimes. Environmental Science and Pollution Research 24,
596 6634-6647.

597 Bakker, P.A.H.M., Berendsen, R.L., Doornbos, R.F., Wintermans, P.C.A., Pieterse,
598 C.M.J., 2013a. The rhizosphere revisited: root microbiomics. Frontiers in
599 Plant Science 4.

600 Bakker, M.G., Otto-Hanson, L., Lange, A.J., Bradeen, J.M., Kinkel, L.L., 2013b.
601 Plant monocultures produce more antagonistic soil *Streptomyces* communities
602 than high-diversity plant communities. Soil Biology and Biochemistry 65,
603 304-312.

604 Bakker, M.G., Schlatter, D.C., Otto-Hanson, L., Kinkel, L.L., 2014. Diffuse
605 symbioses: roles of plant-plant, plant-microbe and microbe-microbe
606 interactions in structuring the soil microbiome. Molecular Ecology 23,
607 1571-1583.

608 Baldrian, P., 2006. Fungal laccases-occurrence and properties. FEMS Microbiology
609 Reviews 30, 215-242.

610 Bokulich, N.A., Mills, D.A., 2013. Improved selection of internal transcribed
611 spacer-specific primers enables quantitative, ultra-high-throughput profiling of
612 fungal communities. Applied and Environmental Microbiology 79,
613 2519-2526.

614 Broz, A.K., Broeckling, C.D., De-la-Peña, C., Lewis, M.R., Greene, E., Callaway,
615 R.M., Sumner, L.W., Vivanco, J.M., 2010. Plant neighbor identity influences
616 plant biochemistry and physiology related to defense. BMC Plant Biology 10,

617 115.

618 Broadbent, A., Stevens, C.J., Peltzer, D.A., Ostle, N.J., Orwin, K.H., 2018.

619 Belowground competition drives invasive plant impact on native species

620 regardless of nitrogen availability. *Oecologia* 186, 577-587.

621 Carrara, J.E., Walter, C.A., Hawkins, J.S., Peterjohn, W.T., Averill, C., Brzostek, E.R.,

622 2018. Interactions among plants, bacteria, and fungi reduce extracellular

623 enzyme activities under long-term N fertilization. *Global Change Biology*

624 online.

625 Castelli, J.P., Casper, B.B., 2003. Intraspecific AM fungal variation contributes to

626 plant-fungal feedback in a serpentine grassland. *Ecology* 84, 323-336.

627 Chen, L.C., Wang, S.L., Wang, P., Kong, C.H., 2014. Autoinhibition and soil

628 allelochemical (cyclic dipeptide) levels in replanted Chinese fir

629 (*Cunninghamia lanceolata*) plantations. *Plant and Soil* 374, 793-801.

630 Chen, Y., Ma, S., Sun, J., Wang, X., Cheng, G., Lu, X., 2017. Chemical diversity and

631 incubation time affect non-additive responses of soil carbon and nitrogen

632 cycling to litter mixtures from an alpine steppe soil. *Soil Biology and*

633 *Biochemistry* 109, 124-134.

634 Ciavatta, C., Govi, M., Antisari, L.V., Sequi, P., 1991. Determination of organic

635 carbon in aqueous extracts of soils and fertilizers. *Communications in Soil*

636 *Science and Plant Analysis* 22, 795-807.

637 Cline, L.C., Zak, D.R., 2015. Soil microbial communities are shaped by plant-driven

638 changes in resource availability during secondary succession. *Ecology* 96,

639 3374-3385.

640 Constancias, F., Saby, N.P.A., Terrat, S., Dequiedt, S., Horrigue, W., Nowak, V.,
641 Guillemain, J.-P., Biju-Duval, L., Chemidlin Prevost-Boure, N., Ranjard, L.,
642 2015. Contrasting spatial patterns and ecological attributes of soil bacterial and
643 archaeal taxa across a landscape. *Microbiologyopen* 4, 518-531.

644 Cox, F., Barsoum, N., Lilleskov, E.A., Bidartondo, M.I., 2010. Nitrogen availability
645 is a primary determinant of conifer mycorrhizas across complex environmental
646 gradients. *Ecology Letters* 13, 1103-1113.

647 Deng, Q., Cheng, X., Yang, Y., Zhang, Q., Luo, Y., 2014. Carbon-nitrogen interactions
648 during afforestation in central China. *Soil Biology and Biochemistry* 69,
649 119-122.

650 Dietrich, P., Buchmann, T., Cesarz, S., Eisenhauer, N., Roscher, C., 2017.
651 Fertilization, soil and plant community characteristics determine soil microbial
652 activity in managed temperate grasslands. *Plant and Soil* 419, 189-199.

653 Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial
654 communities. *Proceedings of the National Academy of Sciences of the United*
655 *States of America* 103, 626-631.

656 Fierer, N., Lauber, C.L., Ramirez, K.S., Zaneveld, J., Bradford, M.A., Knight, R.,
657 2012. Comparative metagenomic, phylogenetic and physiological analyses of
658 soil microbial communities across nitrogen gradients. *ISME Journal* 6,
659 1007-1017.

660 Gunina, A., Smith, A.R., Godbold, D.L., Jones, D.L., Kuzyakov, Y., 2017. Response

661 of soil microbial community to afforestation with pure and mixed species.
662 Plant and Soil 412, 357-368.

663 Guo, Q., Li, J., Zhang, Y., Zhang, J., Lu, D., Korpelainen, H., Li, C., 2016.
664 Species-specific competition and N fertilization regulate non-structural
665 carbohydrate contents in two *Larix* species. Forest Ecology and Management
666 364, 60-69.

667 Guo, Q., Zhang, Y., Wang, D., Zhang, Y., Korpelainen, H., Li, C., 2017. Influence of
668 soil qualities on intra- and interspecific competition dynamics of *Larix*
669 *kaempferi* and *L. olgensis*. Environmental and Experimental Botany 135,
670 96-105.

671 Guo, Q., Song, H., Kang, J., Korpelainen, H., Li, C., 2018. Different responses in
672 leaf-level physiology to competition and facilitation under different soil types
673 and N fertilization. Environmental and Experimental Botany 150, 69-78.

674 Hättenschwiler, S., Aeschlimann, B., Coûteaux, M.M., Roy, J., Bonal, D., 2008. High
675 variation in foliage and leaf litter chemistry among 45 tree species of a
676 neotropical rainforest community. New Phytologist 179, 165-175.

677 Harrison, K.A., Bardgett, R.D., 2010. Influence of plant species and soil conditions on
678 plant-soil feedback in mixed grassland communities. Journal of Ecology 98,
679 384-395.

680 Helfrich, M., Ludwig, B., Thoms, C., Gleixner, G., Flessa, H., 2015. The role of soil
681 fungi and bacteria in plant litter decomposition and macroaggregate formation
682 determined using phospholipid fatty acids. Applied Soil Ecology 96, 261-264.

683 Hendriks, M., Ravenek, J.M., Smit-Tiekstra, A.E., van der Paauw, J.W., de Caluwe, H.,
684 van der Putten, W.H., de Kroon, H., Mommer, L., 2015. Spatial heterogeneity
685 of plant-soil feedback affects root interactions and interspecific competition.
686 *New Phytologist* 207, 830-840.

687 Hodge, A., Fitter, A.H., 2013. Microbial mediation of plant competition and
688 community structure. *Functional Ecology* 27, 865-875.

689 Keymer, D.P., Lankau, R.A., 2017. Disruption of plant-soil-microbial relationships
690 influences plant growth. *Journal of Ecology* 105, 816-827.

691 Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013.
692 Development of a dual-index sequencing strategy and curation pipeline for
693 analyzing amplicon sequence data on the MiSeq Illumina sequencing platform.
694 *Applied and Environmental Microbiology* 79, 5112-5120.

695 Kozovits, A.R., Matyssek, R., Blaschke, H., Göttlein, A., Grams, T.E.E., 2005.
696 Competition increasingly dominates the responsiveness of juvenile beech and
697 spruce to elevated CO₂ and/or O₃ concentrations throughout two subsequent
698 growing seasons. *Global Change Biology* 11, 1387-1401.

699 Kumar, M., Männistö, M.K., van Elsas, J.D., Nissinen, R.M., 2016. Plants impact
700 structure and function of bacterial communities in Arctic soils. *Plant and Soil*
701 399, 319-332.

702 Kunstler, G., Falster, D., Coomes, D.A., Hui, F., Kooyman, R.M., Laughlin, D.C.,
703 Poorter, L., Vanderwel, M., Vieilledent, G., Wright, S.J., Aiba, M., Baraloto,
704 C., Caspersen, J., Cornelissen, J.H.C., Gourlet-Fleury, S., Hanewinkel, M.,

705 Herault, B., Kattge, J., Kurokawa, H., Onoda, Y., Peñuelas, J., Poorter, H.,
706 Uriarte, M., Richardson, S., Ruiz-Benito, P., Sun, I.F., Ståhl, G., Swenson,
707 N.G., Thompson, J., Westerlund, B., Wirth, C., Zavala, M.A., Zeng, H.,
708 Zimmerman, J.K., Zimmermann, N.E., Westoby, M., 2015. Plant functional
709 traits have globally consistent effects on competition. *Nature* 529, 204.

710 Kuzyakov, Y., Xu, X., 2013. Competition between roots and microorganisms for
711 nitrogen: mechanisms and ecological relevance. *New Phytologist* 198,
712 656-669.

713 Leroy, C., Jauneau, A., Martinez, Y., Cabin-Flaman, A., Gibouin, D., Orivel, J.,
714 Séjalon-Delmas, N., 2017. Exploring fungus–plant N transfer in a tripartite
715 ant–plant–fungus mutualism. *Annals of Botany* 120, 417-426.

716 Li, J., Guo, Q., Zhang, J., Korpelainen, H., Li, C., 2016. Effects of nitrogen and
717 phosphorus supply on growth and physiological traits of two *Larix* species.
718 *Environmental and Experimental Botany* 130, 206-215.

719 Lovelock, C.E., Ewel, J.J., 2005. Links between tree species, symbiotic fungal
720 diversity and ecosystem functioning in simplified tropical ecosystems. *New
721 Phytologist* 167, 219-228.

722 Manea, A., Leishman, M.R., 2011. Competitive interactions between native and
723 invasive exotic plant species are altered under elevated carbon dioxide.
724 *Oecologia* 165, 735-744.

725 McKane, R.B., Johnson, L.C., Shaver, G.R., Nadelhoffer, K.J., Rastetter, E.B., Fry, B.,
726 Giblin, A.E., Kielland, K., Kwiatkowski, B.L., Laundre, J.A., Murray, G.,

727 2002. Resource-based niches provide a basis for plant species diversity and
728 dominance in arctic tundra. *Nature* 415, 68-71.

729 McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible
730 interactive analysis and graphics of microbiome census data. *PLoS One* 8,
731 e61217.

732 Morrison, E.W., Frey, S.D., Sadowsky, J.J., van Diepen, L.T.A., Thomas, W.K.,
733 Pringle, A., 2016. Chronic nitrogen additions fundamentally restructure the
734 soil fungal community in a temperate forest. *Fungal Ecology* 23, 48-57.

735 Mortimer, P., Gui, H., Xu, J., Zhang, C., Barrios, E., Hyde, K., 2015. Alder trees
736 enhance crop productivity and soil microbial biomass in tea plantations.
737 *Applied Soil Ecology* 96, 25-32.

738 Nazaries, L., Tottey, W., Robinson, L., Khachane, A., Al-Soud, W.A., Sørensen, S.,
739 Singh, B.K., 2015. Shifts in the microbial community structure explain the
740 response of soil respiration to land-use change but not to climate warming.
741 *Soil Biology and Biochemistry* 89, 123-134.

742 Nie, Y., Wang, M., Zhang, W., Ni, Z., Hashidoko, Y., Shen, W., 2018. Ammonium
743 nitrogen content is a dominant predictor of bacterial community composition
744 in an acidic forest soil with exogenous nitrogen enrichment. *Science of The*
745 *Total Environment* 624, 407-415.

746 O'Hehir, J., Nambiar, E., 2010. Productivity of three successive rotations of *P.*
747 *radiata* plantations in South Australia over a century. *Forest Ecology and*
748 *Management* 259, 1857-1869.

749 Ochoa-Hueso, R., Eldridge, D.J., Delgado-Baquerizo, M., Soliveres, S., Bowker,
750 M.A., Gross, N., Le Bagousse-Pinguet, Y., Quero, J.L., García-Gómez, M.,
751 Valencia, E., Arredondo, T., Beinticenco, L., Bran, D., Cea, A., Coaguila, D.,
752 Dougill, A.J., Espinosa, C.I., Gaitán, J., Guuroh, R.T., Guzman, E., Gutiérrez,
753 J.R., Hernández, R.M., Huber-Sannwald, E., Jeffries, T., Linstädter, A., Mau,
754 R.L., Monerris, J., Prina, A., Pucheta, E., Stavi, I., Thomas, A.D., Zaady, E.,
755 Singh, B.K., Maestre, F.T., 2018. Soil fungal abundance and plant functional
756 traits drive fertile island formation in global drylands. *Journal of Ecology* 106,
757 242-253.

758 Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J.,
759 Suggests, M., 2007. The vegan package. *Community Ecology Package* 10:
760 631-637.

761 Paul, M., Catterall, C.P., Pollard, P.C., Kanowski, J., 2010. Recovery of soil
762 properties and functions in different rainforest restoration pathways. *Forest*
763 *Ecology and Management* 259, 2083-2092.

764 Pierik, R., Mommer, L., Voisenek, L.A.C.J., 2013. Molecular mechanisms of plant
765 competition: neighbour detection and response strategies. *Functional Ecology*
766 27, 841-853.

767 Pivato, B., Bru, D., Busset, H., Deau, F., Matejicek, A., Philippot, L., Moreau, D.,
768 2017. Positive effects of plant association on rhizosphere microbial
769 communities depend on plant species involved and soil nitrogen level. *Soil*
770 *Biology and Biochemistry* 114, 1-4.

771 Richards, A.E., Forrester, D.I., Bauhus, J., Scherer-Lorenzen, M., 2010. The influence
772 of mixed tree plantations on the nutrition of individual species: a review. *Tree*
773 *Physiology* 30, 1192-1208.

774 Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B.,
775 Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres,
776 B., Thallinger, G.G., Van Horn, D.J., Weber, C.F., 2009. Introducing mothur:
777 Open-source, platform-independent, community-supported software for
778 describing and comparing microbial communities. *Applied and Environmental*
779 *Microbiology* 75, 7537-7541.

780 Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S.,
781 Huttenhower, C., 2011. Metagenomic biomarker discovery and explanation.
782 *Genome Biology* 12, R60.

783 Suz, L., Kallow, S., Reed, K., Bidartondo, M., Barsoum, N., 2017. Pine mycorrhizal
784 communities in pure and mixed pine-oak forests: Abiotic environment trumps
785 neighboring oak host effects. *Forest Ecology and Management* 406, 370-380.

786 Treseder, K.K., Allen, M.F., 2002. Direct nitrogen and phosphorus limitation of
787 arbuscular mycorrhizal fungi: a model and field test. *New Phytologist* 155,
788 507-515.

789 Treseder, K.K., Allen, E.B., Egerton-Warburton, L.M., Hart, M.M., Klironomos, J.N.,
790 Maherali, H., Tedersoo, L., 2018. Arbuscular mycorrhizal fungi as mediators
791 of ecosystem responses to nitrogen deposition: A trait-based predictive
792 framework. *Journal of Ecology* 106, 480-489.

793 van der Putten, W.H., Bradford, M.A., Brinkman, E.P., van de Voorde, T.F.J., Veen,
794 G.F., 2016. Where, when and how plant-soil feedback matters in a changing
795 world. *Functional Ecology* 30, 1109-1121.

796 Wilson, S.D., Tilman, D., 1991. Component of plant competition along an
797 experimental gradient of nitrogen availability. *Ecology* 72, 1050-1065.

798 Xia, Z.-C., Kong, C.-H., Chen, L.-C., Wang, P., Wang, S.-L., 2016. A broadleaf
799 species enhances an autotoxic conifers growth through belowground chemical
800 interactions. *Ecology* 97, 2283-2292.

801 Xiong, W., Zhao, Q., Zhao, J., Xun, W., Li, R., Zhang, R., Wu, H., Shen, Q., 2015.
802 Different continuous cropping spans significantly affect microbial community
803 membership and structure in a vanilla-grown soil as revealed by deep
804 pyrosequencing. *Microbial Ecology* 70, 209-218.

805 Yao, M., Rui, J., Li, J., Dai, Y., Bai, Y., Hedeneč, P., Wang, J., Zhang, S., Pei, K., Liu,
806 C., Wang, Y., He, Z., Frouz, J., Li, X., 2014. Rate-specific responses of
807 prokaryotic diversity and structure to nitrogen deposition in the *Leymus*
808 *chinensis* steppe. *Soil Biology and Biochemistry* 79, 81-90.

809 Yao, M., Rui, J., Niu, H., Heděneč, P., Li, J., He, Z., Wang, J., Cao, W., Li, X., 2017.
810 The differentiation of soil bacterial communities along a precipitation and
811 temperature gradient in the eastern Inner Mongolia steppe. *CATENA* 152,
812 47-56.

813 Yin, H., Wheeler, E., Phillips, R.P., 2014. Root-induced changes in nutrient cycling in
814 forests depend on exudation rates. *Soil Biology and Biochemistry* 78,

815 213-221.

816 Yuan, X., Knelman, J.E., Gasarch, E., Wang, D., Nemergut, D.R., Seastedt, T.R.,
817 2016. Plant community and soil chemistry responses to long-term nitrogen
818 inputs drive changes in alpine bacterial communities. *Ecology* 97, 1543-1554.

819 Zhu, C., Tian, G., Luo, G., Kong, Y., Guo, J., Wang, M., Guo, S., Ling, N., Shen, Q.,
820 2018. N-fertilizer-driven association between the arbuscular mycorrhizal
821 fungal community and diazotrophic community impacts wheat yield.
822 *Agriculture Ecosystems and Environment* 254, 191-201.

823

824 **Table 1** Effects of plant-plant interactions and N fertilization on soil properties (two-way ANOVA).

		pH	AP	SOM	TN	NH ₄ ⁺	NO ₃ ⁻	C _{Mic}	N _{Mic}	P _{Mic}
N	F	56.674	1.087	10.246	0.173	0.120	0.024	3.358	0.455	1.507
	P	<0.000	0.305	0.003	0.680	0.732	0.012	0.076	0.505	0.229
Interactions	F	8.297	0.140	26.386	48.904	11.358	7.143	12.470	1.686	0.528
	P	0.007	0.711	<0.000	<0.000	0.002	0.012	0.001	0.203	0.473
N* Interactions	F	0.002	0.301	0.119	1.460	0.003	0.936	1.103	0.153	0.153
	P	0.967	0.587	0.732	0.236	0.960	0.341	0.301	0.699	0.698

825 AP: available phosphorus, SOM: soil organic matter, TN: total nitrogen, C_{Mic}: soil microbial biomass, N_{Mic}: soil carbon nitrogen biomass and

826 P_{Mic}: soil carbon phosphorus biomass.

827 **Table 2** Summaries of PERMANOVA tests with 999 permutations based on Bray-Curtis distances.

Taxonomy	Experiment	Tree species or interactions		N treatment	
		PERMANOVA	Betadisper	PERMANOVA	Betadisper
Bacteria	Single-plant	ns	ns	ns	ns
	Double-plant	*	***	**	ns
	KO vs KK	**	ns	ns	ns
	KK vs OO	**	ns	ns	ns
	KO vs OO	ns	ns	ns	ns
	Plantations	***	ns	–	–
	KO vs KK	*	ns	–	–
	KK vs OO	**	ns	–	–
	KO vs OO	ns	ns	–	–
	Fungi	Single-plant	*	ns	ns
Double-plant		**	ns	***	ns
KO vs KK		*	ns	ns	ns
KK vs OO		**	ns	ns	ns
KO vs OO		ns	ns	ns	ns
Plantations		***	ns	–	–
KO vs KK		**	ns	–	–
KK vs OO		**	ns	–	–
KO vs OO		**	ns	–	–

828 When a plant was grown in isolation, we analyzed the effect of species, and when grown under intra- or
829 inter-specific interactions, we analyzed the effect of plant-plant interactions. KK and OO refer to intra-specific
830 interactions of *Larix kaempferi* and *L. olgensis*, respectively. KO refers to inter-specific interactions of the two
831 species. In the plantations, KK and OO refer to monoculture plantations of *Larix kaempferi* and *L. olgensis*,
832 respectively. KO refers to a mixed plantation of the two species.

833

834

835

836

837

838

839

840

841

842

843 **Figure legends**

844

845 **Figure 1** Neighbor effect index (NEI) of *L. kaempferi* and *L. olgensis* in different
846 plant-plant interactions. KK and OO refer to *L. kaempferi* and *L. olgensis* in
847 intra-specific plant-plant interactions, respectively. K/KO and O/KO refer to *L.*
848 *kaempferi* and *L. olgensis* in inter-specific plant-plant interactions, respectively. N-
849 and N+ refer to soil without and with N fertilization, respectively. Negative NEI
850 values indicate that a plant suffers negative impacts from its neighbor. Tukey's HSD
851 tests are conducted for multiple comparisons.

852

853 **Figure 2** Taxonomic composition of bacterial communities in the rhizosphere soil at
854 the phylum level under N fertilization. (a) Rhizosphere soil sampled from single-plant
855 pots (without plant-plant interactions); (b) rhizosphere soil sampled from pots with
856 two plants, where KK, KO and OO refer to *L. kaempferi* + *L. kaempferi*, *L. kaempferi*
857 + *L. olgensis* and *L. olgensis* + *L. olgensis*, respectively. N- and N+ refer to pots
858 without and with N fertilization, respectively. For bacteria, only phyla with relative
859 abundance over 1% are shown.

860

861 **Figure 3** Taxonomic composition of fungal communities in the rhizosphere soil at the
862 phylum level under N fertilization. (a) Rhizosphere soil sampled from single-plant
863 pots (without plant-plant interactions); (b) rhizosphere soil sampled from pots with
864 two plants, where KK, KO and OO refer to *L. kaempferi* + *L. kaempferi*, *L. kaempferi*

865 + *L. olgensis* and *L. olgensis* + *L. olgensis*, respectively. N- and N+ refer to pots
866 without and with N fertilization, respectively. For fungi, only phyla with relative
867 abundance over 1% are shown.

868

869 **Figure 4** Effects of different plant-plant interactions on the alpha diversity (Shannon
870 index) of rhizosphere soil bacteria under N fertilization. (a) Rhizosphere soil sampled
871 from single-plant pots (without plant-plant interactions); (b) rhizosphere soil sampled
872 from pots with two plants (intra- or inter-specific interactions). KK, KO and OO refer
873 to *L. kaempferi* + *L. kaempferi*, *L. kaempferi* + *L. olgensis* and *L. olgensis* + *L.*
874 *olgensis*, respectively. N- and N+ refer to pots without and with N fertilization in the
875 pot experiment, respectively. Tukey's HSD tests were conducted for multiple
876 comparisons.

877

878 **Figure 5** Effects of different plant-plant interactions on the alpha diversity (Shannon
879 index) of rhizosphere soil fungi under N fertilization. (a) Rhizosphere soil sampled
880 from single-plant pots (without plant-plant competition); (b) rhizosphere soil sampled
881 from pots with two plants (intra- or inter-specific plant interactions). KK, KO and OO
882 refer to *L. kaempferi* + *L. kaempferi*, *L. kaempferi* + *L. olgensis* and *L. olgensis* + *L.*
883 *olgensis*, respectively. N- and N+ refer to pot experiment without and with N
884 fertilization in the pot experiment, respectively. Tukey's HSD tests were conducted
885 for multiple comparisons.

886

887 **Figure 6** Bacterial (a and c) and fungal (b and d) taxa with different abundance
888 changes in single-plant pots between N-treated (N+) and control (N-) soil, irrespective
889 of plant species (a and b, Class: N treatment; Subclass: Species) and ii) between plant
890 species, irrespective of N treatment (a and b, Class: Species; Subclass: N treatment),
891 as detected by LEfSe analysis. The taxa with the absolute LDA scores over 3 and *P*
892 values less than 0.05 are shown.

893

894 **Figure 7** Bacterial (a) and fungal (b) taxa with different abundance changes in
895 two-plant pots between N-treated (N+) and control (N-) soil, irrespective of
896 plant-plant interactions (Class: N treatment; Subclass: Plant-plant interactions), as
897 detected by LEfSe analysis. The taxa with the absolute LDA scores over 3 and *P*
898 values less than 0.05 are shown.

899


900 **Figure 8** Bacterial (a) and fungal (b) taxa with different abundance changes in
901 two-plant pots between intra- and interspecific plant interactions, irrespective of N
902 treatment (Class: Plant-plant interactions, Subclass: N treatment), as detected by
903 LEfSe analysis. The taxa with the absolute LDA scores over 3 and *P* values less than
904 0.05 are shown. KK, KO and OO refer to plant-plant interactions *L. kaempferi* + *L.*
905 *kaempferi*, *L. kaempferi* + *L. olgensis* and *L. olgensis* + *L. olgensis*, respectively.

906

907

908


909 **Figure 1**



910

911

912 **Figure 2**



913

914

915

916

917

918

919

920

921

922

923


924

925

926

927

928 **Figure 3**



929

930

931

932

933

934

935

936

937

938

939


940

941

942

943

944 **Figure 4**



945

946

947

948

949

950

951

952

953

954

955


956

957

958


959

960 **Figure 5**




961

962 **Figure 6**



979 **Figure 7**



980

981

982

983

984

985

986

987

988

989

