



Plasma 25-hydroxyvitamin D concentration and metabolic syndrome among middle-aged and elderly Chinese individuals

Citation

Lu, Ling, Zhijie Yu, An Pan, Frank B. Hu, Oscar H. Franco, Huaixing Li, Xiaoying Li, Xilin Yang, Yan Chen, and Xu Lin. 2009. Plasma 25-hydroxyvitamin D concentration and metabolic syndrome among middle-aged and elderly Chinese individuals. *Diabetes Care* 32(7): 1278-1283.

Published Version

doi:10.2337/dc09-0209

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:4879212>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Plasma 25-Hydroxyvitamin D Concentration and Metabolic Syndrome Among Middle-Aged and Elderly Chinese Individuals

LING LU, MSC¹
 ZHIJIE YU, PHD¹
 AN PAN, MSC¹
 FRANK B. HU, MD, PHD²
 OSCAR H. FRANCO, MD, DSC, PHD³

HUAXING LI, PHD¹
 XIAOYING LI, MD⁴
 XILIN YANG, PHD⁵
 YAN CHEN, PHD¹
 XU LIN, MD, PHD¹

OBJECTIVE — To evaluate the association between 25-hydroxyvitamin D [25(OH)D] and metabolic syndrome in the Chinese population.

RESEARCH DESIGN AND METHODS — Plasma 25(OH)D was measured in a cross-sectional sample of 1,443 men and 1,819 women aged 50–70 years from Beijing and Shanghai. Metabolic syndrome was defined according to the updated National Cholesterol Education Program Adult Treatment Panel III criteria for Asian Americans. Fasting plasma glucose, insulin, lipid profile, A1C, and inflammatory markers were measured.

RESULTS — The geometric mean of plasma 25(OH)D was 40.4 nmol/l, and percentages of vitamin D deficiency [25(OH)D <50 nmol/l] and insufficiency [50 ≤ 25(OH)D <75 nmol/l] were 69.2 and 24.4%, respectively. Compared with the highest 25(OH)D quintile (≥57.7 nmol/l), the odds ratio for metabolic syndrome in the lowest quintile (≤28.7 nmol/l) was 1.52 (95% CI 1.17–1.98, $P_{\text{trend}} = 0.0002$) after multiple adjustment. Significant inverse associations also existed between 25(OH)D and individual metabolic syndrome components plus A1C. Moreover, we observed significant inverse associations of 25(OH)D with fasting insulin and the insulin resistance index (homeostasis model assessment of insulin resistance [HOMA-IR]) in overweight and obese individuals (BMI ≥24 kg/m²) but not in their normal-weight counterparts (test for interaction: $P = 0.0363$ and 0.0187 for insulin and HOMA-IR, respectively).

CONCLUSIONS — Vitamin D deficiency is common in the middle-aged and elderly Chinese population, and a low 25(OH)D level is significantly associated with an increased risk of having metabolic syndrome and insulin resistance. Prospective studies and randomized clinical trials are warranted to determine the role of 25(OH)D in the development of metabolic syndrome and related metabolic diseases.

Diabetes Care 32:1278–1283, 2009

Vitamin D deficiency is now recognized as a worldwide concern (1). A growing body of evidence suggests that 25-hydroxyvitamin D [25(OH)D], a generally accepted indicator of vitamin D status, is inversely associated with adiposity, glucose homeostasis, lipid profiles, and blood pressure along with its classic

role in calcium homeostasis and bone metabolism (1–6). Even though the underlying mechanism has not been well understood, vitamin D appears to exert effects through direct modulation of gene expression via vitamin D receptors (VDRs) (1) and through regulation of extra- and intracellular calcium (1,7).

Metabolic syndrome, a constellation of cardiometabolic disease risk factors, has become a global epidemic (8). Several epidemiologic studies (5,6,9,10) have suggested that 25(OH)D status is inversely associated with metabolic syndrome in western populations, although data for morbidly obese individuals are inconsistent (11,12). Nevertheless, evidence from the Asian population is limited. Because of ethnic differences in vitamin D metabolism and its nutritional status indicated by previous studies (3,13), it is not clear whether the findings from western populations could be extrapolated directly to Asian individuals. With rapid nutrition and lifestyle transitions in the last 20 years, metabolic syndrome has become one of the most widespread health problems in Asian countries (8). However, little is known regarding whether vitamin D deficiency plays an important role in the heightened prevalence of metabolic syndrome and other metabolic disorders among Asian individuals. Therefore, the aim of our study was to evaluate the plasma 25(OH)D concentration and its association with metabolic syndrome among Chinese individuals aged 50–70 years.

RESEARCH DESIGN AND METHODS

The present study is a part of the Nutrition and Health of Aging Population in China (NHAPC) project, which is a population-based cross-sectional study among noninstitutionalized Chinese individuals aged 50–70 years in Beijing (latitude 40° north) and Shanghai (latitude 31° north), China. The details of the study design have been described previously (14). In brief, the study was conducted simultaneously in both

From the ¹Key Laboratory of Nutrition and Metabolism, Institute for Nutritional Sciences, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences and Graduate School of the Chinese Academy of Sciences, Shanghai, China; the ²Departments of Nutrition and Epidemiology, Harvard School of Public Health, Boston, Massachusetts; the ³Health Sciences Research Institute, University of Warwick, Coventry, U.K.; the ⁴Shanghai Clinical Center for Endocrine and Metabolic Diseases, Shanghai Institute of Endocrinology and Metabolism, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China; and the ⁵Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong SAR, China.

Corresponding author: Xu Lin, xlin@sibs.ac.cn.

Received 4 February 2009 and accepted 4 April 2009.

Published ahead of print at <http://care.diabetesjournals.org> on 14 April 2009. DOI: 10.2337/dc09-0209.

L.L. and Z.Y. contributed equally to this work.

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

geographic locations from April to June 2005. In each city, one rural and two urban districts were selected. A total of 3,289 eligible participants (1,458 men and 1,831 women) were recruited. After excluding those who did not have adequate blood samples for 25(OH)D measurement ($n = 27$), 3,262 individuals were eligible for the present analysis. The study was approved by the institutional review board of the Institute for Nutritional Sciences, and all participants provided informed consent.

Data collection

In a home interview, a standardized questionnaire was used by trained health workers to collect information such as age, sex, geographic location (Beijing/Shanghai), residential region (urban/rural), visit date (April or May/June), education level (≤ 6 , 7–9, or ≥ 10 years in school), smoking (current, former, or never), alcohol drinking (yes/no), and self-reported diabetes, hypertension, dyslipidemia, coronary heart disease (CHD), stroke, and medication use. Physical activity level was classified as low, moderate, or high according to the International Physical Activity Questionnaire (short last 7-day format) scoring protocol with minor modification (14). According to participants' responses to the corresponding questions, family history of cardiovascular disease (CVD) or diabetes was classified as yes or no.

After the home interview, all participants were invited to attend a physical examination after an overnight fast. Measurements of weight, height, waist circumference, and blood pressure have been described previously (14). BMI was calculated as weight in kilograms divided by the square of height in meters and categorized as normal weight ($< 24.0 \text{ kg/m}^2$) or overweight or obesity ($\geq 24.0 \text{ kg/m}^2$), according to the criteria for Chinese individuals (15).

Laboratory methods

Peripheral venous blood samples were collected in tubes containing liquid EDTA and centrifuged at 4°C. After being frozen, the samples were shipped on dry ice to the Institute for Nutritional Sciences and stored at -80°C until analysis. Plasma glucose, triglycerides, and HDL cholesterol were measured enzymatically on an automatic analyzer (Hitachi 7080) with reagents purchased from Wako Pure Chemical Industries (Osaka, Japan). A1C was quantified from resolved erythrocytes

with an automated immunoassay (Tinaquant Hemoglobin A1C II; Roche Diagnostics, Indianapolis, IN), which was standardized according to the Diabetes Control and Complications Trial/National Glycohemoglobin Standardization Program. Plasma high-sensitive C-reactive protein (CRP) was measured by a particle-enhanced immunoturbidimetric assay (Ultrasensitive CRP kit; Orion Diagnostica, Espoo, Finland). Interleukin-6 (IL-6) was measured using a high-sensitivity ELISA (Quantikine HS IL-6 Immunoassay; R&D Systems, Minneapolis, MN). Fasting insulin was determined by radioimmunoassay (Linco Research, St. Charles, MO). The insulin resistance index (homeostasis model assessment of insulin resistance [HOMA-IR]) was calculated using updated homeostasis model assessment methods (<http://www.dtu.ox.ac.uk/>). The plasma 25(OH)D concentration was assayed with a radioimmunoassay kit (DiaSorin, Stillwater, MN). Vitamin D nutritional status was assessed as "sufficiency" ($\geq 75 \text{ nmol/l}$), "insufficiency" [$50 \leq 25(\text{OH})\text{D} < 75 \text{ nmol/l}$], or "deficiency" ($< 50 \text{ nmol/l}$) (1). All of the intra- and interassay coefficients of variation were $< 13\%$.

Definition of the metabolic syndrome

Metabolic syndrome was defined using the updated National Cholesterol Education Program Adult Treatment Panel III criteria for Asian Americans (16) as presentation of three or more of the following components: 1) waist circumference $\geq 90 \text{ cm}$ for men or $\geq 80 \text{ cm}$ for women; 2) triglycerides $\geq 1.7 \text{ mmol/l}$; 3) HDL cholesterol $< 1.03 \text{ mmol/l}$ for men or $< 1.30 \text{ mmol/l}$ for women; 4) blood pressure $\geq 130/85 \text{ mmHg}$ or current use of antihypertensive medications; and 5) fasting glucose $\geq 5.6 \text{ mmol/l}$ or previously diagnosed type 2 diabetes or using oral hypoglycemic agents or insulin.

Statistical methods

The difference in plasma 25(OH)D concentrations between participants with and without metabolic syndrome was tested by the general linear model with controls for age, sex, geographic location, residential region, and visit date. A logistic regression model was used to evaluate the odds ratios (ORs) and 95% CIs of having metabolic syndrome for each quintile of 25(OH)D compared with the highest quintile with adjustment for potential confounders that were suggested in pre-

vious studies (3,5,6,9,17). In addition, effects of inflammatory factors on the association between 25(OH)D and metabolic syndrome were tested by models adjusted for CRP and IL-6. Tests of linear trend across increasing quintiles of 25(OH)D were performed by assigning the median value to each quintile and treating it as a continuous variable. A multiple linear regression model was used to assess the association of 25(OH)D (continuous) with fasting glucose, A1C, waist circumference, triglycerides, HDL cholesterol, blood pressure, fasting insulin, and HOMA-IR. A likelihood ratio test was used to test the potential modifying effect of sex and obesity status (< 24 or $\geq 24 \text{ kg/m}^2$). When appropriate, natural log-transformed values were used for the analyses. Statistical inference was made with $P < 0.05$ (two-sided). The statistical analyses were performed using Stata 9.2 (StataCorp, College Station, TX).

RESULTS— The geometric mean of 25(OH)D was 40.4 nmol/l in our study participants. The percentages of vitamin D deficiency, insufficiency, and sufficiency were 69.2, 24.4, and 6.4%, respectively. Across 25(OH)D quintiles, subjects in lower 25(OH)D quintiles were more likely to be northern (Beijing) and urban residents, to have a physical examination in April, to have higher educational levels, and to have a family history of CVD and diabetes. They also tended to have an adverse cardiometabolic profile. In addition, these subjects were less physically active and were less likely to be male and alcohol drinkers than those in higher 25(OH)D quintiles (Table 1).

Plasma 25(OH)D concentrations were lower in participants with metabolic syndrome than in those without metabolic syndrome (39.3 vs. 41.6 nmol/l, $P < 0.0001$). The risk of having metabolic syndrome increased progressively across the highest to the lowest quintiles of 25(OH)D with the ORs of 1.31 (95% CI 1.03–1.66), 1.36 (95% CI 1.07–1.73), 1.67 (95% CI 1.31–2.13), and 1.62 (95% CI 1.26–2.08), respectively ($P_{\text{trend}} < 0.0001$) (Table 2) after adjustment for age, sex, geographic location, residential region, and visit date (model 1). Further controls for education, behavioral factors, self-reported CHD, and stroke and family history of CVD and diabetes (model 2) did not materially change the inverse associations. After additional adjustment for CRP and IL-6 (model 3), the ORs attenuated slightly, but remained signifi-

Table 1—Characteristics of study participants according to 25(OH)D quintiles

	Quintiles of 25(OH)D				
	Q1: ≤28.7 nmol/l	Q2: 28.8–36.8 nmol/l	Q3: 36.9–45.5 nmol/l	Q4: 45.6–57.6 nmol/l	Q5: ≥57.7 nmol/l
<i>n</i>	652	653	652	653	652
Age (years)	58.8 ± 6.3	58.1 ± 6.0	58.8 ± 5.9	58.2 ± 5.9	59.1 ± 5.9
Male sex	233 (35.7)	252 (38.6)	279 (42.8)	311 (47.6)	368 (56.4)
Urban residents	423 (64.9)	384 (58.8)	367 (56.3)	260 (39.8)	189 (29.0)
Residents of Beijing	466 (71.5)	419 (64.2)	331 (50.8)	238 (36.5)	168 (25.8)
Visit date (April)	433 (66.4)	389 (59.6)	334 (51.2)	245 (37.5)	183 (28.1)
Smoking					
Never	395 (60.6)	422 (64.6)	423 (64.9)	409 (62.6)	376 (57.7)
Former	67 (10.3)	51 (7.8)	63 (9.7)	59 (9.0)	87 (13.3)
Current	190 (29.1)	180 (27.6)	166 (25.5)	185 (28.3)	189 (29.0)
Alcohol drinking, yes	167 (25.6)	183 (28.0)	189 (29.0)	193 (29.6)	196 (30.1)
Physical activity					
Low	63 (9.7)	40 (6.1)	47 (7.2)	50 (7.7)	43 (6.6)
Moderate	310 (47.6)	307 (47.0)	294 (45.1)	228 (34.9)	230 (35.3)
High	279 (42.8)	306 (46.9)	311 (47.7)	375 (57.4)	379 (58.1)
Education					
0–6 years	201 (30.8)	223 (34.2)	240 (36.8)	317 (48.6)	371 (56.9)
7–9 years	278 (42.6)	254 (38.9)	249 (38.2)	201 (30.8)	177 (27.2)
≥10 years	173 (26.5)	176 (27.0)	163 (25.0)	135 (20.7)	104 (16.0)
Hypertension*	373 (57.2)	372 (57.0)	367 (56.3)	329 (50.4)	344 (52.8)
Self-reported CHD†	67 (10.6)	54 (8.4)	48 (7.6)	29 (4.6)	28 (4.4)
Self-reported stroke‡	33 (5.1)	32 (4.9)	23 (3.5)	24 (3.7)	19 (2.9)
Family history of diabetes	105 (16.1)	89 (13.6)	86 (13.2)	93 (14.2)	74 (11.4)
Family history of CVD	179 (27.5)	154 (23.6)	152 (23.3)	149 (22.8)	97 (14.9)
BMI (kg/m ²)	24.9 ± 3.8	24.9 ± 3.8	24.6 ± 3.6	24.3 ± 3.2	23.5 ± 3.3
Waist circumference (cm)	85.2 ± 10.5	84.7 ± 10.7	84.2 ± 10.8	83.2 ± 10.1	81.3 ± 10.2
Fasting glucose (mmol/l)	5.52 (5.08–6.09)	5.46 (5.09–6.00)	5.46 (5.09–5.98)	5.42 (4.99–5.88)	5.28 (4.96–5.72)
A1C (%)	5.76 (5.46–6.22)	5.75 (5.48–6.14)	5.73 (5.46–6.10)	5.78 (5.47–6.11)	5.68 (5.39–6.02)
Insulin (pmol/l)§	87.0 (61.2–123.0)	87.0 (60.0–120.6)	82.2 (61.2–113.4)	80.7 (59.4–106.8)	73.8 (55.2–99.3)
HOMA-IR	1.67 (1.18–2.30)	1.66 (1.17–2.26)	1.59 (1.18–2.12)	1.52 (1.14–2.02)	1.40 (1.07–1.88)
Triglycerides (mmol/l)	1.15 (0.83–1.82)	1.17 (0.82–1.80)	1.15 (0.80–1.71)	1.06 (0.73–1.61)	0.92 (0.65–1.38)
HDL cholesterol (mmol/l)	1.25 ± 0.34	1.27 ± 0.33	1.27 ± 0.33	1.27 ± 0.33	1.32 ± 0.33
CRP (mg/l)	0.79 (0.36–1.92)	0.74 (0.35–1.68)	0.67 (0.33–1.49)	0.60 (0.32–1.37)	0.59 (0.29–1.25)
IL-6 (pg/ml)¶	1.13 (0.73–1.86)	1.07 (0.73–1.65)	1.00 (0.62–1.49)	1.06 (0.69–1.57)	0.94 (0.61–1.49)

Data are means ± SD, *n* (%), or median (interquartile range). *Blood pressure ≥140/90 mmHg, or current use of anti-hypertensive medications. †77 participants were excluded with a missing value for self-reported CHD. ‡10 participants were excluded with a missing value for self-reported stroke. §4 participants were excluded with a missing value of fasting insulin. ||23 participants were excluded with a missing value for HOMA-IR. ¶89 participants were excluded with a missing value for IL-6.

cant ($P_{\text{trend}} = 0.0002$). The association appeared to be stronger in men than in women (data not shown), although the interaction for sex was not statistically significant ($P = 0.1745$). When study participants were dichotomized with the cutoff point of 50 nmol/l, the risk of having metabolic syndrome was 27% higher in those with an 25(OH)D concentration <50 nmol/l (OR 1.27; 95% CI 1.06–1.53) than in those with a concentration ≥50 nmol/l after controlling for all the covariates in model 3.

In both simple and multiple adjusted linear regression analyses, 25(OH)D was inversely associated with fasting glucose,

A1C, triglycerides, waist circumference, and diastolic blood pressure and positively associated with HDL cholesterol (Table 3). Further adjustment for BMI (not for waist circumference) did not substantially alter the associations. Sex-specific interactions were observed in the associations of 25(OH)D with triglycerides and HDL cholesterol (test for interaction: $P = 0.0034$ and 0.0027 for triglycerides and HDL cholesterol, respectively), and both associations were only significant in men but not in women.

There were negative associations of 25(OH)D with fasting insulin and HOMA-IR in the total population ($P =$

0.0114 and 0.0044 for insulin and HOMA-IR, respectively), after controlling for potential confounding factors (Table 3). We further observed a modifying effect of obesity status on these associations (test for interaction: $P = 0.0363$ and 0.0187 for insulin and HOMA-IR, respectively). In stratified analyses, 25(OH)D was negatively associated with fasting insulin and HOMA-IR only in participants with BMI ≥24 kg/m² ($P = 0.0001$ and $P < 0.0001$ for insulin and HOMA-IR, respectively) but not in those with BMI <24 kg/m² ($P = 0.5948$ and 0.5874 for insulin and HOMA-IR, respectively).

Table 2—Adjusted ORs (95% CI) of having metabolic syndrome according to quintiles of plasma 25(OH)D concentrations

	Quintiles of 25(OH)D					<i>P</i> _{trend}
	Q5: ≥57.7 nmol/l	Q4: 45.6–57.6 nmol/l	Q3: 36.9–45.5 nmol/l	Q2: 28.8–36.8 nmol/l	Q1: ≤28.7 nmol/l	
No. cases/participants	193/652	248/653	279/652	326/653	335/652	
Model 1	1.00 (reference)	1.31 (1.03–1.66)	1.36 (1.07–1.73)	1.67 (1.31–2.13)	1.62 (1.26–2.08)	<0.0001
Model 2	1.00 (reference)	1.31 (1.02–1.66)	1.38 (1.08–1.77)	1.76 (1.37–2.26)	1.64 (1.27–2.13)	<0.0001
Model 3	1.00 (reference)	1.28 (1.00–1.64)	1.33 (1.03–1.71)	1.71 (1.32–2.21)	1.52 (1.17–1.98)	0.0002

Model 1: adjusted for age, sex, geographic location (Beijing/Shanghai), residential region (urban/rural), and visit date (April or May/June). Model 2: further adjusted for education (≤6, 7–9, or ≥10 years in school), physical activity (low, moderate, or high), smoking (current, former, or never), alcohol drinking (yes/no), family history of CVD (yes/no) and diabetes (yes/no), and self-reported CHD (yes/no) and stroke (yes/no). Model 3: further adjusted for inflammatory factors (CRP and IL-6).

When overweight and obese participants were divided into 25(OH)D quintiles, the multivariate-adjusted geometric mean of fasting insulin was 13.86 pmol/l lower (*P* = 0.0003) and HOMA-IR was 0.26 lower (*P* = 0.0002) for the participants in the highest quintile compared with those

in the lowest. Excluding those with diagnosed diabetes and CVD yielded similar results patterns (data not shown).

CONCLUSIONS— We observed that ~94% of our study participants had 25(OH)D deficiency or insufficiency. A

lower 25(OH)D concentration was associated with increased risk of having metabolic syndrome and its individual components. In addition, poor 25(OH)D status was also significantly associated with increased insulin resistance, especially among those who were overweight or obese.

Table 3—Adjusted regression coefficients of 25(OH)D with components of metabolic syndrome, A1C, fasting insulin, and HOMA-IR

	Model 1		Model 2		Model 3		Model 4	
	β (SEM)	<i>P</i>	β (SEM)	<i>P</i>	β (SEM)	<i>P</i>	β (SEM)	<i>P</i>
All (<i>N</i> = 3,262)								
Fasting glucose (mmol/l)	−0.17 (0.08)	0.0277	−0.17 (0.08)	0.0337	−0.18 (0.08)	0.0262	−0.17 (0.08)	0.0281
A1C (%)	−0.21 (0.05)	<0.0001	−0.20 (0.05)	<0.0001	−0.21 (0.05)	<0.0001	−0.21 (0.05)	<0.0001
Fasting insulin (pmol/l)*	−0.08 (0.02)	0.0004	−0.07 (0.02)	0.0036	−0.06 (0.02)	0.0088	−0.06 (0.02)	0.0114
HOMA-IR†	−0.09 (0.02)	0.0002	−0.07 (0.02)	0.0014	−0.07 (0.02)	0.0035	−0.06 (0.02)	0.0044
Waist circumference (cm)	−1.61 (0.46)	0.0005	−1.64 (0.47)	0.0005	−1.46 (0.48)	0.0023	—	—
Triglycerides (mmol/l)‡	−0.13 (0.03)	<0.0001	−0.12 (0.03)	<0.0001	−0.11 (0.03)	<0.0001	−0.10 (0.03)	<0.0001
HDL cholesterol (mmol/l)§	0.07 (0.01)	<0.0001	0.07 (0.02)	<0.0001	0.06 (0.02)	<0.0001	0.06 (0.01)	<0.0001
Diastolic blood pressure (mmHg)	−1.29 (0.48)	0.0072	−1.32 (0.49)	0.0074	−1.25 (0.50)	0.0122	−1.15 (0.48)	0.0169
Systolic blood pressure (mmHg)	−1.37 (0.95)	0.1486	−1.52 (0.98)	0.1185	−1.28 (0.99)	0.1964	−1.08 (0.95)	0.2577
Sex								
Men (<i>n</i> = 1,443)								
Triglycerides (mmol/l)	−0.19 (0.04)	<0.0001	−0.19 (0.04)	<0.0001	−0.19 (0.04)	<0.0001	−0.19 (0.04)	<0.0001
HDL cholesterol (mmol/l)	0.10 (0.02)	<0.0001	0.10 (0.02)	<0.0001	0.10 (0.02)	<0.0001	0.10 (0.02)	<0.0001
Women (<i>n</i> = 1,819)								
Triglycerides (mmol/l)	−0.07 (0.04)	0.0574	−0.04 (0.04)	0.2554	−0.03 (0.04)	0.4546	−0.02 (0.03)	0.5907
HDL cholesterol (mmol/l)	0.03 (0.02)	0.0841	0.03 (0.02)	0.1182	0.02 (0.02)	0.3121	0.02 (0.02)	0.4327
Obesity status								
BMI <24 kg/m ² (<i>n</i> = 1,526)								
Fasting insulin (pmol/l)	−0.04 (0.03)	0.2653	−0.03 (0.03)	0.4596	−0.02 (0.04)	0.5948	—	—
HOMA-IR	−0.03 (0.03)	0.2977	−0.02 (0.03)	0.4530	−0.02 (0.03)	0.5874	—	—
BMI ≥24 kg/m ² (<i>n</i> = 1,736)								
Fasting insulin (pmol/l)	−0.14 (0.03)	<0.0001	−0.13 (0.03)	<0.0001	−0.12 (0.03)	0.0001	—	—
HOMA-IR	−0.15 (0.03)	<0.0001	−0.13 (0.03)	<0.0001	−0.13 (0.03)	<0.0001	—	—

25(OH)D, fasting insulin, HOMA-IR, and triglycerides were log-transformed before analysis. Model 1: adjusted for age, sex (not for sex-stratified analysis), geographic location (Beijing/Shanghai), residential region (urban/rural), and visit date (April or May/June). Model 2: further adjusted for education (≤6, 7–9, or ≥10 years in school), physical activity (low, moderate, or high), smoking (current, former, or never), alcohol drinking (yes/no), family history of CVD (yes/no) and diabetes (yes/no), and self-reported CHD (yes/no) and stroke (yes/no). For fasting glucose, A1C, fasting insulin, and HOMA-IR, hypertension (yes/no) was included in the model. Model 3: further adjusted for inflammatory factors (CRP and IL-6, continuous variables). Model 4: further adjusted for BMI (not for waist circumference and obesity status-stratified analysis). *4 participants were excluded with a missing value of fasting insulin; *P* for interaction between 25(OH)D and obesity status = 0.0363. †23 participants were excluded with a missing value of HOMA-IR; *P* for interaction between 25(OH)D and obesity status = 0.0187. ‡*P* for interaction between 25(OH)D and sex = 0.0034. §*P* for interaction between 25(OH)D and sex = 0.0027.

Although two studies reported that the VDR gene polymorphism was not associated with metabolic syndrome (18,19), the inverse association between 25(OH)D and metabolic syndrome found in our study is consistent with that from three previous cross-sectional reports in American (9,10) and British adults (6). These three studies (6,9,10) were also performed in the general population with large sample sizes; two American studies (9,10) included subjects aged >20 years and the other one (6) comprised participants aged 45 years. In these studies (6,9,10), 25(OH)D concentrations were relatively higher than in our study. A cohort study by Forouhi et al. (5) showed that the baseline concentration of 25(OH)D was inversely associated with an increased metabolic syndrome risk α score. However, a study in southern California residents (17) did not show a significant association between 25(OH)D and metabolic syndrome, which may be attributed to extraordinary high levels of 25(OH)D (mean concentration was 108.9 nmol/l in men and 101.6 nmol/l in women). Whether there is a threshold or a specific range for the association between 25(OH)D and metabolic abnormalities merits further investigation.

Elevated CRP and IL-6 levels have been found to be positively associated with metabolic syndrome (8) and inversely correlated with 25(OH)D in our study and other studies (20). Thus, we further explored whether the observed inverse association between 25(OH)D and metabolic syndrome was mediated by CRP and/or IL-6. In our study, the association was slightly attenuated but remained significant after we control for CRP and IL-6. Our result was in accordance with the data from the Third National Health and Nutrition Examination Survey (NHANES III) (9) in which 25(OH)D was inversely associated with metabolic syndrome independent of CRP levels.

25(OH)D status in our study was shown to be inversely associated with A1C and individual metabolic syndrome features such as fasting glucose, waist circumference, diastolic blood pressure, and triglycerides and positively associated with HDL cholesterol. However, unlike some of the previous studies (6,9,10), we have noticed that plasma 25(OH)D was associated with triglycerides and HDL cholesterol only in men but not in women. These sex-specific associations may partly explain the stronger

25(OH)D–metabolic syndrome association in men than in women in our study. Although the mechanistic role of vitamin D deficiency in the pathogenesis of dyslipidemia is not well understood, vitamin D supplementation was reported to attenuate the beneficial effect of hormone replacement therapy on serum lipid levels (21). Thus, the potential sex-specific effects of vitamin D on metabolic disorders need to be elucidated further.

Insulin resistance is a major underlying mechanism for the metabolic syndrome. Experimental studies have suggested that vitamin D may exert its beneficial effects by stimulating the expression of insulin receptor to improve insulin responsiveness for glucose transport or by controlling calcium influx, which is essential for the insulin-mediated intracellular process in insulin-responsive tissues (7). Most epidemiologic studies (3,5,19), but not all (18), suggested that serum 25(OH)D or the VDR polymorphism was associated with insulin sensitivity. Moreover, data from NHANES III documented an ethnic difference in the association between 25(OH)D and insulin resistance (4), with a stronger association in Caucasians than in African Americans, who have much lower levels of 25(OH)D concentrations. Interestingly, in our population, with 25(OH)D levels similar to those of African Americans, 25(OH)D was also negatively associated with fasting insulin and HOMA-IR, and the associations were stronger among overweight and obese subjects.

Obesity is known to be associated with decreased bioavailability of vitamin D, which is sequestered in body fat (2). In fact, Forouhi et al. (5) reported a significant interaction between 25(OH)D and BMI on the risk for a 10-year increase in HOMA-IR. In addition, the release of free fatty acids from adipose tissue can induce insulin resistance, whereas 1,25-dihydroxyvitamin D has been shown to counteract the free fatty acid–induced insulin resistance (22). The stronger association of vitamin D with insulin resistance among the overweight and obese participants suggests that adequate vitamin D status is more important for the prevention of insulin resistance and metabolic syndrome in these individuals.

Our data suggested that vitamin D deficiency was common in middle-aged and elderly Chinese individuals [i.e., 69.2% participants with 25(OH)D \leq 50 nmol/l]. Indeed, poor vitamin D status in middle

and older Chinese individuals was also reported previously in two small bone-related studies conducted in Beijing (23) and Shenyang (24), respectively. Although little is known regarding to what extent the high prevalence of vitamin D deficiency in our population could be explained by certain environmental and/or genetic factors, plausible explanations include 1) the skin of elderly individuals is less efficient for forming vitamin D₃, the major precursor of 25(OH)D (25), 2) use of vitamin D supplements is rare among older Chinese individuals (24), 3) unlike in the United States and other western countries, few vitamin D–fortified foods are available in China, and 4) our study participants were from a relatively high latitude, especially those in Beijing (latitude 40° north), who had lower 25(OH)D levels than their Shanghai counterparts (median 35.6 nmol/l in Beijing vs. 47.6 nmol/l in Shanghai). Hence, attention should be paid to improving vitamin D nutritional status in the Chinese population, particularly for those who are older, female, obese, less physically active, and living in high latitudes and urban areas.

To our knowledge, this is the first study to investigate the distribution and association of 25(OH)D levels with metabolic syndrome in Chinese individuals. The major strength of this study is that we used the data from a large representative sample of middle-aged and older men and women living in the two largest municipalities in China. Our analyses took into account many potential covariates that might confound the observed associations. Furthermore, we completed the field study within a relatively short period, which minimized the seasonal variation in the biomarkers. Nevertheless, several limitations should be acknowledged. A cause-effect relationship between 25(OH)D and metabolic syndrome cannot be inferred because of the cross-sectional nature of the study design. Although data on sun exposure and vitamin D supplementation were not available, we used a direct measure of vitamin D status, which reflects cumulative sun exposure and dietary vitamin D intake. In addition, because we did not measure serum calcium and parathyroid hormone, we could not determine whether the association of 25(OH)D with metabolic syndrome was partly mediated by calcium or secondary hyperparathyroidism. However, data from NHANES 2003–2004 (10) and the Medical Research Council Ely Prospective Study (5) suggested that the associations

between 25(OH)D and insulin resistance and metabolic syndrome were independent of calcium and parathyroid hormone.

In summary, our findings suggested that reduced 25(OH)D is associated with an increased risk of having metabolic syndrome and adverse values for metabolic syndrome components. The association of 25(OH)D with insulin resistance is stronger in overweight and obese individuals than in normal-weight individuals. Because there is evidence of ethnic variations in the 25(OH)D effect and limited data on the association of 25(OH)D with metabolic syndrome in Asians, our data provide novel insights into the nature of this association among Asians. Moreover, because of the high prevalence of metabolic syndrome and vitamin D deficiency in the middle-aged and elderly Chinese population, our results may have important public health implications. Increased sun exposure and use of vitamin D supplements or fortified foods are simple and inexpensive means to prevent vitamin D deficiency and related health problems. Nevertheless, the benefits of vitamin D on metabolic syndrome and related diseases such as type 2 diabetes need to be confirmed in future prospective studies and clinical trials.

Acknowledgments— This study was funded by the Chief Scientist Program of Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences (SIBS2008006), the Knowledge Innovation Program Project of the Chinese Academy of Sciences (KSCX1-YW-02, KSCX2-YW-R-73, and KSCX2-YW-R-116), the Ministry of Science and Technology of China (973 Program, Grant 2006CB503902), and the Shanghai-Unilever Research Development Fund (CH-2006-0941).

No potential conflicts of interest relevant to this article were reported.

References

- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266–281
- Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* 2000;72:690–693
- Chiu KC, Chu A, Go VL, Saad MF. Hypovitaminosis D is associated with insulin resistance and β cell dysfunction. *Am J Clin Nutr* 2004;79:820–825
- Scragg R, Sowers M, Bell C. Serum 25-hydroxyvitamin D, diabetes, and ethnicity in the Third National Health and Nutrition Examination Survey. *Diabetes Care* 2004;27:2813–2818
- Forouhi NG, Luan J, Cooper A, Boucher BJ, Wareham NJ. Baseline serum 25-hydroxy vitamin D is predictive of future glycemic status and insulin resistance: the Medical Research Council Ely Prospective Study 1990–2000. *Diabetes* 2008;57:2619–2625
- Hypponen E, Boucher BJ, Berry DJ, Power C. 25-hydroxyvitamin D, IGF-1, and metabolic syndrome at 45 years of age: a cross-sectional study in the 1958 British Birth Cohort. *Diabetes* 2008;57:298–305
- Pittas AG, Lau J, Hu FB, Dawson-Hughes B. The role of vitamin D and calcium in type 2 diabetes: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2007;92:2017–2029
- Cornier MA, Dabelea D, Hernandez TL, Lindstrom RC, Steig AJ, Stob NR, Van Pelt RE, Wang H, Eckel RH. The metabolic syndrome. *Endocr Rev* 2008;29:777–822
- Ford ES, Ajani UA, McGuire LC, Liu S. Concentrations of serum vitamin D and the metabolic syndrome among U.S. adults. *Diabetes Care* 2005;28:1228–1230
- Reis JP, von Muhlen D, Miller ER 3rd: Relation of 25-hydroxyvitamin D and parathyroid hormone levels with metabolic syndrome among US adults. *Eur J Endocrinol* 2008;159:41–48
- Botella-Carretero JI, Alvarez-Blasco F, Villafuella JJ, Balsa JA, Vazquez C, Escobar-Morreale HF. Vitamin D deficiency is associated with the metabolic syndrome in morbid obesity. *Clin Nutr* 2007;26:573–580
- Hjelmsaeth J, Hofso D, Aasheim ET, Jenssen T, Moan J, Hager H, Roislien J, Bollerslev J. Parathyroid hormone, but not vitamin D, is associated with the metabolic syndrome in morbidly obese women and men: a cross-sectional study. *Cardiovasc Diabetol* 2009;8:7
- Awumey EM, Mitra DA, Hollis BW, Kumar R, Bell NH. Vitamin D metabolism is altered in Asian Indians in the southern United States: a clinical research center study. *J Clin Endocrinol Metab* 1998;83:169–173
- Ye X, Yu Z, Li H, Franco OH, Liu Y, Lin X. Distributions of C-reactive protein and its association with metabolic syndrome in middle-aged and older Chinese people. *J Am Coll Cardiol* 2007;49:1798–1805
- Zhou BF. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults—study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci* 2002;15:83–96
- Grundey SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith SC Jr, Spertus JA, Costa F. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005;112:2735–2752
- Reis JP, von Muhlen D, Kritiz-Silverstein D, Wingard DL, Barrett-Connor E. Vitamin D, parathyroid hormone levels, and the prevalence of metabolic syndrome in community-dwelling older adults. *Diabetes Care* 2007;30:1549–1555
- Tworowska-Bardzinska U, Lwow F, Kubicka E, Laczmanski L, Jedrzejuk D, Dunajska K, Milewicz A. The vitamin D receptor gene *Bsm1* polymorphism is not associated with anthropometric and biochemical parameters describing metabolic syndrome in postmenopausal women. *Gynecol Endocrinol* 2008;24:514–518
- Oh JY, Barrett-Connor E. Association between vitamin D receptor polymorphism and type 2 diabetes or metabolic syndrome in community-dwelling older adults: the Rancho Bernardo Study. *Metabolism* 2002;51:356–359
- Dobnig H, Pilz S, Scharnagl H, Renner W, Seelhorst U, Wellnitz B, Kinkeldei J, Boehm BO, Weihrauch G, Maerz W. Independent association of low serum 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D levels with all-cause and cardiovascular mortality. *Arch Intern Med* 2008;168:1340–1349
- Heikkinen AM, Tuppurainen MT, Niskanen L, Komulainen M, Penttila I, Saarikoski S. Long-term vitamin D3 supplementation may have adverse effects on serum lipids during postmenopausal hormone replacement therapy. *Eur J Endocrinol* 1997;137:495–502
- Zhou QG, Hou FF, Guo ZJ, Liang M, Wang GB, Zhang X. 1,25-Dihydroxyvitamin D improved the free fatty-acid-induced insulin resistance in cultured C2C12 cells. *Diabetes Metab Res Rev* 2008;24:459–464
- Xue Y. Serum levels of 25-hydroxyvitamin D in normal Beijing subjects. *China Prev Med* 1991;25:177–179
- Yan L, Zhou B, Wang X, D'Ath S, Laidlaw A, Laskey MA, Prentice A. Older people in China and the United Kingdom differ in the relationships among parathyroid hormone, vitamin D, and bone mineral status. *Bone* 2003;33:620–627
- Holick MF, Matsuoka LY, Wortsman J. Age, vitamin D, and solar ultraviolet. *Lancet* 1989;2:1104–1105