

## **Plasticity of GABAergic synapses in the neonatal rat hippocampus**

**J.-L. Gaiarsa \***

*Institut de Neurobiologie de la Méditerranée (INMED), Institut National de la Santé et de la Recherche  
Médicale (INSERM) Unité 29,  
Marseille, France*

*Received: February 13, 2004; Accepted: March 10, 2004*

- **Introduction**
- **Developing inhibitory synapses undergoes long-term changes in synaptic strength**
- **Long-term plasticity is generated by physiologically relevant stimuli**
- **Long-term plasticity and the establishment of appropriate inhibitory synaptic connections**
- **Conclusion and perspectives**

### **Abstract**

While the development and plasticity of excitatory synaptic connections have been studied into detail, little is known about the development of inhibitory synapses. As proposed for excitatory synapses, recent studies have indicated that activity-dependent forms of synaptic plasticity, such as long-term potentiation and long-term depression, may play a role in the establishment of functional inhibitory synaptic connections. Here, I review these different forms of plasticity and focus on their possible role in the developing neuronal network.

**Keywords:** synaptic plasticity • GABA • glycine • calcium

---

\* Correspondence to: Prof. Jean-Luc GAIARSA  
Institut de Neurobiologie de la Méditerranée (INMED),  
Institut National de la Santé et de la Recherche Médicale (INSERM)

Unité 29, route de Luminy, B.P. 13, Marseille, France.  
Tel.: +33 (4)-91 82 81 29  
E-mail: gaiarsa@inmed.univ-mrs.fr

## Introduction

The development of highly organized structures in the CNS is a complex process that will determine the functional integrity at adult stage. Neuronal circuits are constructed over an extended period, both before and after birth, during which both activity-independent and activity-dependent mechanisms interact. During the last two decades, investigations have revealed that synaptic activity plays a role in shaping several aspects of circuits development including cell migration and differentiation, axonal path finding and the establishment of functional synaptic connections. The mechanisms by which synaptic activity participates to the refinement of initially coarse patterns of synaptic connections is presently not fully understood, but several studies have proposed that long-term changes in synaptic efficacy, such as long-term depression or potentiation (LTD and LTP) might participate in this process [1, 2].

Most of our understandings on activity-dependent maturation of synaptic connections and the possible contribution of long-term changes in synaptic efficacy in this process largely derived from studies on excitatory synaptic transmission [3-5]. However, several lines of evidence support the idea that spontaneous neuronal activity is also necessary for the full development of the inhibitory circuitry. In the auditory system, the topographic organization of glycinergic projections is achieved through synapses elimination [6], a process involving activity-dependent mechanisms [7]. In organotypic slice cultures of postnatal hippocampus [8] and cerebellum [9], synaptic activity regulates the density of inhibitory synapses. Moreover, sensory deprivations from birth lead to profound alterations in the number of GABA-immunopositive synaptic contacts or neurons in rat somatosensory [10] and visual [11-13] cortex. These changes provide the morphological substrate for the alterations in the spontaneous GABAergic synaptic activity observed both *in vitro* [14, 15] and *in vivo* [16] following manipulations of neuronal activity. The mechanisms underlying activity-dependent maturation of inhibitory synapses are unknown. However, it is tempting to speculate that they are similar to those required for long-term plasticity, as proposed for excitatory synapses.


The validity of this hypothesis however depends on several assumptions that are discussed in the present review.

## Developing inhibitory synapses undergoes long-term changes in synaptic strength

The major aim of this review is to discuss the possible role of long-term synaptic plasticity in the establishment of functional inhibitory synapses in the developing brain. However, the characterization of their induction and expression processes could help to understand the relevance of any form of synaptic plasticity. In this chapter, I will therefore briefly summarize the minimal requirements necessary for the induction and expression of long-term plasticity at developing GABAergic and glycinergic synapses.

Both LTP and LTD of inhibitory synapses have been described in different developing brain regions including the lateral superior olive [17], the cortex [18] and the hippocampus [19]. In all these structures, attention had been paid to demonstrate that GABAergic and glycinergic synapses themselves undergo long-term changes in synaptic efficacy. Not surprisingly, several mechanisms have been reported to contribute to the induction and maintenance of long-term plasticity at developing inhibitory synapses. A rise in intracellular  $Ca^{2+}$  concentration appears to be the common trigger for inducing the synaptic plasticity, although the source of  $Ca^{2+}$  and the underlying consequences on inhibitory synaptic efficacy may differ from one structure to another (reviewed in [20]). The rise in intracellular  $Ca^{2+}$  concentration can be produced by the opening of voltage-gated calcium channels (VGCCs) [21] or NMDA-gated channels [22, 23], or by the release of  $Ca^{2+}$  from  $InsP_3$ -sensitive stores [24] or  $Ca^{2+}$ -induced  $Ca^{2+}$  release stores [25]. In the neonatal rat hippocampus, the same conditioning protocol, i.e. high frequency stimulation, can lead to either LTP or LTD of GABAergic synapses (Fig. 1). The polarity of the plasticity induced depends on whether or not NMDA receptors have been activated during the conditioning protocol [19]. Given that both forms of plasticity require a postsynaptic rise in calcium [19], these observations suggest that the direction of synaptic changes is determined by the source of the calcium influx (*i.e.* VGCCs for LTP [21] versus NMDA for LTD [23]) (Fig. 1).

The mechanisms underlying the expression of long-term plasticity at inhibitory synapses have been



**Fig. 1 Long-term changes in GABAergic synaptic efficacy in the developing rat hippocampus.** During the first postnatal week of life, direct stimulation of the GABAergic interneurons, in the presence of ionotropic glutamatergic receptor antagonist, induces a depolarizing GABAergic potential. High frequency stimulation induces either a long-term potentiation (LTP<sub>GABA-A</sub>) or depression (LTD<sub>GABA-A</sub>) of GABAergic synaptic potentials. Both LTP and LTD required a membrane depolarization, mediated by the activation of GABA<sub>A</sub> receptors. This depolarization induces the opening of voltage gated Ca<sup>2+</sup> channels (VGCCs) and allows the removal of Mg<sup>2+</sup>-block from NMDA channels. The calcium influx through VGCCs leads to an LTP of GABAergic synapses, while the calcium influx through NMDA channels lead to an LTD.

investigated. Long-term changes in the strength of synaptic efficacy can be accounted for by at least three non exclusive mechanisms: (i) modifications in the number or properties of receptors at functional synapses, (ii) modifications in the probability of transmitter release, and (iii) modifications in the number of functional synapses though either pre- or post-synaptic mechanisms. In the neonatal rat hippocampus, long-term plasticity of GABAergic synapses is expressed as modifications in the number of functional synapses or changes in the probability of transmitter release. Thus GABAergic synaptic plasticity is associated with a modification in the frequency, but not amplitude, of quantal inhibitory postsynaptic currents (IPSCs) [21, 23] and in the coefficient of variation of evoked IPSCs [23]. If the induction has a postsynaptic origin while the locus of expression is presynaptic, then a retrograde messenger should travel backward across the synapses to modulate the probability of

transmitter release. Retrograde messengers have been reported to participate in long-lasting plasticity of GABAergic synapses in the adult hippocampus [26, 27]. Whether similar synaptic feedback also contributes to the induction of long-term plasticity at developing inhibitory synapses remained to be determined.

### Long-term plasticity is generated by physiologically relevant stimuli

In most studies tetanic stimulation has been used to induce long-term changes in the efficacy of inhibitory synaptic transmission. Although such conditioning protocol provides a good tool to study the mechanisms by which plasticity is triggered and expressed, it is less obvious that inhibitory synapses are stimulated at such rates with the exception of

ripples and sharp waves [28]. Therefore, one always has to take into account the relevance of the conditioning protocol to physiological conditions to gain insight into the role that this plasticity may serve.

There are now compelling evidences suggesting that conditioning protocols relevant to physiological or pathological conditions can induce plasticity at GABAergic and glycinergic synapses. The first demonstration was provided by Korn and collaborators, in the goldfish [29]. They showed that auditory stimulation can trigger an LTP of inhibitory synapses *in vivo* that shares similar properties with the LTP induced by tetanic stimulation [30-32]. Although less direct, other evidences have been provided in the neonatal rat hippocampus. Thus, a  $Ca^{2+}$ -dependent LTP at GABAergic synapses can be induced by repeated bursts of action potential applied at a low frequency to CA3 pyramidal cells [21, 33]. This conditioning protocol is clearly related to the sort of activity that neonatal CA3 pyramidal cells experience *in vivo* [34]. In neonates, burst of action potentials can result from endogenous bursting properties of CA3 pyramidal cells [35] or from the network-driven synaptic activity present at early stages of development [36]. To further address the relevance of the conditioning protocol, it will be of interest in future studies to determine whether action potentials generated by the ongoing spontaneous activity are also able to trigger long-term plasticity at GABAergic and glycinergic synapses.

### **Long-term plasticity and the establishment of appropriate inhibitory synaptic connections**

Activity-dependent long-term changes in inhibitory synaptic efficacy are believed to be an important phenomenon in learning and memory formation, and pathological states of neuronal excitability in the adult central nervous system (reviewed in [37]). A number of studies now suggest that long-term changes in synaptic efficacy, such as long-term depression or potentiation, might also play a crucial role in the establishment of functional inhibitory synapses in the developing brain.

First, in all developing structures where inhibitory synaptic plasticity has been described,

the induction is restricted to a limited period of development. Moreover, although the period during which the plasticity can be induced differ in these structures, this period always closely matches the period of functional maturation of neuronal network as shown in the visual cortex [18], the hippocampus [19, 33] and the auditory system [17]. In the neonatal rat hippocampus, for instance, the GABAergic synaptic plasticity induced by postsynaptic firing of pyramidal cells is restricted to the first postnatal week of life [33]. This developmental window coincides with the period of functional maturation of GABAergic synapses [38].

Second, the conditioning protocol leading to long-term changes in synaptic plasticity also leads to functional maturation of inhibitory synapses. The developing rat hippocampus is an interesting model to test this hypothesis. Thus, at birth the hippocampal pyramidal cells are synaptically “silent”: that is, they show neither spontaneous, nor evoked synaptic responses. The synaptic maturation follows a sequence with the GABAergic synapses being functional before the glutamatergic one [39, 40]. In our study, we have recorded such “silent” neurons, and asked whether a relevant conditioning protocol can trigger the functional maturation of GABAergic synapses. Indeed, repeated bursts of action potentials, leading to long-term plasticity of GABAergic synapses when applied to “active” cells”, also lead to the appearance of evoked and spontaneous GABAergic synaptic activity when applied to “silent” cells [33].

Finally, to further strengthen the link between activity-dependent maturation and long-term plasticity, it is necessary to show that the same cellular mechanisms are involved in both phenomena. In this context, neurotrophins and related Trk receptor-coupled protein tyrosine kinases (PTKs) have been implicated in synapse development and plasticity. In cerebellar and hippocampal cultures, activity deprivation decreases the number of GABAergic synapses, an effect reversed by the application of brain-derived neurotrophic factor (BDNF) or neurotrophin-4 [8, 41]. Moreover, overexpression of BDNF [42] or chronic treatment with different neurotrophins [43-45] accelerates the functional maturation of GABAergic synapses. Several studies have also shown that neurotrophins can modulate the efficacy of inhibitory synaptic transmission [46-48]. Interestingly, it was reported that NT-3 enhances



GABAergic synaptic transmission during the development period when GABA<sub>A</sub> receptors activation induced a depolarization and a postsynaptic Ca<sup>2+</sup> rise, but not later when GABA leads to a membrane hyperpolarisation [49]. Neurotrophins may therefore be the signal linking long-term plasticity and activity-dependent maturation of inhibitory synapses. In agreement with this hypothesis, a recent study suggests that Trk receptors participate in the induction of long-term depression of glycinergic synapses in the developing auditory brain stem [50]. Similarly, in the developing neonatal rat hippocampus, our data show that the long-term potentiation of GABAergic synapses induced by the postsynaptic firing of CA3 pyramidal neurons is prevented by the application of PTK inhibitors (Gaiarsa and Gubellini, unpublished results).

## Conclusion and perspectives

The data reviewed here indicate that inhibitory synapses undergo long-term change in synaptic efficacy in the developing brain, and that this form of plasticity might be instrumental in the establishment of appropriate synaptic connections. In future studies, a key issue to further link long-term plasticity and activity-dependent maturation of synaptic transmission will be to characterize the synaptic activity underlying functional maturation of inhibitory synapses, and to determine whether this activity is able to induce long-term plasticity at inhibitory synapses. A hallmark of developing network is the presence of spontaneous patterned synaptic activity (for review see [51]). In spite of differences in the mechanisms underlying their generation, these patterned activities share several interesting features with regard to activity-dependent maturation of the neuronal networks. They are present during a restricted period of development corresponding to the period of synaptic maturation. In the neonatal rat hippocampus, this activity disappeared at a time when long-term plasticity can no longer be induced by postsynaptic firing of CA3 pyramidal cells. Moreover, they are always associated with postsynaptic variation in intracellular Ca<sup>2+</sup> concentration, although the mechanisms leading to this Ca<sup>2+</sup> transient differ from one structure to another. These activities may therefore represent

the physiological pattern of activity leading to the functional maturation of inhibitory synapses through LTP/LTD like mechanisms. Further studies are needed to directly address this hypothesis.

## References

1. **Shatz C.J.**, Impulse activity and the patterning of connections during CNS development, *Neuron*, **5**: 745-756, 1990
2. **Constantine-Paton M., Cline H.T.**, LTP and activity-dependent synaptogenesis: the more alike they are, the more different they become, *Curr.Opin.Neurobiol.*, **8**: 139-148, 1998
3. **Lo Y.J., Poo M.M.**, Activity-dependent synaptic competition in vitro: heterosynaptic suppression of developing synapses. *Science*, **254**: 1019-1021, 1991
4. **Lo Y.J., Poo M.M.**, Heterosynaptic suppression of the developing neuromuscular synapses in culture. *J.Neurosci.*, **14**: 4684-4693, 1994
5. **Kirkwood A., Lee H.K., Bear M.F.**, Co-regulation of long-term potentiation and experience-dependent synaptic plasticity in visual cortex by age and experience, *Nature*, **375**: 328-331, 1995
6. **Sanes D.H., Siverls V.**, The development and specificity of inhibitory axonal arborizations in the lateral superior olive, *J.Neurobiol.*, **22**: 837-854, 1991
7. **Sanes D.H., Tackacs C.** Activity-dependent refinement of inhibitory connections. *Eur. J. Neurosci.*, **5**: 570-574, 1993
8. **Marty S., Wehrle R., Sotelo C.**, Neuronal activity and brain-derived neurotrophic factor regulate the density of inhibitory synapses in organotypic slice cultures of postnatal hippocampus. *J. Neurosci.*, **20**: 8087-8095, 2000
9. **Seil F.J., Drake-Baumann R.**, Reduced cortical inhibitory synaptogenesis in organotypic cerebellar cultures developing in the absence of neuronal activity., *J Comp.Neurol.*, **342**: 366-377, 1994
10. **Micheva K.D., Beaulieu C.**, An anatomical substrate for experience-dependent plasticity of the rat barrel field cortex, *Proc.Natl.Acad.Sci.U.S.A.*, **92**: 11834-11838, 1995
11. **Hendry S.H., Jones E.G.**, Activity-dependent regulation of GABA expression in the visual cortex of adult monkey, *Neuron*, **1**: 701-712, 1988
12. **Benevento L.A., Bakkum B.W., Cohen R.S.**, Gamma-aminobutyric acid and somatostatin immunoreactivity in the visual cortex of normal and dark-reared rats, *Brain Res.*, **689**: 172-182, 1995
13. **Morales B., Choi S.Y., Kirkwood A.**, Dark rearing alters the development of GABAergic transmission in visual cortex, *J.Neurosci.*, **22**: 8084-8090, 2002
14. **Rutherford L.C., DeWan A., Lauer H.M., Turrigiano G.G.**, Brain-derived neurotrophic factor mediates the activity-dependent regulation of inhibition in neocortical cultures., *J.Neurosci.*, **17**: 4527-4535, 1997
15. **Galante M., Nistri A., Ballerini L.**, Opposite changes in synaptic activity of organotypic rat spinal cord cultures

- after chronic block of AMPA/kainate or glycine and GABAA receptors, *J.Physiol.*, **523**: 639-651, 2000
16. **Aamodt M.A., Shi J., Colonnese M.T., Veras W., Constantine-Paton M.**, Chronic NMDA exposure accelerated development of GABAergic inhibition in the supri- or colliculus. *J.Neurophysiol.*, **83**: 1580-1591, 2000
  17. **Kotak V.C., Sanes D.H.**, Long-lasting inhibitory synaptic depression is age- and calcium-dependent., *J.Neurosci.*, **20**: 5820-5826, 2000
  18. **Komatsu Y.**, Age-dependent long-term potentiation of inhibitory synaptic transmission in rat visual cortex. *J.Neurosci.*, **14**: 6488-6499, 1994
  19. **McLean H.A., Caillard O., Ben-Ari Y., Gaïarsa J.-L.**, Bidirectional plasticity expressed by GABAergic synapses in the neonatal rat hippocampus, *J.Physiol.(Lond.)*, **496**: 471-477, 1996
  20. **Gaïarsa J.L., Caillard O., Ben Ari Y.**, Long-term plasticity at GABAergic and glycinergic synapses: mechanisms and functional significance, *Trends Neurosci.*, **25**: 564, 2002
  21. **Caillard O., Ben-Ari Y., Gaïarsa J.-L.**, Long-term potentiation of GABAergic synaptic transmission in neonatal rat hippocampus, *J.Physiol.(Lond.)*, **518**: 109-119, 1999
  22. **Komatsu Y., Iwakiri M.**, Long-term modification of inhibitory synaptic transmission in developing visual cortex, *Neuroreport*, **4**: 907-910, 1993
  23. **Caillard O., Ben-Ari Y., Gaïarsa J.-L.**, Mechanisms of induction and expression of long-term depression at GABAergic synapses in neonatal rat hippocampus, *J. Neurosci.*, **19**: 7568-7577, 1999
  24. **Komatsu Y.**, GABA<sub>B</sub> receptors, monoamine receptors, and postsynaptic inositol trisphosphate-induced Ca<sup>2+</sup> release are involved in the induction of long-term potentiation at visual cortical inhibitory synapses, *J. Neurosci.*, **16**: 6342-6352, 1996
  25. **Caillard O., Ben-Ari Y., Gaïarsa J.-L.**, Activation of presynaptic and postsynaptic ryanodine-sensitive calcium stores is required for the induction of long-term depression at GABAergic synapses in the neonatal rat hippocampus, *J. Neurosci.*, **20**: 1-5, 2000
  26. **Kang J., Jiang L., Goldman S.A., Nedergaard M., Kang J.** Astrocyte-mediated potentiation of inhibitory synaptic transmission. *Nature Neuroscience*, **1**: 683-692, 1998
  27. **Chevalyere V., Castillo P.E.**, Heterosynaptic LTD of hippocampal GABAergic synapses: a novel role of endocannabinoids in regulating excitability. *Neuron*, **38**: 461-472, 2003
  28. **Freund T., Buzsaki G.**, Interneurons of the hippocampus. *Hippocampus*, **6**: 345-470, 1996
  29. **Oda Y., Kawasaki H., Morita M., Korn H., Matsui H.**, Inhibitory long-term potentiation underlies auditory conditioning of goldfish escape behaviour. *Nature*, **394**: 182-185, 1998
  30. **Korn H., Oda Y., Faber D.S.**, Long-term potentiation of inhibitory circuits and synapses in the central nervous system. *Proc.Natl.Acad.Sci.USA*, **89**: 440-443, 1992
  31. **Oda Y., Charpier S., Murayama Y., Suma C., Korn H.**, Long-term potentiation of glycinergic inhibitory synaptic transmission. *J. Neurophysiol.*, **74**: 1056-1074, 1995.
  32. **Charpier S., Behrends J.C., Triller A., Faber D.S., Korn H.**, "Latent" inhibitory connections become functional during activity-dependent plasticity, *Proc. Natl. Acad. Sci.U.S.A.*, **92**: 117-120, 1995.
  33. **Gubellini P., Ben-Ari Y., Gaïarsa J.-L.**, Activity- and age-dependent GABAergic synaptic plasticity in the developing rat hippocampus, *Eur. J. Neurosci.*, **14**: 1937-1946, 2001.
  34. **Leinekugel X., Khazipov R., Cannon R., Hirase H., Ben-Ari Y., Buzsáki G.**, Correlated burst of activity in the neonatal rat hippocampus *in vivo*, *Science*, **296**: 2049-2052, 2002
  35. **Menendez de la Prida L., Sanchez-Andres J.V.**, Heterogenous populations of cells mediate spontaneous synchronous bursting in the developing hippocampus though a frequency-dependent mechanism, *Neuroscience*, **97**: 227-241, 2000.
  36. **Ben-Ari Y., Cherubini E., Corradetti R., Gaïarsa J.-L.**, Giant synaptic potentials in immature rat CA3 hippocampal neurones, *J.Physiol.(Lond.)*, **416**: 303-325, 1989
  37. **Gaïarsa J.L., Caillard O., Ben Ari Y.**, Long-term plasticity at GABAergic and glycinergic synapses: mechanisms and functional significance, *Trends Neurosci.*, **25**: 564, 2002
  38. **Ben-Ari Y., Khazipov R., Leinekugel X., Caillard O., Gaïarsa J.-L.**, GABA<sub>A</sub>, NMDA and AMPA receptors: a developmentally regulated 'ménage a trois', *Trends Neurosci.*, **20**: 523-529, 1997
  39. **Tyzio R., Represa A., Jorquera I., Ben-Ari Y., Gozlan H., Aniksztejn L.**, The establishment of GABAergic and glutamatergic synapses on CA1 pyramidal neurons is sequential and correlates with the development of the apical dendrite., *J. Neurosci.*, **19**: 10372-10382, 1999
  40. **Hennou S., Khalilov I., Diabira D., Ben Ari Y., Gozlan H.**, Early sequential formation of functional GABAA and glutamatergic synapses on CA1 interneurons of the rat foetal hippocampus., *Eur. J. Neurosci.*, **16**: 197-208, 2002
  41. **Seil F.J., Drake-Baumann R.**, TrkB receptor ligands promote activity-dependent inhibitory synaptogenesis. *J. Neurosci.*, **20**: 5367-5373, 2000.
  42. **Aguado F., Carmona M.A., Pozas E., Aguilo A., Martinez-Guijarro F.J., Alcantara S., Borrell V., Yuste R., Ibanez C.F., Soriano E.**, BDNF regulates sponaneous correlated activity at early developmental stages by increasing synaptogenesis and expression of K<sup>+</sup>/Cl<sup>-</sup> cotransport, *Development*, **130**: 1287-1280, 2003
  43. **Marty S., Berninger B., Carroll P., Thoenen H.**, GABAergic stimulation regulates the phenotype of hippocampal interneurons through the regulation of brain-derived neurotrophic factor, *Neuron*, **16**: 565-570, 1996
  44. **Vicario-Abejon C., Collin C., McKay R.D., Segal M.**, Neurotrophins induce formation of functional excitatory and inhibitory synapses between cultured hippocampal neurons. *J. Neurosci.*, **18**: 7256-7271, 1998
  45. **Schwyzler L., Mateos J.M., Abegg M., Rietschin L., Heeb L., Thompson S.M., Luthi A., Gahwiler B.H., McKinney R.A.**, Physiological and morphological plasticity induced by chronic treatment with NT-3 or NT-4/5 in hippocampal slice cultures, *Eur. J. Neurosci.*, **16**: 1939-1948, 2002

46. **Tanaka T., Saito H., Matsuki N.**, Inhibition of GABA<sub>A</sub> synaptic responses by brain-derived neurotrophic factor (BDNF) in rat hippocampus, *J. Neurosci.*, **17**: 2959-2966, 1997
47. **Li Y.X., Zhang Y., Lester H.A., Schuman E.M., Davidson N.** Enhancement of neurotransmitter release induced by brain-derived neurotrophic factor in cultured hippocampal neurons. *J. Neurosci.*, **18**: 10231-10240, 1998
48. **MacLean Bolton M., Pittman A.J., Lo D.C.**, Brain-derived neurotrophic factor differentially regulates excitatory and inhibitory synaptic transmission in hippocampal neurons, *J. Neurosci.*, **20**: 3221-3232, 2000
49. **Gao B.X., Van den Pol A.N.**, Neurotrophin-3 potentiates GABAergic synaptic transmission in cultured developing hypothalamic neurones of the rat, *J. Physiol. (Lond.)*, **518**: 81-95, 1999
50. **Kotak V.C., Dimattina C., Sanes D.H.**, GABA<sub>B</sub> and Trk receptor signaling mediates long-lasting inhibitory synaptic depression., *J. Neurophysiol.*, **86**: 536-540, 2001
51. **Ben-Ari Y.**, Developing networks play similar melody. *Trends Neurosci.*, **24**: 354-360, 2001