

PMRD: plant microRNA database

Zhenhai Zhang¹, Jingyin Yu¹, Daofeng Li², Zuyong Zhang¹, Fengxia Liu¹, Xin Zhou¹,
Tao Wang², Yi Ling¹ and Zhen Su^{1,*}

¹State Key Laboratory of Plant Physiology and Biochemistry and ²State Key Laboratory for Agricultural Biotechnology, College of Biological Sciences, China Agricultural University, Beijing 100193, China

Received July 4, 2009; Revised August 27, 2009; Accepted September 16, 2009

ABSTRACT

MicroRNAs (miRNA) are ~21 nucleotide-long non-coding small RNAs, which function as post-transcriptional regulators in eukaryotes. miRNAs play essential roles in regulating plant growth and development. In recent years, research into the mechanism and consequences of miRNA action has made great progress. With whole genome sequence available in such plants as *Arabidopsis thaliana*, *Oryza sativa*, *Populus trichocarpa*, *Glycine max*, etc., it is desirable to develop a plant miRNA database through the integration of large amounts of information about publicly deposited miRNA data. The plant miRNA database (PMRD) integrates available plant miRNA data deposited in public databases, gleaned from the recent literature, and data generated in-house. This database contains sequence information, secondary structure, target genes, expression profiles and a genome browser. In total, there are 8433 miRNAs collected from 121 plant species in PMRD, including model plants and major crops such as *Arabidopsis*, rice, wheat, soybean, maize, sorghum, barley, etc. For *Arabidopsis*, rice, poplar, soybean, cotton, medicago and maize, we included the possible target genes for each miRNA with a predicted interaction site in the database. Furthermore, we provided miRNA expression profiles in the PMRD, including our local rice oxidative stress related microarray data (LC Sciences miRPlants_10.1) and the recently published microarray data for poplar, *Arabidopsis*, tomato, maize and rice. The PMRD database was constructed by open source technology utilizing a user-friendly web interface, and multiple search tools. The PMRD is freely available at <http://bioinformatics.cau.edu.cn/PMRD>. We expect PMRD to be a useful tool for scientists in

the miRNA field in order to study the function of miRNAs and their target genes, especially in model plants and major crops.

INTRODUCTION

MicroRNAs (microRNA) are ~21 nucleotide-long endogenous non-coding small RNAs and function as post-transcriptional regulators in eukaryotes. The processing of miRNA is well studied and is comprised of a discrete series of steps (1). miRNA genes are transcribed and excised into miRNAs, then miRNA is recruited by RISC (RNA-induced silencing complex, including Argonaute and other proteins) which combines with mRNA to inhibit or degrade the target mRNA (1) and thus translation is interrupted.

MiRNAs play essential roles in regulating plant growth and development (2). Research into miRNA function on target genes has progressed at a very rapid rate in plants. As examples, Archak *et al.* (3) discovered that miRNAs may participate in diverse functions including transcription, catalysis, binding and transporter activity. Some miRNAs related to abiotic stress responses such as cold stress and nutrient deprivation were identified in *Arabidopsis thaliana* using transcriptome analysis (4,5); Lu *et al.* (6) analyzed miRNA regulatory roles in the response of *Populus trichocarpa* to the stressful environment incurred over their long-term growth; Morin *et al.* conducted comparative analyses on the conservation of miRNAs between *Pinus contorta* and *Oryza sativa* and discovered that important RNA silencing processes were highly developed in the earliest spermatophytes (7).

Recently, more and more miRNAs have been identified in plant genomes. Jones-Rhoades *et al.* (8) developed comparative genomic approaches to systematically identify both miRNAs and target genes, which enlarged the miRNAs family and the number of target genes in *A. thaliana*. Using high-throughput sequencing, Rajagopalan *et al.* (9) identified 38 new miRNAs in

*To whom correspondence should be addressed. Tel: +86 10 62731380; Fax: +86 10 62731214; Email: zhensu@cau.edu.cn

A. thaliana. Using a similar approach, a large number of miRNAs were recently identified in rice (10). Using support vector machine classification based on intra-genomic matching of potential miRNAs and their targets, Lindow *et al.* (11) predicted ~1200, ~2100 and ~2500 miRNA candidate genes in *A. thaliana*, *O. sativa* and *P. trichocarpa*, respectively. Using EST analysis, Zhang *et al.* (12,13) identified numerous new miRNAs in many species. Finally, Ramanjulu Sunkar and Guru Jagadeeswaran (14) identified miRNAs in a large number of plant species.

With the discovery of a large number of miRNAs and their target genes, some public resources have been constructed: miRBase (<http://microrna.sanger.ac.uk/>), provides a set of precursor and mature miRNAs discovered in many plants (15); ASRP (<http://asrp.cgrb.oregonstate.edu/>) lists miRNAs and their target genes in *Arabidopsis* (16); CSRDB (<http://sundarlab.ucdavis.edu/smrnas/>) is a collection of miRNAs identified in maize and rice (17); Rfam (<http://rfam.sanger.ac.uk/>) provides secondary structures of miRNA precursors in many plant species (18,19). However, there are limitations to the current databases in this field. For example, miRBase only provides miRNA sequence data and annotation in 25 plant species; Rfam only contains miRNA precursor sequences from less than 50 plant species; the ASRP integrates miRNAs, precursor, target genes and genome browser only for *Arabidopsis*. It is necessary to develop a plant miRNA database through integration of the large amount of information about plant miRNAs available to the public. For this purpose, we developed a plant microRNA database (PMRD) to integrate the available data pertinent to plant miRNAs available from public resources. Uniquely, we provided miRNA expression profiles in the PMRD, including our local generated rice oxidative stress microarray data and the recent published microarray data generated from poplar, *Arabidopsis*, tomato, maize and rice.

DATABASE CONTENT

miRNA data collection

The majority of our data was gleaned from miRBase, Rfam and literature published in recent years. The number of mature miRNAs that we immediately retrieved from the miRBase was 1931, 73 miRNAs came from Rfam, and other miRNAs were from the literature. As shown in Table 1, we integrated 8433 non-redundant miRNA sequences and their original resources. Annotation of miRNAs was mainly dependent on information present in miRBase, but in some particular instances, the annotation included information about the corresponding author who discovered the miRNA. Detailed information concerning the variety of miRNAs in PMRD is presented in Table 1.

Recently in PMRD, there are 1427, 2540 and 2780 mature miRNA entries for *Arabidopsis*, rice and poplar, respectively, while in miRBase the entries for these three species are 207, 415 and 237, respectively. The other miRNAs we curated from these three species mainly

come from Lindow *et al.* (11). There are 166 soybean entries in PMRD, of which 79 sequences came from miRBase (15,20–22), and the others were from Zhang *et al.* (13,23). For wheat we deposited 85 miRNAs in PMRD, of which 32 came from miRBase (15,24) and the others were from Yao *et al.* (24), Zhang *et al.* (13) and Wei *et al.* (25). As for maize, 207 entries were created in PMRD, including 98 from miRBase (13,15,20,26–28) and 109 from Zhang *et al.* (13,29). For cotton we deposited 53 mature sequences in PMRD, in which 13 items were from miRBase (15,30), and the others were from Qiu *et al.* (31) and Zhang *et al.* (32). In total, there were 8433 miRNAs taken from 121 plant species integrated into PMRD.

miRNA expression profiling

In our PMRD database, we had already added data concerning miRNA expression profiling for several public miRNA expression data sets including maize miRNA-seq results (33), poplar cold stress-responsive miRNAs (6) (sample data shown in Supplementary Figure S1), tomato miRNA expression in leaf tissue (34), *Arabidopsis* stress (cold, salt and osmotic stresses) regulated miRNAs (35), maize miRNAs responding to salt stress in roots (36), miRNA expression profiles generated from *Arabidopsis* grown at different temperatures (GEO: GSE11535), and rice miRNAs responding to drought stress (GEO: GSE5986), etc.

For example, in the data set for poplar cold stress, there were 75 probes for 168 miRNAs, and a total of 21 samples including seven time points collected in triplicate. As shown in Supplementary Figure S1, the ratio represented the fold-change of poplar ptc-miR474b expression between cold treatment and control. We highlighted time points demonstrating a significant change in miRNA expression using $P \leq 0.01$ as the cut-off.

In addition, we added our local rice oxidative stress related microarray data into PMRD. In order to investigate differences in expression levels of miRNAs in response to oxidative stress in rice, we conducted a series of experiments using the miRPlants_10.1 small RNA chip (LC Sciences, Houston, TX). One-week-old 9311 (rice Indica variety) seedling plants were incubated in a solution containing 10 μ M methyl viologen (MV) (paraquat, an herbicide that induces oxidative stresses in plants) using water as a mock-treatment control, for 24 h. A total of four samples were examined in these experiments: 9311-CK01 and 9311-CK02 were controls and 9311-MV01 and 9311-MV02 were two samples treated with MV. Quantile normalization (37) was used to analyze the miRNA expression data. Within the total of 653 probes related to 1773 miRNAs representing 14 species in the chip, 18 probes showed significant differential expression ($P \leq 0.05$, listed in Table 2) between MV treatment and mock, including 12 probes that were up-regulated and 6 probes that were down-regulated after MV treatment. Detailed relative expression levels of each probe are presented in the Expression page in the PMRD database. In this article, we present the example of osa-miR159a as Supplementary Figure S2.

Table 1. Species and sources for raw miRNA data

Species	Number of miRNAs			References
	Total	Experimental	Computational	
<i>Aegilops speltoides</i>	1		1	(12,13)
<i>Aegilops tauschii</i>	2		2	(19)
<i>Alliaria petiolata</i>	1		1	(19)
<i>Allium cepa</i>	7		7	(12,13)
<i>Annona cherimola</i>	1		1	(19)
<i>Antirrhinum majus</i>	1		1	(19)
<i>Arabidopsis arenosa</i>	1		1	(19)
<i>Arabidopsis cebennensis</i>	1		1	(19)
<i>Arabidopsis halleri</i>	1		1	(19)
<i>Arabidopsis thaliana</i>	1427	207	1220	(2,4,8,9,11–13,16,41–58)
<i>Arabis hirsuta</i>	1		1	(19)
<i>Arachis hypogaea</i>	1		1	(59)
<i>Australopyrum velutinum</i>	1		1	(19)
<i>Avena sativa</i>	1		1	(19)
<i>Barbarea vulgaris</i>	1		1	(19)
<i>Boechera stricta</i>	1		1	(14)
<i>Brachypodium distachyon</i>	12	12		(25)
<i>Brachypodium sylvaticum</i>	1		1	(19)
<i>Brassica napus</i>	45	45		(50,60,61)
<i>Brassica oleracea</i>	9	1	8	(14,50,62)
<i>Brassica rapa</i>	20	20		(14,50,62)
<i>Camelina microcarpa</i>	1		1	(19)
<i>Capsella rubella</i>	1		1	(19)
<i>Capsicum annuum</i>	3		3	(12,13)
<i>Carica papaya</i>	2	1	1	(14,63)
<i>Chlamydomonas reinhardtii</i>	84	84		(64,65)
<i>Citrus clementina</i>	1		1	(14)
<i>Citrus sinensis</i>	5		5	(12,13)
<i>Citrus × paradisi × Poncirus trifoliata</i>	4		4	(14)
<i>Conringia orientalis</i>	1		1	(19)
<i>Corylus avellana</i>	1		1	(19)
<i>Cycas rumphii</i>	1		1	(12,13)
<i>Descurainia sophia</i>	1		1	(19)
<i>Dictyostelium discoideum</i>	2	2		(66)
<i>Erysimum cheiri</i>	1		1	(19)
<i>Eschscholzia californica</i>	1		1	(12,13)
<i>Festuca arundinacea</i>	1		1	(14)
<i>Festuca glaucescens × Lolium multiflorum</i>	1		1	(19)
<i>Fourraea alpina</i>	1		1	(19)
<i>Glycine clandestina</i>	3		3	(12,13,22,23)
<i>Glycine max</i>	166	80	86	(12,13,20–23,59)
<i>Glycine soja</i>	3		3	(22,23)
<i>Gossypium arboreum</i>	2		2	(12,13)
<i>Gossypium herbaceum</i>	4		4	(32)
<i>Gossypium hirsutum</i>	53		53	(30–32)
<i>Gossypium raimondii</i>	11		11	(12,13,30,32)
<i>Hedyotis centranthoides</i>	3		3	(12,13)
<i>Hedyotis terminalis</i>	1		1	(12,13)
<i>Helianthus annuus</i>	3		3	(12,13)
<i>Henrardia persica</i>	1		1	(19)
<i>Heterantherum piliferum</i>	1		1	(19)
<i>Hordeum vulgare</i>	1		1	(14)
<i>Hordeum vulgare subsp. Spontaneum</i>	1		1	(12,13)
<i>Hordeum vulgare subsp. Vulgare</i>	16		16	(12,13)
<i>Ipomoea batatas</i>	1		1	(12,13)
<i>Ipomoea nil</i>	5		5	(12,13)
<i>Lactuca sativa</i>	3		3	(12,13)
<i>Liriodendron tulipifera</i>	2		2	(12,13)
<i>Lotus japonicus</i>	8		8	(12,13,22,59)
<i>Lupinus luteus</i>	1		1	(12,13)
<i>Malcolmia maritima</i>	1		1	(19)
<i>Malus × domestica</i>	3		3	(12,13)
<i>Manihot esculenta</i>	1		1	(14)
<i>Medicago truncatula</i>	76	36	40	(13,20,22,59,67–69)
<i>Nasturtium officinale</i>	1		1	(19)
<i>Nicotiana benthamiana</i>	4		4	(12,13)
<i>Nicotiana sylvestris</i>	1		1	(19)
<i>Nicotiana tabacum</i>	1		1	(12,13)

Table 1. Continued

Species	Number of miRNAs			References
	Total	Experimental	Computational	
<i>Nuphar advena</i>	3		3	(12,13)
<i>Oenocarpus bataua</i>	1		1	(19)
<i>Oryza barthii</i>	1		1	(19)
<i>Oryza brachyantha</i>	1		1	(19)
<i>Oryza nivara</i>	1		1	(19)
<i>Oryza rufipogon</i>	6		6	(19)
<i>Oryza sativa</i>	2540	269	2271	(2,3,6–8,10–13,28,52–55,69–78)
<i>Panicum virgatum</i>	1		1	(14)
<i>Pennisetum ciliare</i>	1		1	(14)
<i>Pennisetum glaucum</i>	1		1	(12,13)
<i>Peridictyon sanctum</i>	1		1	(19)
<i>Persea americana</i>	1		1	(12,13)
<i>Petunia x hybrida</i>	1		1	(19)
<i>Phaseolus coccineus</i>	2		2	(12,13)
<i>Phaseolus vulgaris</i>	2		2	(14,59)
<i>Phleum pratense</i>	1		1	(19)
<i>Physochloa patens</i>	282	281	1	(2,12,13,70,79–82)
<i>Picea glauca</i>	5		5	(12,13)
<i>Picea sitchensis</i>	3		3	(12,13)
<i>Pinus taeda</i>	40	24	16	(6,7,12,13,77)
<i>Populus tremula</i>	4		4	(12,13)
<i>Populus tremula x Populus tremuloides</i>	6		6	(14)
<i>Populus tremuloides</i>	7		7	(12,13)
<i>Populus trichocarpa</i>	2780	77	2703	(2,6,11,13,83,84)
<i>Populus trichocarpa x Populus deltoides</i>	5		5	(12–14)
<i>Prunus armeniaca</i>	2		2	(12,13)
<i>Prunus persica</i>	3		3	(14)
<i>Prunus salicina</i>	1		1	(19)
<i>Ricinus communis</i>	24		24	(19)
<i>Rorippa indica</i>	1		1	(19)
<i>Saccharum officinarum</i>	32		32	(14,20)
<i>Saccharum sp</i>	2		2	(12,13)
<i>Schedonorus arundinaceus</i>	1		1	(12,13)
<i>Secale cereale</i>	1		1	(12,13)
<i>Selaginella moellendorffii</i>	64	64		(79)
<i>Sesamum indicum</i>	2		2	(12,13)
<i>Sibara virginica</i>	1		1	(19)
<i>Solanum demissum</i>	1		1	(19)
<i>Solanum lycopersicum</i>	37	21	16	(12–14,34,85)
<i>Solanum tuberosum</i>	14		14	(12–14)
<i>Sorghum bicolor</i>	76		76	(12–14,20,27,86)
<i>Sorghum propinquum</i>	2		2	(12,13)
<i>Theobroma cacao</i>	1		1	(12,13)
<i>Thlaspi arvense</i>	1		1	(19)
<i>Trifolium repens</i>	1		1	(19)
<i>Triticum aestivum</i>	85	71	14	(12,13,24,25)
<i>Triticum monococcum</i>	2		2	(19)
<i>Triticum turgidum</i>	2		2	(12,13)
<i>Triticum urartu</i>	2		2	(19)
<i>Vigna unguiculata</i>	2		2	(22,59)
<i>Vitis vinifera</i>	142		142	(13,22,87)
<i>Zea mays</i>	207	2	205	(12,13,20,26–29)
<i>Zinnia elegans</i>	3		3	(12,13)

The curation of miRNAs that are not included in miRBase or that have not yet been named was performed using a special format that includes information about the corresponding author. An example is osa-miRf10000-akr, a rice miRNA discovered by Anders Krogh's group (11).

miRNA target gene, genome browser and stem-loop information

In order to construct a collection of miRNA target genes, we mainly focused on *Arabidopsis*, rice, poplar, soybean, cotton, *Medicago* and maize. The target genes collection

was curated from literature published in recent years. We also listed the location of interaction sites between miRNAs and target genes in *Arabidopsis* and rice [predicted by psRNATarget server (38,39)] in our database. Furthermore, we added information concerning miRNA action on target genes in *Arabidopsis*, rice and soybean

Species		Arabidopsis thaliana				
Precursor region	Precursor	ath-MIR168a [miRBase] stem-loop information stem loop structure computed using [Vienna RNA Package]				
	Sequence	CACCAUCGGGUCUGGAUUCGCUUGGUGCAGGUCGGGAACCAAUUCGGCUGACACAGCCUCGUGACUUUUAAACUUUUUUGGUUUG UGAGCAGGGAUUGGAUCCCGCCUUGCAUCAACUGAAUCGGAUCCCGAGGUG				
	Location	chr4: 10578646--10578783 [-] Genome Browser				
	Promoter	500bp 1kbp 3kbp				
	MIRNA family	MIRfam168				
Mature region	microRNA	ath-miR168a				
	Sequence	UCGCUUGGUCAGGUCGGAA				
	Evidence	experimental: cloned [44,45], Northern [44], 454 [2,46], MPSS [46] Rice microRNA expression profiling under oxidase stress microRNAs expression in tomato leaf tissue				
Expression	Microarray-based analysis of stress-regulated microRNAs in Arabidopsis thaliana Differential expression of miRNAs in response to salt stress in maize roots Expression profiles of MicroRNA of Arabidopsis grown at different temperature					
Targets region	Targets	Genes	Alignment	Expection	Target Description	
		AT1G48410.1	ath-miR168a: 21 AAGGGCUGGACGUGGUUCGCU 1 * * *	3	Symbols: AGO1 AGO1 (ARGONAUT1) chr1:17889766-17896254 REVERSE	
		AT1G48410.2	ath-miR168a: 21 AAGGGCUGGACGUGGUUCGCU 1 * * *	3	Symbols: AGO1 AGO1 (ARGONAUT1) chr1:17889766-17896254 REVERSE	
		AT3G04380.1	ath-miR168a: 21 AAGGGCUGGACGUGGUUCGCU 2 = * * *	3	Symbols: SUV4 SUV4; histone-lysine N-methyltransferase chr3:1161393-1164935 FORWARD	
		AT3G04380.2	ath-miR168a: 21 AAGGGCUGGACGUGGUUCGCU 2 = * * *	3	Symbols: SUV4 SUV4 chr3:1161507-1164771 FORWARD	
		Arabidopsis thaliana TAIR7, cDNA, released 04/25/2007 [Predicted through web-service: psRNAtarget]				
		[1] Gustafson AM, Allen E, Givan S, Smith D, Carrington JC, Kasschau KD.(2005). ASRP: The Arabidopsis Small RNA Project Database. Nucleic Acids Res 33, D637-40. [NCBI]				
References region	References	[2] Ramya Rajagopalan, Herve Vaucheret, Jerry Trejo and David P. Bartel.(2006). A diverse and evolutionarily fluid set of microRNAs in Arabidopsis thaliana. Genes & Dev. 20: 3407-3425. [NCBI]				
		[5] Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP. Prediction of Plant MicroRNA Targets. Cell, Vol. 110, 513-520, August 23, 2002. [NCBI]				
		[44] Reinhart BJ, Weinstein EG, Rhoades MW, Bartel B, Bartel DP. MicroRNAs in plants. Genes Dev. 16:1616-1626 (2002). [NCBI]				
		[45] Xie Z, Allen E, Fahlgren N, Calamari A, Givan SA, Carrington JC. Expression of Arabidopsis MIRNA genes. Plant Physiol. 138:2145-2154(2005). [NCBI]				
		[46] Lu C, Kulkarni K, Souret FF, MuthuVallippan R, Tej SS, Poethig RS, Henderson IR, Jacobsen SE, Wang W, Green PJ, Meyers BC. MicroRNAs and other small RNAs enriched in the Arabidopsis RNA-dependent RNA polymerase-2 mutant. Genome Res. 16:1276-1288(2006). [NCBI]				

Figure 1. Sample page for detailed information regarding each unique miRNA. Contents of this page are divided into four parts (miRNA ath-MIR168a was used as an example): the first section concerns miRNA precursor, including sequence, stem-loop information, miRNA family and location in the whole genome; the second section is about the mature miRNA, including sequence, evidence and expression patterns; the third section concerns target genes: for *Arabidopsis* and rice, the target genes and sites were predicted by psRNA target server (38) in default parameters; for poplar, soybean, cotton, medicago and maize, the listed genes were collected from the references. The last section includes the related references. The miRNA family and the evidence were based on the definition from miRBase, Rfam and the references.

into our local genome browser. We also provided the chromosomal location, secondary structure and dot matrix [generated by RNAfold (40)] of miRNA precursor sequences.

ACCESS TO THE DATABASE

The PMRD website was constructed using Hypertext Markup Language (HTML), perl CGI (<http://www.perl.com>), and the MySQL 4.0 (<http://www.mysql.com>) database engine. The detailed architecture of the

database tables is presented in Supplementary Figure S3. These tables are divided into three sections: the first section concerns miRNAs, the second section is about target genes, and the last is a presentation of miRNA expression profiles. The whole database is run on a server managed by the LINUX operating system.

The PMRD database web interface enables users to view and analyze each miRNA through either the Browse page or Search page. Detail information page (shown as Figure 1) displays basic information for each unique miRNA. The evidence for each entry is either

Table 2. Differentially expressed rice miRNAs after methyl viologen (MV) treatment

Probe ID	Log ₂ ratio (MV/mock)	P-value	Target_Seq
ptc-miR474c	2.17	3.67E-06	CAAAAGCUGUUGGGUUUGGCUGGG
ptc-miR474b	2.11	4.13E-05	CAAAAGUUGUUGGGUUUGGCUGGG
ptc-miR474a	2.08	1.49E-05	CAAAAGUUGCUGGGUUUGGCUGGG
tae-miR1125	1.66	1.12E-02	AACCAACGAGACCAACUCGCGGCGG
ath-miR404	1.37	2.51E-05	AUUAACGCUGGCGGUUGCGGCAGC
ath-miR854a	0.80	7.00E-04	GAUGAGGAUAGGGAGGAGGAG
osa-miR169b	0.77	4.30E-02	CAGCCAAGGAUGACUUGCCGG
ppt-miR903	0.76	4.31E-02	GCUACUUCGGCGGGACAAGAGC
osa-miR529b	0.68	2.29E-03	GAGAAGAGAGAGAGUACAGC
osa-miR820a	0.66	7.61E-03	CGGCCUCGUGGAUGGACCAGG
ppt-miR900-5p	0.65	4.60E-02	UCCCAGGUACAAGAACACAGC
ppt-miR395	0.60	5.00E-03	CUGAAGCGUUUGGGGGAAG
osa-miR159f	-0.76	5.86E-04	CUUGGAUUGAAGGGAGCUCUA
osa-miR396a	-0.81	2.72E-02	UCCACAGCUUUCUUGAACUG
osa-miR159a	-0.83	1.19E-04	UUUGGAUUGAAGGGAGCUCUG
osa-miR397b	-0.99	6.99E-03	UUAUUGAGUGCAGCGUUGAUG
tae-miR1120	-1.08	2.36E-02	ACAUCUUAUUAUUGAGACGGAG
osa-miR396c	-1.22	1.36E-03	UCCACAGCUUUCUUGAACUU

experimental or computational. We listed the methods used for experimentally demonstrated miRNAs and the programs applied for computational prediction. Within the total 8433 miRNAs in PMRD, 1297 of them were experimentally demonstrated and 7136 of them were from computational predictions (the detailed number shown in Table 1). Users can view a miRNA display in the local Genome Browser (shown as Supplementary Figure S4) in which users can extract information about *Arabidopsis*, rice and soybean. Through PMRD, users can also inspect miRNA expression profiles, target gene list and predicted interaction sites between miRNA and cDNA. In addition, the secondary structure of miRNA precursor could be predicted by using the prediction tool in our PMRD database. All miRNA sequences are available for downloading.

Currently, we have collected thousands of miRNA sequences from *Arabidopsis*, rice and poplar in our database, but for other species such as soybean, maize, wheat, cotton and sorghum, etc., the identified miRNAs are limited. Since this field is progressing rapidly due to new deep sequencing technologies, many more plant miRNAs will be discovered, as well as their target genes and expression profiles. The PMRD will be updated and maintained in order to integrate the newest plant miRNA data.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

The authors thank Ms Wenyang Xu for critical suggestions. They also thank Dongxia Yao and Qiang Wei for their help on taxonomic browse.

FUNDING

Ministry of Science and Technology of China grants (2008AA02Z312 and 2006CB100105). Funding for open access charge: Ministry of Science and Technology of China (2008AA02Z312 and 2006CB100105).

Conflict of interest statement. None declared.

REFERENCES

- Bartel,D.P. (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*, **116**, 281–297.
- Jones-Rhoades,M.W., Bartel,D.P. and Bartel,B. (2006) MicroRNAs and their regulatory roles in plants. *Annu. Rev. Plant Biol.*, **57**, 19–53.
- Archak,S. and Nagaraju,J. (2007) Computational prediction of rice (*Oryza sativa*) miRNA targets. *Genomics Proteomics Bioinformatics*, **5**, 196–206.
- Sunkar,R., Chinnusamy,V., Zhu,J. and Zhu,J.K. (2007) Small RNAs as big players in plant abiotic stress responses and nutrient deprivation. *Trends Plant Sci.*, **12**, 301–309.
- Zhou,X., Wang,G., Sutoh,K., Zhu,J.K. and Zhang,W. (2008) Identification of cold-inducible microRNAs in plants by transcriptome analysis. *Biochim. Biophys. Acta*.
- Lu,S., Sun,Y.H. and Chiang,V.L. (2008) Stress-responsive microRNAs in *Populus*. *Plant J.*, **55**, 131–151.
- Morin,R.D., Aksay,G., Dolgosheina,E., Ebhardt,H.A., Magrini,V., Mardis,E.R., Sahinalp,S.C. and Unrau,P.J. (2008) Comparative analysis of the small RNA transcriptomes of *Pinus contorta* and *Oryza sativa*. *Genome Res.*, **18**, 571–584.
- Jones-Rhoades,M.W. and Bartel,D.P. (2004) Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Mol. Cell*, **14**, 787–799.
- Rajagopalan,R., Vaucheret,H., Trejo,J. and Bartel,D.P. (2006) A diverse and evolutionarily fluid set of microRNAs in *Arabidopsis thaliana*. *Genes Dev.*, **20**, 3407–3425.
- Sunkar,R., Zhou,X., Zheng,Y., Zhang,W. and Zhu,J.K. (2008) Identification of novel and candidate miRNAs in rice by high throughput sequencing. *BMC Plant Biol.*, **8**, 25.
- Lindow,M., Jacobsen,A., Nygaard,S., Mang,Y. and Krogh,A. (2007) Intragenomic matching reveals a huge potential for miRNA-mediated regulation in plants. *PLoS Comput. Biol.*, **3**, e238.

12. Zhang, B., Pan, X., Cannon, C.H., Cobb, G.P. and Anderson, T.A. (2006) Conservation and divergence of plant microRNA genes. *Plant J.*, **46**, 243–259.
13. Zhang, B.H., Pan, X.P., Wang, Q.L., Cobb, G.P. and Anderson, T.A. (2005) Identification and characterization of new plant microRNAs using EST analysis. *Cell Res.*, **15**, 336–360.
14. Sunkar, R. and Jagadeeswaran, G. (2008) In silico identification of conserved microRNAs in large number of diverse plant species. *BMC Plant Biol.*, **8**, 37.
15. Griffiths-Jones, S., Grocock, R.J., van Dongen, S., Bateman, A. and Enright, A.J. (2006) miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.*, **34**, D140–144.
16. Gustafson, A.M., Allen, E., Givan, S., Smith, D., Carrington, J.C. and Kasschau, K.D. (2005) ASRP: the Arabidopsis Small RNA Project Database. *Nucleic Acids Res.*, **33**, D637–640.
17. Johnson, C., Bowman, L., Adai, A.T., Vance, V. and Sundaresan, V. (2007) CSRDB: a small RNA integrated database and browser resource for cereals. *Nucleic Acids Res.*, **35**, D829–833.
18. Gardner, P.P., Daub, J., Tate, J.G., Nawrocki, E.P., Kolbe, D.L., Lindgreen, S., Wilkinson, A.C., Finn, R.D., Griffiths-Jones, S., Eddy, S.R. *et al.* (2009) Rfam: updates to the RNA families database. *Nucleic Acids Res.*, **37**, D136–140.
19. Griffiths-Jones, S., Bateman, A., Marshall, M., Khanna, A. and Eddy, S.R. (2003) Rfam: an RNA family database. *Nucleic Acids Res.*, **31**, 439–441.
20. Dezulian, T., Palatnik, J., Huson, D. and Weigel, D. (2005) Conservation and divergence of microRNA families in plants. *Genome Biol.*, **6**, P13.
21. Subramanian, S., Fu, Y., Sunkar, R., Barbazuk, W.B., Zhu, J.K. and Yu, O. (2008) Novel and nodulation-regulated microRNAs in soybean roots. *BMC Genomics*, **9**, 160.
22. Wang, Y., Li, P., Cao, X., Wang, X., Zhang, A. and Li, X. (2009) Identification and expression analysis of miRNAs from nitrogen-fixing soybean nodules. *Biochem. Biophys. Res. Commun.*, **378**, 799–803.
23. Zhang, B., Pan, X. and Stellwag, E.J. (2008) Identification of soybean microRNAs and their targets. *Planta*, **229**, 161–182.
24. Yao, Y., Guo, G., Ni, Z., Sunkar, R., Du, J., Zhu, J.K. and Sun, Q. (2007) Cloning and characterization of microRNAs from wheat (*Triticum aestivum* L.). *Genome Biol.*, **8**, R96.
25. Wei, B., Cai, T., Zhang, R., Li, A., Huo, N., Li, S., Gu, Y.Q., Vogel, J., Jia, J., Qi, Y. *et al.* (2009) Novel microRNAs uncovered by deep sequencing of small RNA transcriptomes in bread wheat (*Triticum aestivum* L.) and *Brachypodium distachyon* (L.) Beauv. *Funct. Integr. Genomics.*, **9**, 532–541.
26. Juarez, M.T., Kui, J.S., Thomas, J., Heller, B.A. and Timmermans, M.C. (2004) microRNA-mediated repression of rolled leaf1 specifies maize leaf polarity. *Nature*, **428**, 84–88.
27. Maher, C., Timmermans, M., Stein, L. and Ware, D. (2004) Identifying microRNAs in plant genomes. *Proc. IEEE CSB*, **6**, 718–723.
28. Xue, L.J., Zhang, J.J. and Xue, H.W. (2009) Characterization and expression profiles of miRNAs in rice seeds. *Nucleic Acids Res.*, **37**, 916–930.
29. Zhang, B., Pan, X. and Anderson, T.A. (2006) Identification of 188 conserved maize microRNAs and their targets. *FEBS Lett.*, **580**, 3753–3762.
30. Khan Barozai, M.Y., Irfan, M., Yousaf, R., Ali, I., Qaisar, U., Maqbool, A., Zahoor, M., Rashid, B., Hussain, T. and Riazuddin, S. (2008) Identification of micro-RNAs in cotton. *Plant Physiol. Biochem.*, **46**, 739–751.
31. Qiu, C.X., Xie, F.L., Zhu, Y.Y., Guo, K., Huang, S.Q., Nie, L. and Yang, Z.M. (2007) Computational identification of microRNAs and their targets in *Gossypium hirsutum* expressed sequence tags. *Gene*, **395**, 49–61.
32. Zhang, B., Wang, Q., Wang, K., Pan, X., Liu, F., Guo, T., Cobb, G.P. and Anderson, T.A. (2007) Identification of cotton microRNAs and their targets. *Gene*, **397**, 26–37.
33. Wang, X., Elling, A.A., Li, X., Li, N., Peng, Z., He, G., Sun, H., Qi, Y., Liu, X.S. and Deng, X.W. (2009) Genome-Wide and Organ-Specific Landscapes of Epigenetic Modifications and Their Relationships to mRNA and Small RNA Transcriptomes in Maize. *Plant Cell*, **21**, 1053–1069.
34. Zhang, J., Zeng, R., Chen, J., Liu, X. and Liao, Q. (2008) Identification of conserved microRNAs and their targets from *Solanum lycopersicum* Mill. *Gene*, **423**, 1–7.
35. Liu, H.H., Tian, X., Li, Y.J., Wu, C.A. and Zheng, C.C. (2008) Microarray-based analysis of stress-regulated microRNAs in *Arabidopsis thaliana*. *RNA*, **14**, 836–843.
36. Ding, D., Zhang, L., Wang, H., Liu, Z., Zhang, Z. and Zheng, Y. (2008) Differential expression of miRNAs in response to salt stress in maize roots. *Ann. Botany*, **103**, 29–38.
37. Hu, J. and He, X. (2007) Enhanced quantile normalization of microarray data to reduce loss of information in gene expression profiles. *Biometrics*, **63**, 50–59.
38. Zhang, Y. (2005) miRU: an automated plant miRNA target prediction server. *Nucleic Acids Res.*, **33**, W701–W704.
39. Enright, A.J., John, B., Gaul, U., Tuschl, T., Sander, C. and Marks, D.S. (2003) MicroRNA targets in *Drosophila*. *Genome Biol.*, **5**, R1.
40. Schuster, P., Fontana, W., Stadler, P.F. and Hofacker, I.L. (1994) From sequences to shapes and back: a case study in RNA secondary structures. *Proceedings*, **255**, 279–284.
41. Park, W., Li, J., Song, R., Messing, J. and Chen, X. (2002) CARPEL FACTORY, a Dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in *Arabidopsis thaliana*. *Curr. Biol.*, **12**, 1484–1495.
42. Adai, A., Johnson, C., Mlotshwa, S., Archer-Evans, S., Manocha, V., Vance, V. and Sundaresan, V. (2005) Computational prediction of miRNAs in *Arabidopsis thaliana*. *Genome Res.*, **15**, 78–91.
43. Palatnik, J.F., Allen, E., Wu, X., Schommer, C., Schwab, R., Carrington, J.C. and Weigel, D. (2003) Control of leaf morphogenesis by microRNAs. *Nature*, **425**, 257–263.
44. Xie, Z., Allen, E., Fahlgren, N., Calamar, A., Givan, S.A. and Carrington, J.C. (2005) Expression of Arabidopsis MIRNA genes. *Plant Physiol.*, **138**, 2145–2154.
45. Arteaga-Vazquez, M., Caballero-Perez, J. and Vielle-Calzada, J.P. (2006) A family of microRNAs present in plants and animals. *Plant Cell*, **18**, 3355–3369.
46. Franco-Zorrilla, J.M., Del Toro, F.J., Godoy, M., Perez-Perez, J., Lopez-Vidriero, I., Oliveros, J.C., Garcia-Casado, G., Llave, C. and Solano, R. (2009) Genome-wide identification of small RNA targets based on target enrichment and microarray hybridizations. *Plant J.*, **59**, 840–850.
47. German, M.A., Pillay, M., Jeong, D.H., Hetawal, A., Luo, S., Janardhanan, P., Kannan, V., Rymarquis, L.A., Nobuta, K., German, R. *et al.* (2008) Global identification of microRNA-target RNA pairs by parallel analysis of RNA ends. *Nat. Biotechnol.*, **26**, 941–946.
48. Fahlgren, N., Howell, M.D., Kasschau, K.D., Chapman, E.J., Sullivan, C.M., Cumbie, J.S., Givan, S.A., Law, T.F., Grant, S.R., Dangel, J.L. *et al.* (2007) High-throughput sequencing of Arabidopsis microRNAs: evidence for frequent birth and death of MIRNA genes. *PLoS ONE*, **2**, e219.
49. Allen, E., Xie, Z., Gustafson, A.M. and Carrington, J.C. (2005) microRNA-directed phasing during trans-acting siRNA biogenesis in plants. *Cell*, **121**, 207–221.
50. Kutter, C., Schob, H., Stadler, M., Meins, F. Jr and Si-Ammour, A. (2007) MicroRNA-mediated regulation of stomatal development in *Arabidopsis*. *Plant Cell*, **19**, 2417–2429.
51. Lu, C., Kulkarni, K., Souret, F.F., MuthuValliappan, R., Tej, S.S., Poethig, R.S., Henderson, I.R., Jacobsen, S.E., Wang, W., Green, P.J. *et al.* (2006) MicroRNAs and other small RNAs enriched in the *Arabidopsis* RNA-dependent RNA polymerase-2 mutant. *Genome Res.*, **16**, 1276–1288.
52. Reinhart, B.J., Weinstein, E.G., Rhoades, M.W., Bartel, B. and Bartel, D.P. (2002) MicroRNAs in plants. *Genes Dev.*, **16**, 1616–1626.
53. Sunkar, R. and Zhu, J.K. (2004) Novel and stress-regulated microRNAs and other small RNAs from *Arabidopsis*. *Plant Cell*, **16**, 2001–2019.
54. Wang, X.J., Reyes, J.L., Chua, N.H. and Gaasterland, T. (2004) Prediction and identification of Arabidopsis thaliana microRNAs and their mRNA targets. *Genome Biol.*, **5**, R65.
55. Rhoades, M.W., Reinhart, B.J., Lim, L.P., Burge, C.B., Bartel, B. and Bartel, D.P. (2002) Prediction of plant microRNA targets. *Cell*, **110**, 513–520.

56. Baulcombe, D. (2004) RNA silencing in plants. *Nature*, **431**, 356–363.
57. Mette, M.F., van der Winden, J., Matzke, M. and Matzke, A.J. (2002) Short RNAs can identify new candidate transposable element families in Arabidopsis. *Plant Physiol.*, **130**, 6–9.
58. Zhou, X., Wang, G. and Zhang, W. (2007) UV-B responsive microRNA genes in Arabidopsis thaliana. *Mol. Syst. Biol.*, **3**, 103.
59. Jagadeeswaran, G., Zheng, Y., Li, Y.F., Shukla, L.I., Matts, J., Hoyt, P., Macmill, S.L., Wiley, G.B., Roe, B.A., Zhang, W. et al. (2009) Cloning and characterization of small RNAs from Medicago truncatula reveals four novel legume-specific microRNA families. *New Phytol.*, **184**, 85–98.
60. Wang, L., Wang, M.B., Tu, J.X., Helliwell, C.A., Waterhouse, P.M., Dennis, E.S., Fu, T.D. and Fan, Y.L. (2007) Cloning and characterization of microRNAs from Brassica napus. *FEBS Lett.*, **581**, 3848–3856.
61. Xie, F.L., Huang, S.Q., Guo, K., Xiang, A.L., Zhu, Y.Y., Nie, L. and Yang, Z.M. (2007) Computational identification of novel microRNAs and targets in Brassica napus. *FEBS Lett.*, **581**, 1464–1474.
62. He, X.F., Fang, Y.Y., Feng, L. and Guo, H.S. (2008) Characterization of conserved and novel microRNAs and their targets, including a TuMV-induced TIR-NBS-LRR class R gene-derived novel miRNA in Brassica. *FEBS Lett.*, **582**, 2445–2452.
63. Porter, B.W., Aizawa, K.S., Zhu, Y.J. and Christopher, D.A. (2008) Differentially expressed and new non-protein-coding genes from a Carica papaya root transcriptome survey. *Plant Sci.*, **174**, 38–50.
64. Zhao, T., Li, G., Mi, S., Li, S., Hannon, G.J., Wang, X.J. and Qi, Y. (2007) A complex system of small RNAs in the unicellular green alga Chlamydomonas reinhardtii. *Genes Dev.*, **21**, 1190–1203.
65. Molnar, A., Schwach, F., Studholme, D.J., Thuenemann, E.C. and Baulcombe, D.C. (2007) miRNAs control gene expression in the single-cell alga Chlamydomonas reinhardtii. *Nature*, **447**, 1126–1129.
66. Hinas, A., Reimegard, J., Wagner, E.G., Nellen, W., Ambros, V.R. and Soderbom, F. (2007) The small RNA repertoire of Dictyostelium discoideum and its regulation by components of the RNAi pathway. *Nucleic Acids Res.*, **35**, 6714–6726.
67. Zhou, Z.S., Huang, S.Q. and Yang, Z.M. (2008) Bioinformatic identification and expression analysis of new microRNAs from Medicago truncatula. *Biochem. Biophys. Res. Commun.*, **374**, 538–542.
68. Szittyá, G., Moxon, S., Santos, D.M., Jing, R., Fevereiro, M.P., Moulton, V. and Dalmay, T. (2008) High-throughput sequencing of Medicago truncatula short RNAs identifies eight new miRNA families. *BMC Genomics*, **9**, 593.
69. Guddeti, S., Zhang, D.C., Li, A.L., Leseberg, C.H., Kang, H., Li, X.G., Zhai, W.X., Johns, M.A. and Mao, L. (2005) Molecular evolution of the rice miR395 gene family. *Cell Res.*, **15**, 631–638.
70. Arazi, T., Talmor-Neiman, M., Stav, R., Riese, M., Huijser, P. and Baulcombe, D.C. (2005) Cloning and characterization of micro-RNAs from moss. *Plant J.*, **43**, 837–848.
71. Sunkar, R., Girke, T., Jain, P.K. and Zhu, J.K. (2005) Cloning and characterization of microRNAs from rice. *Plant Cell*, **17**, 1397–1411.
72. Li, Y., Li, W. and Jin, Y.X. (2005) Computational identification of novel family members of microRNA genes in Arabidopsis thaliana and Oryza sativa. *Acta Biochim Biophys. Sin.*, **37**, 75–87.
73. Zhu, Q.H., Spriggs, A., Matthew, L., Fan, L., Kennedy, G., Gubler, F. and Helliwell, C. (2008) A diverse set of microRNAs and microRNA-like small RNAs in developing rice grains. *Genome Res.*, **18**, 1456–1465.
74. Lu, C., Jeong, D.H., Kulkarni, K., Pillay, M., Nobuta, K., German, R., Thatcher, S.R., Maher, C., Zhang, L., Ware, D. et al. (2008) Genome-wide analysis for discovery of rice microRNAs reveals natural antisense microRNAs (nat-miRNAs). *Proc. Natl Acad. Sci. USA*, **105**, 4951–4956.
75. Lacombe, S., Nagasaki, H., Santi, C., Duval, D., Piegu, B., Bangratz, M., Breitler, J.C., Guiderdoni, E., Brugidou, C., Hirsch, J. et al. (2008) Identification of precursor transcripts for 6 novel miRNAs expands the diversity on the genomic organisation and expression of miRNA genes in rice. *BMC Plant Biol.*, **8**, 123.
76. Liu, B., Li, P., Li, X., Liu, C., Cao, S., Chu, C. and Cao, X. (2005) Loss of function of OsDCL1 affects microRNA accumulation and causes developmental defects in rice. *Plant Physiol.*, **139**, 296–305.
77. Lu, S., Sun, Y.H., Amerson, H. and Chiang, V.L. (2007) MicroRNAs in loblolly pine (Pinus taeda L.) and their association with fusiform rust gall development. *Plant J.*, **51**, 1077–1098.
78. Luo, Y.C., Zhou, H., Li, Y., Chen, J.Y., Yang, J.H., Chen, Y.Q. and Qu, L.H. (2006) Rice embryogenic calli express a unique set of microRNAs, suggesting regulatory roles of microRNAs in plant post-embryonic development. *FEBS Lett.*, **580**, 5111–5116.
79. Axtell, M.J., Snyder, J.A. and Bartel, D.P. (2007) Common functions for diverse small RNAs of land plants. *Plant Cell*, **19**, 1750–1769.
80. Fattash, I., Voss, B., Reski, R., Hess, W.R. and Frank, W. (2007) Evidence for the rapid expansion of microRNA-mediated regulation in early land plant evolution. *BMC Plant Biol.*, **7**, 13.
81. Talmor-Neiman, M., Stav, R., Frank, W., Voss, B. and Arazi, T. (2006) Novel micro-RNAs and intermediates of micro-RNA biogenesis from moss. *Plant J.*, **47**, 25–37.
82. Cho, S.H., Addo-Quaye, C., Coruh, C., Arif, M.A., Ma, Z., Frank, W. and Axtell, M.J. (2008) Physcomitrella patens DCL3 is required for 22-24 nt siRNA accumulation, suppression of retrotransposon-derived transcripts, and normal development. *PLoS Genet.*, **4**, e1000314.
83. Tuskan, G.A., Difazio, S., Jansson, S., Bohlmann, J., Grigoriev, I., Hellsten, U., Putnam, N., Ralph, S., Rombauts, S., Salamov, A. et al. (2006) The genome of black cottonwood, Populus trichocarpa (Torr. & Gray). *Science (New York, NY)*, **313**, 1596–1604.
84. Lu, S., Sun, Y.H., Shi, R., Clark, C., Li, L. and Chiang, V.L. (2005) Novel and mechanical stress-responsive MicroRNAs in Populus trichocarpa that are absent from Arabidopsis. *Plant Cell*, **17**, 2186–2203.
85. Moxon, S., Jing, R., Szittyá, G., Schwach, F., Rusholme Pilcher, R.L., Moulton, V. and Dalmay, T. (2008) Deep sequencing of tomato short RNAs identifies microRNAs targeting genes involved in fruit ripening. *Genome Res.*, **18**, 1602–1609.
86. Bedell, J.A., Budiman, M.A., Nunberg, A., Citek, R.W., Robbins, D., Jones, J., Flick, E., Rholting, T., Fries, J., Bradford, K. et al. (2005) Sorghum genome sequencing by methylation filtration. *PLoS Biol.*, **3**, e13.
87. Jaillon, O., Aury, J.M., Noel, B., Policriti, A., Clepet, C., Casagrande, A., Choisne, N., Aubourg, S., Vitulo, N., Jubin, C. et al. (2007) The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature*, **449**, 463–467.