



***Poaceascoma aquaticum* sp. nov. (Lentitheciaceae), a new species from submerged bamboo in freshwater**

ZONG-LONG LUO^{1,2}, ALI H. BAHKALI³, XIAO-YING LIU^{1,4}, RUNGTIWA PHOOKAMSAK², YONG-CHANG ZHAO⁵, DE-QUN ZHOU⁶, HONG-YAN SU^{1*,6} & KEVIN D. HYDE²

¹ College of Agriculture & Biology, Dali University, Dali, 671003, Yunnan, PR China.

² Centre of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand.

³ College of Science, Botany and Microbiology Department, King Saud University, Riyadh 1145, Saudi Arabia.

⁴ College of Basic Medicine, Dali University, Dali, Yunnan 671000, China.

⁵ Institute of Biotechnology and Germplasmic Resources, Yunnan Academy of Agricultural Sciences, Kunming, 650223, PR China.

⁶ Faculty of Environmental Sciences & Engineering, Kunming University of Science & Technology, Kunming 650500, Yunnan, PR China.

Corresponding author: Hong-Yan Su, email address: suhongyan16@163.com

Abstract

Poaceascoma aquaticum sp. nov. (Lentitheciaceae) was found on decaying bamboo submerged in a freshwater stream in northern Thailand. The new species is introduced based on its unique morphology and phylogeny. *Poaceascoma aquaticum* is characterized by black, semi-immersed or erumpent, papillate ascomata, bitunicate, cylindrical to cylindrical-clavate, short pedicellate asci and filiform, pale brown to brown ascospores. Phylogenetic analyses of combined LSU, SSU and RPB2 sequence data shows that *Poaceascoma aquaticum* forms a robust clade with *P. helicoides* in the family Lentitheciaceae (Pleosporales). *Poaceascoma aquaticum* is described, illustrated and compared with related taxa.

Key words: Aquatic fungi, Lentitheciaceae, Phylogeny, Taxonomy

Introduction

Freshwater ascomycetes are an ecological assemblage that occur on submerged or partially submerged woody substrates in freshwater habitats, including both sexual and asexual species (Shearer 1993, 2001, Hyde *et al.* 2015). These taxa play a key role in the degradation of organic carbon (Wong *et al.* 1998; Gulis *et al.* 2006; Krauss *et al.* 2011). Both sexual and asexual morphs of ascomycetes as well as a few basidiomycetes have been found on various substrates in freshwater in lentic (lakes, ponds) and lotic (rivers, streams, creeks, peat swamps) habitats (Wong *et al.* 1999, Cai *et al.* 2003a, b, Jones & Choeyklin 2008, Hyde *et al.* 2015). The freshwater fungi of the Asian region have been particularly well-studied with collections made from Australia, China, Malaysia, Philippines and Thailand (Nawawi 1985, Kuthubutheen 1987, Hyde 1995, Sivichai *et al.* 2000, Ho *et al.* 2001, 2002, Cai *et al.* 2003a, b, Goh and Tsui 2003, Luo *et al.* 2004, Sivichai & Jones 2004, Jones *et al.* 2007, Hu *et al.* 2010, Kurniawati *et al.* 2010); and the data are summarized by Hyde *et al.* (2015). Despite this, collections in unstudied streams will often result in the discovery of new species.

This manuscript is part of a study on the taxonomy and diversity of microfungi on substrates in freshwater, along a north-south latitudinal gradient from China through to New Zealand (Hyde *et al.* 2015). We have previously introduced several new taxa (Luo *et al.* 2015, Su *et al.* 2015a, b, Liu *et al.* 2015a). In this paper, we introduce a new species, *Poaceascoma aquaticum*, in the family Lentitheciaceae (Pleosporales). The new species is defined based on its unique morphological characters as well as support from phylogenetic analyses of combined LSU, SSU and RPB2 sequence data.

Materials and methods

Isolation and morphology

Decaying bamboo was collected in November 2013 from a freshwater stream in Chiang Rai Province, Thailand and returned to the laboratory in plastic Ziploc bags. The samples were incubated in plastic boxes lined with moistened tissue paper at room temperature for one week. The samples were processed and examined following the methods described by Taylor and Hyde (2003). Morphological observations were made under a Nikon SMZ-171 dissecting microscope and captured by using a Cannon EOS 600D camera on a Nikon Eclipse 80i compound microscope.

Single ascospore isolations were made to obtain pure cultures as described in Chomnunti *et al.* (2014). The pure cultures are deposited in Mae Fah Luang University Culture Collection (MFLUCC) and Dali University Culture Collection (DLUCC). Herbarium specimens are deposited at the herbarium of Mae Fah Luang University (MFLU) and the herbaria of Kunming Institute of Botany, Chinese Academy of Sciences (HKAS). Faces of fungi numbers are as detailed in Jayasiri *et al.* (2015).

DNA extraction, PCR amplification and sequencing

Total genomic DNA was extracted from fresh fungal mycelium grown on PDA at 25°C. The EZ gene™ Fungal gDNA kit (GD2416) was used to extract DNA according to the manufacturer's instructions. The primer pair LROR and LR7 was used to amplify partial large subunits of the nuclear ribosomal RNA gene (LSU) (Vilgalys & Hester 1990). The PCR thermal cycle program for LSU, SSU amplification were as follows: initially 95 °C for 3 minutes, followed by 35 cycles of denaturation at 95°C for 30 seconds, annealing at 50 °C for 40 seconds, elongation at 72 °C for 90 seconds, and final extension at 72°C for 10 minutes. The PCR thermal cycle program for the partial RNA polymerase second largest subunit (*RPB2*) was followed as initially 95 °C for 5 mins, followed by 40 cycle of denaturation at 95 °C for 1 mins, annealing at 52 °C for 2 mins, elongation at 72 °C for 90 seconds, and finalextension at 72°C for 10 mins. PCR products were purified using minicolumns, purification resin and buffer according to the manufacturer's protocols (Amersham product code: 27–9602–01). The PCR products were observed on 1% agarose electrophoresis gels stained with ethidium bromide. Purification and sequencing of PCR products were carried at Shanghai Sangon Biological Engineering Technology and Services Co., Ltd (Shanghai, P.R. China).

Phylogenetic analysis

The sequences generated from this study were analyzed with sequence data from related taxa and others from families in the suborder Massarineae (Pleosporales) as obtained from GenBank using the dataset in Phookamsak *et al.* (2015). The consensus sequences were then initially aligned using MAFFT v.7 (<http://mafft.cbrc.jp/alignment/server/>) (Katoh & Standley 2013) and further improved using Bioedit v.5.0.6 (Hall 2001) and ClustalX v. 1.83 (Thompson *et al.* 1997) to allow maximum alignment and maximum sequence similarity.

A maximum likelihood analysis was performed using RAXMLGUI v. 1.3 (Silvestro & Michalak 2011). The optimal ML tree search was conducted with 100 separate runs, using the default algorithm of the program from a random starting tree for each run. The final tree was selected among suboptimal trees from each run by comparing likelihood scores under the GTR+GAMMA substitution model.

Maximum-parsimony analyses were performed using the heuristic search option with 1000 random taxa additions and tree bisection and reconnection (TBR) as the branch-swapping algorithm. All characters were unordered and of equal weight and gaps were treated as missing data. Maxtrees were unlimited, branches of zero length were collapsed and all multiple, equally parsimonious trees were saved. Clade stability was assessed using a bootstrap (BT) analysis with 1000 replicates, each with 10 replicates of random stepwise addition of taxa (Hillis & Bull 1993).

Bayesian analyses were performed by using PAUP v.4.0b10 (Swofford 2002) and MrBayes v3.0b4 (Ronquist & Huelsenbeck 2003). The model of evolution was estimated by using MrModeltest 2.2 (Nylander 2004). Posterior probabilities (PP) (Rannala & Yang 1996) were performed by Markov Chain Monte Carlo Sampling (BMCMC) in MrBayes v. 3.0b4 (Liu *et al.* 2012). Six simultaneous Markov Chains were run for 1 m generations and trees were sampled every 100th generation (Resulting 10000 total trees) (Cai *et al.* 2006). The first 2000 trees representing the burn-in phase of the analyses were discarded and the remaining 8000 (post burning) trees used for calculating posterior probabilities (PP) in the majority rule consensus tree (Cai *et al.* 2006, Liu *et al.* 2012).

Trees were viewed in Treeview (Page 1996). Sequences derived in this study are deposited in GenBank (Table 1).

TABLE 1. Isolates used in this study and their GenBank accession numbers. The ex-type strains are in bold; the newly generated sequences are indicated in red.

Taxon	Culture/voucher	GenBank Accession Number		
		LSU	SSU	RPB2
<i>Bambusicola bambusae</i> ^T	MFLUCC 11-0614	JX442035	JX442039	KP761718
<i>Bambusicola irregulispota</i> ^T	MFLUCC 11-0437	JX442036	JX442040	KP761719
<i>Bambusicola massarinia</i> ^{T/Ts}	MFLUCC 11-0389	JX442037	JX442041	KP761716
<i>Bambusicola splendida</i> ^T	MFLUCC 11-0439	JX442038	JX442042	KP761717
<i>Bimuria novae-zelandiae</i> ^{T/Ts}	CBS 107.79	AY016356	AY016338	DQ470917
<i>Corynespora cassiicola</i>	CBS 100822	GU301808	GU296144	GU371742
<i>Corynespora smithii</i>	CABI 5649b	GU323201	–	GU371783
<i>Deniquelata barringtoniae</i> ^{T/Ts}	MFLUCC 11-0422	JX254655	JX254656	–
<i>Didymosphaeria rubi-ulmifolii</i> ^T	CBS 100299	JX496124	AY642523	–
<i>Falciformispora lignatilis</i> ^{Ts}	BCC 21117	GU371826	GU371834	–
<i>Falciformispora lignatilis</i> ^{Ts}	BCC 21118	GU371827	GU371835	–
<i>Helicascus nypae</i>	BCC 36752	GU479789	GU479755	GU479827
<i>Kalmusia scabrispora</i> ^T	NBRC 106237	AB524594	AB524453	AB539094
<i>Karstenula rhodostoma</i>	CBS 690.94	GU301821	GU296154	GU371788
<i>Katumotoa bambusicola</i> ^{T/Ts}	MAFF 239641	AB524595	AB524454	AB539095
<i>Keissleriella cladophila</i> ^T	CBS 104.55	GU301822	GU296155	–
<i>Keissleriella dactylis</i> ^T	MFUCC 13-0751	KP197668	KP197666	–
<i>Keissleriella poagensis</i> ^T	CBS 136767	KJ869170	–	–
<i>Keissleriella trichophoricola</i> ^T	CBS 136770	KJ869171	–	–
<i>Lentithecium aquaticum</i> ^T	CBS 123099	GU301823	GU296156	–
<i>Lentithecium arundinaceum</i>	CBS 619.86	GU301824	GU296157	–
<i>Lentithecium fluviatile</i> ^{Ts}	CBS 122367	GU301825	GU296158	–
<i>Massarina cisti</i>	CBS 266.62	FJ795447	FJ795490	FJ795464
<i>Massarina eburnea</i>	CBS 473.64	GU301840	GU296170	GU371732
<i>Melanomma pulvis-pyrius</i> ^{T/Ts}	CBS 124080	GU456323	GU456302	GU456350
<i>Montagnula opulenta</i>	CBS 168.34	DQ678086	AF164370	DQ677984
<i>Morosphaeria ramunculicola</i>	JK 5304B	GU479794	GU479760	GU479831
<i>Murilentithecium clematidis</i> ^{T/Ts}	MFLUCC 14-0561	KM408758	KM408760	–
<i>Neottiosporina paspali</i>	CBS 331.37	EU754172	EU754073	GU371779
<i>Ophiosphaerella sasicola</i>	MAFF 239644	AB524599	AB524458	–
<i>Palmiascoma gregariascomum</i> ^{T/Ts}	MFLUCC 11-0175	KP744495	KP753958	–
<i>Paraconiothyrium minitans</i>	CBS 122788	EU754173	EU754074	GU371776
<i>Paraphaeosphaeria michotii</i> ^{T/Ts}	MFLUCC 13-0349	KJ939282	KJ939285	–
<i>Phaeodothis winteri</i>	CBS 182.58	GU301857	GU296183	–
<i>Poaceascoma aquaticum</i> ^{T/Ts}	MFLUCC 14-0048	KT324690	KT324691	KT373846
<i>Prosthemium canba</i>	JCM 16966	AB553760	AB553646	–
<i>Setoseptoria phragmitis</i> ^{T/Ts}	CBS 114802	KF251752	–	–
<i>Stagonospora macropycnidia</i>	CBS 114202	GU301873	GU296198	–
<i>Stagonospora paludosa</i> ^{T/Ts}	CBS135088	KF251760	–	KF252262
<i>Trematosphaeria pertusa</i> ^{T/Ts}	CBS 122368	FJ201990	FJ201991	FJ795476
<i>Trematosphaeria pertusa</i> ^{Ts}	CBS 122371	FJ201992	FJ201993	GU371801
<i>Tingoldiogo graminicola</i> ^{T/Ts}	JCM 16485	AB521743	AB521726	–

Abbreviations: **BCC**: BIOTEC Culture Collection, Bangkok, Thailand; **CABI**: International Mycological Institute, CABI-Bioscience, Egham, Basingstoke, Hampshire, UK; **CBS**: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; **JCM**: The Japan Collection of Microorganisms, Japan; **MAFF**: Ministry of Agriculture, Forestry and Fisheries, Japan; **MFLUCC**: Mae Fah Luang University Culture Collection, Chiang Rai, Thailand; **NBRC**: NITE Biological Resource Centre, Japan; Culture and specimen abbreviations: **JK**: J. Kohlmeier; **T**: ex-type/ex-epitype isolates; **Ts**: type species.

Results

Phylogeny

Combined analyses of LSU, SSU and RPB2 sequence data were used to determine the taxonomic placement of our strain. The dataset comprised 45 taxa with *Melanomma pulvis-pyrius* (CBS 124080) as the outgroup taxon and the manually adjusted dataset comprised 2998 characters including gaps. Phylogenetic analyses obtained from maximum likelihood (RAxML), maximum parsimony (MP) and Bayesian analyses showed similar topologies and were not significantly different. The best scoring RAxML tree was selected to represent the relationships among taxa and is shown in Figure 1.

The phylogenetic analyses obtained from maximum likelihood (RAxML), maximum parsimony (MP) and Bayesian analyses gave similar results for related families in the suborder Massarinae (Pleosporales) as in previous studies (Hyde *et al.* 2013, Singtripop *et al.* 2015, Wanasinghe *et al.* 2014, Wijayawardene *et al.* 2014, Phookamsak *et al.* 2015). *Poaceasca aquaticum* clusters with *P. helicoides* with high support values (99% MP, 99% ML and 1.00 PP), basal to *Stagonospora macropycnidia* (CBS 114202), *Setoseptoria phragmitis* which clustered with *Lentithecium arundinaceum* (95% MP, 97% ML and 1.00 PP) and may be the asexual morph of *Lentithecium*.




FIGURE 1. Phylogram generated from maximum likelihood analysis (RAxML) based on combined LSU, SSU and RPB2 sequenced data of the families in the suborder Massarinae. Bootstrap support values for maximum likelihood (ML, red) and maximum parsimony (MP, green) equal to or greater than 70% are given above the nodes. The values of the Bayesian posterior probabilities from MCMC analyses (BYPP, blue) equal or higher than 95% are given below the nodes. The tree is rooted to *Melanomma pulvis-pyrius* (CBS 124080). Newly generated sequences are indicated in red-gray.

Taxonomy

Poaceascoma aquaticum Z.L. Luo & K.D. Hyde, *sp. nov.* **FIGURE 2.**

Index Fungorum: IF551351

Facesoffungi number: FoF00910

Etymology: With reference to the habitat of this taxon.

Holotype: MFLU 15–0075




FIGURE 2. *Poaceascoma aquaticum* (MFLU 15–0075, holotype). **a, b.** Ascomata on submerged wood. **c.** Pseudoparaphyses. **d.** Sections through ascomata. **e.** Section of peridium. **f–i.** Asci. **j.** Ascospores. **k.** Germinating ascospore. **l.** Colony on MEA. Scale bars: b = 500 μ m, c = 20 μ m, d, f–j = 100 μ m, e = 50 μ m, k = 20 μ m, l = 30 mm.

Saprobic on decaying bamboo submerged in freshwater. **Sexual morph:** *Ascomata* 280–380 µm high, 310–500 µm diam, scattered, solitary, semi-immersed or erumpent through host tissue, uniloculate, globose to subglobose, glabrous, black, papilla, visible as raised, dark spots on host surface, ostiole central, with periphyses. *Peridium* 23–40 µm wide, of equal thickness, composed of several layers of pseudoparenchymatous cells, outer layer comprising several layers of pale black cells, arranged in a *textura angularis*, inner layer comprising several layers of hyaline, flattened cells, arranged in a *textura prismatica*. *Hamathecium* composed of numerous, 2–4.5 µm wide, filamentous, broad, cellular pseudoparaphyses, with distinct septa, embedded in a mucilaginous matrix. *Asci* 184.5–210.5 × 12–16 µm (\bar{x} = 197.5 × 14 µm, n = 20), 8-spored, bitunicate, fissionate, cylindrical to cylindrical-clavate, pedicellate, apically rounded with an ocular chamber. *Ascospores* 214–231 × 4.2–5.2 µm (\bar{x} = 223 × 4.7 µm, n = 20), fasciculate, arranged spirally in ascus, filiform, widest at the rounded apex and tapering towards the rounded base, pale brown to brown, 21–36-septate, not constricted at the septa, smooth-walled. **Asexual morph:** Undetermined.

Material examined:—THAILAND. Chiang Rai Province: (N 20°3'39", E 99°52'29"), saprobic on decaying bamboo submerged in a stream, November 2013, Z.L. Luo ZL-26 (MFLU 15–0075!, **holotype**); ex-type culture, MFLUCC 14-0048, DLUCC; *ibid.* (HKAS 86450, **isotype**).

Notes:—The scolecosporous *Poaceascoma* species was collected from decaying culms of bamboo submerged in a freshwater stream in northern Thailand. There are many genera forming filiform ascospores in Pleosporales, such as *Leptospora*, *Ophiobolus* and *Ophiosphaerella* (Shoemaker 1976, Boonmee *et al.* 2011, Zhang *et al.* 2012, Wijayawardene *et al.* 2014, Hyde *et al.* 2013, Ariyawansa *et al.* 2014, 2015). Phookamsak *et al.* (2015) introduced a new genus *Poaceascoma* in Lentitheciaceae, Pleosporales to accommodate the species associated with Poaceae which formed setose ascoma and filiform ascospores. *Poaceascoma aquaticum* resembles *P. helicoides* in having uni-loculate, immersed ascomata, a peridium of *textura angularis* to *textura prismatica* and filiform, multi-septate ascospores arranged spirally in the ascus. However, *P. aquaticum* differs from *P. helicoides* in having a thinner, pale black to hyaline peridium without setae, wider and longer asci (\bar{x} = 197.5 × 14 µm, versus 173.5 × 9.4 µm) and ascospores (\bar{x} = 223 × 4.7 µm, versus 178.1 × 2.4 µm). The phylogenetic data also confirmed them as distinct taxa (Figure 1).

Discussion

Zhang *et al.* (2012) introduced the family Lentitheciaceae in the order Pleosporales to accommodate massarina-like species, typified by *Lentithecium* with *L. fluviatile* as the type species. Species of the Lentitheciaceae occur on herbaceous plants such as *Phragmites* (*Lentithecium fluviatile* (Aptroot & Van Ryck.) K.D. Hyde *et al.*, *L. arundinaceum* (Sowerby) K.D. Hyde *et al.*, *Tingoldiagio graminicola* K. Hirayama & Kaz. Tanaka) and on submerged wood (*Lentithecium aquaticum* Yin. Zhang *et al.*) in freshwater environments (Zhang *et al.* 2012). *Katumotoa bambusicola* Kaz. Tanaka & Y. Harada and *Ophiosphaerella sasicola* (Nagas. & Y. Otani) Shoemaker & C.E. Babc. are bambusicolous species known from Japan (Nagasawa & Otani 1977; Tanaka & Harada 2005). Hyde *et al.* (2013) included *Katumotoa*, *Keissleriella*, *Lentithecium*, *Setoseptoria* and *Tingoldiagio* in Lentitheciaceae based on multigene phylogenetic analyses. Phookamsak *et al.* (2015) introduced a new genus *Poaceascoma* to accommodate the ophiosphaeriella-like taxa in Lentitheciaceae and mentioned that the scolecosporous fungi were polyphyletic in the class Dothideomycetes. Presently, *Katumotoa*, *Keissleriella*, *Lentithecium*, *Setoseptoria*, *Tingoldiagio* and *Poaceascoma* are accepted in the family (Hyde *et al.* 2013, Liu *et al.* 2015b, Phookamsak *et al.* 2015). We included all of these genera of Lentitheciaceae in the phylogenetic tree (Figure. 1) to determine the placement of our taxon. Several clades in Lentitheciaceae were not well-resolved, which may be due to limited sequence data in GenBank, however the genus *Poaceascoma* is strongly supported.

Acknowledgements

We would like to thank the National Natural Science Foundation of China (Project ID: 31460015, 31360013, 31160160) and middle younger academic leaders of candidate's projects in Yunnan Province (2012HB-042) for financial and laboratory support. The authors extend their sincere appreciations to the Deanship of Scientific Research at King Saud University for its funding this Prolific Research Group (PRG-1436-09). We thank the National Research Council of Thailand (grant for Dothideomycetes No: 58201020010) for supporting studies on Dothideomycetes. Zonglong Luo thanks Shaun Pennycook (Landcare Research, New Zealand) to help in checking the taxon name. We thank Shu-

Hong Li (Institute of Biotechnology and Gerplamic Resources, Yunnan Academy of Agricultural Sciences, China) for assistance in molecular work.

References

- Ariyawansa, H.A., David, L.H., Hyde, K.D., Jones, E.B.G., Maharachchikumbura, S.S.N., Manamgoda, D.S., Thambugala, K.M., Udayanga D., Camporesi, E., Daranagama, A., Jayawardena, R., Liu, J.K., McKenzie, E.H.C., Phookamsak, R., Senanayake, I.C., Shivas, R.G., Tian, Q. & Xu, J.-C. (2014) Epitypification and neotypification: guidelines with appropriate and inappropriate examples. *Fungal Diversity* 69: 57–91.
<http://dx.doi.org/10.1007/s13225-014-0315-4>
- Ariyawansa, H.A., Thambugala, K.M., Manamgoda, D.S., Jayawardena, R., Camporesi, E., Boonmee, S., Wanasinghe, D.N., Phookamsak, R., Hongsanan, S., Singtripop, C., Chukeatirote, E., Kang, J.C., Jones, E.B.G. & Hyde, K.D. (2015) Towards a natural classification and backbone tree for *Pleosporaceae*. *Fungal Diversity* 71: 85–139.
<http://dx.doi.org/10.1007/s13225-015-0323-z>
- Boonmee, S., Zhang, Y., Chomnunti, P., Chukeatirote, E., Tsui, C.K., Bahkali, A.H. & Hyde, K.D. (2011) Revision of lignicolous Tubeufiaceae based on morphological reexamination and phylogenetic analysis. *Fungal Diversity* 51(1): 63–102.
<http://dx.doi.org/10.1007/s13225-011-0147-4>
- Cai, L., Jeewon, R. & Hyde, K.D. (2006) Phylogenetic investigations of Sordariaceae based on multiple gene sequences and morphology. *Mycological Research* 110:137–150.
<http://dx.doi.org/10.1016/j.mycres.2005.09.014>
- Cai, L., Zhang, K.Q. & Hyde, K.D. (2003a) Freshwater ascomycetes. In: Tsui CKM, Hyde KD (eds) *Freshwater Mycology. Fungal Diversity Research Series* 10: 275–324.
- Cai, L., Zhang, K.Q., McKenzie, E.H.C. & Hyde, K.D. (2003b) Freshwater fungi from bamboo and wood submerged in the Liput River in the Philippines. *Fungal Diversity* 13:1–12.
- Chomnunti, P., Hongsanan, S., Aguirre-Hudson, B., Tian, Q., Peršoh, D., Dhimi, M.K., Alias, A.S., Xu, J.C., Liu, X.Z., Stadler, M. & Hyde, K.D. (2014) The sooty moulds. *Fungal Diversity* 66(1):1–36.
<http://dx.doi.org/10.1007/s13225-014-0278-5>
- Goh, T.K. & Tsui, C.K.M. (2003) Key to common dematiaceous hyphomycetes from freshwater. In: Hyde KD, Tsui CKM (eds) *Freshwater Mycology. Fungal Diversity Research Series* 10: 325–343.
- Gulis, V., Kuehn, K.A. & Suberkropp, K. (2006) *The role of fungi in carbon and nitrogen cycles in freshwater ecosystems*. Fungi in Biogeochemical Cycles. Cambridge University Press, Cambridge.
- Hall, T. (2001) BioEdit version 5.0. 6. North Carolina State University, Department of Microbiology, Raleigh, North Carolina 192.
- Hillis, D.M. & Bull, J.J. (1993) An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* 42: 182.
<http://dx.doi.org/10.1093/sysbio/42.2.182>
- Ho, W.H., Hyde, K.D. & Hodgkiss, I.J. (2001) Fungal communities on submerged wood from streams in Brunei, Hong Kong and Malaysia. *Mycological Research* 105: 1492–1501.
<http://dx.doi.org/10.1017/S095375620100507X>
- Ho, W.H., Yanna, Hyde, K.D. & Hodgkiss, I.J. (2002) Seasonality and sequential occurrence of fungi on woodsubmerged in Tai Po Kau Forest Stream, Hong Kong. *Fungal Diversity* 10: 21–43. Available from: http://www.fungaldiversity.org/fdp/sfdp/FD_10_21-43.pdf
- Hu, D.M., Cai, L., Chen, H., Bahkali, A.H. & Hyde, K.D. (2010) Four new freshwater fungi associated with submerged wood from Southwest Asia. *Sydowia* 62: 191–203.
- Hyde, K.D. (1995) Tropical Australia freshwater fungi. VII. New genera and species of ascomycetes. *Nova Hedwigia* 61: 119–140.
- Hyde, K.D., Jones, E.B.G., Liu, J.K., Ariyawansa, H., Boehm, E., Boonmee, S., Braun, U., Chomnunti, P., Crous, P.W., Dai, D.Q., Diederich, P., Dissanayake, A., Doilom, M., Doveri, F., Hongsanan, S., Jayawardena, R., Lawrey, J.D., Li, Y.M., Liu, Y.X., Lücking, R., Monkai, J., Muggia, L., Nelsen, M.P., Pang, K.L., Phookamsak, R., Senanayake, I.C., Shearer, C.A., Suetrong, S., Tanaka, K., Thambugala, K.M., Wijayawardene, N.N., Wikee, S., Wu, H.X., Zhang, Y., Aguirre-Hudson, B., Alias, S.A., Aptroot, A., Bahkali, A., Bezerra, J.L., Bhat, D.J., Camporesi, E., Chukeatirote, E., Gueidan, C., Hawksworth, D.L., Hirayama, K., Hoog, S.D., Kang, J.C., Knudsen, K., Li, W.J., Li, X.H., Liu, Z.Y., Mapook, A., Mckenzie, E.H.C., Miller, A.N., Mortimer, P.E., Phillips, A.J.L., Raja, H.A., Scheuer, C., Schumm, F., Taylor, J.E., Tian, Q., Tibpromma, S., Wanasinghe, D.N., Wang, Y., Xu, J.C., Yacharoen, S., Yan, J.Y. & Zhang, M. (2013) Families of Dothideomycetes. *Fungal Diversity* 63: 1–313.

<http://dx.doi.org/10.1007/s13225-013-0263-4>

- Hyde, K.D., Fryar, S., Tian, Q., Bahkali, A.H. and Xu, J.C. (2015) Lignicolous freshwater fungi along a north-south latitudinal gradient in the Asian/Australian region; can we predict the affects of global warming on biodiversity and function? *Fungal Ecology* 19: 190–200.
- <http://dx.doi.org/10.1016/j.funeco.2015.07.002>
- Jayasiri, S.C., Hyde, K.D., Ariyawansa, H.A., Bhat, J., Buyck, B., Cai L., Dai Y.C., Abd-Elsalam, K.A., Ertz, D., Hidayat, I., Jeewon, R., Jones, E.B.G., Bahkali, A.H., Karunarathna, S.C., Liu, J.K., Luangsa-ard, J.J., Lumbsch, H.T., Maharachchikumbura, S.S.N., McKenzie, E.H.C., Moncalvo, J.M., Ghobad-Nejhad, M., Nilsson, H., Pang, K.L., Pereira, O.L., Phillips, A.J.L., Raspé, O., Rollins, A.W., Romero, A.I., Etayo, J., Selçuk, F., Stephenson, S.L., Suetrong, S., Taylor, J.E., Tsui, C.K.M., Vizzini, A., Abdel-Wahab, M.A., Wen, T.C., Boonmee, S., Dai, D.Q., Daranagama, D.A., Dissanayake, A.J., Ekanayaka, A.H., Fryar, S.C., Hongsanan, S., Jayawardena, R.S., Li, W.J., Perera, R.H., Phookamsak, R., Silva, N.I., Thambugala, K.M., Tian, Q., Wijayawardene, N.N., Zhao, R.L., Zhao, Q., Kang, J.C., & Promputtha, I. (2015). The Faces of Fungi database: fungal names linked with morphology, phylogeny and human impacts. *Fungal diversity* 74 (1): 3–18.
- <http://dx.doi.org/10.1007/s13225-015-0351-8>
- Jones, E.B.G. & Choeyklin, R. (2008) Ecology of marine and freshwater basidiomycetes. In: Boddy L, FranklandJC, van West P (eds) Ecology of saprotrophic basidiomycetes. *Academic Press*: 301–324.
- [http://dx.doi.org/10.1016/S0275-0287\(08\)80018-5](http://dx.doi.org/10.1016/S0275-0287(08)80018-5)
- Jones, E.B.G., Alias, S.A. & Nawawi, A. (2007) New fungi described from Malaysia. In: Jones, E.B.G., Hyde, K.D. & Vikineswary, S. (Eds.) *Malaysian fungal diversity*. pp. 377–419.
- Katoh, K. & Standley, D.M. (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30: 772–780.
- <http://dx.doi.org/10.1093/molbev/mst010>
- Kirk, P.M., Cannon, P.F., Minter, D.W. & Stalpers, J.A. (2008) *Ainsworth & Bisby's dictionary of the fungi, 10th edn*. CABI, Wallingford
- Krauss, G.J., Sole, M., Krauss, G., Schlosser, D., Wesenberg, D. & Baerlocher, F. (2011) Fungi in freshwaters: ecology, physiology and biochemical potential. *FEMS Microbiology Reviews* 35: 620–651.
- <http://dx.doi.org/10.1111/j.1574-6976.2011.00266.x>
- Kurniawati, E., Zhang, H., Chukeatirote, E., Sulistyowati, L., Moslem, M.A. & Hyde, K.D. (2010) Diversity of freshwater ascomycetes in freshwater bodies at Amphoe Mae Chan, Chiang Rai. *Cryptogamie Mycologie* 31: 323–331.
- Kuthubutheen, A.J. (1987) A new species of Phalangiospora and further observations on *P. constricta* from Malaysia. *Transactions of the British Mycological Society* 89: 414–420.
- [http://dx.doi.org/10.1016/S0007-1536\(87\)80135-3](http://dx.doi.org/10.1016/S0007-1536(87)80135-3)
- Liu, J.K., Phookamsak, R., Doilom, M., Wikee, S., Li, Y.M., Ariyawansa, H., Boonmee, S., Chomnunti, P., Dai, D.Q., Bhat, J.D., Romero, A.I., Zhuang, W.Y., Monkai, J., Jones, E.B.G., Chukeatirote, E., KO-KO, W.T., Zhao, Y.C., Wang, Y. & Hyde, K.D. (2012) Towards a natural classification of *Botryosphaerales*. *Fungal Diversity* 57: 149–210.
- <http://dx.doi.org/10.1007/s13225-012-0207-4>
- Liu, J.K., Hyde, K.D., Jones, E.B.G., Ariyawansa, H.A., Bhat, J.D., Boonmee, S., Maharachchikumbura, S.S.N., Mckenzie, E.H.C., Phookamsak, R., Phukhamsakda, C., Shenoy, B.D., Abdel-Wahab M.A., Buyck, B., Chen J., Chethana, W.T., Singtripop, C., Dai, D.Q., Dai, Y.C., Daranagama, D.A., Dissanayake, A.J., Doilom, M., D'Souza, M.J., Fan, X.L., Goonasekara, I.D., Hirayama, K., Hongsanan, S., Jayasiri, S.C., Jayawardena, R.S., Karunarathana, S.C., Li, W.J., Mapook, A., Norphanphoun, C., Pang, K.L., Perera, R.H., Perošoh, D., Pinruan, U., Senanayake, I.C., Somrithipol, S., Suethrong, S., Tanaka, K., Thambugala, K.M., Tian Q., Tibpromma, S., Udayanag, A.D., Wijayawardene, N.N., Wanasinghe, D., Wistrassameewong, K., Zeng, X.Y., Abdel-Aziz, F.A., Adamčík, S., Bahkali, A.H., Boonyuen, N., Bulgakov, T., Callac, P., Chomnunti, P., Greiner, K., Hashimoto, A., Hofstetter, V., Kang, J.C., Lewis, D., Li, X.L., Liu, X.X., Liu, Z.Y., Matsumura, M., Mortimer, P.E., Rambold, G., Randrianjohany, E., Sato, G., Srindrasutdhi, V., Tian C.M., Verbeke, A., Vonbrackel, W., Wang, Y., Wen, T.C., Xu, J.C., Yan, J.Y., Zhao, R.L. & Camporesi, E. (2015) Fungal diversity notes 1–110: taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity* 72 (1): 1–197.
- <http://dx.doi.org/10.1007/s13225-015-0324-y>
- Liu, X.Y., Udayanga, D., Luo, Z.L., Chen, L.J., Zhou, D.Q., Su, H.Y. & Hyde, K.D. (2015a) Backbone tree for *Chaetothyriales* with four new species of *Minimelanolocus* from aquatic habitats. *Fungal Biology* 119 (11): 1046–1062.
- <http://dx.doi.org/10.1016/j.funbio.2015.08.005>
- Lumbsch, H.T. & Huhndorf, S.M. (2010) Outline of Ascomycota–2009. *Fieldiana Life and Earth Sciences* 1: 1–60.
- <http://dx.doi.org/10.3158/1557.1>
- Luo, J., Yin, J.F., Cai, L., Zhang, K.Q. & Hyde, K.D. (2004) Freshwater fungi in Lake Dianchi, a heavily polluted lake in Yunnan, China. *Fungal Diversity* 16: 93–112.
- Luo, Z.L., Maharachchikumnura, S.S.N., Liu, X.Y., Li, S.H., Chen, L.J., Su, H.Y., Zhou, D.Q. & Hyde, K.D. (2015) *Annulatascus*

- saprophyticus* sp. nov. and *Pseudoannulatascus* gen. nov. to accommodate *Annulatascus biatriisporus* (Annulatascales Sordariomycetes) from Thailand. *Phytotaxa* 239 (2): 174–182.
<http://dx.doi.org/10.11646/phytotaxa.239.2.6>
- Nagasawa, E. & Otani, Y. (1977) Some species of the bambusicolous ascomycetes from Japan I. *Rep Tottori Mycol Instit* 15: 38–42.
- Nawawi, A. (1985) Basidiomycetes with branched water-borne conidia. *Botanical Journal of the Linnean Society* 91: 51–60.
<http://dx.doi.org/10.1111/j.1095-8339.1985.tb01134.x>
- Nylander, J.A.A. (2004) *MrModeltest* 2.0. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Phookamsak, R., Li, W.J., Dai, D.Q., Singtripop, C. & Hyde, K.D. (2015) *Poaceascoma* gen et sp. nov., a new genus with scolecospores in *Lentitheciaceae*. *Cryptogamie Mycologie* 36 (2): 225–236.
<http://dx.doi.org/10.7872/crym/v36.iss2.2015.225>
- Phookamsak, R., Liu, J.K., McKenzie, E.H.C., Manamgoda, D.S., Chatpapamon, C., Ariyawansa, H., Thambugala, K.M., Dai, D.Q., Camporesi, E., Chukeatiro, E., Wijayawardene, N.N., Mortimer, P.E., Xu, J.C. & Hyde, K.D. (2014) Revision of Phaeosphaeriaceae. *Fungal Diversity* 68: 159–238.
<http://dx.doi.org/10.1007/s13225-014-0308-3>
- Rannala, B. & Yang, Z. (1996) Probability distribution of molecular evolutionary trees: a new method of phylogenetic inference. *Journal of Molecular Evolution* 43: 304–31.
<http://dx.doi.org/10.1007/BF02338839>
- Ronquist, F. & Huelsenbeck, J.P. (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19 (12): 1572.
<http://dx.doi.org/10.1093/bioinformatics/btg180>
- Shearer, C.A. (1993) The freshwater ascomycetes. *Nova Hedwigia* 56: 1–33.
- Shearer, C.A., Anderson, J.L. & Pringle, C.M. (2001) Is there a pantropical freshwater ascomycota? Mycological society of America, annual meeting, Salt Lake City, Utah. *Phytopathology* 91: S123.
- Shoemaker, R.A. (1976) Canadian and some extralimital *Ophiobolus* species. *Canadian Journal of Botany* 54: 2365–2404.
<http://dx.doi.org/10.1139/b76-252>
- Silvestro, D. & Michalak, I. (2011). RAXML-GUI: a graphical front-end for RAXML. *Organisms Diversity and Evolution* 12: 335–337.
<http://dx.doi.org/10.1007/s13127-011-0056-0>
- Singtripop, C., Camporesi, E., Ariyawansa, H.A., Wanasinghe, D.N., Boonmee, S., Mortimer, P.E., Xu, J.C. & Hyde, K.D. (2015) *Keissleriella dactylidis*, sp. nov., from *Dactylis glomerata* and its phylogenetic placement. *Scienceasia* 41 (2015): 295–304.
- Sivichai, S. & Jones, E.B.G. (2004) *Stauriella* gen. nov., proposed for a new lignicolous basidiomycetous anamorph from freshwater in Thailand. *Sydowia* 56: 131–136.
- Sivichai, S., Jones, E.B.G. & Hywel-Jones, N.L. (2000) Fungal colonisation of wood in a freshwater stream at KhaoYai National Park, Thailand. *Fungal Diversity* 5: 71–88.
- Su, H.Y., Udayanga, D., Luo, Z.L., Manamgoda, D.S., Zhao, Y.C., Yang, J., Liu, X.Y., Mckenzie, E.H.C. & Hyde, K.D. (2015) Hyphomycetes from aquatic habitats in Southern China: Species of *Curvularia* (Pleosporaceae) and *Phragmocephala* (Melannomataceae). *Phytotaxa* 226 (3): 201–216.
<http://dx.doi.org/10.11646/phytotaxa.226.3.1>
- Su, H.Y., Luo, Z.L., Bhat, J.D., Bahkali, A.H., Mckenzie, E.H.C., Zhou, D.Q. & Hyde, K.D. (In press) *Helminthosporium velutinum* and *H. aquaticum* sp. nov. from aquatic habitats in Yunnan Province, China. *Phytotaxa*.
- Swofford D.L. (2002) PAUP: phylogenetic analysis using parsimony, version 4.0 b10. Sunderland, MA: Sinauer Associates.
- Tanaka, K. & Harada, Y. (2005) Bambusicolous fungi in Japan (6): *Katumotoa*, a new genus of phaeosphaeriaceous ascomycetes. *Mycoscience* 46: 313–318.
<http://dx.doi.org/10.1007/S10267-005-0251-Y>
- Taylor, J.E. & Hyde, K.D. (2003) Microfungi of tropical and temperate palms. *Fungal Diversity Research Series* 12: 1–459.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F. & Higgins, D.G. (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25: 4876–4882.
<http://dx.doi.org/10.1093/nar/25.24.4876>
- Vilgalys, R. & Hester, M. (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172 (8): 4238–4246.
- Wanasinghe, D.N., Jones, E.B.G., Camporesi, E., Boonmee, S., Ariyawansa, H.A., Wijayawardene, N.N. & Hyde, K.D. (2014) An Exciting Novel Member of Lentitheciaceae in Italy from *Clematis Vitalba*. *Cryptogamie Mycologie* 35 (4): 323–337.
<http://dx.doi.org/10.7872/crym.v35.iss4.2014.323>
- Wijayawardene, N.N., Crous, P.W., Kirk, P.M., Hawksworth, D.L., Boonmee, S., Braun, U., Dai, D.Q., D'souza, M.J., Diederich, P., Dissanayake, A., Doilom, M., Hongsanan, S., Jones, E.B.G., Groenewald, J.Z., Jayawardena, R., Lawrey, J.D., Liu, J.K., Lücking,

- R., Madrid, H., Manamgoda, D.S., Muggia, L., Nelsen, M.P., Phookamsak, R., Suetrong, S., Tanaka, K., Thambugala, K.M., Wanasinghe, D.N., Wikee, S., Zhang, Y., Aptroot, A., Ariyawansa, H.A., Bahkali, A.L., Bhat, D.J., Gueidan, C., Chomnunti, P., Hoog, G.S.D., Knudsen, K., Li, W.J., McKenzie, E.H.C., Miller, A.N., Phillips, A.J.L., Piątek, M., Raja, H.A., Shivas, R.S., Slippers, B., Taylor, J.E., Tian, Q., Wang, Y., Woudenberg, J.H.C., Cai, L., Jaklitsch, W.M. & Hyde, K.D. (2014) Naming and outline of Dothideomycetes–2014 including proposals for the protection or suppression of generic names. *Fungal Diversity* 69: 1–55.
<http://dx.doi.org/10.1007/s13225-014-0309-2>
- Wong, M.K.M., Goh, T.K., Hodgkiss, I.J., Hyde, K.D., Ranghoo, V.M., Tsui, C.K.M., Ho, W.H., Wong, W.S.W. & Yuen, T.K. (1998) Role of fungi in freshwater ecosystems. *Biodiversity and Conservation* 7: 1187–1206.
<http://dx.doi.org/10.1023/A:1008883716975>
- Wong, S.W., Hyde, K.D. & Jones, E.B.G. (1999) Ultrastructure studies on freshwater ascomycetes, *Fluminicola bipolaris* gen. et sp. nov. *Fungal Diversity* 2: 189–197.
- Zhang, Y., Crous, P.W., Schoch, C.L. & Hyde, K.D. (2012) Pleosporales. *Fungal Diversity* 53: 1–221.
<http://dx.doi.org/10.1007/s13225-011-0117-x>