

## Polyacetylene Glycosides from *Gymnaster koraiensis*

JinYoung PARK,<sup>a</sup> ByungSun MIN,<sup>b</sup> HyunJu JUNG,<sup>a</sup> YoungHo KIM,<sup>a</sup> HyeongKyu LEE,<sup>b</sup> and KiHwan BAE<sup>\*,a</sup>

<sup>a</sup> College of Pharmacy, Chungnam National University; Taejon 305–764, Korea; and <sup>b</sup> Immunomodulator Research Laboratory, Korea Research Institute of Bioscience & Biotechnology; P.O. Box 115, Yusong, Taejon 305–333, Korea.

Received December 13, 2001; accepted February 5, 2002

Two polyacetylene glycosides, gymnasterkoreasides A and B, were isolated from the roots of *Gymnaster koraiensis*. Their structures were elucidated to be (3*R*)-8-decene-4,6-diyne-1,3-diol 1-*O*- $\beta$ -D-glucopyraside and (3*R*)-8-decene-4,6-diyne-1,3-diol 1-*O*- $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyraside on the basis of spectroscopic analysis including COSY, HMQC, and HMBC experiments, as well as chemical methods, which confirmed the determination of a chiral center by the modified Mosher's method.

**Key words** *Gymnaster koraiensis*; polyacetylene glycoside; modified Mosher's method

*Gymnaster koraiensis* (NAKAI) KITAMURA (Compositae) is an endemic species in Korea. Earlier and more recently, we isolated eight polyacetylenes from a CH<sub>2</sub>Cl<sub>2</sub>-soluble fraction of the roots of this plant, which showed significant cytotoxicity against L1210 tumor cells with ED<sub>50</sub> values of 0.12–3.28  $\mu$ g/ml.<sup>1</sup> As part of our continuing research to find pharmacologically active constituents from *G. koraiensis*, we have isolated two new polyacetylene glycosides, called gymnasterkoreasides A (**1**) and B (**2**), from a BuOH-soluble fraction of the roots of this plant. This paper describes the isolation and structure elucidation of two new C<sub>10</sub>-acetylenic glycosides isolated from *G. koraiensis*.

### Results and Discussion

The BuOH-soluble fraction of 80% ethanol extract of the roots of *G. koraiensis* was subjected to repeated column chromatography on silica gel, Sephadex LH-20, and preparative HPLC to give two compounds (**1**, **2**) (Chart 1).

Gymnasterkoreaside A (**1**), bright yellow oil, has the molecular formula C<sub>16</sub>H<sub>22</sub>O<sub>7</sub> based on high-resolution fast atom bombardment mass spectrometry (HR-FAB-MS) with a molecular ion peak at *m/z* 349.1260 [M+Na]<sup>+</sup>. The UV spectrum showed typical absorptions at  $\lambda_{\max}$  239, 252, 266, and 281 nm for an ene-diyne chromophore.<sup>2</sup> The IR spectrum of **1** showed the presence of a conjugated triple bond (2245 cm<sup>-1</sup>), a conjugated double bond (1660 cm<sup>-1</sup>), and one or more hydroxyl groups (3350 cm<sup>-1</sup>). The proton nuclear magnetic resonance (<sup>1</sup>H-NMR) spectrum showed signals for a methyl at  $\delta$  1.80 (dd, *J*=6.8, 1.8 Hz), an oxymethylene at  $\delta$  4.64 (t, *J*=6.7 Hz), an anomeric proton at  $\delta$  4.26 (d, *J*=7.8 Hz), and two olefinic protons at  $\delta$  6.31 (dq, *J*=15.8, 6.8 Hz) and 5.57 (dq, *J*=15.8, 1.8 Hz) (Table 1). The <sup>13</sup>C-NMR measurements, aided by distortionless enhancement by polarization transfer (DEPT) and <sup>1</sup>H-detected multiple quantum coherence (HMQC) spectra, revealed the presence of one methyl at  $\delta$  18.9, two olefinic carbons at  $\delta$  145.1 and 110.5, four quaternary carbons at  $\delta$  83.7, 78.1, 72.4, and 69.7, and an anomeric carbon at  $\delta$  104.5.

Analysis of the <sup>1</sup>H–<sup>1</sup>H shift correlation spectroscopy (COSY), HMQC, and heteronuclear multiple-bond correlations (HMBC) spectra of **1** allowed its structural fragments to be determined. In the COSY spectrum, two terminal spin systems began with a methyl at  $\delta$  1.80 and hydroxymethylene at  $\delta$  3.73 and 3.98, respectively. The former proton cou-

pled to an olefinic proton ( $\delta$  6.31) that was further coupled to an other olefinic proton ( $\delta$  5.57). The latter protons correlated to methylene protons ( $\delta$  1.96) and an oxymethylene ( $\delta$  4.64), indicating the end of the spin system (Chart 1). In addition, partial structures were linked in the HMBC experiments. Long-range correlations among  $\delta_{\text{H}}$  4.64 (H-3) and  $\delta_{\text{C}}$  83.7 (C-4)/69.7 (C-5),  $\delta_{\text{H}}$  5.57 (H-8) and  $\delta_{\text{C}}$  72.4 (C-6), and  $\delta_{\text{H}}$  6.31 (H-9) and  $\delta_{\text{C}}$  78.1 (C-7) confirmed that a conjugated diyne was connected to C-3 and C-8. The C-8,9 double bond was found to be *E*, as evidenced by their vicinal coupling constant (*J*<sub>8,9</sub>=15.8 Hz). From these data, the aglycone was determined to be 8-decene-4,6-diyne-1,3-diol.

The presence of glucose in **1** was confirmed by the signals of anomeric proton at  $\delta$  4.26 in the <sup>1</sup>H-NMR spectrum and six carbon signals at  $\delta$  104.54, 75.06, 78.09, 71.56, 77.90, and 62.70 in the <sup>13</sup>C-NMR spectrum.<sup>3</sup> The anomeric configuration of the glucose moiety was determined to be  $\beta$  on the basis of the *J*<sub>H-H</sub> value (7.8 Hz) of the anomeric proton in the <sup>1</sup>H-NMR spectrum. Acid hydrolysis of **1** with 4N HCl–diox-




Chart 1. Structures of Compounds Isolated from the Roots of *G. koraiensis*

\* To whom correspondence should be addressed. e-mail: baekh@cnu.ac.kr




Fig. 1. Chemical Shift Difference for the (*S*)-MTPA Ester (**1c**) and (*R*)-MTPA Ester (**1b**) in ppm

ane produced a sugar component, which was determined to be *D*-glucose by GLC of its pertrimethylsilated *L*-cysteine methyl ester derivative.<sup>4</sup> The linkage of glucose was determined by HMBC, which showed a correlation between the signals at  $\delta_{\text{H}}$  3.73/3.98 (H<sub>2</sub>-1 of the aglycone) and  $\delta_{\text{C}}$  104.54 (Glc-C-1'), indicating glycosylation at C-1.

Determination of the absolute configuration at C-3 of **1** was examined with the modified Mosher's method.<sup>5-7</sup> Compound **1a**, obtained by enzyme hydrolysis of **1**, was treated with (+)- and (-)- $\alpha$ -methoxy- $\alpha$ -trifluoromethylphenylacetate (MTPA) chlorides in the presence of 4-dimethylaminopyridine (DMPA) to give (*R*)- and (*S*)-MTPA esters (**1b, c**). In the <sup>1</sup>H-NMR spectrum of (*S*)-MTPA ester (**1c**), proton signals assigned to H<sub>2</sub>-1 and H<sub>2</sub>-2 were observed at a lower field than those in the (*R*)-MTPA ester (**1b**), while signals due to H-8 and H-9 in the former ester were shifted to a higher field than those in the latter ester (Fig. 1). Therefore the absolute configuration at C-3 was concluded to be *3R*. On the basis of the above findings, gymnasterkoreaside A (**1**) was determined to be (*3R*)-8-decene-4,6-diyne-1,3-diol 1-*O*- $\beta$ -*D*-glucopyranoside.

Gymnasterkoreaside B (**2**) was isolated as optically active oil,  $[\alpha]_{\text{D}} -78^{\circ}$ , and its molecular formula was established to be C<sub>21</sub>H<sub>30</sub>O<sub>11</sub> by HR-FAB-MS. On acid hydrolysis, **2** yielded glucose and apiose. Its UV spectrum was similar to that of **1**, indicating the presence of a conjugated diyne. The <sup>1</sup>H- and <sup>13</sup>C-NMR spectra were similar to those of **1**, except for the signals of sugar components. In its <sup>1</sup>H-NMR spectrum, **2** exhibited signals for two anomeric protons at  $\delta$  4.25 (d, *J*=7.7 Hz) and 4.99 (d, *J*=2.3 Hz). Detailed analysis of the 2D-NMR (COSY, HMQC, and HMBC) revealed that its aglycone was the same as that of **1**. Long-range correlation between the anomeric proton at  $\delta$  4.99 and the carbon signal at  $\delta$  68.49 (C-6') of the glucose observed in the HMBC spectrum suggested the second sugar moiety to be an ether-linked one at C-6'. This was confirmed by comparing the <sup>13</sup>C-NMR spectrum of **2** with that of **1**, in which a downfield shift of the C-6' (5.8 ppm) signal due to glucose moiety was observed. The ether-linked sugar was determined to be *D*-apiose by acid hydrolysis of **2** followed by GLC analysis of its sugar derivative (see Experimental). By comparing the NMR spectra of **2** with those of **1** and analysis of its 2D-NMR, the <sup>1</sup>H- and <sup>13</sup>C-NMR signals were assigned as shown in Table 1. The structure of **2** therefore was confirmed to be (*3R*)-8-decene-4,6-diyne-1,3-diol 1-*O*- $\beta$ -*D*-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -*D*-glucopyranoside.

#### Experimental

Optical rotations were measured with a JASCO DIP-370 Digital polarimeter. UV spectra were recorded on a Milton Roy Spectronic 3000 spectrophotometer. IR spectra were determined on an IR Report-100 spectropho-

Table 1. <sup>1</sup>H- and <sup>13</sup>C-NMR Spectral Data of **1** and **2** in MeOH-*d*<sub>4</sub>

	<b>1</b>		<b>2</b>	
	<sup>1</sup> H <sup>a</sup>	<sup>13</sup> C <sup>b</sup>	<sup>1</sup> H <sup>a</sup>	<sup>13</sup> C <sup>b</sup>
1a	3.73 (dd, 10.1, 6.3)	66.8	3.73 (dd, 6.6, 4.0)	67.0
1b	3.98 (dd, 10.1, 6.3)		3.95 (dd, 6.6, 4.0)	
2	1.96 (m)	38.9	1.96 (m)	39.0
3	4.64 (t, 6.7)	60.2	4.64 (t, 6.7)	60.2
4		83.7		83.7
5		69.7		69.7
6		72.4		72.0
7		78.1		78.1
8	5.57 (qd, 15.8, 1.8)	110.5	5.58 (qd, 15.8, 1.7)	110.5
9	6.31 (qd, 15.8, 6.8)	145.1	6.33 (qd, 15.8, 6.9)	145.1
10	1.80 (dd, 6.8, 1.8)	18.9	1.81 (dd, 6.9, 1.7)	18.8
glc-1'	4.26 (d, 7.8)	104.5	4.25 (d, 7.7)	104.6
-2'	3.16 (dd, 8.9, 7.8)	75.1	3.16 (dd, 8.9, 7.9)	75.1
-3'	3.34 (t, 8.9)	78.9	3.32 (t, 8.9)	78.0
-4'	3.29 (t, 8.9)	71.6	3.28 (t, 8.9)	71.6
-5'	3.26 (m)	77.9	3.39 (m)	76.8
-6'a	3.67 (dd, 12.4, 5.8)	62.7	3.62 (dd, 11.6, 5.9)	68.5
-6'b	3.98 (dd, 12.4, 5.8)		3.98 (dd, 11.6, 5.8)	
api-1''			4.99 (d, 2.3)	111.0
-2''			3.90 (d, 2.3)	78.0
-3''				80.6
-4''a			3.76 (d, 9.7)	75.0
-4''b			3.96 (d, 9.7)	
-5''			3.60 (s)	65.7

a) 300 MHz. b) 75 MHz.

tometer (JASCO). FAB-MS spectra were measured on an Autospec Mass spectrometer (Micromass). NMR spectra were recorded on a Bruker NMR DRX300, 600 spectrometer, with the chemical shift being represented in parts per million with tetramethylsilane as an internal standard. Preparative HPLC was performed on Shimadzu LC-10AD pump, CTO-10A column oven, and SPD-10AV UV-detector. GC-MS was carried out on a Shimadzu GC-17A and JEOL Automass system II mass detector.

**Plants Materials** Roots of *G. koraiensis* (NAKAI) KITAMURA were collected in May 1997 at Gurae, Chunnam province, Korea. A voucher specimen (CNU96003) is deposited in the herbarium of the College of Pharmacy, Chungnam National University, Taejeon, Korea.

**Extraction and Isolation** The dried root (4.8 kg) was extracted with 80% aqueous EtOH (700 g). The EtOH extract was suspended in H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub> and BuOH successively to give the CH<sub>2</sub>Cl<sub>2</sub>-soluble fraction (140 g) and BuOH-soluble fraction (123 g). The BuOH-soluble fraction was chromatographed on a silica gel column with a stepwise gradient of CHCl<sub>3</sub> and MeOH as eluent to give eight fractions (Fr. 1-8). Repeated column chromatography of Fr. 6 on silica gel (CHCl<sub>3</sub>/MeOH/H<sub>2</sub>O, 5:1:0:0.5) and RP-C<sub>18</sub> column chromatography (58% aq. MeOH) afforded **1** (160 mg). Column chromatography of Fr. 7 using silica gel (CHCl<sub>3</sub>/MeOH/H<sub>2</sub>O, 5:1:0:0.5) and Sephadex LH-20 (100% MeOH), followed by preparative HPLC on RP-C<sub>18</sub> (250 $\times$ 10 mm, 45% aq. MeOH, flow rate 1.5 ml/min) gave **2** (20 mg, retention time 17.9 min).

Compound **1**: Bright yellow oil.  $[\alpha]_{\text{D}}^{20} -28^{\circ}$  (*c*=1, MeOH); UV  $\lambda_{\text{max}}$  nm (MeOH, log  $\epsilon$ ): 239 (0.92), 252 (1.17), 266 (1.32), 281 (1.21); IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3350 (OH), 2930, 2245 (C $\equiv$ C), 1660 (C=C), 1080 (C-O); HR-FAB-MS *m/z*: 349.1260 [M+Na]<sup>+</sup> (Calcd for [C<sub>16</sub>H<sub>22</sub>O<sub>7</sub>+Na]<sup>+</sup> 349.1264); <sup>1</sup>H- and <sup>13</sup>C-NMR spectral data: see Table 1.

Compound **2**: Bright yellow oil.  $[\alpha]_{\text{D}}^{20} -78^{\circ}$  (*c*=1, MeOH); UV  $\lambda_{\text{max}}$  nm (MeOH, log  $\epsilon$ ): 239 (0.12), 252 (0.39), 266 (0.54), 281 (0.47); IR (KBr)  $\nu_{\text{max}}$  cm<sup>-1</sup>: 3380 (OH), 2345 (C $\equiv$ C), 1650 (C=C), 1060 (C-O); HR-FAB-MS *m/z*: 481.1682 [M+Na]<sup>+</sup> (Calcd for [C<sub>21</sub>H<sub>30</sub>O<sub>11</sub>+Na]<sup>+</sup> 481.1686); <sup>1</sup>H- and <sup>13</sup>C-NMR spectral data: see Table 1.

**Enzymatic Hydrolysis of 1** Naringinase (200 mg) was added to a suspension of **1** (30 mg) in 50 mM acetate buffer (pH 5.5) and the mixture was stirred at 37 $^{\circ}$ C for 5 h. The reaction mixture was extracted with EtOAc (10 ml $\times$ 3) and the organic layer was evaporated to dryness. The residue was chromatographed on silica gel eluted with hexane-acetone (3:1) to give 8-decene-4,6-diyne-1,3-diol (**1a**, 8 mg) as bright yellow oil. <sup>1</sup>H-NMR (CHCl<sub>3</sub>):  $\delta$  6.30 (1H, dq, *J*=15.8, 6.9 Hz, H-2), 5.50 (1H, dq, *J*=15.8, 1.7 Hz, H-3),

4.68 (1H, t,  $J=6.3$  Hz, H-8), 3.79 (2H, t,  $J=5.6$  Hz, H-10), 1.94 (2H, m, H-9), 1.77 (3H, dd,  $J=6.9, 1.7$  Hz, H-1).

**(R)-MTPA Ester of 1a** (+)-MTPA chloride (15 mg) and DMAP (10 mg) in pyridine (0.2 ml) was added to a solution of **1a** (2.0 mg) in  $\text{CCl}_4$  (0.2 ml). After stirring at room temperature for 12 h, the mixture was poured into water (10 ml) and extracted with  $\text{CHCl}_3$  (10 ml $\times$ 2). The  $\text{CHCl}_3$  extract was concentrated *in vacuo* and purified with preparative thin-layer chromatography (hexane/acetone, 5:1) to give an (R)-MTPA ester (**1b**, 1.5 mg) as a colorless oil.  $^1\text{H-NMR}$  ( $\text{CHCl}_3$ ):  $\delta$  6.38 (1H, m, H-9), 5.58 (1H, t,  $J=6.9$  Hz, H-3), 5.52 (1H, ddd,  $J=15.8, 1.8, 0.7$  Hz, H-8), 4.46 (1H, m, H-1a), 4.35 (1H, m, H-1b), 3.51 (3H, d,  $J=1.2$  Hz,  $\text{MTPA-OCH}_3$ ), 3.52 (3H, d,  $J=1.0$  Hz,  $\text{MTPA-OCH}_3$ ), 2.26 (2H, m, H-2), 1.84 (3H, dd,  $J=6.9, 1.8$  Hz, H-10).

**(S)-MTPA Ester of 1a** (-)-MTPA chloride (15 mg) and DMAP (10 mg) in pyridine (0.2 ml) was added to a solution of **1a** (2.0 mg) in  $\text{CCl}_4$  (0.2 ml). Work-up as described above gave an (S)-MTPA ester (**1c**, 1.5 mg) as a colorless oil.  $^1\text{H-NMR}$  ( $\text{CHCl}_3$ ):  $\delta$  6.39 (1H, m, H-9), 5.61 (1H, t,  $J=6.9$  Hz, H-3), 5.53 (1H, ddd,  $J=15.8, 1.8, 0.7$  Hz, H-8), 4.34 (1H, m, H-1a), 4.20 (1H, m, H-1b), 3.57 (3H, d,  $J=1.2$  Hz,  $\text{MTPA-OCH}_3$ ), 3.53 (3H, d,  $J=1.0$  Hz,  $\text{MTPA-OCH}_3$ ), 2.20 (2H, m, H-2), 1.84 (3H, dd,  $J=6.9, 1.8$  Hz, H-10).

**Determination of Sugars in 1 and 2** Each sample (2 mg) was refluxed with 4N HCl-dioxane (1:1, 2 ml) for 2 h. The mixture was extracted with EtOAc (5 ml $\times$ 3). The residual water layer was neutralized with Amberlite MB-3 and dried to give a residue. The residue was dissolved with pyridine (1 ml), to which 0.1M L-cysteine methyl ester hydrochloride in pyridine (2 ml) was added. The mixture was stored at 60 °C for 1.5 h. After the reac-

tion mixture was dried *in vacuo*, the residue was trimethylsilylated with hexamethyldisilazane-trimethylchlorosilane (0.1 ml) at 60 °C for 1 h. The mixture was partitioned between hexane and  $\text{H}_2\text{O}$  (0.3 ml each), and the hexane extract was analyzed by GC-MS. In the acid hydrolysate of **1** and **2**, D-glucose and D-apiose were confirmed by comparison of the retention times of their derivatives with those of D-glucose, L-glucose, and D-apiose derivatives prepared in a similar way, which showed retention times of 21.30, 22.00, and 17.30 min, respectively.

**Acknowledgments** This work has been partially supported by a grant from the KOSEF (R01-1999-00155) in Korea. We are grateful to KBSI for  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectral measurements.

#### References

- 1) Jung H., Min B., Park J., Kim Y., Lee H., Bae K., *J. Nat. Prod.*, in press (2002).
- 2) Tsukamoto S., Kato H., Hirota H., Fusetani N., *J. Nat. Prod.*, **60**, 126—130 (1997).
- 3) Wang C. Z., Yu D. Q., *Phytochemistry*, **48**, 711—717 (1998).
- 4) Min B. S., Nakamura N., Miyashiro H., Kim Y. H., Hattori M., *Chem. Pharm. Bull.*, **48**, 194—200 (2000).
- 5) Dale J. A., Mosher H. S., *J. Am. Chem. Soc.*, **95**, 512—519 (1973).
- 6) Ohitani I., Kusumi T., Kashman Y., Kakisawa H., *J. Am. Chem. Soc.*, **113**, 4092—4096 (1991).
- 7) Min B. S., Gao J. J., Nakamura N., Hattori M., *Chem. Pharm. Bull.*, **48**, 1026—1033 (2000).