

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

**Pakistan
Journal of Biological Sciences**

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

Polyculture of Carp, Tilapia and Pangas Using Low Cost Inputs

¹M.A.K. Azad, ¹M.R. Rahman, ¹Z. Rahman, ²M.A. Kader, ¹M.M. Haque and ¹M.J. Alam

¹Department of Fisheries Management,

²Department of Aquaculture, Faculty of Fisheries,
Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

Abstract: An experiment was conducted to evaluate the water quality, growth and production of fishes at different species compositions in polyculture using tilapia (*Oreochromis niloticus*), pangas (*Pangasius hypophthalmus*), mrigal (*Cirrhinus mrigala*) and silver carp (*Hypophthalmichthys molitrix*). The experiment was conducted for a period of 90 days from September 8, to December 7, 2002 at the Fisheries Faculty Field Laboratory of Bangladesh Agricultural University, Mymensingh. There were three treatments each with three replications. In all the treatments, the stocking density was 100 fish/decimal at different species compositions. All the ponds were subjected to same regime of feeding and fertilization. Fortnightly random sampling was done to adjust the feeding rate. In this experiment, mrigal showed the highest SGR (% per day) value (3.74) in T₂. Tilapia showed the second highest (3.65) SGR (% per day) in T₁. Specific growth rate (% per day) of pangas was comparatively lower in all treatments. Mean survival rates of various species ranged from 90.63 to 91.10%. Comparatively, tilapia showed the highest survival rate where pangas showed the lowest. T₁ showed the highest survival rate but there was no significant variation among the treatments. Weight gain (g) of tilapia, pangas, mrigal and silver carp were 89.18, 30.61, 85.75 and 91.71 g in T₁, 63.95, 37.24, 86.86 and 65.63 g in T₂ and 70.84, 47.65, 70.42 and 72.87 g in T₃, respectively. The highest and the lowest weight gain were found in T₁. Silver carp attained maximum weight gain (91.71 g) in T₁ where pangas showed the lowest (30.61 g) in the same treatment. Among the treatments, the highest production was found in T₁ (1974.02 kg/ha/90 days) which was significantly higher than other treatments. The production in T₂ and T₃ were 1405.59 and 1522.01 kg/ha/90 days but there was no significant variation between T₂ and T₃. Considering the above, pangas showed the lowest growth performance and production in all treatments compare with other species. From the findings of the present study, it can be said that polyculture of pangas with tilapia is not suitable, but it may be suitable with carps and for this further research is necessary.

Key words: Polyculture, carp, tilapia, pangas, low-cost feed

INTRODUCTION

Polyculture or composite culture is the system in which fast growing compatible species of different feeding habits are stocked in different proportions in the same ponds^[1]. The basic principles of the polyculture; species of different feeding habits are, culture in the same pond to avoid food competition and best utilization of natural foods of different habits without any harm to each other. It is a fact that, polyculture may produce an expected result if the fish with different feeding habits are stocked in proper ratios and combinations^[2].

Selection of species plays an important role for any cultural practices. The better utilization of different strata and zones of a pond three or more species must be

stocked. In our country, suitable or common combinations of fish for composite culture are rui (*Labeo rohita*), catla (*Catla catla*) and mrigal (*Cirrhinus mrigala*). Sometimes calbasu (*Labeo calbasu*) has been included in polyculture. In recent years, some exotic fish species draw an attention to farmers for their better growth performance. Fish culturist introduced several exotic carps in order to obtain more production at minimum cost and the shortest possible period. They are now being cultured combindly with the local carps. These include silver carp (*Hypophthalmichthys molitrix*), grass carp (*Ctenopharyngodon idella*), common carp (*Cyprinus carpio*), raj punti (*Puntius gonionotus*) etc. Among those, silver carp is the most popular species for it's fast growth, lucrative size, good taste and high market demand.

Corresponding Author: Mohammad Rafiqur Rahman, C/O Md. Noor Khan, Scientific Officer, Freshwater Station, Bangladesh Fisheries Research Institute, Mymensingh-2201, Bangladesh
Tel: 091 55695 E-mail: mrafiqu@yaho.com

Now-a-days, pangas (*Pangasius hypophthalmus*) and tilapia (*Oreochromis niloticus*) have also been introduced in fresh water ponds for polyculture. Once a time, *P. hypophthalmus* was also a popular table fish in our country and farmers were economically benefited from this industry. But in recent years, this fishery is being depleted partly because of increasing feed cost, decreasing feed quality, lack of proper management, unavailability of low cost supplementary feeds and socio-economic constraints. So, the production cost became high enough compared to market value and the farmers are discouraged from pangas culture day by day. For this purpose, it is urgent need to find out a suitable culture technique as well as polyculture technique of pangas with local carps to get the sustainable production using low cost feed and fertilizer.

Tilapia is an excellent fish for growing in the shallow and seasonal ponds in a country like Bangladesh^[3-5] because Bangladesh enjoys very suitable climate and ecological conditions for culture of warm-water species.

That is why, the present study was undertaken to evaluate the growth performance of fishes (tilapia, pangas, mrigal and silver carp) in polyculture at different species combinations and to determine the suitable species combinations of both Indian major carp and exotic fishes on the basis of their ecological relationships.

MATERIALS AND METHODS

Description of the study area and duration: The experiment was conducted for a period of 90 days from September 8 to December 7, 2002. The experiment was carried out three treatments each with three replications, situated in the Field Laboratory of the Faculty of Fisheries, Bangladesh Agricultural University, Mymensingh. The area of each pond was 200 m² (5 decimal) of which 18.5 m length and 13.5 m width with an average depth of 1.2 m.

Pond preparation: During the experiment, all the ponds were freed from predatory and undesirable species by repeated netting. Aquatic weeds were removed from the ponds manually by day labour. All the ponds were limed at the rate of 1 kg/decimal and manured with cow dung at the rate of 10 kg/decimal. One week after liming, the ponds were fertilized with urea and TSP at the rate of 100 g/decimal to stimulate the productivity of the ponds.

Stocking of fish: Two weeks after fertilization, the ponds were stocked with the fingerlings of GIFT tilapia (*Oreochromis niloticus*), silver carp (*Hypophthalmichthys molitrix*), mrigal (*Cirrhinus mrigala*) and pangas

(*Pangasius hypophthalmus*). Before stocking, all fishes were measured in length and weight of fingerlings by centimeter scale and electric balance. In all the treatments, GIFT tilapia, silver carp, mrigal and pangas were stocked in different species composition at the rate of 100/decimal.

Fertilization and supplementary feeding: Both inorganic and organic fertilizers were used during the period of experiment. Each pond was treated with urea and TSP at the rate of 100 g/decimal daily. On the other hand, all the ponds were fertilized with organic fertilizer (cow dung) at the rate of 5 kg/decimal at 7 days interval. Everyday, rice bran (60%) and Mustard Oil Cake (MOC, 40%) was used as supplementary feed. The feeds were calculated at a rate of 5-7% of the body weight. Feeds were applied twice a day, half in the morning (9:00 am) and the rest in the afternoon (4:00 pm).

Growth study of fishes: To record the growth of the fishes and adjust the feeding rate, 10% of fishes (tilapia and pangas) from each pond were sampled fortnightly by using a seine net and then the length (cm) and weight (g) of each species were recorded separately. Weight of fishes in each sampling was measured by using a digital electronic balance (Denver Instrument, Model XP-3000). Weight Gained (g) and Specific Growth Rate (SGR) were used to evaluate the growth of fishes.

Water quality parameters: Physico-chemical factors viz., temperature (°C), Dissolved Oxygen (DO), transparency (cm), depth (cm) and pH were measured daily and Nitrate-nitrogen (NO₃-N), Phosphate-phosphorus (PO₄-P), chlorophyll-a (µg L⁻¹) and total alkalinity (mg L⁻¹), were measured fortnightly.

Plankton population: Water samples were collected from all ponds fortnightly for qualitative and quantitative study of both phytoplankton and zooplankton. About 10 L of water samples were collected from different depth and area of each pond and was then passed through plankton net of 25 µm mesh size. For the quantitative and qualitative study of plankton, the plankton samples were analyzed in Water Quality and pond dynamics Laboratory, Faculty of Fisheries, Bangladesh Agricultural University, Mymensingh.

Planktons were expressed as cells per liter of water of each pond. The quantitative analysis of phytoplankton and zooplankton was done according to Stirling^[6] and qualitative analysis was done according to Needham and Needham^[7], Bellinger^[8].

Harvesting of fish: At the end of the experiment all the fishes were harvested by netting repeatedly with a fine meshed seine net from each pond and then counted species wise. Then the final growth gained by each species was recorded by measuring the length (cm) and weight (g) of individual fish.

Estimation of survival rate and production of fishes: The survival rate of fishes of each species for each treatment and replication were calculated from the number of fish of each species harvested from each replication at the end of the experiment. The gross and net production for each treatment and replication was determined by multiplying the average gained in weight (g) of each species of fish by the total number of fishes of the same species survived at the end of the experiment. This was done separately for each replication and treatment.

Statistical analysis: Data were analyzed statistically to observe the performances of fishes in different species combinations. Using Analysis of Variance (ANOVA) compared the mean values of growth, survival and yields obtained. The mean values were compared according to Duncan's Multiple Range Test^[9].

RESULTS AND DISCUSSION

Water quality parameters: The results of water quality parameters in different ponds such as temperature, transparency, water depth, pH, dissolved oxygen, phosphate-phosphorus, nitrate-nitrogen, total alkalinity and chlorophyll-a monitored during the study period are shown in Table 1.

The water quality parameters measured in different treatments throughout the experimental period were found to be more or less similar and all of them were within the acceptable range for fish culture. Fish culturists are more conscious about the maintenance of optimum condition of water quality parameters. The ranges of water temperature measured were 24.0 to 35.0°C. The present findings agree with the findings of Mollah and Haque^[10], Rahman *et al.*^[11], Wahab *et al.*^[12], Azim *et al.*^[13], Hasan^[14], Kohinoor *et al.*^[15] and Paul^[16]. Dewan *et al.*^[17] reported a surface water temperature ranged from 30.2 to 34.0°C in polyculture with Indian and Chinese carps. Ali *et al.*^[18] also observed temperature range of 20.5-30.5°C in pond water. These findings are also more or less similar to that of the present study. Dissolved Oxygen (DO) content were varied from 4.1 to 8.9 mg L⁻¹. The highest (8.9 mg L⁻¹) and lowest (4.1 mg L⁻¹) dissolved oxygen content were recorded in T₁. The fluctuation in DO value

might be due to alteration in the rate photosynthesis in the pond and due to rate of dissolved oxygen consumption by fish through respiration. Wahab *et al.*^[12] recorded dissolved oxygen ranging from 2.2 to 7.1 mg L⁻¹. in nine ponds of BAU campus, Mymensingh. Nirod^[19] recorded dissolved oxygen ranged from 3.4 to 8.79 mg L⁻¹, Paul^[16] found dissolved oxygen 0.85 to 7.85 mg L⁻¹, while Kohinoor^[20] measured dissolved oxygen 2 to 7.4 mg L⁻¹ in the research ponds of BAU campus, Mymensingh. From the above findings, it is concluded that the oxygen content of the experimental ponds were within the good productive range. The ranges of water pH in different treatments varied from 6.18 to 9.16 which indicate good productive conditions of the ponds. The optimum pH range for carp polyculture in pond is 6.5-9.0^[17,21,22]. The weekly variation recorded in pH might reflect the change in plankton abundance. Mean pH value recorded in different months in different treatments in the present study were in the suitable range. In the present study, transparency ranged from 16.0-37.5 cm which was near the findings of Kohinoor^[20] who recorded transparency values ranging from 15 to 58 cm. Wahab *et al.*^[22] found

Table 1: Mean values (± SD, n=9) with ranges (in parenthesis) of water quality parameters of different experimental ponds during the experimental period under three treatments

Water quality parameters	T ₁	T ₂	T ₃
Temperature (°C)	30.32±2.70 (35-25)	30.22±2.86 (35-24)	30.21±2.87 (34.5-25)
Transparency (cm)	25.67±3.09 (36.0-18.5)	25.26±3.56 (37.5-16.0)	25.18±3.74 (37.0-16.5)
Water depth (cm)	107.53±7.69 (131.0-93.0)	103.68±7.76 (126.0-89.0)	107.27±10.52 (132.0-85.0)
pH	7.89±0.53 (9.16-6.77)	7.82±0.54 (9.1-6.18)	7.82±0.53 (9.14-6.24)
Dissolved oxygen (mg L ⁻¹)	6.81±0.99 (8.9-4.1)	6.63±0.96 (8.8-4.5)	6.65±0.96 (8.7-4.3)
Total alkalinity (mg L ⁻¹)	110.14±39.65 (176.0-60.0)	109.95± 42.56 (199.0-61.0)	110.52±45.33 (194.0-63.0)
PO ₄ -P (mg L ⁻¹)	0.99±0.76 (3.2-0.23)	1.42±1.01 (3.2-0.15)	1.35±1.02 (3.6-0.19)
NO ₃ -N (mg L ⁻¹)	0.62±0.54 (2.1-0.2)	0.76±0.55 (2.3-0.2)	0.85±0.7 (3.5-0.3)
Chlorophyll-a (µg L ⁻¹)	207.29±72.23 (377.23-117.81)	175.56±69.62 (292.74-82.11)	194.25±83.00 (379.61-73.78)

Table 2: Group wise mean (± SD) plankton cell (×103) count per litre of water in different treatments during the study period.

Plankton-group	T ₁	T ₂	T ₃
Phytoplankton			
Bacillariophyceae	10.39±4.87	15.33±22.13	13.37±5.63
Chlorophyceae	19.22±8.99	26.11±12.77	31.88±11.14
Cyanophyceae	35.16±17.89	31.85±24.74	44.67±49.12
Euglenophyceae	8.34±4.52	10.99±3.07	11.23±6.63
Total	73.11±36.27	84.28±62.71	101.15±72.52
Zooplankton			
Crustacea	11.26±2.10	15.13±0.69	16.45±1.76
Rotifera	1.78±0.52	3.06±0.14	3.24±0.36
Total	13.04±2.62	18.19±0.83	19.69±2.12

Table 3: Growth, survival and production of fish in different treatments during the study period

Parameters	T ₁	T ₂	T ₃	±S.E.
Initial weight (g)				
Tilapia	4.18a	3.88a	3.77a	±0.31
Pangas	5.61a	4.33a	4.45a	±0.40
Mrigal	4.18b	4.28ab	4.34a	±0.03
Silver carp	6.07a	5.93a	5.99a	±0.09
Final weight (g)				
Tilapia	93.36a	67.83b	74.61b	±3.56
Pangas	36.22b	41.57ab	52.10a	±3.77
Mrigal	89.92a	94.47a	74.76a	±7.96
Silver carp	97.78a	71.53c	78.86b	±1.61
Weight gain (g)				
Tilapia	89.18a	63.95b	70.84b	±3.83
Pangas	30.61b	37.24ab	47.65a	±3.89
Mrigal	85.75a	86.86a	70.42a	±8.54
Silver carp	91.71a	65.60c	72.87b	±1.53
SGR (% per day)				
Tilapia	3.65a	3.16b	3.32ab	±0.12
Pangas	2.08b	2.49ab	2.74a	±0.15
Mrigal	3.36b	3.74a	3.08b	±0.10
Silver carp	3.09a	2.77b	2.85b	±0.04
Survival rate (%)				
Tilapia	92.45a	90.76a	91.26a	±0.92
Pangas	87.87a	88.56a	88.85a	±0.91
Mrigal	92.45a	90.88a	90.75a	±0.43
Silver carp	92.70a	90.60a	91.54a	±0.71
Production (kg/treatment/90 days)				
Tilapia	28.88a	18.89b	19.39b	±1.34
Pangas	0.67c	1.64b	3.16a	±0.20
Mrigal	1.91a	1.97a	1.60a	±0.19
Silver carp	8.5a	5.95c	6.66b	±0.18
Total production	39.96a	28.45b	30.81b	±1.55
(kg/treatment/90 days)				
Total production	7.99a	5.69b	6.16b	±0.31
(kg/decimal/90 days)				
Total production	0.1974a	0.1406b	0.1522b	±76.43
(kg/ha/90 days)				

Figures in the same row with same superscripts are not significantly different ($p > 0.05$)

transparency depth ranging from 15-75 cm in polyculture pond. Rahman^[23] concluded that the transparency of productive water bodies should be 40 cm or less. The observed range of water transparency was more or less similar with the findings of Wahab *et al.*^[12] and Paul^[16]. Total alkalinity values depending upon the location, season, plankton population, nature of bottom deposits etc.^[21]. Boyd^[24] advocated that the total alkalinity should be more than 20 ppm in fertilized ponds as production increases with the increase in total alkalinity. The range of total alkalinity during the study period in different treatments ranged from 60-199 mg L⁻¹ which was more or less similar to the results of Wahab *et al.*^[12], Nirod^[19], Paul^[17] and Kohinoor^[20]. Therefore, it might be concluded that this range was within the suitable range for fish culture. Phosphorus is a major nutrient and very essential for phytoplankton, the primary producer of waters. Phosphate-phosphorus (PO₄-P) in the present study was found to range from 0.15 to 3.6 mg L⁻¹. Wahab *et al.*^[12] found the highest concentration of phosphate-

phosphorus 0.09 to 5.2 mg L⁻¹ in nine experimental ponds of BAU campus, Mymensingh. So, the content of Phosphate-phosphorus in the present study was suitable for fish culture. Nitrate is extremely important as a nutrient in supplying nitrogen for protein synthesis. The ranges of nitrate-nitrogen measured in the present study were 0.2 to 3.5 mg L⁻¹. Bhuiyan^[25] reported that the range of NO₃-N from 0.06 to 0.1 ppm is suitable range for fish culture. Alikunhi^[26] also stated a concentrated of 0.06 ppm nitrate was good for fish cultivation. The mean values of nitrate concentration in the present study were higher than those of the above statements. The present findings are also more or less similar to the findings of Mollah and Haque^[10]. The value of chlorophyll-a (µg L⁻¹) indirectly express the abundance of phytoplankton in a water body. Khatrai^[27] found a positive relationship between phytoplankton and chlorophyll-a. The chlorophyll-a values showed a wide variation during the experimental period which might be due to the abundance of phytoplankton by pond fertilization. In the present study, chlorophyll-a values ranged between 73.78 and 379.61 µg L⁻¹ which indicate that all ponds under three treatments were highly productive and suitable for fish culture. Comparatively higher chlorophyll-a value was found in T₃ which due to the presence of higher concentration of phytoplankton in pond water. Several others^[16,19,20,28] have recorded more or less similar values of chlorophyll-a.

Plankton population: The group-wise mean abundance of plankton observed in three treatments is shown in Table 2. Phytoplankton population was mainly composed of Bacillariophyceae, Chlorophyceae, Cyanophyceae and Euglenophyceae whereas the zooplankton population included Crustaceans and Rotifers. Total phytoplankton was 73.11×10³, 84.28×10³ and 101.15×10³ cells L⁻¹ and zooplakton were 13.04×10³, 18.19×10³ and 19.69×10³ cells L⁻¹ in T₁, T₂ and T₃, respectively. The mean abundance of phytoplankton 73.11 to 101.15×10³ cells L⁻¹ and zooplankton concentration 13.04 to 19.69×10³ cells L⁻¹ which were more or less similar to that of Wahab and Ahmed^[29] who estimated mean phytoplankton population of 17.72×10⁴, 9.26×10⁴ and 13.87×10⁴ cells L⁻¹ from three sets of ponds, respectively. Wahab *et al.*^[22] also recorded phytoplankton numbers ranging from 2 to 8×10⁵ cells L⁻¹ and zooplankton between 2 to 3.2×10⁴ cells L⁻¹. Application of cattle manure might have enhanced the production of plankton in the present study.

Growth performance of fish: The growth performance of different species of fish viz., Tilapia, pangas, mrigal and silver carp in terms of initial weight, final weight, weight

gain, specific growth rate (SGR % per day), survival rate and total production are shown in Table 3.

There was no significant variation in initial weight of various species among the treatments. Mean final weight of different fish species varied significantly among the treatments. Mean final weight ranged between 67.83 and 93.36 g in tilapia, 36.22 and 52.10 g in pangas, 74.76 and 91.14 g in mrigal and 71.53 and 97.78 g in silver carp. The mean final weight of tilapia (93.36 g) in T₁ was significantly ($p < 0.05$) higher than the other treatments (Table 3). However there was no significant variation in mean final weight of tilapia between T₂ and T₃. The mean final weight of pangas was significantly different ($p < 0.05$) between the T₁ (36.22 g) and T₃ (52.10 g) but there was no significant variation in final weight of pangas between T₁ (36.22 g) and T₂ (41.57 g) and also between T₂ (41.57 g) and T₃ (52.10 g). There was no significant variation ($p > 0.05$) in final weight of mrigal among the treatments T₂ (94.47 g), T₁ (89.92 g) and T₃ (74.76 g), respectively. The mean final weight of silver carp was significantly different ($p < 0.05$) among the treatments. It grew best in T₁ (97.78 g) which was significantly different ($p < 0.05$) from T₂ (71.53 g) and T₃ (78.86 g). Again significant differences were found between T₂ (71.53 g) and T₃ (78.86 g).

Weight gain of all the four species of fishes varied considerably among the treatments. Weight gain (g) of tilapia, pangas, mrigal and silver carp were 89.18, 30.61, 85.75 and 91.71 g in T₁, 63.95, 37.24, 86.86 and 65.63 g in T₂ and 70.84, 47.65, 70.42 and 72.87 g in T₃, respectively. From the results, it is evident that the highest weight gain (g) was in T₁ followed by T₂ and T₃ due to proper utilization of both natural and supplementary feed in all stages by the fishes and also due to good water quality conditions. Among the treatments the weight gain of tilapia was significantly ($p < 0.05$) higher in T₁ (89.18 g) than other treatments. This might be due to the fact that tilapia is an omnivorous fish and it properly utilized the supplementary (Rice bran and Mustard oil cake) and natural food (Phytoplankton and zooplankton), as the ponds were fertilized with inorganic (Urea and TSP) and organic (cow dung) fertilizer daily and weekly. On the other hand, the water quality conditions were also found better in T₁. However, there was no significant variation between the weight gain of tilapia in T₂ (63.95 g) and T₃ (70.84 g). The weight gain of pangas was found comparatively lower in all the treatments. This might be due to the fact that it could not compete with other fishes for food and space. Weight gain of pangas was found significantly ($p < 0.05$) higher in T₃ (47.65 g) compared to the T₁ (30.61g). There was no significant variation between the weight gain of pangas in T₃ and T₂. Similarly, there was no significant variation between T₂ and T₁.

There was no significant variation in weight gain of mrigal among the T₂ (86.86 g), T₁ (85.75 g) and T₃ (70.42 g), respectively and expected weight gain was found in all the treatments. The reasons were mrigal being a bottom feeder, proper utilizing of bottom feeds and no competitor of it in all treatments. The mean weight gain of silver carp was significantly different ($p < 0.05$) among the treatments. The highest weight gain of silver carp was found in T₁ (91.71 g) and the lowest in T₂ (65.60 g). This might be silver carp being a planktivorous fish and utilized phytoplankton properly in T₁ as the lowest amount of phytoplankton was present in T₁. Phytoplankton were used highly by the silver carp, so the amount was found low.

The specific growth rate of tilapia in different treatments ranged between 3.16 and 3.65%. The significantly ($p < 0.05$) higher SGR value was recorded in T₁ (3.65%) than the T₂ (3.16%). However, no significant variation in SGR was found between T₁ (3.65%) and T₃ (3.32%) and T₃ and T₂, respectively. The SGR (% per day) in pangas varied between 2.08 and 2.74 in different treatments. Pangas in T₃ showed significantly ($p < 0.05$) the highest SGR values (2.74%) while pangas in T₁ showed the lowest (2.08%). There was no significant variation between T₃ and T₂ and T₁. Mrigal showed higher SGR values compared to other species of fish in the present study. The SGR values range between 3.08 and 3.74%. Mrigal in T₂ showed significantly ($p < 0.05$) the highest SGR values (3.74%) than the rest of the treatments. However, there was no significant variation between the SGR values of mrigal in T₁ and T₃. The SGR value of silver carp in different treatments ranged from 2.77 to 3.10% and the highest value was recorded in T₁ (3.10%) and the lowest in T₂ (2.77%). SGR of silver carp in T₁ was significantly higher ($p < 0.05$) than those in T₃ and T₂. No significant variation was found between the SGR values in T₃ and T₂. In the present study, the fortnightly average specific growth rate (SGR % per day) of fishes was found to increase more or less rapidly at the beginning of the experiment and then slowly after October till the end of the experiment. Relatively slower SGR (% per day) towards the end of the experiment might be associated with the winter season. Among all the species, mrigal showed the highest SGR (% per day) value (3.74) in T₁ both by net increase and percentage of increase in weight than the rest of the species. Tilapia showed the second highest SGR (% per day) among the fishes under different treatments and the highest value was found in T₁. Specific growth rate (SGR % per day) of silver carp and pangas were comparatively lower than mrigal and tilapia. The highest value of silver carp was found in T₁ and lowest in T₂. On the other hand, the highest SGR

value of pangas was found in T₃ where the lowest in T₁. SGR values obtained in the present study are similar to that obtained by Hossain *et al.*^[30], Salimullah^[31] and Pandey *et al.*^[32] fed supplemental feed (mustard oil cake and rice bran, 1: 1) at the rate of 5% of total body weight daily. SGR of fish fed on high protein and energy diet show higher value, but fish fed on supplemental feeds made on-farm could show SGR value between 3-4% per day^[33].

The survival rate (%) of different fish species in different treatments was fairly high. This might be due to application of both fertilization and supplementary feed. There was no significant variation of survival rate (%) of different fish species among the treatments. The survival rate of the fishes was in the range of 90.63-91.10 % in different treatments. T₁ showed the highest survival rate (91.10%) where T₃ showed the lowest (90.63%). In general, mrigal showed the highest (91.79%) and pangas showed the lowest (88.43%) survival rate among the treatments. In the present study, mean survival rates of various fish were recorded 91.10, 90.63 and 90.76% in T₁, T₂ and T₃, respectively. Similar types of survival rates were observed by Mostaque^[34], who recorded the survival rates of 86 to 95% in a polyculture system in BAU ponds. The survival rate of the present study were higher than the survival rates reported by Wahab *et al.*^[35] for Indian major carps in polyculture where supplemental feed (mustard oil cake + rice bran, 2: 1) was given. Lakshmanan *et al.*^[36] observed similar survival rate of 80% with seven species composite culture of Indian and Chinese carps in which ponds were fertilized with both organic and inorganic manures at short interval and fish were fed daily with a mixture of mustard oil cake and rice bran. There was no significant difference of survival of each fish species among the treatments. The highest survival rate was seen in T₁ followed by T₂ and T₃. Among all the species relatively higher survival rate was recorded in tilapia which was closely followed by mrigal and silver carp. The lowest survival rate was recorded in pangas. This might be due to the fact that it could not compete with other fishes for food and space.

Production of fish: Species-wise and total production of fish in terms of kg/treatment has been shown in Table 3. Total production ranged between 28.45 and 39.96 during the experimental period of 90 days. From the Table 3, it revealed that all of the fish species showed significantly ($p < 0.05$) higher production in T₁ (7.99 kg/decimal/90 days) but there was no significant variation between T₂ (5.69 kg/decimal/90 days) and T₃ (6.16 kg/decimal/90 days). In terms of kg/ha/90 days, the production range was 1406-1974 kg/ha/90 days. The highest production of tilapia was found in T₁ (28.88 kg/treatment/90 days). In T₂

and T₃ the production was 18.89 and 19.39 kg/treatment 90 days. In pangas, the highest production was found in T₃ (3.16 kg/treatment/90 days) where in T₁ and T₂ the production were 0.67 and 1.64 kg/treatment/90 days. The production of mrigal in different treatments was 1.91, 1.97 and 1.60 kg/treatment/90 days. In case of silver carp, T₁ showed the highest production (8.50 kg/treatment/90 days). The production in T₂ and T₃ was 5.95 and 6.66 kg/treatment/90 days, respectively.

The production of tilapia (kg/decimal/90 days) in T₁ was significantly ($p < 0.05$) different from the other treatments. However, there was no significant ($p > 0.05$) variation between the T₃ and T₂. In pangas, the production (kg/decimal/90 days) in T₃ was significantly ($p < 0.05$) different than the other treatments. On the other hand, significant ($p < 0.05$) variation was found between T₂ and treatment T₁. The production (kg/decimal/90 days) in case of mrigal, no significant differences were found among the T₁, T₂ and T₃.

Among the three treatments, the highest production of fish was recorded in T₁ which was subsequently followed by the production of T₂ and T₃. The reason behind the highest production in T₁ might be the role of both supplemental and natural food especially phytoplankton and zooplankton and its abundance throughout the study period. Another reason might be the value of individual final weight and survival were found to be averagely higher and better species combination in this treatment than the rest of the treatments. Tang^[37], Sinha and Gupta^[38,39] have reported that different species in polyculture pond occupy different niches with their complimentary feeding habits, fully utilizing all the natural foods in the ponds, thus increasing the total fish production from the pond. Lakshmanan *et al.*^[36] obtained good result in polyculture with Chinese and Indian major carps by stocking in varying proportions in different densities. The fish production recorded by them varied from 2230-4209 kg ha⁻¹/year. Good results were also obtained by Singh *et al.*^[40] from polyculture experiment using silver carp, grass carp, common carp together with Indian major carps and the yield recorded by them was 6196 kg/ha/year.

Jhingran^[41], Chaudhuri^[42], Mathew *et al.*^[43] and Gupta *et al.*^[44] also reported good results from the polyculture of Indian carps with exotic species and the yield recorded by them were 7000-9000, 7444.8, 10183 and 4917 kg/ha/year, respectively. Whereas, Dunseth and Smitherman^[45] find out the effectiveness of polyculture using silver carp and bighead carp in catfish grow out ponds. They noted the yield of catfishes reduce with the combination of carps.

However, the fish productions recorded in the present study were 1974, 1406 and 1522 kg/ha/90 days in treatments 1, 2 and 3, respectively. Mazid *et al.*^[46] found a gross production of 3,600 kg/ha/year from composite culture of Indian major carps and Chinese carps in Bangladesh. Wahab *et al.*^[12] also recorded 5,294 to 5,670 kg/ha/year productions in the polyculture of carps with silver carp. The total productions of fish to the present experiment were within the range of good production when compared with the above findings.

Among the fish species, expected growth performance of mrigal was found in all treatments. The highest production was found in T₂ and lowest in T₃, but there was no significant variation among the treatments. The reasons were bottom feeds were properly utilized and no competitor of it in bottom layer in all treatments.

Growth and production of tilapia and silver carp were better in T₁ than the other treatments. This might be due to the fact, that in T₁, they properly utilized both natural and artificial feed and successfully compete with other species for food and space. The water quality condition of T₁ was also better which might be the reason.

Pangas was experimentally introduced in this study to observe the growth and production compare with tilapia, silver carp and mrigal. But expected production was not found in any treatment. This might be due to its failure to compete with other species for food and space. So, polyculture of pangas with tilapia, mrigal and silver carp is not suggested. Therefore, more studies are required to establish a suitable species composition of pangas polyculture to observe its growth and production.

The result of the present study demonstrated that stocking density of 100 fish/decimal at different species combination of tilapia, pangas, mrigal and silver carp in T₁ is optimum for polyculture using low-cost feed and fertilizer.

REFERENCES

1. Jhingran, V.G., 1975. Aquaplosion Lecture, Summer Int. Intensive Freshwater Fish culture, Cuttack, Organized by centre. Inland Fish Res. Inst., Barrackpore (ICAR), pp: 12.
2. Halver, J.E., 1984. Special Methods in Pen Fish Husbandry. Akademiai Nyomda, Budapest, pp: 146.
3. Hussain, M.G., M.A. Rahman, M. Akteruzzaman and A.H.M. Kohinoor, 1987. A study on the production of *O. niloticus* in semi-intensive system in Bangladesh. Bangladesh J. Fish., 11: 59-65.
4. Kohinoor, A.H.M., M. Akteruzzaman and M.S. Shah, 1993. Red tilapia production potential under low input management in Bangladesh. Paper presented at the Third Fisheries Forum, 26-30 October, World Trade Center, Singapore.
5. Gupta, M.V., M. Akteruzzaman, A.H.M. Kohinoor and M.S. Shah, 1994. Nile tilapia (*O. niloticus*) culture under different feeding and fertilization regimes. In: Pullin, R.S.V., J. Lazer, M. Legendre and J.B. Amon Kothias (Eds.). The third International Symposium on Tilapia in Aquaculture. ICLARM Conf. Proc., pp: 41.
6. Stirling, H.P., 1985. Chemical and Biological Methods of Water Analysis for Aquaculturists. Institute of Aquaculture, University of Stirling, Scotland, pp: 119.
7. Needham, J.G. and P.R. Needham, 1963. A Guide to Study of Freshwater Biology. 5th Edn., Holden-day Inc., San Francisco, pp: 106.
8. Bellinger, E.G., 1992. A Key to Common Algae: Freshwater Estuarine and Some Coastal Species. The Institute of Water and Environmental Management, London, UK.
9. Gomez, K.A. and A.A. Gomez, 1984. Statistical Procedures for Agricultural Research. 2nd Ed. John Wiley and Sons, New York, pp: 680.
10. Mollah, M.F.A. and A.K.M.A. Haque, 1978. Studies on monthly variations of plankton in relation to the physico-chemical conditions of water and bottom soil of two ponds. II. Zooplankton. Bangladesh J. Fish., 1: 99-103.
11. Rahman, M.S., M.Y. Chowdhury, A.K.M.A. Haque and M.S. Haq, 1982. Limnological studies of four ponds. Bangladesh J. Fish., 2-5: 25-35.
12. Wahab, M.A., Z.F. Ahmed, M.A. Islam and S.M. Rahmatullah, 1995. Effect of introduction of common carp, *Cyprinus carpio* (L), of the pond ecology and growth of fish in polyculture. Aquaculture Research., 26: 619-628.
13. Azim, M.E., G.S. Talukder, M.A. Wahab, M.M. Haque and M.S. Haq, 1995. Effect of liming and maintenance of total hardness levels on fish production in fertilized ponds. Progress. Agric., 6: 7-14.
14. Hasan, M.A., 1998. Development of carp polyculture techniques with small indigenous fish species mola *Amblypharyngodon mola*, chela *Chela cachius*, punti *Puntius sophore*. MS dissertation, Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh, Bangladesh, pp: 71.
15. Kohinoor, A.H.M., M.L. Islam, M.A. Wahab and S. H. Thilsted, 1998. Effect of mola (*Amblypharyngodon mola* Ham.) on the growth and production of carps in polyculture. Bangladesh J. Fish., 2: 119-126.

16. Paul, S., 1998. Comparison Between Carp Polyculture System with Silver Carp (*Hypophthalmichthys molitrix*) and with Small Indigenous fish mola (*Amblypharyngodon mola*) MS Thesis, Bangladesh Agricultural University, Mymensingh, Bangladesh, pp: 85.
17. Dewan, S., M.A. Wahab, M.C.M. Beveridge, M.H. Rahman and B.K. Sarker, 1991. Food selection, electivity and dietary overlap among planktivorous Chinese and Indian major carps fry and fingerlings grown in extensively managed, rain-fed ponds in Bangladesh. *Aquaculture and Fisheries Management*, 22: 277-294.
18. Ali, S., A.K. Rahman, A.R. Patwary and K.H.R. Islam, 1982. Studies on the diurnal variations in physico-chemical factors and zooplankton in a freshwater pond. *Bangladesh J. Fish.*, 2-5: 15-23.
19. Nirod, D.B., 1997. Effect on stocking density on the growth and production of mola (*Amblypharyngodon mola*). MS Thesis, Bangladesh Agricultural University, Mymensingh, Bangladesh, pp: 75.
20. Kohinoor, A.H.M., 2000. Development of culture technology of three small indigenous fish mola (*Amblypharyngodon mola*), punti (*Puntius sophore*) and chela (*Chela cachius*) with notes on some aspects of their biology. Ph.D. Thesis, Bangladesh Agricultural University, Mymensingh, Bangladesh, pp: 363.
21. Jhingran, V.G. and P.K. Talwar, 1991. *Inland Fisheries of India and Adjacent Countries*. Vol. 1. Oxford and IBH Publishing Co. Pvt. Ltd. Calcutta, pp: 541.
22. Wahab, M.A., Z.F. Ahmed, M.S. Haq and M. Begum, 1994. Compatibility of silver carp in the polyculture of crprinid fishes. *Progress. Agric.*, 5: 221-227.
23. Rahman, M.S., 1992. *Water Quality Management In Aquaculture*. BRAC Prokashana, Dhaka 1212, pp: 84.
24. Boyd, C.E., 1982. *Water Quality Management for Pond Fish Culture*. Elsevier Sci. Publ. Co. Amsterdam-Oxford New York, pp: 318.
25. Bhuiyan, B.R., 1970. Physico-chemical qualities of some ancient tanks of Sibsagar. *Assam. Environmental Health*, 12: 129-134.
26. Alikunhi, K.H., 1957. Fish culture in India. *Fm. Bull. India Coun. Agri. Res.*, 20: 144.
27. Khatrai, T.C., 1984. Seasonal variation in the ecosystem of the Lakhotia lake in Rajasthan. *India J. Fish.*, 31: 122-129.
28. Rahman, M.M., 2000. Effect of addition of calbasu *Labeo calbasu* in the periphyton based aquaculture system of rohu and catla. MS. Thesis, Bangladesh Agricultural University, Mymensingh, Bangladesh, pp: 82.
29. Wahab, M.A. and Z.F. Ahmed, 1992. Effect of planktivorous carp species combination on food organisms and electivity indices in the fish ponds. *Progress. Agric.*, 2: 21-30.
30. Hossain, M.A., M. Ahmed and M.N. Islam, 1997. Mixed culture of fishes in seasonal ponds through fertilization and feeding. *Bangladesh J. Fish. Res.*, 1: 9-18.
31. Salimullah, M., 1997. A comparative study on the growth of fishes under polyculture system with supplementary and fertilization. M.S. Thesis. Bangladesh Agricultural University, Mymensingh, Bangladesh, pp: 72.
32. Pandey, B.N., R. Perween, R. Yasmin and A.K. Khan, 1992. Effect of supplementary feeding on the growth of Indian major carp, *Catla catla* (Ham). *J. Freshwat. Biol.*, 4: 261-265.
33. De Silva, S.S. and F.B. Davy, 1992. Fish nutrition research for semi-intensive culture system in Asia. *Asian Fish. Sci.*, 5: 129-144.
34. Mostaque, A., 1995. Studies on the mixed culture of fishes in seasonal ponds through fertilization and feeding. MS. Thesis, Faculty of Fisheries, Bangladesh Agricultural University, Mymensingh, Bangladesh.
35. Wahab, M.A., M. Begum and Z.F. Ahmed, 1991. The effects of silver carp introduction in the polyculture of major Indian carps. *Proc. BAU Res. Prog.*, 5: 429-437.
36. Lakshmanan, M.A.V., K.K. Sukumaran, D.S. Murty, D.P. Chakraborty and M.T. Philipose, 1971. Preliminary observation on intensive fish farming in freshwater ponds by the composite culture of Indian and exotic species. *J. Inland Fish. Soc.*, 3: 1-21.
37. Tang, Y.A., 1970. Evaluation of balance between fishes and available fish foods in multispecies fish culture ponds in Taiwan. *Trans. Am. Fish. Soc.*, 99: 708-718.
38. Sinha, V.R.P. and M.V. Gupta, 1974. Composite culture of Indian and exotic fisheries at Kalyani. *Indian Fmg*, 24: 29-31.
39. Sinha, V.R.P. and M.V. Gupta, 1975. On the growth of grass carp, *Ctenopharyngodon idella* (Val.) in composite fish culture at Kalyani, West Bengal (India). *Aquaculture*, 5: 283-290.
40. Singh, S.B. K.K. Sukumaran, P.C. Chakraborty and M.M. Bagchi, 1972. Observations of composite culture of exotic carps. *J. Inland Fish. Soc., India*, 4: 38-50.
41. Jhingran, V.G., 1976. System of polyculture of polyculture in inland waters in India. *J. Fish. Res. Board can.*, 33: 905-910.

42. Chaudhuri, H., N.G.S. Rao, G.N. Saha, M. Rout and D.R. Ranaujia, 1978. Record fish production through intensive fish culture in a farmers ponds. *J. Inland Fish. Soc. India*, 10: 19-27.
43. Mathew, P.M., B.K. Singh and D.P. Chakrabarty, 1988. A case study stocking of fry in six species composite fish culture. *J. Maharashtra Agric. Univ.*, 13: 79-82.
44. Gupta, V.K., J.P. Sharma and J.B. Srivastava, 1990. Polyculture of Indian and exotic carps using cattle manurer with and without supplementary feed. *Inland Fish. Res. Inst.*, pp: 430-446.
45. Dunseth, D.R. and R.O. Smitherman, 1977. Polyculture of catfish, tilapia and silver carp. *Proc. Ann. Comm. Fish. Workshop*, 6: 25-26
46. Mazid, M.A., M. Zaher, N.N. Begum, M.Z. Ali and F. Nahar, 1997. Formulation of cost effective feeds from locally available ingredients for carp polyculture systems for increased production. *Aquaculture*, 151: 71-78.