

POLYHARMONIC HOMOGENIZATION, ROUGH POLYHARMONIC SPLINES AND SPARSE SUPER-LOCALIZATION

HOUMAN OWHADI¹, LEI ZHANG² AND LEONID BERLYAND³

Abstract. We introduce a new variational method for the numerical homogenization of divergence form elliptic, parabolic and hyperbolic equations with arbitrary rough (L^∞) coefficients. Our method does not rely on concepts of ergodicity or scale-separation but on compactness properties of the solution space and a new variational approach to homogenization. The approximation space is generated by an interpolation basis (over scattered points forming a mesh of resolution H) minimizing the L^2 norm of the source terms; its (pre-)computation involves minimizing $\mathcal{O}(H^{-d})$ quadratic (cell) problems on (super-)localized sub-domains of size $\mathcal{O}(H \ln(1/H))$. The resulting localized linear systems remain sparse and banded. The resulting interpolation basis functions are biharmonic for $d \leq 3$, and polyharmonic for $d \geq 4$, for the operator $-\operatorname{div}(a\nabla\cdot)$ and can be seen as a generalization of polyharmonic splines to differential operators with arbitrary rough coefficients. The accuracy of the method ($\mathcal{O}(H)$ in energy norm and independent from aspect ratios of the mesh formed by the scattered points) is established *via* the introduction of a new class of higher-order Poincaré inequalities. The method bypasses (pre-)computations on the full domain and naturally generalizes to time dependent problems, it also provides a natural solution to the inverse problem of recovering the solution of a divergence form elliptic equation from a finite number of point measurements.

Mathematics Subject Classification. 41A15, 34E13, 35B27.

Received August 3, 2013.

Published online March 11, 2014.

1. INTRODUCTION

Consider the partial differential equation

$$\begin{cases} -\operatorname{div}\left(a(x)\nabla u(x)\right) = g(x) & x \in \Omega; g \in L^2(\Omega), a(x) = \{a_{ij} \in L^\infty(\Omega)\} \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

Keywords and phrases. Homogenization, polyharmonic splines, localization.

¹ Corresponding author. California Institute of Technology, Computing and Mathematical Sciences, MC 9-94 Pasadena, CA 91125, USA. owhadi@caltech.edu

² Shanghai Jiaotong University, Institute of Natural Sciences and Department of Mathematics, Key Laboratory of Scientific and Engineering Computing (Shanghai Jiao Tong University), Ministry of Education, 800 Dongchuan Road, Shanghai 200240, P.R. China. lzhang2012@sjtu.edu.cn

³ Pennsylvania State University, Department of Mathematics, University Park, PA, 16802, USA. berlyand@math.psu.edu

where Ω is a bounded subset of \mathbb{R}^d with piecewise Lipschitz boundary which satisfies an interior cone condition and a is symmetric, uniformly elliptic on Ω and with entries in $L^\infty(\Omega)$. It follows that the eigenvalues of a are uniformly bounded from below and above by two strictly positive constants, denoted by $\lambda_{\min}(a)$ and $\lambda_{\max}(a)$. More precisely, for all $\xi \in \mathbb{R}^d$ and almost all $x \in \Omega$,

$$\lambda_{\min}(a)|\xi|^2 \leq \xi^T a(x)\xi \leq \lambda_{\max}(a)|\xi|^2. \quad (1.2)$$

In this paper, as in [15, 69, 70], we are interested in the homogenization of (1.1) (and its parabolic and hyperbolic analogues in Sect. 8), but not in the classical sense, *i.e.*, that of asymptotic analysis [14] or that of G or H -convergence [37, 65, 80] in which one considers a sequence of operators $-\operatorname{div}(a_\epsilon \nabla \cdot)$ and seeks to characterize the limit of solutions. We are interested in the homogenization of (1.1) in the sense of “numerical homogenization,” *i.e.*, that of the approximation of the solution space of (1.1) by a finite-dimensional space. This approximation is not based on concepts of scale separation and/or of ergodicity but on strong compactness properties, *i.e.*, the fact that the unit ball of the solution space is strongly compactly embedded into $\mathcal{H}_0^1(\Omega)$ if source terms (g) are in L^2 (see [70], Appendix B).

In other words if (1.1) is “excited” by source terms $g \in L^2$ then the solution space can be approximated by a finite dimensional sub-space (of $\mathcal{H}_0^1(\Omega)$) and the question becomes “how to approximate” this solution space. The approximate solution space introduced in this paper is constructed as follows (see Sect. 2):

- (1) Select a finite subset of points $(x_i)_{i \in \mathcal{N}}$ in Ω with $\mathcal{N} := \{1, \dots, N\}$.
- (2) Identify the nodal interpolation basis $(\phi_i)_{i \in \mathcal{N}}$ as minimizers of $\int_\Omega |\operatorname{div}(a \nabla \phi)|^2$ subject to $\phi_i(x_j) = \delta_{i,j}$ (where $\delta_{i,j}$ is the Kronecker delta defined by $\delta_{i,i} = 1$ and $\delta_{i,j} = 0$ for $i \neq j$).
- (3) Identify the approximate solution space as the interpolation space generated by the elements ϕ_i .

Note that we construct here an interpolation basis with N elements, which interpolates a given function using its values at the nodes $(x_i)_{i \in \mathcal{N}}$ (analogously to *e.g.*, the classical interpolation polynomials of degree $N - 1$).

Our motivation in identifying the interpolation elements through the minimization step (2) lies in observation that the origin of the compactness of the solution space lies in the higher integrability of g . In Section 2 we will show that one of the properties of the interpolation basis ϕ_i is that $\sum_i w_i \phi_i$ minimizes $\int_\Omega |\operatorname{div}(a \nabla w)|^2$ over all functions w interpolating the points x_i (*i.e.* such that $w(x_i) = w_i$). The error of the interpolation and the accuracy of the resulting finite element method is established in Section 3 through the introduction of a new class of higher order Poincaré inequalities.

The accuracy of our method (as a function of the locations of the interpolation points) will depend only on the mesh norm of the discrete set $(x_i)_{i \in \mathcal{N}}$, *i.e.*, the constant H defined by

$$H := \sup_{x \in \Omega} \min_{i \in \mathcal{N}} |x - x_i| \quad (1.3)$$

Our method is meshless in its formulation and error analysis and the density of the interpolation points $(x_i)_{i \in \mathcal{N}}$ can be increased near points of the domain where high accuracy is required. In the fully discrete formulation of our method (Sect. 9), a will be chosen to be piecewise constant on a tessellation \mathcal{T}_h (of Ω) of resolution $h \ll H$ and the points x_i will be a subset of interior nodes of \mathcal{T}_h . More precisely, in numerical applications where a is represented *via* constant values on a fine mesh \mathcal{T}_h of resolution h , H will be the resolution of a coarse mesh \mathcal{T}_H having interior nodes x_i (the nodes of \mathcal{T}_H will be nodes of \mathcal{T}_h but triangles of \mathcal{T}_H may not be unions of triangles of \mathcal{T}_h) and the accuracy of our method will depend only on H (*i.e.*, the accuracy will be independent from the regularity of \mathcal{T}_H and the aspect ratios of its elements (tetrahedra for $d = 3$, triangles for $d = 2$)).

In Section 4 we show that our method also provides a natural solution to the (inverse) problem of recovering u from the (partial) measurements of its (nodal) values at the sites x_i . Although we restrict, in this paper, our attention to $d \leq 3$, we show, in Section 5, how our method can be generalized to $d \geq 4$ by requiring that $g \in L^{2m}(\Omega)$ (for $(d - 1)/2 \leq m \leq d/2$) and obtaining the elements ϕ_i as minimizers of $\int_\Omega |\operatorname{div}(a \nabla \phi)|^{2m}$. The resulting elements are $2m$ -harmonic in the sense that they satisfy $(\operatorname{div}(a \nabla \cdot))^{2m} \phi_i = 0$ in $\Omega \setminus \{x_j : j \in \mathcal{N}\}$.

In that sense our method can be seen as a polyharmonic formulation of homogenization and a generalization of polyharmonic splines [24, 43] to PDEs with rough coefficients (see Sect. 6, the basis elements ϕ_i can be seen as “rough polyharmonic splines”). In Section 7 we show how the computation of the basis elements ϕ_i can be localized to sub-domains $\Omega_i \subset \Omega$ and obtain *a posteriori* error estimates. Those *a posteriori* error estimates show that if the localized elements ϕ_i^{loc} decay at an exponential rate away from x_i (which we observe in our numerical experiments in Sect 9) then the accuracy of the method remains $\mathcal{O}(H)$ in \mathcal{H}^1 -norm for sub-domains Ω_i of size $\mathcal{O}(H \ln(1/H))$, which we refer to as the *super-localization property*. We defer the proof and statement of *a priori* error estimates to a sequel work. We give a short review of numerical homogenization in Section 10 and we discuss connections with classical homogenization in Section 10.4.

We discuss and compare the accuracy and computational cost of our method in Section 10.3. The linear systems to be solved for the identification of the (super-)localized basis ϕ_i^{loc} are sparse and banded (by virtue of this property, the computational cost of our method is minimal).

2. VARIATIONAL FORMULATION AND PROPERTIES OF THE INTERPOLATION BASIS

2.1. Identification of the interpolation basis

Let V be the set of functions $u \in \mathcal{H}_0^1(\Omega)$ such that $\text{div}(a\nabla u) \in L^2(\Omega)$. Observe that V is the solution space of (1.1) for $g \in L^2(\Omega)$. By [36, 81], solutions of (1.1) are Hölder continuous functions over Ω provided that their source terms g belong to $L^{d/2+\varepsilon}(\Omega)$ for some $\varepsilon > 0$. It follows that if the dimension of the physical space is less than or equal to three ($d \leq 3$) then elements of V are Hölder continuous functions over Ω . For the sake of simplicity, we will restrict our presentation to $d \leq 3$ and show in Section 5 how the method and results of this paper can be generalized to $d \geq 4$.

Let $\|\cdot\|_V$ be the norm on V defined by

$$\|u\|_V := \|\text{div}(a\nabla u)\|_{L^2(\Omega)}. \tag{2.1}$$

It is easy to check that $(V, \|\cdot\|_V)$ is a complete linear space; in particular, it is closed under the norm $\|\cdot\|_V$.

Let $(x_i, i = 1, \dots, N)$ be a finite subset of points in Ω . Write $\mathcal{N} := \{1, \dots, N\}$ and define H as in (1.3) (in practical applications, $H > 0$ will be a parameter determined by available computing power and desired accuracy). For each $i \in \{1, \dots, N\}$ define

$$V_i := \{\phi \in V \mid \phi(x_i) = 1 \text{ and } \phi(x_j) = 0, \text{ for } j \in \{1, \dots, N\} \text{ such that } j \neq i\} \tag{2.2}$$

and consider the following optimization problem over V_i :

$$\begin{cases} \text{Minimize } \int_{\Omega} |\text{div}(a\nabla \phi)|^2 \\ \text{Subject to } \phi \in V_i. \end{cases} \tag{2.3}$$

We will now show that (2.3) is a well posed (strictly convex) quadratic optimization problem and we will identify its unique minimizer ϕ_i .

Remark 2.1. Observe that ϕ_i depends only on a , Ω and the locations of the points $(x_i)_{i \in \mathcal{N}}$. Note that we do not require the distribution of those points be uniform or regular (in particular, the density of points x_i can be adapted to the structure of a or increased in locations where higher accuracy is desired). In particular, in the construction of ϕ_i , we do not use any tessellation, we just use the set of points $(x_i)_{i \in \mathcal{N}}$.

Let $G(x, y)$ be the Green’s function of (1.1). Recall that G is symmetric and that it satisfies (for $y \in \Omega$)

$$\begin{cases} -\text{div}(a(x)\nabla G(x, y)) = \delta(x - y) & x \in \Omega; \\ G(x, y) = 0 \text{ for} & x \in \partial\Omega, \end{cases} \tag{2.4}$$

where $\delta(x - y)$ is the Dirac distribution centered at y . For $i, j \in \{1, \dots, N\}$ define

$$\Theta_{i,j} := \int_{\Omega} G(x, x_i) G(x, x_j) dx. \quad (2.5)$$

Lemma 2.2. *The integrals (2.5) are finite, well defined and for all $i, j \in \{1, \dots, N\}$, $\Theta_{i,j}$ is bounded by a constant depending only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω . The $N \times N$ matrix Θ is symmetric positive definite. Furthermore for all $l \in \mathbb{R}^N$,*

$$l^T \Theta l = \|v\|_{L^2(\Omega)}^2 \quad (2.6)$$

where v is the solution of

$$\begin{cases} -\operatorname{div}(a(x)\nabla v(x)) = \sum_{j=1}^N l_j \delta(x - x_j) & \text{for } x \in \Omega; \\ v(x) = 0 & \text{for } x \in \partial\Omega. \end{cases} \quad (2.7)$$

Proof. By Theorem 1.1. of [42], for $d \geq 3$, the Green's function is bounded by

$$G(x, y) \leq K_d |x - y|^{2-d} \quad (2.8)$$

where K_d depends only on $d, \lambda_{\min}(a)$ and $\lambda_{\max}(a)$. For $d = 2$, we have (see for instance [84] and references therein)

$$G(x, y) \leq K_2 \left(1 + \ln \frac{\operatorname{diam}(\Omega)}{|x - y|} \right) \quad (2.9)$$

where K_2 depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω . For $d = 1$, the Green function is trivially bounded by a constant depending only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω . It follows from these observations that for $d = 1, 2, 3$

$$\int_{\Omega} (G(x_i, y))^2 dy \leq K \quad (2.10)$$

where the constant K is finite and depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω . We deduce that the integrals in (2.5) (and hence Θ) are well defined.

Let $l \in \mathbb{R}^N$, write

$$v(x) := \sum_{i=1}^N G(x, x_i) l_i. \quad (2.11)$$

Observe that v is the solution of (2.7) and that $\|v\|_{L^2(\Omega)}^2 = l^T \Theta l$. Then, it follows that Θ is symmetric positive definite. Indeed if Θ is not positive definite, then there would exist a non zero vector $l \in \mathbb{R}^N$ such that $\Theta l = 0$. This would imply $\|v\|_{L^2(\Omega)} = 0$ which is a contradiction since the equation $-\operatorname{div}(a(x)\nabla v(x)) = \sum_{j=1}^N l_j \delta(x - x_j)$ has a non zero solution. \square

Let P be the $N \times N$ symmetric definite positive matrix defined as the inverse of Θ , i.e.

$$P := \Theta^{-1} \quad (2.12)$$

Note that Lemma 2.2 (i.e. the well-posedness and invertibility of Θ) guarantees the existence of P .

Define

$$\tau(x, y) := \int_{\Omega} G(x, z) G(z, y) dz. \quad (2.13)$$

Note that τ is symmetric and (see proof of Lem. 2.2) that for each $y \in \Omega$, $x \rightarrow \tau(x, y)$ is a well defined element of V (in particular, it is Hölder continuous on Ω). Note also that for $i, j \in \mathcal{N}$ we have $\tau(x_i, x_j) = \Theta_{i,j}$ and that τ is the fundamental solution of $(\operatorname{div}(a\nabla\cdot))^2$ in the sense that for $y \in \Omega$

$$\begin{cases} \operatorname{div} \left(a\nabla \left(\operatorname{div} (a\nabla\tau(x, y)) \right) \right) = \delta(x - y) & \text{for } x \in \Omega \\ \tau(x, y) = \operatorname{div} (a\nabla\tau(x, y)) = 0 & \text{for } x \in \partial\Omega. \end{cases} \quad (2.14)$$

Proposition 2.3. V_i is a non-empty closed affine subspace of V . Problem (2.3) is a strictly convex optimization problem over V_i . The unique minimizer of (2.3) is

$$\phi_i(x) := \sum_{j=1}^N P_{i,j} \tau(x, x_j). \quad (2.15)$$

Remark 2.4. It is important to note that in practical (numerical) applications each element ϕ_i would be obtained by solving the quadratic optimization problem (2.3) rather than through the representation formula (2.15).

Proof. Let us first prove that $\phi_i \in V_i$. First observe that for all $j \in \{1, \dots, N\}$,

$$-\operatorname{div} \left(a(x)\nabla\tau(x, x_j) \right) = G(x, x_j) \quad (2.16)$$

and $\tau(x, x_j) = 0$ on $\partial\Omega$. Noting that $\|\operatorname{div}(a(x)\nabla\tau(x, x_j))\|_{L^2(\Omega)}^2 = \Theta_{j,j}$ we deduce from Lemma 2.2 that $\tau(x, x_j) \in V$. We conclude from (2.15) that $\phi_i \in V$. Now observing that

$$\Theta_{i,j} = \tau(x_i, x_j) \quad (2.17)$$

we deduce (using the fact that $P \cdot \Theta$ is the $N \times N$ identity matrix) that

$$\phi_i(x_j) = (P \cdot \Theta)_{i,j} = \delta_{i,j}. \quad (2.18)$$

We conclude that $\phi_i \in V_i$ which implies that V_i is non empty (it is easy to check that it is a closed affine sub-space of V).

Now let us prove that problem (2.3) is a strictly convex optimization problem over V_i . Let $v, w \in V_i$ such that $v \neq w$. Write for $\lambda \in [0, 1]$,

$$f(\lambda) := \int_{\Omega} \left| \operatorname{div} (a\nabla(v + \lambda(w - v))) \right|^2. \quad (2.19)$$

and we need to show that $f(\lambda)$ is a strictly convex function. Observing that

$$f(\lambda) = \int_{\Omega} \left| \operatorname{div}(a\nabla v) \right|^2 + 2\lambda \int_{\Omega} (\operatorname{div}(a\nabla v))(\operatorname{div}(a\nabla(w - v))) + \lambda^2 \int_{\Omega} \left| \operatorname{div}(a\nabla(w - v)) \right|^2 \quad (2.20)$$

and noting that $\int_{\Omega} \left| \operatorname{div}(a\nabla(w - v)) \right|^2 > 0$ (otherwise one would have $v = w$ given that v and w are both equal to zero on $\partial\Omega$) we deduce that f is strictly convex in λ . We conclude that (see, for example, [32], p. 35, Prop. 1.2) that Problem (2.3) is a strictly convex optimization problem over V_i and that it admits a unique minimizer in V_i .

Let us now prove that ϕ_i is the minimizer of (2.3). Let $v \in V_i$ with $v \neq \phi_i$. Write $I(v) := \int_{\Omega} \left| \operatorname{div}(a\nabla v) \right|^2$. We have

$$I(v) = I(\phi_i) + I(v - \phi_i) + 2J(\phi_i, v - \phi_i) \quad (2.21)$$

with

$$J(\phi_i, v - \phi_i) = \int_{\Omega} (\operatorname{div}(a\nabla\phi_i))(\operatorname{div}(a\nabla(v - \phi_i))). \quad (2.22)$$

Note that (as before) $I(v - \phi_i) > 0$ if $v \neq \phi_i$ and using (2.16) and (2.15) we obtain that

$$-\operatorname{div}(a(x)\nabla\phi_i(x)) := \sum_{j=1}^N P_{i,j} G(x, x_j) \quad (2.23)$$

which leads to

$$J(\phi_i, v - \phi_i) = \sum_{j=1}^N P_{i,j} (v(x_j) - \phi_i(x_j)) = 0 \quad (2.24)$$

where in the last equality we have used $v(x_j) - \phi_i(x_j) = 0$ for all $j \in \{1, \dots, N\}$. It follows that $I(v) > I(\phi_i)$ which implies that ϕ_i is the minimizer of (2.3). \square

2.2. Variational properties of the interpolation basis

Write V_0 the subset of V defined by

$$V_0 := \{v \in V : v(x_i) = 0, \forall i \in \{1, \dots, N\}\} \quad (2.25)$$

For $u, v \in V$, define the (scalar) product $\langle \cdot, \cdot \rangle$ by

$$\langle u, v \rangle := \int_{\Omega} (\operatorname{div}(a\nabla u))(\operatorname{div}(a\nabla v)) \quad (2.26)$$

The following theorem shows that the functions ϕ_i generate a linear space interpolating elements of V with minimal $\langle \cdot, \cdot \rangle$ -norm (see (2.28)).

Theorem 2.5. *It holds true that*

- The basis ϕ_i is orthogonall to V_0 with respect to the product $\langle \cdot, \cdot \rangle$, i.e.

$$\langle \phi_i, v \rangle = 0, \quad \forall i \in \{1, \dots, N\} \text{ and } \forall v \in V_0 \quad (2.27)$$

- $\sum_i w_i \phi_i$ is the unique minimizer of

$$\langle w, w \rangle = \int_{\Omega} (\operatorname{div}(a\nabla w))^2 \quad (2.28)$$

over all $w \in V$ such that $w(x_i) = w_i$.

- For all $i \in \{1, \dots, N\}$ and for all $v \in V$,

$$\langle \phi_i, v \rangle = \sum_{j=1}^N P_{i,j} v(x_j) \quad (2.29)$$

- For all $i, j \in \{1, \dots, N\}$,

$$\langle \phi_i, \phi_j \rangle = P_{i,j} \quad (2.30)$$

Proof. Equation (2.27) trivially follows from (2.29). (2.29) is a direct consequence of (2.23). (2.29) leads to $\langle \phi_i, v \rangle = 0$ for $v \in V_0$. To prove that $w = \sum_i w_i \phi_i$ is the unique minimizer of (2.28) over $W := \{w \in V \mid w(x_i) = w_i \text{ for all } i \in \{1, \dots, N\}\}$ we simply observe that if w' is another element of W then $w' - w$ belongs to V_0 and for $w' \neq w$ we have (from (2.27), using $\langle w, w' - w \rangle = 0$) that

$$\langle w', w' \rangle = \langle w, w \rangle + \langle w' - w, w' - w \rangle > \langle w, w \rangle. \tag{2.31}$$

For (2.30), using (2.29) and writing I_N the $N \times N$ identity matrix we have

$$\langle \phi_i, \phi_j \rangle = \sum_{k=1}^N P_{ik} \phi_j(x_k) = (P \cdot I_N)_{i,j} = P_{i,j}. \tag{2.32}$$

□

Remark 2.6. It is easy to check that the orthogonality (2.27) implies that ϕ_i solves

$$\begin{cases} \operatorname{div} (a \nabla (\operatorname{div} (a \nabla \phi_i))) = 0 & \text{on } \Omega \setminus \{x_j \mid j \in \mathcal{N}\} \\ \phi_i = \operatorname{div} (a \nabla \phi_i) = 0 & \text{on } \partial \Omega \\ \phi_i(x_j) = \delta_{i,j}. \end{cases} \tag{2.33}$$

However, (2.33) alone cannot be used to uniquely identify ϕ_i as (2.15). This is due to the fact that the solution of (2.33) may not be unique. As a counter-example consider $\Omega = (-1, 1)$, $x_1 = 0$, $\mathcal{N} = \{1\}$ and $a = I_d$. For $\alpha \in \mathbb{R}$ define

$$\begin{cases} \psi_\alpha(x) := \frac{\alpha-1}{2}x^3 + \frac{3}{2}(\alpha-1)x^2 + \alpha x + 1 & \text{on } [-1, 0] \\ \psi_\alpha(x) := \frac{\alpha+1}{2}x^3 - \frac{3}{2}(\alpha+1)x^2 + \alpha x + 1 & \text{on } [0, 1] \end{cases} \tag{2.34}$$

then for any $\alpha \in \mathbb{R}$, ψ_α is a solution (2.33). Using the variational formulation of ϕ_i it is possible to show that to enforce the uniqueness of the solution of (2.33) we also need to require that (see Prop. 7.6), for some $c \in \mathbb{R}^N$,

$$\operatorname{div} (a \nabla (\operatorname{div} (a \nabla \phi_i))) = \sum_{j=1}^N c_j \delta(x - x_j) \text{ on } \Omega \tag{2.35}$$

which in dimension one is equivalent to the continuity of $\operatorname{div}(a \nabla \phi_i)$ across the coarse nodes $(x_j)_{j \in \mathcal{N}}$. Note that (2.35) implies that for all $v \in V_0$

$$\int_{\Omega} \operatorname{div}(a \nabla \phi_i) \operatorname{div}(a \nabla v) = \int_{\Omega} v \operatorname{div} (a \nabla (\operatorname{div} (a \nabla \phi_i))) = 0. \tag{2.36}$$

3. FROM A HIGHER ORDER POINCARÉ INEQUALITY TO THE ACCURACY OF THE INTERPOLATION BASIS

In this section we introduce a new higher order Poincaré inequality and derive the accuracy of the interpolation space computed in (2.3). This new class can be thought of as a generalization of Sobolev inequalities for functions with scattered zeros (see [56, 58–60, 66]) to operators with rough coefficients.

3.1. A Higher order Poincaré inequality

We will first present the new Poincaré inequality (Thm. 3.3). Analogous inequalities when $a = I_d$ can be found in Theorem 1.1 of [66] and in [26] but the presence of the L^∞ matrix a renders the proof much more difficult and requires a new strategy based on the following lemma.

Lemma 3.1. *Let $d \leq 3$ and B_1 be the open ball of center 0 and radius 1. There exists a finite strictly positive constant $C_{d,\lambda_{\min}(a),\lambda_{\max}(a)}$ such that for all $v \in \mathcal{H}^1(B_1)$ such that $\operatorname{div}(a\nabla v) \in L^2(B_1)$ it holds true that*

$$\|v - v(0)\|_{L^2(B_1)}^2 \leq C_{d,\lambda_{\min}(a),\lambda_{\max}(a)} \left(\|\nabla v\|_{L^2(B_1)}^2 + \|\operatorname{div}(a\nabla v)\|_{L^2(B_1)}^2 \right) \tag{3.1}$$

Proof. The proof is per absurdum. Note that since $d \leq 3$ the assumptions $v \in \mathcal{H}^1(B_1)$ and $\operatorname{div}(a\nabla v) \in L^2(B_1)$ imply the Hölder continuity of v in B_1 . Assume that (3.1) does not hold. Then there exists a sequence v_n and a sequence a'_n whose maximum and minimum eigenvalues are uniformly bounded by $\lambda_{\min}(a)$ and $\lambda_{\max}(a)$ (we need to introduce that sequence because we want the constant in (3.1) to depend only $d, \lambda_{\min}(a), \lambda_{\max}(a)$) such that

$$\|v_n - v_n(0)\|_{L^2(B_1)}^2 > n \left(\|\nabla v_n\|_{L^2(B_1)}^2 + \|\operatorname{div}(a'_n \nabla v_n)\|_{L^2(B_1)}^2 \right) \tag{3.2}$$

Letting $w_n = \frac{v_n - v_n(0)}{\|v_n - v_n(0)\|_{L^2(B_1)}}$ we obtain that $w_n(0) = 0$, $\|w_n\|_{L^2(B_1)} = 1$ and

$$\|\nabla w_n\|_{L^2(B_1)}^2 + \|\operatorname{div}(a'_n \nabla w_n)\|_{L^2(B_1)}^2 < \frac{1}{n} \tag{3.3}$$

Since

$$\|w_n\|_{\mathcal{H}^1(B_1)} < 1 + \frac{1}{n} \leq 2 \tag{3.4}$$

it follows that there exists a subsequence w_{n_j} and a $w \in \mathcal{H}^1(B_1)$ such that $w_{n_j} \rightharpoonup w$ weakly in $\mathcal{H}^1(B_1)$ and $\nabla w_{n_j} \rightharpoonup \nabla w$ weakly in $L^2(B_1)$. Using $\|\nabla w_n\|_{L^2(B_1)} \leq 1/n$ we deduce that $\nabla w = 0$ which implies that w is a constant in B_1 . Since by the Rellich–Kondrachov theorem the embedding $\mathcal{H}^1(B_1) \subset L^2(B_1)$ is compact it follows from (3.4) that $w_{n_j} \rightarrow w$ strongly in $L^2(B_1)$ which (using $\|w_n\|_{L^2(B_1)} = 1$) implies that $\|w\|_{L^2(B_1)} = 1$. Now (3.4) together with the fact that $\|\operatorname{div}(a'_n \nabla w_n)\|_{L^2(B_1)}^2$ is uniformly bounded and that $d \leq 3$ implies that w_n is uniformly Hölder continuous on $B(0, \frac{1}{2})$ (see for instance [82]). This implies that w is continuous in $B(0, \frac{1}{2})$ and that $w(0) = 0$. This contradicts the fact that w is a constant in B_1 with $\|w\|_{L^2(B_1)} = 1$. \square

We also need the following lemma.

Lemma 3.2. *Let $f \in \mathcal{H}_0^1(\Omega)$ and define Ω_{2H} as the set of points in Ω at distance at most $2H$ from $\partial\Omega$. There exists a constant $C_{d,\Omega}$ such that*

$$\|f\|_{L^2(\Omega_{2H})} \leq C_{d,\Omega} H \|\nabla f\|_{L^2(\Omega_{2H})} \tag{3.5}$$

Proof. The proof is analogous to Poincaré inequality. We observe that $u = 0$ on $\partial\Omega$. Recalling that Ω is piecewise Lipschitz continuous we cover $\partial\Omega \cup \Omega_{2H}$ with a collection of charts and express $u(x)$ as an integral from the boundary (where $u = 0$) to x of $\nabla u(y)$ along a path of length bounded by a multiple of H . \square

Theorem 3.3. *Let $f \in V_0$ (and $d \leq 3$). It holds true that*

$$\|\nabla f\|_{L^2(\Omega)} \leq CH \|\operatorname{div}(a\nabla f)\|_{L^2(\Omega)} \tag{3.6}$$

where the constant C depends only $d, \lambda_{\min}(a), \lambda_{\max}(a)$ and Ω .

Proof. We have by integration by parts and the Cauchy–Schwartz inequality,

$$\begin{aligned} \int_{\Omega} (\nabla f)^T a \nabla f &= - \int_{\Omega} f (\operatorname{div}(a\nabla f)) \\ &\leq \|f\|_{L^2(\Omega)} \|\operatorname{div}(a\nabla f)\|_{L^2(\Omega)}. \end{aligned} \tag{3.7}$$

Therefore, using Young's inequality we obtain that for any $\alpha > 0$,

$$\int_{\Omega} (\nabla f)^T a \nabla f \leq \frac{1}{2\alpha H^2} \|f\|_{L^2(\Omega)}^2 + \frac{\alpha H^2}{2} \|\operatorname{div}(a \nabla f)\|_{L^2(\Omega)}^2. \quad (3.8)$$

Now it follows from the definition (1.3) that there exists an index set $\mathcal{N}' \subset \mathcal{N}$ such that: (i) for all $i \in \mathcal{N}'$, $B(x_i, H) \subset \Omega$ (where $B(x_i, H)$ is the ball of center x_i and radius H) (ii) $\cup_{i \in \mathcal{N}'} B(x_i, H) \cup \Omega_{2H} = \Omega$ (iii) and that any $x \in \Omega$ is contained in at most K balls of $(B(x_i, H))_{i \in \mathcal{N}'}$ where K is a finite number depending only on the dimension d . It follows from (ii) that

$$\|f\|_{L^2(\Omega)}^2 \leq \sum_{i \in \mathcal{N}'} \|f\|_{L^2(B(x_i, H))}^2 + \|f\|_{\Omega_{2H}}^2. \quad (3.9)$$

Observing that $f(x_i) = 0$ we obtain from Lemma 3.1 and scaling that for each $i \in \mathcal{N}'$

$$\|f\|_{L^2(B(x_i, H))}^2 \leq C_{d, \lambda_{\min}(a), \lambda_{\max}(a)} \left(H^2 \|\nabla f\|_{L^2(B(x_i, H))}^2 + H^4 \|\operatorname{div}(a \nabla f)\|_{L^2(B(x_i, H))}^2 \right) \quad (3.10)$$

Therefore (3.10) and (3.5) imply that

$$\begin{aligned} \|f\|_{L^2(\Omega)}^2 &\leq C_{d, \Omega} H^2 \|\nabla f\|_{L^2(\Omega_{2H})}^2 \\ &+ C_{d, \lambda_{\min}(a), \lambda_{\max}(a)} \sum_{i \in \mathcal{N}'} \left(H^2 \|\nabla f\|_{L^2(B(x_i, H))}^2 + H^4 \|\operatorname{div}(a \nabla f)\|_{L^2(B(x_i, H))}^2 \right). \end{aligned} \quad (3.11)$$

Using (iii) we deduce that

$$\begin{aligned} \|f\|_{L^2(\Omega)}^2 &\leq C_{d, \Omega, \lambda_{\min}(a), \lambda_{\max}(a)} K \\ &\left(H^2 \|\nabla f\|_{L^2(\Omega)}^2 + H^4 \|\operatorname{div}(a \nabla f)\|_{L^2(\Omega)}^2 \right) \end{aligned} \quad (3.12)$$

Combining (3.12) with (3.8) we obtain that

$$\int_{\Omega} (\nabla f)^T a \nabla f \leq \frac{\alpha H^2}{2} \|\operatorname{div}(a \nabla f)\|_{L^2(\Omega)}^2 + \frac{1}{\alpha} C_{d, \Omega, \lambda_{\min}(a), \lambda_{\max}(a)} \left(\|\nabla f\|_{L^2(\Omega)}^2 + H^2 \|\operatorname{div}(a \nabla f)\|_{L^2(\Omega)}^2 \right). \quad (3.13)$$

which concludes the proof by taking $\alpha = \frac{1}{2\lambda_{\min}(a)} C_{d, \Omega, \lambda_{\min}(a), \lambda_{\max}(a)}$. \square

3.2. Accuracy of the interpolation basis

Now let us use (3.6) to estimate the interpolation error.

Corollary 3.4. *Let $u \in V$ be the solution of (1.1) and $u^{\text{in}}(x) := \sum_{i=1}^N u(x_i) \phi_i(x)$. It holds true that*

$$\|u - u^{\text{in}}\|_{\mathcal{H}_0^1(\Omega)} \leq CH \|g\|_{L^2(\Omega)} \quad (3.14)$$

where the constant C depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω .

Proof. Observing that $u - u^{\text{in}} \in V_0(\Omega)$ we deduce from Theorem 3.3 that

$$\int_{\Omega} |\nabla(u - u^{\text{in}})|^2 \leq CH^2 \int_{\Omega} \left| g - \sum_i u(x_i) \operatorname{div}(a \nabla \phi_i) \right|^2 dx \quad (3.15)$$

$$\leq CH^2 \left(\int_{\Omega} g^2 dx + \int_{\Omega} \left(\sum_i u(x_i) \operatorname{div}(a \nabla \phi_i) \right)^2 dx \right) \quad (3.16)$$

$$\leq CH^2 \left(\int_{\Omega} g^2 dx + \int_{\Omega} (\operatorname{div}(a \nabla u))^2 dx \right) \quad (3.17)$$

$$\leq CH^2 \|g\|_{L^2(\Omega)}^2 \quad (3.18)$$

Note that in the third inequality, we have used the Lemma 2.5, which states that the linear combination of ϕ_i minimizes $\|\cdot\|_V$ among all the functions in $V(\Omega)$ sharing the same values at the nodes $\{x_i\}$. Note also that through this paper, for the sake of clarity, we only keep track of the dependence of C and not its precise numerical value (we will, for instance, write $2C/\lambda_{\min}(a)$ as C to avoid cumbersome expressions). \square

3.3. Accuracy of the FEM with elements ϕ_i

The following theorem shows that the finite element method with elements ϕ_i achieves the optimal (see [15]) convergence rate $\mathcal{O}(H)$ in \mathcal{H}^1 norm in its approximation of (1.1).

Theorem 3.5. *Let u be the solution of equation (1.1), and u^H be the finite element solution of (1.1) over the linear space spanned by the elements $\{\phi_i, i = 1, \dots, N\}$ identified in (2.3). It holds true that*

$$\|u - u^H\|_{\mathcal{H}_0^1(\Omega)} \leq CH\|g\|_{L^2(\Omega)} \quad (3.19)$$

where the constant C depends only on $\lambda_{\min}(a)$ and $\lambda_{\max}(a)$.

Proof. The theorem is a straightforward consequence of Corollary 3.4 and of the fact that u^H minimizes the (squared) distance $\int_{\Omega} (\nabla u - \nabla u^H)^T a (\nabla u - \nabla u^H)$ over the linear space spanned by the elements ϕ_i . \square

Remark 3.6. Recall that the convergence rate of a FEM applied to (1.1) with piecewise linear elements can be arbitrarily bad [11]. Recall also that the convergence rate of Theorem 3.5 is optimal [15] (this is related to the Kolmogorov n -widths [61, 74, 85], which measure how accurately a given set of functions can be approximated by linear spaces of dimension n in a given norm).

4. RECOVERING u FROM PARTIAL MEASUREMENTS

In this section we will show that our method provides a natural solution to the inverse problem of recovering u from partial measurements. Consider the problem (1.1). In this inverse problem one is given the following (incomplete) information: $a(x)$ is known, g is unknown but we know that $\|g\|_{L^2(\Omega)} \leq M$, u is unknown but its values are known on a finite set of points $\{x_j : j \in \mathcal{N}\}$ (through site measurements). The inverse problem is then to recover u (accurately in the \mathcal{H}^1 norm). A solution of this inverse problem can be obtained by first computing the minimizers of (2.3) (or their localized version (7.4)) and approximate u with $u^{\text{in}}(x) := \sum_{i=1}^N u(x_i)\phi_i(x)$. Corollary 3.4 can then be used to bound the accuracy of the recovery by

$$\|u - u^{\text{in}}\|_{\mathcal{H}_0^1(\Omega)} \leq CHM \quad (4.1)$$

where the constant C depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω . Note that the method is meshless.

5. GENERALIZATION TO $d \geq 4$

Although for the sake of simplicity we have restricted our presentation to $d \leq 3$, the method and results of this paper can be generalized to $d \geq 4$ by introducing the space V^m defined as the set of functions $u \in \mathcal{H}_0^1(\Omega)$ such that $(\text{div}(a\nabla\cdot))^m u \in L^2(\Omega)$ where m is an integer $m \geq 1$ and $(\text{div}(a\nabla\cdot))^m u$ is the m -iterate of the operator $(\text{div}(a\nabla\cdot))$, i.e.

$$(\text{div}(a\nabla\cdot))^1 u := (\text{div}(a\nabla u)) \text{ and } (\text{div}(a\nabla\cdot))^m u := (\text{div}(a\nabla\cdot))^{m-1} (\text{div}(a\nabla u)). \quad (5.1)$$

Introducing $\|\cdot\|_{V^m}$ as the norm on V^m defined by

$$\|u\|_{V^m} := \left\| (\text{div}(a\nabla\cdot))^m u \right\|_{L^2(\Omega)}. \quad (5.2)$$

The results of this paper can be generalized to $d \geq 4$ by choosing m such that $(d - 1)/2 \leq m \leq d/2$, assuming that $g \in L^{2m}(\Omega)$ and using the interpolation basis defined as the minimizer of

$$\begin{cases} \text{Minimize } \|\phi\|_{V^m} \\ \text{Subject to } \phi \in V_i^m. \end{cases} \tag{5.3}$$

where

$$V_i^m := \{\phi \in V^m \mid \phi(x_i) = 1 \text{ and } \phi(x_j) = 0, \text{ for } j \in \{1, \dots, N\} \text{ such that } j \neq i\}. \tag{5.4}$$

Note that the solutions of (5.3) are $2m$ -harmonic in the sense that they satisfy

$$(\operatorname{div}(a\nabla\cdot))^{2m} \phi(x) = 0 \text{ for } x \neq x_j. \tag{5.5}$$

Note that the solutions of (2.3) are bi-harmonic on $\Omega \setminus \{x_j : j \in \mathcal{N}\}$.

For $d = 1$ one can also obtain an accurate interpolation basis by minimizing $\int_{\Omega} \nabla\phi a\nabla\phi$ subject to the pointwise interpolation constraints used in (5.4). By doing so one would obtain basis elements ϕ_i that are harmonic in $\Omega \setminus \{x_j : j \in \mathcal{N}\}$ and recover a numerical homogenization method based on “harmonic coordinates” [8, 69].

Note also that the variation formulation (5.3) does not require the operator $L := \operatorname{div}(a\nabla\cdot)$ to be self-adjoint (but solely depends on the minimization of the L^2 norm of $L^{2m}\phi$).

6. ROUGH POLYHARMONIC SPLINES

6.1. The elements ϕ_i as rough polyharmonic splines

The interpolation basis ϕ_i constructed in this work can be seen as a generalization of polyharmonic splines to differential operators with rough coefficients. More precisely, as polyharmonic splines lead to an accurate interpolation of smooth functions (solutions of the laplace operator), the “rough polyharmonic splines” introduced in this paper lead to an accurate interpolation of solutions of differential operators with rough coefficients. Note also that as polyharmonic splines are m -harmonic in the sense that they satisfy $\Delta^m\phi = 0$ away from the coarse nodes, the solutions of (5.3) are $2m$ -harmonic in the sense that they satisfy $(\operatorname{div}(a\nabla\cdot))^{2m} \phi(x) = 0$ away from those coarse nodes. This generalization is challenging in several major ways due to the lack of regularity of coefficients. In particular, the Fourier analysis toolkit is no longer available and the simple task of enforcing clamped boundary conditions (trivial with smooth coefficients) becomes a significant difficulty with rough coefficients: consider, for instance, the problem of finding a function $\phi \in \mathcal{H}^1(B(0, 1))$ that satisfies zero Neumann and Dirichlet boundary conditions on $\partial B(0, 1)$ and such that $\operatorname{div}(a\nabla\phi) \in L^2(B(0, 1))$. We will now give a short review of polyharmonic splines.

6.2. Polyharmonic splines: a short review

Let X be a discrete (possibly finite) set of points of \mathbb{R}^d . Let u be a real valued function defined on X . Polyharmonic splines have emerged through the problem of interpolating u (from its known values on X) to a real function ϕ defined on \mathbb{R}^d (with $\phi(x_j) = u(x_j)$ for $x_j \in X$). Let m be an integer strictly greater than $d/2$. The idea behind polyharmonic splines is to seek the most “regular” interpolation by selecting ϕ to be a minimizer of the semi-norm

$$\left(\int_{\mathbb{R}^d} \sum_{\alpha \in \mathbb{N}^d, |\alpha|=m} c_{\alpha} \left(\frac{\partial^{\alpha} u}{\partial x^{\alpha}}(x) \right)^2 dx \right)^{\frac{1}{2}} \tag{6.1}$$

over all functions ϕ in $\mathcal{H}^m(\mathbb{R}^d)$ interpolating u on X , where the parameters c_{α} are positive coefficients usually [25, 78] chosen to be equal to $\frac{m!}{\alpha!}$ to ensure the rotational invariance of the semi-norm (those coefficients are also sometimes chosen to be equal to one [76, 77]). Writing Δ the Laplace operator on \mathbb{R}^d ($\Delta\phi = \sum_{i=1}^d \frac{\partial^2 \phi}{\partial x_i^2}$) and Δ^m the m -iterate of ϕ ($\Delta^1 = \Delta$ and $\Delta^k\phi = \Delta(\Delta^{k-1}\phi)$) the solutions of (6.1) satisfy $\Delta^m\phi = 0$ on $\mathbb{R}^d \setminus X$ and are

therefore called m-harmonic splines. A Polyharmonic spline of order m is therefore also commonly defined [48] as a tempered distribution ϕ on \mathbb{R}^d that is $2m - d - 1$ continuously differentiable and such that $\Delta^m \phi = 0$ on $\mathbb{R}^d \setminus X$.

Polyharmonic splines can be represented *via* weighted sums of shifted fundamental solutions of Δ , (*i.e.* they can be written [24] $\phi = \sum_{x_j \in X} c_j \tau(x - x_j) + p_{m-1}(x)$ where τ is the fundamental solution of Δ^m ($\Delta^m \tau(x) = \delta(x)$) and p_{m-1} is a polynomial of degree at most $m - 1$).

When X forms a regular lattice of \mathbb{R}^d the resulting splines are referred to as polyharmonic cardinal splines. In this case (which has been extensively studied [52–54, 79]) it can be shown [75–77] that the basis elements allowing for the interpolation of u can be expressed as $\Delta_d^m \tau$ (where τ is the fundamental solutions of Δ^m and Δ_d is the finite difference discretization of Δ on \mathcal{Z}^d).

Polyharmonic splines can be traced back to the seminal work of Harder and Desmarais [43] on the interpolation of functions of two variables based on the minimization of a quadratic functional corresponding to the bending energy of a thin plate (we refer to [20] for a review) and the extension of this idea to higher dimensions in the seminal work of Duchon [24–26] (built on earlier work by Atteia [7]). We also refer to the related work of Schoenberg ($d = 1$) on cardinal spline interpolation [79], to [51] for L^∞ radial basis functions interpolation error estimates and to [57] for the rapid decay (locality) of polyharmonic splines.

7. LOCALIZATION OF THE INTERPOLATION BASIS

7.1. Introduction of the localized basis

For each $i \in \{1, \dots, N\}$ let Ω_i be an open subset of Ω containing x_i . Let $\mathcal{N}(\Omega_i)$ be the set of indices corresponding to the interior nodes of the coarse mesh contained in Ω_i , *i.e.*

$$\mathcal{N}(\Omega_i) := \{j \in \{1, \dots, N\} \mid x_j \in \Omega_i\}. \quad (7.1)$$

Define

$$V(\Omega_i) := \{\phi \in \mathcal{H}_0^1(\Omega) \mid \phi = 0 \text{ on } \partial\Omega_i \cup (\Omega \setminus \Omega_i) \text{ and } \operatorname{div}(a\nabla\phi) \in L^2(\Omega_i)\} \quad (7.2)$$

and

$$V_i(\Omega_i) := \{\phi \in V(\Omega_i) \mid \phi(x_i) = 1 \text{ and } \phi(x_j) = 0, \text{ for } j \in \mathcal{N}(\Omega_i) \text{ such that } j \neq i\} \quad (7.3)$$

Consider the following optimization problem over $V_i(\Omega_i)$:

$$\begin{cases} \text{Minimize } \int_{\Omega_i} |\operatorname{div}(a\nabla\phi)|^2 \\ \text{Subject to } \phi \in V_i(\Omega_i). \end{cases} \quad (7.4)$$

The proof of the following proposition is identical to that of Proposition 2.3 (V_i is simply replaced by $V_i(\Omega_i)$).

Proposition 7.1. $V_i(\Omega_i)$ is a non-empty closed convex affine subspace of $V(\Omega_i)$ under the norm $\|v\|_{V(\Omega_i)}^2 := \int_{\Omega_i} (\operatorname{div}(a\nabla v))^2$. Problem (7.4) is a strictly convex (quadratic) optimization problem over $V_i(\Omega_i)$ with a unique minimizer denoted by ϕ_i^{loc} .

Remark 7.2. As in Proposition 2.3, the unique minimizer of (7.4) can be represented using the Green's function of the operator $-\operatorname{div}(a\nabla \cdot)$ with Dirichlet boundary condition on $\partial\Omega_i$, *i.e.*, we have for $x \in \Omega_i$

$$\phi_i^{\text{loc}}(x) := \sum_{j \in \mathcal{N}(\Omega_i)} P_{i,j}^{i,\text{loc}} \tau^{i,\text{loc}}(x, x_j) \quad (7.5)$$

with

$$\tau^{i,\text{loc}}(x, y) := \int_{\Omega_i} G^{i,\text{loc}}(x, z) G^{i,\text{loc}}(z, y) \, dy \quad (7.6)$$

where $G^{i,\text{loc}}(x, y)$ is the Green's function of $-\text{div}(a\nabla)$ in Ω_i with zero Dirichlet boundary condition, *i.e.* the solution of (for $y \in \Omega_i$)

$$\begin{cases} -\text{div}(a(x)\nabla G^{i,\text{loc}}(x, y)) = \delta(x - y) & \text{for } x \in \Omega_i; \\ G^{i,\text{loc}}(x, y) = 0 & \text{for } x \in \partial\Omega_i, \end{cases} \quad (7.7)$$

and, writing $N_i := |\mathcal{N}(\Omega_i)|$ the number of coarse nodes x_j that are contained in Ω_i , $P^{i,\text{loc}}$ is the $N_i \times N_i$ positive definite symmetric matrix, defined over $\mathcal{N}(\Omega_i)$, by $P^{i,\text{loc}} = (\Theta^{i,\text{loc}})^{-1}$ where $\Theta^{i,\text{loc}}$ is the $N_i \times N_i$ positive definite symmetric matrix, defined over $\mathcal{N}(\Omega_i)$, by

$$\Theta_{k,j}^{i,\text{loc}} := \int_{\Omega_i} G^{i,\text{loc}}(x, x_k) G^{i,\text{loc}}(x, x_j) \, dx. \quad (7.8)$$

Remark 7.3. It is important to note that in practical (numerical) applications each (localized) element ϕ_i^{loc} is obtained by solving one (local, *i.e.* over Ω_i) quadratic optimization problem (7.4) rather than through the representation formula (7.5) (which would require solving N_i elliptic problems in Ω_i corresponding to the values of the local Green's function).

7.2. Accuracy of the localized basis as a function of the norm of the difference between the global and the localized basis

We introduce in this subsection lemmas that will be used to control the accuracy of the local basis $(\phi_i^{\text{loc}})_{i \in \mathcal{N}}$ in approximating solutions of (1.1) (*i.e.*, the error associated with replacing the global basis $(\phi_i)_{i \in \mathcal{N}}$ with a localized version $(\varphi_i)_{i \in \mathcal{N}}$). Although $(\varphi_i)_{i \in \mathcal{N}}$ is arbitrary in the following lemma, it is intended to be a localized version of the global basis.

Lemma 7.4. *Let $(\varphi_i)_{i \in \mathcal{N}}$ be N elements of $\mathcal{H}_0^1(\Omega)$. For $c \in \mathbb{R}^N$, $c \neq 0$, define*

$$v(x) := \sum_{i=1}^N c_i \phi_i(x) \quad \text{and} \quad w(x) := \sum_{i=1}^N c_i \varphi_i(x). \quad (7.9)$$

It holds true that

$$\frac{\|v - w\|_{\mathcal{H}_0^1(\Omega)}}{\|\text{div}(a\nabla v)\|_{L^2(\Omega)}} \leq CN \max_{i \in \mathcal{N}} \|\phi_i - \varphi_i\|_{\mathcal{H}_0^1(\Omega)} \quad (7.10)$$

where the constant C is finite and depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω .

Proof. We have, using (2.30),

$$\|v - w\|_{\mathcal{H}_0^1(\Omega)} = \left(\frac{c^T M c}{c^T P c} \right)^{\frac{1}{2}} \|\text{div}(a\nabla v)\|_{L^2(\Omega)} \quad (7.11)$$

where M is the $N \times N$ matrix defined by

$$M_{i,j} := \int_{\Omega} (\nabla \phi_i - \nabla \varphi_i)^T (\nabla \phi_j - \nabla \varphi_j) \quad (7.12)$$

and P is defined in (2.12). Note that since Θ is symmetric positive definite, there exists a symmetric positive definite $N \times N$ matrix T such that $\Theta = T^2$. Recalling that $P = \Theta^{-1}$, we obtain

$$\sup_{c \in \mathbb{R}^N, c \neq 0} \frac{c^T M c}{c^T P c} = \lambda_{\max}(TMT) \leq \lambda_{\max}(M) \lambda_{\max}(\Theta) \quad (7.13)$$

Using (2.5) we have for $c \in \mathbb{R}^N$

$$c^T \Theta c = \int_{\Omega} \left(\sum_{i=1}^N G(x, x_i) c_i \right)^2 dx. \quad (7.14)$$

It follows from the Cauchy–Schwartz inequality that

$$c^T \Theta c \leq \left(\sum_{i=1}^N c_i^2 \right) \left(\sum_{i=1}^N \int_{\Omega} (G(x, x_i))^2 dx \right). \quad (7.15)$$

Using (2.10), we conclude that

$$\lambda_{\max}(\Theta) \leq NK \quad (7.16)$$

where the constant K is finite and depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω . To summarize we have obtained that

$$\frac{\|v - w\|_{\mathcal{H}_0^1(\Omega)}}{\|\operatorname{div}(a \nabla v)\|_{L^2(\Omega)}} \leq CN^{1/2} (\lambda_{\max}(M))^{1/2}. \quad (7.17)$$

We now need to control $\lambda_{\max}(M)$. Since M is symmetric and positive its maximum eigenvalue is less or equal than its trace. It follows that,

$$\lambda_{\max}(M) \leq \sum_{i=1}^N M_{i,i} \leq N \max_i M_{i,i}. \quad (7.18)$$

Combining (7.18) with (7.17) we deduce (7.10). \square

The following lemma bounds the accuracy of the localized basis $(\phi_i^{\text{loc}})_{i \in \mathcal{N}}$ in approximating solutions of (1.1). In particular, it shows that if $\max_{i \in \mathcal{N}} \|\phi_i - \phi_i^{\text{loc}}\|_{\mathcal{H}_0^1(\Omega)}$ is sufficiently small then the accuracy of the localized basis is similar to that of the global basis.

Lemma 7.5. *Let u be the solution of equation (1.1), and $u^{H,\text{loc}}$ be the finite element solution of (1.1) over the linear space spanned by the elements $\{\phi_i^{\text{loc}}, i = 1, \dots, N\}$ identified in (7.4). It holds true that*

$$\|u - u^{H,\text{loc}}\|_{\mathcal{H}_0^1(\Omega)} \leq C \|g\|_{L^2(\Omega)} \left(H + N \max_{i \in \mathcal{N}} \|\phi_i - \phi_i^{\text{loc}}\|_{\mathcal{H}_0^1(\Omega)} \right) \quad (7.19)$$

where the constant C depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω .

Proof. Let u^{in} be the interpolation of the solution of (1.1) over the elements $\{\phi_i, i = 1, \dots, N\}$ identified in (2.3). Let $u^{\text{in,loc}}$ be the interpolation of the solution of (1.1) over the elements $\{\phi_i^{\text{loc}}, i = 1, \dots, N\}$ identified in (7.4). Using the triangle inequality we have

$$\|u - u^{\text{in,loc}}\|_{\mathcal{H}_0^1(\Omega)} \leq \|u - u^{\text{in}}\|_{\mathcal{H}_0^1} + \|u^{\text{in}} - u^{\text{in,loc}}\|_{\mathcal{H}_0^1(\Omega)}. \quad (7.20)$$

Using Corollary 3.4 and Lemma 7.4 we deduce that

$$\|u - u^{\text{in,loc}}\|_{\mathcal{H}_0^1(\Omega)} \leq CH \|g\|_{L^2(\Omega)} + CN \max_{i \in \mathcal{N}} \|\phi_i - \phi_i^{\text{loc}}\|_{\mathcal{H}_0^1(\Omega)} \|\operatorname{div}(a \nabla u^{\text{in}})\|_{L^2(\Omega)}. \quad (7.21)$$

Using the minimization property given in Theorem 2.5 we have

$$\|\operatorname{div}(a \nabla u^{\text{in}})\|_{L^2(\Omega)} \leq \|\operatorname{div}(a \nabla u)\|_{L^2(\Omega)} = \|g\|_{L^2(\Omega)}. \quad (7.22)$$

We deduce that

$$\|u - u^{\text{in,loc}}\|_{\mathcal{H}_0^1(\Omega)} \leq C \|g\|_{L^2(\Omega)} \left(H + N \max_{i \in \mathcal{N}} \|\phi_i - \phi_i^{\text{loc}}\|_{\mathcal{H}_0^1(\Omega)} \right). \quad (7.23)$$

We conclude the proof using the fact that $u^{H,\text{loc}}$ minimizes $\int_{\Omega} (\nabla u - \nabla v) a (\nabla u - \nabla v)$ over all v in the linear space spanned by the elements $\{\phi_i^{\text{loc}}, i = 1, \dots, N\}$. \square

7.3. Identification of the difference between the global and the localized basis

In this subsection we will represent $\phi_i - \phi_i^{\text{loc}}$ as the solution of a fourth order PDE. We will assume that $\partial\Omega_i \cap \Omega$ is at some strictly positive distance from the set of coarse nodes, *i.e.* writing

$$\delta_i := \inf_{x \in \partial\Omega_i \cap \Omega, j \in \mathcal{N}} \|x - x_j\| \quad (7.24)$$

we assume that $\delta_i > 0$. Although the construction of the localized basis φ_i^{loc} does not require δ_i to be strictly positive our proof of the accuracy of the localized basis will require that δ_i is uniformly bounded from below by some arbitrary power of H (*i.e.*, CH^k for some $C > 0$ and $k \geq 1$). More precisely, our (final and simplified) *a posteriori* error estimates (Thm. 7.15) will be given under the assumption that $\min_{i \in \mathcal{N}} \delta_i \geq H_{\min}/10$ where

$$H_{\min} := \min_{i, j \in \mathcal{N}} \|x_i - x_j\|. \quad (7.25)$$

and that the assumption that $\max_{i \in \mathcal{N}} \delta_i \leq H \leq 1$ (we also make these assumptions to simplify our expressions, note that there is no loss of generality here since Ω can be re-scaled to have a diameter bounded by 1).

Let $f \in H^{-1}(\Omega)$ and consider the system of equations with unknowns $c \in \mathbb{R}^N$ and $\chi \in V_0$ (where V_0 is defined in (2.25))

$$\begin{cases} \operatorname{div}(a\nabla(\operatorname{div}(a\nabla\chi))) = f + \sum_{i=1}^N c_i \delta(x - x_i) & \text{on } \Omega \\ \chi = \operatorname{div}(a\nabla\chi) = 0 & \text{on } \partial\Omega \\ \chi = 0 & \text{on } \{x_j : j \in \mathcal{N}\}. \end{cases} \quad (7.26)$$

Proposition 7.6. *Equation (7.26) admits a unique solution $\chi \in V_0$ given by*

$$\chi(x) = \chi^1(x) + \chi^2(x) \quad (7.27)$$

where

$$\chi^1(x) = \int_{\Omega} \tau(x, y) f(y) \, dy \quad (7.28)$$

and

$$\chi^2(x) = - \sum_{k=1}^N \sum_{j=1}^N \tau(x, x_j) P_{j,k} \chi^1(x_k). \quad (7.29)$$

Proof. Noting that τ is the fundamental solution of (2.14) we obtain that (for $c \in \mathbb{R}^N$) χ is the unique solution of

$$\begin{cases} \operatorname{div}(a\nabla(\operatorname{div}(a\nabla\chi))) = f + \sum_{i=1}^N c_i \delta(x - x_i) & \text{on } \Omega \\ \chi = \operatorname{div}(a\nabla\chi) = 0 & \text{on } \partial\Omega \end{cases} \quad (7.30)$$

if and only if

$$\chi(x) = \int_{\Omega} \tau(x, y) f(y) \, dy + \sum_{j=1}^N \tau(x, x_j) c_j. \quad (7.31)$$

The additional constraint

$$\chi = 0 \text{ on } \{x_j\}_{j \in \mathcal{N}} \quad (7.32)$$

is then equivalent to

$$\chi^1(x_i) + \sum_{j=1}^N \tau(x_i, x_j) c_j = 0 \quad (7.33)$$

which (recalling that $\tau(x_i, x_j) = \Theta_{i,j}$ and that $P = \Theta^{-1}$) admits the unique solution $c = -P(\chi^1(x_i))_{i \in \mathcal{N}}$. Note that χ^1 has well defined values at the nodes $(x_j)_{j \in \mathcal{N}}$ since, by [36, 81], it is Hölder continuous over Ω . \square

Consider the equation

$$\begin{cases} \operatorname{div} \left(a \nabla \left(\operatorname{div} (a \nabla \chi) \right) \right) = - \operatorname{div} \left(a \nabla \left(\operatorname{div} (a \nabla \phi_i^{\text{loc}}) \right) \right) + \sum_{i=1}^N c_i \delta(x - x_i) \text{ on } \Omega \\ \chi = \operatorname{div} (a \nabla \chi) = 0 \text{ on } \partial \Omega \\ \chi = 0 \text{ on } \{x_j : j \in \mathcal{N}\}. \end{cases} \tag{7.34}$$

We will show in the proof of Lemma 7.7 that (7.34) admits a unique solution. We will denote that solution by χ_i and obtain (in Lem. 7.7) a representation formula for it.

Lemma 7.7. *It holds true that*

$$\phi_i = \phi_i^{\text{loc}} + \chi_i \tag{7.35}$$

where $\chi_i = \chi_i^1 + \chi_i^2$ with (n is the unit surface vector oriented towards the exterior of Ω_i)

$$\chi_i^1(x) := \int_{\partial \Omega_i \cap \Omega} \tau(x, y) n \cdot a \nabla \left(\operatorname{div} (a \nabla \phi_i^{\text{loc}}) \right) dy - \int_{\partial \Omega_i \cap \Omega} G(x, y) n \cdot a \nabla \phi_i^{\text{loc}} dy \tag{7.36}$$

and

$$\chi_i^2(x) := - \sum_{k=1}^N \sum_{j=1}^N \tau(x, x_j) P_{j,k} \chi_i^1(x_k). \tag{7.37}$$

Proof.

Initial proof. Let \bar{f} be an arbitrary element of $\mathcal{H}^{-1}(\Omega)$ and $\bar{\chi}$ the solution of (7.30) and (7.32). We have (using $\phi_i - \phi_i^{\text{loc}} = 0$ on $\{x_j : j \in \mathcal{N}\}$)

$$\int_{\Omega} (\phi_i - \phi_i^{\text{loc}}) \bar{f} = \int_{\Omega} (\phi_i - \phi_i^{\text{loc}}) \operatorname{div} \left(a \nabla \left(\operatorname{div} (a \nabla \bar{\chi}) \right) \right). \tag{7.38}$$

Integrating by parts we deduce that

$$\int_{\Omega} (\phi_i - \phi_i^{\text{loc}}) \bar{f} = \int_{\Omega \setminus \partial \Omega_i} \operatorname{div} \left(a \nabla (\phi_i - \phi_i^{\text{loc}}) \right) \operatorname{div} (a \nabla \bar{\chi}) + \int_{\partial \Omega_i \cap \Omega} \operatorname{div} (a \nabla \bar{\chi}) n \cdot a \nabla \phi_i^{\text{loc}}. \tag{7.39}$$

Integrating by parts again (using the continuity of $\operatorname{div} (a \nabla \phi_i^{\text{loc}})$ across $\partial \Omega_i \cap \Omega$ and the fact that $\bar{\chi}$ is null on the nodes x_j) we have

$$\int_{\Omega \setminus \partial \Omega_i} \operatorname{div} \left(a \nabla (\phi_i - \phi_i^{\text{loc}}) \right) \operatorname{div} (a \nabla \bar{\chi}) = \int_{\partial \Omega_i \cap \Omega} \bar{\chi} n \cdot a \nabla \operatorname{div} \left(a \nabla \phi_i^{\text{loc}} \right). \tag{7.40}$$

Replacing $\bar{\chi}$ by the solution expressed in (7.27), *i.e.*

$$\bar{\chi}(x) = \int_{\Omega} \tau(x, y) \bar{f}(y) dy - \sum_{k=1}^N \sum_{j=1}^N \tau(x, x_j) P_{j,k} \int_{\Omega} \tau(x_k, y) \bar{f}(y) dy \tag{7.41}$$

we obtain that

$$\begin{aligned} \int_{\Omega} (\phi_i - \phi_i^{\text{loc}}) \bar{f} &= - \int_{y \in \Omega} \int_{x \in \partial \Omega_i \cap \Omega} G(x, y) \bar{f}(y) n \cdot a \nabla \phi_i^{\text{loc}}(x) dy dx \\ &\quad + \int_{y \in \Omega} \int_{x \in \partial \Omega_i \cap \Omega} \tau(x, y) \bar{f}(y) n \cdot a \nabla \operatorname{div} \left(a \nabla \phi_i^{\text{loc}}(x) \right) dy dx \\ &\quad - \sum_{k=1}^N \sum_{j=1}^N P_{j,k} \int_{y \in \Omega} \int_{x \in \partial \Omega_i \cap \Omega} \tau(x, x_j) \tau(x_k, y) \bar{f}(y) n \cdot a \nabla \operatorname{div} \left(a \nabla \phi_i^{\text{loc}}(x) \right) dy dx \\ &\quad + \sum_{k=1}^N \sum_{j=1}^N P_{j,k} \int_{y \in \Omega} \int_{x \in \partial \Omega_i \cap \Omega} G(x, x_j) \tau(x_k, y) \bar{f}(y) n \cdot a \nabla \phi_i^{\text{loc}}(x) dy dx \end{aligned} \tag{7.42}$$

which leads us to (7.35).

Alternate proof. For the sake of clarity we will now also give an intuitive proof of (7.35) based on a direct application of Proposition 7.6. Note that since ϕ_i^{loc} and $\text{div}(a\nabla\phi_i^{\text{loc}})$ are continuous across $\partial\Omega_i \cap \Omega$ we have (using the representation formulae (2.15) and (7.5)) that

$$\text{div} \left(a\nabla \left(\text{div} (a\nabla(\phi_i - \phi_i^{\text{loc}})) \right) \right) = f + \sum_{i=1}^N c_i \delta(x - x_i) \quad (7.43)$$

where

$$f = n \cdot a\nabla(\text{div}(a\nabla\phi_i^{\text{loc}}))\delta_{\partial\Omega_i \cap \Omega} + \text{div} \left(a\nabla(n \cdot a\nabla\phi_i^{\text{loc}}\delta_{\partial\Omega_i \cap \Omega}) \right) \quad (7.44)$$

and $\delta_{\partial\Omega_i \cap \Omega}$ is the Dirac (surface) distribution on $\partial\Omega_i \cap \Omega$. We conclude by using Proposition 7.6 and integrating by parts. Note that although Proposition 7.6 requires $f \in \mathcal{H}^{-1}(\Omega)$, our initial proof shows that it can still be applied here (with $f \in \mathcal{H}^{-3}(\Omega)$) to show (7.35). Note that our assumption that $\partial\Omega_i \cap \Omega$ (the support of f) is at some strictly positive distance from the coarse nodes $(x_j)_{j \in \mathcal{N}}$ (i.e., $\delta_i > 0$) implies χ_i^1 has well defined values at the coarse nodes $(x_j)_{j \in \mathcal{N}}$ (since $\text{div}(a\nabla\chi_i^1)$ is in \mathcal{H}^1 in a neighborhood of those nodes). \square

7.4. Reverse Poincaré inequality

The following lemma is a generalization of the classical Caccioppoli's inequality (usually referred to as the reversed Poincaré inequality, see [35] and Prop. 1.4.1 of [64]). We will remind its statement and proof for the sake of completeness.

Lemma 7.8. *Let D_1, D_2 be open subsets of D_0 such that $D_2 \subset D_1$ (and such that $D_0 \setminus D_1$ is non-empty). If $v \in \mathcal{H}^1(D_0)$ satisfies $\text{div}(a\nabla v) = 0$ in D_1 , and $v = 0$ on ∂D_0 , then*

$$\|\nabla v\|_{L^2(D_2)} \leq \frac{C}{\text{dist}(D_2, D_0 \setminus D_1)} \|v\|_{L^2(D_1)} + \left(\|\text{div}(a\nabla v)\|_{L^2(D_1)} \|v\|_{L^2(D_1)} \right)^{\frac{1}{2}}. \quad (7.45)$$

Proof. The idea of the proof is as follows, choose η to be a differentiable function on D_0 such that $\eta = 1$ on D_2 , $\eta = 0$ on $D_0 \setminus D_1$, $0 \leq \eta \leq 1$ and $\|\nabla\eta\|_{L^\infty(D_0)} \leq \frac{C}{\text{dist}(D_2, D_0 \setminus D_1)}$. Using the fact that $\eta^2 v = 0$ on ∂D_1 we obtain that

$$\int_{D_1} \nabla(\eta^2 v) a \nabla v = - \int_{D_1} \eta^2 v \text{div}(a \nabla v) \quad (7.46)$$

which leads to

$$\int_{D_1} \eta^2 \nabla v a \nabla v = -2 \int_{D_1} \eta v \nabla \eta a \nabla v - \int_{D_1} \eta^2 v \text{div}(a \nabla v). \quad (7.47)$$

Using the Cauchy–Schwartz inequality, the uniform bound on $\nabla\eta$, and the uniform bound on a we deduce that

$$\int_{D_1} \eta^2 \nabla v a \nabla v \leq \frac{C}{\text{dist}(D_2, D_0 \setminus D_1)} \left(\int_{D_1} \eta^2 \nabla v a \nabla v \right)^{\frac{1}{2}} \|v\|_{L^2(D_1)} + \|\text{div}(a \nabla v)\|_{L^2(D_1)} \|v\|_{L^2(D_1)}. \quad (7.48)$$

Writing $(\int_{D_1} \eta^2 \nabla v a \nabla v)^{\frac{1}{2}}$, we conclude by using the fact that, for positive x, y, z , $x \leq y + z/x$ implies that $x \leq y + \sqrt{z}$ and also using the fact that $\int_{D_2} \nabla v a \nabla v \leq \int_{D_1} \eta^2 \nabla v a \nabla v$. \square

7.5. Bound on the maximum eigenvalue of P

In this subsection we will provide an upper bound on the maximum eigenvalue of P as defined in (2.12).

Proposition 7.9. *Let Θ be defined as in (2.5), P as in (2.12), and H_{\min} as in (7.25). It holds true that*

$$\lambda_{\max}(P) \leq C \frac{N}{H_{\min}^4}. \quad (7.49)$$

Proof. The definition (2.12) implies that the maximum eigenvalue of P is the inverse of the minimum eigenvalue of Θ as defined in (2.5). Let $l \in \mathbb{R}^N$ such that $\|l\| = 1$. We wish to bound from below $l^T \Theta l$. Write

$$v(x) := \sum_{j=1}^N G(x, x_j) l_j \quad (7.50)$$

and observe that

$$l^T \Theta l = \|v\|_{L^2(\Omega)}^2. \quad (7.51)$$

Since $\|l\| = 1$, there exists $i \in \mathcal{N}$ such that $|l_i| \geq 1/\sqrt{N}$. Let B_ρ be the intersection of Ω with the ball of center x_i and radius ρ . Let $\rho = H_{\min}/8$. Observing that v is harmonic in $B_{7\rho} \setminus B_\rho$ we have using Caccioppoli's inequality (see Lem. 7.8) that

$$\|\nabla v\|_{L^2(B_{5\rho} \setminus B_{4\rho})} \leq \frac{C}{\rho} \|v\|_{L^2(B_{6\rho} \setminus B_{3\rho})}. \quad (7.52)$$

Using Green's identity and (7.50) we have

$$\int_{\partial B_{5\rho}} n \cdot a \nabla v = \int_{B_{5\rho}} \operatorname{div}(a \nabla v) = -l_i. \quad (7.53)$$

Let η be a smooth function equal to one on $\partial B_{5\rho}$, zero on $B_{4\rho}$ and such that $0 \leq \eta \leq 1$ and $\|\nabla \eta\|_{L^\infty(B_{5\rho})} \leq C/\rho$. Integrating by parts, we have

$$\begin{aligned} \int_{\partial B_{5\rho}} n \cdot a \nabla v &= \int_{\partial B_{5\rho}} \eta n \cdot a \nabla v \\ &= \int_{B_{5\rho} \setminus B_{4\rho}} \nabla \eta a \nabla v. \end{aligned} \quad (7.54)$$

Observing that

$$\left| \int_{B_{5\rho} \setminus B_{4\rho}} \nabla \eta a \nabla v \right| \leq \frac{C}{\rho} \|\nabla v\|_{L^2(B_{5\rho} \setminus B_{4\rho})} \quad (7.55)$$

we obtain from $|l_i| \geq 1/\sqrt{N}$, (7.53) and (7.54) that

$$\frac{1}{\sqrt{N}} \leq \frac{C}{\rho} \|\nabla v\|_{L^2(B_{5\rho} \setminus B_{4\rho})}. \quad (7.56)$$

Combining (7.56) with (7.52) we deduce that

$$C \frac{\rho^2}{\sqrt{N}} \leq \|v\|_{L^2(B_{6\rho} \setminus B_{3\rho})}. \quad (7.57)$$

Combining (7.57) with (7.51) we deduce that

$$l^T \Theta l \geq C \frac{\rho^4}{N} \quad (7.58)$$

which concludes the proof. \square




FIGURE 1. Ω_i and Ω_i^δ as defined in (7.62).

7.6. Pointwise estimates on solutions of elliptic equations with discontinuous coefficients

Write $B(x, \rho)$ the Euclidean ball of center x and radius ρ and $\Omega(x, \rho)$ the intersection of $B(x, \rho)$ with Ω .

Lemma 7.10. *Let $v \in \mathcal{H}^1(\Omega)$ such that $\operatorname{div}(a\nabla v) \in L^2(\Omega)$, $d \leq 3$ and $x_0 \in \Omega$. Then there exists a constant C depending only on $\lambda_{\min}(a)$ and $\lambda_{\max}(a)$ such that*

$$\|v\|_{L^\infty(\Omega(x_0, \rho))} \leq \left(\frac{C}{\rho^d} \int_{\Omega(x_0, 2\rho)} v^2 \right)^{\frac{1}{2}} + C\rho^{2-\frac{d}{2}} \|\operatorname{div}(a\nabla v)\|_{L^2(\Omega(x_0, \rho))}. \tag{7.59}$$

Proof. Lemma 7.10 is classical and we refer to [82] for its proof. The idea is to decompose v as $v = v_1 + v_2$ where v_1 and v_2 satisfy $\operatorname{div}(a\nabla v_1) = \operatorname{div}(a\nabla v)$ and $\operatorname{div}(a\nabla v_2) = 0$ on $\Omega(x_0, \rho)$ with boundary conditions $v_1 = 0$ and $v_2 = v$ on $\partial\Omega(x_0, \rho)$. By the estimates on the L^2 -norm of the Green’s functions given in Lemma 2.2 (see also [36, 81, 82])

$$\|v_1\|_{L^\infty(\Omega(x_0, \rho))} \leq C\rho^{2-\frac{d}{2}} \|\operatorname{div}(a\nabla v)\|_{L^2(\Omega(x_0, \rho))}. \tag{7.60}$$

Since v_2 is harmonic we have (using for instance Thm. 5.1 of [82]),

$$\|v_2\|_{L^\infty(\Omega(x_0, \rho))} \leq C\|v_2\|_{L^2(\Omega(x_0, 2\rho))}. \tag{7.61}$$

□

7.7. Control of the norm of the difference between the global and the localized basis

Write $\operatorname{dist}(x, A)$ the distance from a point x to a set A in Euclidean norm. Define for $\delta > 0$

$$\Omega_i^\delta := \{x \in \Omega_i \mid \operatorname{dist}(x, \partial\Omega_i \cap \Omega) < \delta\}. \tag{7.62}$$

See Figure 1 for an illustration of Ω_i and ω_i^δ when $\partial\Omega_i \cap \partial\Omega \neq \emptyset$.

Lemma 7.11. *We have for $\delta_i \leq 1$*

$$\|\chi_i^1\|_{L^\infty(\Omega)} \leq C\delta_i^{-2} \|\operatorname{div}(a\nabla\phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + C\delta_i^{-\frac{5}{2}} \|\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)}. \tag{7.63}$$

Proof. Let η be a smooth function equal to 1 on $\bar{\Omega}_i^{\frac{\delta_i}{4}}$, 0 on $\Omega_i \setminus \Omega_i^{\frac{\delta_i}{2}}$ and such that $0 \leq \eta \leq 1$ and such that

$$\sup_{x \in \Omega_i} |\nabla \eta(x)| \leq \frac{C}{\delta_i} \quad (7.64)$$

for some constant C independent from H and δ_i . Let χ_i^1 be the function defined in (7.36). Noting that $\eta = 1$ on $\partial\Omega_i \cap \Omega$, $\tau(x, y) = 0$ and $G(x, y) = 0$ for $y \in \partial\Omega$ we have

$$\chi_i^1(x) = \int_{\partial\Omega_i} \tau(x, y) n \cdot \eta a \nabla(\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \, dy - \int_{\partial\Omega_i} G(x, y) n \cdot \eta a \nabla \phi_i^{\text{loc}} \, dy. \quad (7.65)$$

We deduce, after integration by parts and using the fact that $\operatorname{div}(a \nabla(\operatorname{div}(a \nabla \phi_i^{\text{loc}}))) = 0$ on $\Omega_i^{\frac{\delta_i}{2}}$, that

$$\begin{aligned} \chi_i^1(x) &= \int_{\Omega_i} \nabla \tau(x, y) \eta a \nabla(\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \, dy + \int_{\Omega_i} \tau(x, y) \nabla \eta a \nabla(\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \, dy \\ &\quad - \int_{\Omega_i} \nabla G(x, y) \eta a \nabla \phi_i^{\text{loc}} \, dy - \int_{\Omega_i} G(x, y) \nabla \eta a \nabla \phi_i^{\text{loc}} \, dy \\ &\quad - \int_{\Omega_i} G(x, y) \eta \operatorname{div}(a \nabla \phi_i^{\text{loc}}) \, dy. \end{aligned} \quad (7.66)$$

Noting (by integration by parts) that

$$\begin{aligned} - \int_{\Omega_i} \nabla G(x, y) \eta a \nabla \phi_i^{\text{loc}} \, dy &= -\phi_i^{\text{loc}}(x) \eta(x) + \int_{\Omega_i} \nabla G(x, y) a \nabla \eta \phi_i^{\text{loc}} \, dy \\ &\leq |\phi_i^{\text{loc}}(x) \eta(x)| + C \|\nabla \eta \phi_i^{\text{loc}}\|_{L^\infty(\Omega_i^{\frac{\delta_i}{2}})} \|\nabla G(x, y)\|_{L^1(\Omega_i^{\frac{\delta_i}{2}})} \end{aligned} \quad (7.67)$$

and observing that the L^2 norms of $\nabla \tau(x, y)$, $\tau(x, y)$, $G(x, y)$ and the L^1 norm of $\nabla G(x, y)$ are bounded by a constant C depending only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω (see proof of Lem 2.2, Eq. (2.13), [42, 84] and references therein) we deduce that

$$\begin{aligned} \|\chi_i^1\|_{L^\infty(\Omega)} &\leq C \delta_i^{-1} \left\| \nabla(\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right\|_{L^2(\Omega_i^{\frac{\delta_i}{2}})} + C \delta_i^{-1} \|\nabla \phi_i^{\text{loc}}\|_{L^2(\Omega_i^{\frac{\delta_i}{2}})} \\ &\quad + C \|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2(\Omega_i^{\frac{\delta_i}{2}})} + C \delta_i^{-1} \|\phi_i^{\text{loc}}\|_{L^\infty(\Omega_i^{\frac{\delta_i}{2}})}. \end{aligned} \quad (7.68)$$

Since $(\operatorname{div}(a \nabla \phi_i^{\text{loc}}))$ is harmonic on $\Omega_i^{\frac{3\delta_i}{4}}$ we obtain from Caccioppoli's inequality (*i.e.* Lem. 7.8 with $D_0 = \Omega_i$, $D_2 = \Omega_i^{\frac{\delta_i}{2}}$ and $D_1 = \Omega_i^{\frac{3\delta_i}{4}}$) that

$$\left\| \nabla(\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right\|_{L^2(\Omega_i^{\frac{\delta_i}{2}})} \leq C \delta_i^{-1} \|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2(\Omega_i^{\frac{3\delta_i}{4}})}. \quad (7.69)$$

Therefore

$$\|\chi_i^1\|_{L^\infty(\Omega)} \leq C \delta_i^{-2} \|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2(\Omega_i^{\frac{3\delta_i}{4}})} + C \delta_i^{-1} \|\nabla \phi_i^{\text{loc}}\|_{L^2(\Omega_i^{\frac{\delta_i}{2}})} + C \delta_i^{-1} \|\phi_i^{\text{loc}}\|_{L^\infty(\Omega_i^{\frac{\delta_i}{2}})}. \quad (7.70)$$

Similarly, using the reverse Poincaré inequality (Lem. 7.8 with $D_0 = \Omega_i$, $D_2 = \Omega_i^{\frac{\delta_i}{2}}$ and $D_1 = \Omega_i^{\frac{3\delta_i}{4}}$) and Young's inequality ($|ab| \leq \frac{\epsilon}{2}a^2 + \frac{2}{\epsilon}b^2$) we obtain that

$$\|\nabla\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \leq C\delta_i^{-1}\|\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + \delta_i\|\text{div}(a\nabla\phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)}. \quad (7.71)$$

Therefore

$$\|\chi_i^1\|_{L^\infty(\Omega)} \leq C\delta_i^{-2}\|\text{div}(a\nabla\phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + C\delta_i^{-2}\|\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + C\delta_i^{-1}\|\phi_i^{\text{loc}}\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)}. \quad (7.72)$$

Now to control $\|\phi_i^{\text{loc}}\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)}$ note that on $\Omega_i^{\frac{\delta_i}{2}}$, ϕ_i^{loc} can be decomposed as $\phi_i^{\text{loc}} = v_1 + v_2$ where v_1 and v_2 satisfy $\text{div}(a\nabla v_1) = \text{div}(a\nabla\phi_i^{\text{loc}})$ and $\text{div}(a\nabla v_2) = 0$ on $\Omega_i^{\frac{\delta_i}{2}}$ with boundary conditions $v_1 = 0$ and $v_2 = \phi_i^{\text{loc}}$ on $\partial\Omega_i^{\frac{\delta_i}{2}}$. We therefore obtain that

$$\|\phi_i^{\text{loc}}\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \leq \|v_1\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)} + \|v_2\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)}. \quad (7.73)$$

By [36, 81] (one can also use the bounds on the Green's function given in the proof of Lem. 2.2) we have

$$\|v_1\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \leq C\|\text{div}(a\nabla\phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}. \quad (7.74)$$

Since v_2 is harmonic we have,

$$\|v_2\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \leq C\|\phi_i^{\text{loc}}\|_{L^\infty(\partial\Omega_i^{\frac{\delta_i}{2}})}. \quad (7.75)$$

Using Lemma 7.10 we obtain that for $x_0 \in \partial\Omega_i^{\frac{\delta_i}{2}} \cap \Omega_i$ and $\rho = \delta_i/8$,

$$\|\phi_i^{\text{loc}}\|_{L^\infty(B(x_0, \rho) \cap \Omega_i)} \leq \left(\frac{C}{\rho^d} \int_{B(x_0, 2\rho) \cap \Omega_i} (\phi_i^{\text{loc}})^2\right)^{\frac{1}{2}} + C\rho^{2-\frac{d}{2}}\|\text{div}(a\nabla\phi_i^{\text{loc}})\|_{L^2(B(x_0, \rho) \cap \Omega_i)}. \quad (7.76)$$

Combining (7.76) with (7.75) we deduce that

$$\|v_2\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \leq C\delta_i^{-\frac{d}{2}}\|\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + \delta_i^{2-\frac{d}{2}}\|\text{div}(a\nabla\phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)}. \quad (7.77)$$

Using (7.73) we deduce that (for $\delta_i \leq 1$ and $d \leq 3$)

$$\|\phi_i^{\text{loc}}\|_{L^\infty\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \leq C\delta_i^{-\frac{d}{2}}\|\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + C\|\text{div}(a\nabla\phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)}. \quad (7.78)$$

Combining (7.78) and (7.72) we deduce (7.63) for $\delta_i \leq 1$. \square

Lemma 7.12. *Let χ_i^2 be defined as in (7.37). We have*

$$\|\nabla\chi_i^2\|_{L^2(\Omega)} \leq N\|\chi_i^1\|_{L^\infty(\Omega)} \max_{j \in \mathcal{N}} \sqrt{P_{jj}}. \quad (7.79)$$

Proof. We have

$$\chi_i^2(x) := - \sum_{k=1}^N \phi_k(x) \chi_i^1(x_k). \quad (7.80)$$

Using

$$\|\nabla \phi_k\|_{L^2(\Omega)} \leq C \|\operatorname{div}(a \nabla \phi_k)\|_{L^2(\Omega)} \quad (7.81)$$

we deduce that

$$\|\nabla \chi_i^2\|_{L^2(\Omega)} \leq C \sum_{k=1}^N \sqrt{P_{kk}} |\chi_i^1(x_k)|_{L^\infty(\Omega)}. \quad (7.82)$$

□

Lemma 7.13. *We have for $0 < \delta_i \leq 1$*

$$\|\nabla \chi_i^1\|_{L^2(\Omega)} \leq C \delta_i^{-3} \left(\|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + \|\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} \right). \quad (7.83)$$

Proof. Using

$$\int_{\Omega} \nabla v a \nabla v = - \int_{\Omega} v \operatorname{div}(a \nabla v) \quad (7.84)$$

we obtain from (7.36) that

$$\int_{\Omega} \nabla \chi_i^1 a \nabla \chi_i^1 \leq I_1 + I_2 \quad (7.85)$$

with

$$I_1 = \int_{x \in \Omega} \int_{y \in \partial \Omega_i \cap \Omega} \chi_i^1(x) G(x, y) n \cdot a \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \, dy \quad (7.86)$$

and

$$I_2 = - \int_{\partial \Omega_i \cap \Omega} \chi_i^1(y) n \cdot a \nabla \phi_i^{\text{loc}} \, dy. \quad (7.87)$$

Defining η as in the proof of Lemma 7.11 we have (since $\eta = 1$ on $\partial \Omega_i \cap \Omega$ and $\chi_i^1 = 0$ on $\partial \Omega$)

$$I_2 = - \int_{\partial \Omega_i} \chi_i^1(y) \eta n \cdot a \nabla \phi_i^{\text{loc}} \, dy. \quad (7.88)$$

Integrating by parts, we obtain

$$I_2 = - \int_{\Omega_i} \eta \chi_i^1 \operatorname{div}(a \nabla \phi_i^{\text{loc}}) - \int_{\Omega_i} \eta \nabla \chi_i^1 a \nabla \phi_i^{\text{loc}} - \int_{\Omega_i} \chi_i^1 \nabla \eta a \nabla \phi_i^{\text{loc}}. \quad (7.89)$$

Using Young's inequality we obtain

$$\left| \int_{\Omega_i} \eta \nabla \chi_i^1 a \nabla \phi_i^{\text{loc}} \right| \leq \frac{1}{4} \int_{\Omega} \nabla \chi_i^1 a \nabla \chi_i^1 + C \|\nabla \phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}^2. \quad (7.90)$$

Therefore

$$\begin{aligned} |I_2| &\leq \|\chi_i^1\|_{L^\infty(\Omega)} \|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)} + C \delta_i^{-1} \|\chi_i^1\|_{L^\infty(\Omega)} \|\nabla \phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \\ &\quad + C \|\nabla \phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}^2 + \frac{1}{4} \int_{\Omega} \nabla \chi_i^1 a \nabla \chi_i^1. \end{aligned} \quad (7.91)$$

Similarly we have

$$I_1 = \int_{x \in \Omega} \int_{y \in \partial \Omega_i} \chi_i^1(x) G(x, y) \eta n \cdot a \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \, dy \quad (7.92)$$

which, after integration by parts, leads us to

$$\begin{aligned} I_1 &= \int_{x \in \Omega} \int_{y \in \Omega_i} \chi_i^1(x) G(x, y) \eta \operatorname{div} \left(a \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right) \, dy \\ &\quad + \int_{x \in \Omega} \int_{y \in \Omega_i} \chi_i^1(x) G(x, y) \nabla \eta a \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \, dy \\ &\quad + \int_{x \in \Omega} \int_{y \in \Omega_i} \chi_i^1(x) \nabla_y G(x, y) \eta a \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \, dy. \end{aligned} \quad (7.93)$$

Using the fact that $\eta \operatorname{div} \left(a \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right) = 0$ (since the support of η does not contain any coarse nodes), and using the bounds the L^2 -norm of G (see proof of Lem. 2.2) as well as

$$\int_{y \in \Omega_i} \left(\int_{x \in \Omega} \chi_i^1(x) \nabla_y G(x, y) \right)^2 \leq C \|\chi_i^1\|_{L^\infty(\Omega)}^2 \quad (7.94)$$

we obtain that

$$I_1 \leq C(1 + \delta_i^{-1}) \|\chi_i^1\|_{L^\infty(\Omega)} \left\| a \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}. \quad (7.95)$$

Combining (7.85), (7.91) and (7.95), we have obtained that

$$\begin{aligned} \int_{\Omega} \nabla \chi_i^1 a \nabla \chi_i^1 &\leq C \|\chi_i^1\|_{L^\infty(\Omega)} \left\| \operatorname{div}(a \nabla \phi_i^{\text{loc}}) \right\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)} + C \|\nabla \phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}^2 \\ &\quad + C(1 + \delta_i^{-1}) \|\chi_i^1\|_{L^\infty(\Omega)} \left\| \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \\ &\quad + C \delta_i^{-1} \|\chi_i^1\|_{L^\infty(\Omega)} \|\nabla \phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}. \end{aligned} \quad (7.96)$$

Therefore, using Young's inequality, we have obtained that

$$\begin{aligned} \|\nabla \chi_i^1\|_{L^2(\Omega)} &\leq C(1 + \delta_i^{-1}) \|\chi_i^1\|_{L^\infty(\Omega)} + \left\| \operatorname{div}(a \nabla \phi_i^{\text{loc}}) \right\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)} + C \|\nabla \phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)} \\ &\quad + \left\| \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}. \end{aligned} \quad (7.97)$$

We deduce (7.83) by using (7.63), (7.69) and (7.71) to bound $\|\chi_i^1\|_{L^\infty(\Omega)}$, $\|\nabla \phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}$ and $\left\| \nabla (\operatorname{div}(a \nabla \phi_i^{\text{loc}})) \right\|_{L^2\left(\Omega_i^{\frac{\delta_i}{2}}\right)}$ in (7.97). \square

Combining Lemmas 7.7, 7.12 and 7.13 with Proposition 7.9 we deduce the following lemma.

Lemma 7.14. *For $0 < \delta_i \leq 1$, it holds true that*

$$\|\nabla(\phi_i - \phi_i^{\text{loc}})\|_{L^2(\Omega)} \leq CN^2 H_{\min}^{-4} \delta_i^{-3} \left(\left\| \operatorname{div}(a \nabla \phi_i^{\text{loc}}) \right\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} + \|\phi_i^{\text{loc}}\|_{L^2\left(\Omega_i^{\frac{3\delta_i}{4}}\right)} \right). \quad (7.98)$$

7.8. A posteriori error estimates

Theorem 7.15. Let u be the solution of equation (1.1), and $u^{H,\text{loc}}$ be the finite element solution of (1.1) over the linear space spanned by the elements $\{\phi_i^{\text{loc}}, i = 1, \dots, N\}$ identified in (7.4). Assume that $\min_{i \in \mathcal{N}} \delta_i \geq H_{\min}/10$ (where δ_i is defined in (7.24) and H_{\min} in (7.25)) and that $\max_{i \in \mathcal{N}} \delta_i \leq H \leq 1$. It holds true that

$$\|u - u^{H,\text{loc}}\|_{\mathcal{H}_0^1(\Omega)} \leq C \|g\|_{L^2(\Omega)} (H + E) \quad (7.99)$$

where the constant C depends only on $\lambda_{\min}(a)$, $\lambda_{\max}(a)$ and Ω , and

$$E = H_{\min}^{-7-3d} \max_{i \in \mathcal{N}} \left(\|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2(\Omega_i^H)} + \|\phi_i^{\text{loc}}\|_{L^2(\Omega_i^H)} \right) \quad (7.100)$$

where Ω_i^H is defined in (7.62).

Proof.

Theorem 7.15 is a straightforward consequence of Lemmas 7.14 and 7.5 and the inequality $N \leq CH_{\min}^{-d}$. \square

Remark 7.16. Observe that there is no loss of generality in requiring that $\min_{i \in \mathcal{N}} \delta_i \geq H_{\min}/10$ and that $\max_{i \in \mathcal{N}} \delta_i \leq H \leq 1$. Note also that $\min_{i \in \mathcal{N}} \delta_i \geq H_{\min}/10$ corresponds to the condition that the minimum distance between $\partial\Omega_i \cap \Omega$ and the set of coarse nodes is at least $H_{\min}/10$. Although we need this assumption for the simplicity of the proof, numerical experiments suggest that it is not needed for convergence with an optimal rate.

Remark 7.17. Numerical experiments show that $\|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2(\Omega_i^H)} + \|\phi_i^{\text{loc}}\|_{L^2(\Omega_i^H)}$ decays exponentially fast as a function of n_i where n_i is the minimum number of (coarse) edges (*i.e.*, number of layers of triangles) separating x_i from $\partial\Omega_i \cap \Omega$. Hence if n_i is of the order of $\ln(1/H)$ we can derive *a posteriori* error estimates from the posterior observation that $E \leq H$. *A priori* estimates can be derived by bounding the exponential decay of $\|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2(\Omega_i^H)} + \|\phi_i^{\text{loc}}\|_{L^2(\Omega_i^H)}$ as a function of the number of layers of triangles surrounding each node x_i (work in progress). These *a priori* error estimates would guarantee $\|u - u^{H,\text{loc}}\|_{\mathcal{H}_0^1(\Omega)} \leq C \|g\|_{L^2(\Omega)} H$ provided that $\min_{i \in \mathcal{N}} \operatorname{dist}(x_i, \partial\Omega_i \cap \Omega) \geq C^* H \ln \frac{1}{H}$ for some constant C^* depending on $\lambda_{\min}(a)$ and $\lambda_{\max}(a)$. We refer to this property as *super-localization*. Note that each (super-localized) basis function ϕ_i^{loc} is the solution of a sparse/banded/nearly diagonal linear system. This property is essential in keeping the computational cost minimal (see Sect. 10.3). Note also that since Ω_i^H is defined as a layer of thickness H near the portion of the boundary of Ω_i that is contained in Ω (see Fig. 1) Ω_i^H remains at distance $\mathcal{O}(n_i H)$ from x_i even when x_i is close to the boundary of Ω (and there are no “boundary effects” impacting the accuracy of the method).

Note also that because of the exponential decay of $\|\operatorname{div}(a \nabla \phi_i^{\text{loc}})\|_{L^2(\Omega_i^H)} + \|\phi_i^{\text{loc}}\|_{L^2(\Omega_i^H)}$ we do not require H of the same order as H_{\min} . In particular, one can increase interpolation points in local regions where high accuracy is required without loss of accuracy provided that H_{\min} remains bounded from below by a power of H .

8. ON TIME DEPENDENT PROBLEMS

As shown in Sections 4 and 5 of [70] the accuracy of the basis elements ϕ_i and ϕ_i^{loc} remains unchanged when those elements are used to approximate the solutions of the parabolic and hyperbolic equations associated with $-\operatorname{div}(a \nabla)$, *i.e.*, for equations of the form

$$\begin{cases} \partial_t u(x, t) - \operatorname{div} \left(a(x) \nabla u(x, t) \right) = g(x, t) & (x, t) \in \Omega_T; g \in L^2(\Omega_T), \\ u = 0 & \text{on } \partial\Omega_T, \end{cases} \quad (8.1)$$

and

$$\begin{cases} \rho(x) \partial_t^2 u(x, t) - \operatorname{div} \left(a(x) \nabla u(x, t) \right) = g(x, t) & (x, t) \in \Omega_T; g \in L^2(\Omega_T), \\ u = 0 & \text{on } \partial\Omega_T, \\ \partial_t u = 0 & \text{on } \Omega \times \{t = 0\}. \end{cases} \quad (8.2)$$




FIGURE 2. Mesh details.

More precisely the solutions of (8.1) and (8.2) are approximated, *via* finite element methods, using the linear space spanned by the interpolation basis ϕ_i^{loc} as test and solution space. Note that the approximate solutions are of the form

$$u^H(x, t) = \sum_i c_i(t) \phi_i^{\text{loc}}(x). \quad (8.3)$$

We refer to Sections 4 and 5 of [70] for precise statements and proofs.

9. NUMERICAL IMPLEMENTATION AND EXPERIMENTS

9.1. Implementation

Our method is, in its formal description, meshless in the sense that only the locations of the points $(x_j)_{j \in \mathcal{N}}$ are needed to construct the rough polyharmonic splines ϕ_i or ϕ_i^{loc} as solutions of the minimization problem (2.3) or (7.4).

For numerical implementation, we need to formulate and solve the problem in the full discrete setting. Although, for the sake of conciseness we have provided an error analysis only in the continuum setting, this analysis can be extended to the fully discrete setting without major difficulties.

We will now describe the implementation of our method in the discrete setting. Let \mathcal{T}_H be a tessellation of Ω of resolution H with nodes $\{x_i\}$. Assume that $h \ll H$ and that \mathcal{T}_h is obtained by repeated refinements of \mathcal{T}_H . Note that the accuracy of our method is independent from the regularity of \mathcal{T}_H or the aspect ratios of its elements (triangles for $d = 2$).

Let $W_h(\Omega)$ be the set of piecewise linear functions on \mathcal{T}_h with zero Dirichlet boundary condition on $\partial\Omega$. For $\forall u \in W_h(\Omega)$, we can define g_u , which approximates $-\text{div } a \nabla u$ on Ω . g_u is piecewise constant on the dual mesh of \mathcal{T}_h , and can be defined by the following finite volume formulation,

$$g_u(y_i) = \frac{1}{|\mathfrak{V}_i|} \int_{\Omega} \nabla 1_{\mathfrak{V}_i} a \nabla u \quad (9.1)$$

where \mathfrak{V}_i is the dual Voronoi cell associated with the node y_i of \mathcal{T}_h and $1_{\mathfrak{V}_i}$ is the characteristic function which equals to one on \mathfrak{V}_i and zero elsewhere.

Figure 2a shows the primal (finite element) mesh and its dual mesh used for the numerical implementation in this paper: the red dots are the nodes x_i of the triangulation \mathcal{T}_H . In Figure 2a, the red square around each y_i is the Voronoi cell \mathfrak{V}_i associated with the node y_i , and the dual mesh is composed of such cells.

With this definition, the local basis $\phi_{i,h}^{\text{loc}}$ can be readily computed through the variational formulation (9.2).

$$\begin{cases} \text{Minimize } \int_{\Omega} |g_{\phi}|^2 dx \\ \text{Subject to } \begin{cases} \phi \in W_h(\Omega_i), \phi = 0 \text{ on } \partial\Omega_i \cup \Omega \setminus \Omega_i \\ \phi(x_j) = \delta_{ij}, j \in \mathcal{N}. \end{cases} \end{cases} \tag{9.2}$$

9.2. A two dimensional example

In this example we consider $\Omega = (0, 1) \times (0, 1)$ and

$$\begin{aligned} a(x) := & \frac{1}{6} \left(\frac{1.1 + \sin(2\pi x/\epsilon_1)}{1.1 + \sin(2\pi y/\epsilon_1)} + \frac{1.1 + \sin(2\pi y/\epsilon_2)}{1.1 + \cos(2\pi x/\epsilon_2)} + \frac{1.1 + \cos(2\pi x/\epsilon_3)}{1.1 + \sin(2\pi y/\epsilon_3)} \right. \\ & \left. + \frac{1.1 + \sin(2\pi y/\epsilon_4)}{1.1 + \cos(2\pi x/\epsilon_4)} + \frac{1.1 + \cos(2\pi x/\epsilon_5)}{1.1 + \sin(2\pi y/\epsilon_5)} + \sin(4x^2y^2) + 1 \right), \end{aligned} \tag{9.3}$$

where $\epsilon_1 = \frac{1}{5}, \epsilon_2 = \frac{1}{13}, \epsilon_3 = \frac{1}{17}, \epsilon_4 = \frac{1}{31}, \epsilon_5 = \frac{1}{65}$.

Note that a , as defined by (9.3), corresponds to the one used in [69] (Example 1 of Sect. 3) and [63]. The number of coarse nodes is $N = N_c \times N_c$ with $N_c = 32$. The (regular) coarse mesh $\mathcal{T}_H = \mathcal{T}^0$ is obtained by first (uniformly) subdividing Ω into $N_c \times N_c$ rectangles, then partitioning each rectangle into two triangles along the direction $(1, 1)$. We can further refine the mesh to obtain $\mathcal{T}^1, \mathcal{T}^2, \dots, \mathcal{T}^k$. Global equations are solved on a fine triangulation with 66 049 nodes and 131 072 triangles. We compute localized elements ϕ_i^{loc} on localized sub-domains Ω_i^l defined by adding l layers of coarse triangles around x_i . More precisely Ω_i^1 is the union of triangles sharing x_i as a node and Ω_i^{l+1} is the union of Ω_i^l with coarse triangles sharing a node with a triangle contained in Ω_i^l . We refer to Figures 2b, 2c and 2d for an illustration of Ω_i^l with $l = 1, 2, 3$.

We refer to Figure 3 for the results of our numerical experiment. Figure 3a shows ϕ_i (one element of the global basis in dimension two). Figure 3b shows a slice of ϕ_i along the x -axis. Figure 3c shows $\|\phi_i - \phi_i^{\text{loc}}\|$ in L^2, \mathcal{H}^1 , and L^∞ norms in the log scale as a function of the number of layers $l = 1, \dots, 8$ used in the localization. Note that $\|\phi_i - \phi_i^{\text{loc}}\|$ decays exponentially fast with l and that the decay rate is nearly independent from the choice of the norm. Figures 3e and 3f show $\|u - u^{H,\text{loc}}\|_{L^2(\Omega)}$ and $\|u - u^{H,\text{loc}}\|_{\mathcal{H}^1(\Omega)}$ in log-scale as a function of the number of layers $l = 1, \dots, 8$ (defining the size of localized sub-domains) for different values of (coarse) mesh resolution $H = 1/4, 1/8, 1/16, 1/32$. Note that as the number of layers increases, $\|u - u^{H,\text{loc}}\|$ decreases first and then quickly saturates around $O(H)$.

9.3. Wave equation

In this example we consider a and Ω as defined in Section 9.2 and we compute the solutions of wave equation (8.2) up to time $T = 1$. The initial condition is $u(x, 0) = 0$ and $u_t(x, 0) = 0$. The boundary condition is $u(x, t) = 0$, for $x \in \partial\Omega$. The density ρ is uniformly equal to one and we choose $g = \sin(\pi x) \sin(\pi y)$. Figures 4a and 4b show $u(x, T)$ and $u^{H,\text{loc}}(x, T)$ with $T = 1$. $u^{H,\text{loc}}(x, T)$ is computed over the approximate solution space spanned by the localized basis elements ϕ_i^{loc} obtained by choosing Ω_i as the union of 3 layers of triangles around x_i (hence, we choose $l = 3$). The resolution of the fine mesh is $h = 1/80$. The resolution of the coarse mesh is $H = 1/10$. Figure 4c shows approximation error $\|u - u^{H,\text{loc}}(\cdot, T)\|$ in L^2, \mathcal{H}^1 , and L^∞ norms in the log scale as a function of the number of layers $l = 1, \dots, 8$.

9.4. A one dimensional example

In this example we consider $\Omega = (0, 1)$ and

$$a(x) := 1 + \frac{1}{2} \sin \left(\sum_{k=1}^K k^{-\alpha} (\zeta_{1k} \sin(kx) + \zeta_{2k} \cos(kx)) \right) \tag{9.4}$$





FIGURE 3. Two dimensional example (9.3).

where $\{\zeta_{1k}\}$ and $\{\zeta_{2k}\}$ are two independent random vectors with independent entries uniformly distributed in $[-\frac{1}{2}, \frac{1}{2}]$. This example is taken from [45, 63] and it has no scale separation. Note that a has an algebraically decaying (Fourier energy) spectrum, *i.e.*,

$$\langle |\hat{a}(k)|^2 \rangle \simeq |k|^{-\alpha}, \tag{9.5}$$


where $\langle \cdot \rangle$ denotes ensemble averaging, and $\hat{a}(k)$ is the Fourier transform of the coefficients $a(x)$. We choose $K = 20$ in the numerical experiment.

The coarse nodes $(x_j)_{j \in \mathcal{N}}$ correspond to $N = d_c = 80$ points uniformly distributed in $(0, 1)$. More-precisely, those points correspond to the nodes of a coarse mesh of $(0, 1)$ and the distance between two successive points is $H = 1/81$. The fine mesh is then obtained by refining (bisecting) the coarse mesh 3 times. Local bases are



(a) $u(x, T)$ for $T = 1$

(b) $u^{H,loc}(x, T)$ for $T = 1$



(c) $\|u - u^{H,loc}(., T)\|$ in L^2 , \mathcal{H}^1 , and L^∞ in log scale as a function of l

FIGURE 4. Wave equation of Section 9.3.

computed on intervals with l layers, where $l = 1, \dots, 8$. More precisely our localized sub-domains (over which the elements ϕ_i^{loc} are computed) are $\Omega_i = (x_i - l/81, x_i + l/81) \cap \Omega$ for $l = 1, \dots, 8$.

We refer to Figure 5 for the results of our numerical implementation. Figure 5a shows the local basis ϕ_i^{loc} for the node $i = 40$ for various degrees of localization (*i.e.*, for $l = 1, \dots, 8$). Figure 5b shows the global basis in the log scale (*i.e.*, $\log_{10}(10^{-9} + |\phi_{40}|)$) and illustrates the exponential decay of $\phi_i(x)$ away from x_i . Figure 5c shows $\phi_i - \phi_i^{loc}$, the difference between the global and the local basis, at node $i = 40$, for various degrees of localization (*i.e.*, for $l = 1, \dots, 8$), and illustrates the decay of this difference with respect to the number of layers l defining Ω_i . Figure 5d shows $-\text{div}(a\nabla\phi_i)$ for $i = 40$. Figure 5e shows $-\text{div}(a\nabla\phi_i)$ for $i = 40$ in the log




FIGURE 5. One dimensional example (9.4).

scale (*i.e.*, $\log_{10}(10^{-9} + |\operatorname{div} a \nabla \phi_i|)$). Figures 5d and 5e illustrate the exponential decay of $-\operatorname{div}(a \nabla \phi_i)(x)$ away from x_i . Figure 5f shows the matrix $\Theta_{i,j}$, as defined in (2.5), as a function of (i, j) (two dimensional surface plot). Similarly Figure 5g shows the matrix $P_{i,j}$, as defined in (2.12), as a function of (i, j) . Figure 5h shows P in the log scale and illustrates the exponential decay of the entries of P away from the diagonal.

Analogy with polyharmonic splines. Note that as $H \downarrow 0$ the matrix Θ shown in Figure 5f converges towards $\tau(x, y)$ the fundamental solution of the operator $L = (\operatorname{div}(a \nabla \cdot))^2$. Note also that since $P^{-1} = \Theta$, P is, in fact (in analogy with [75–77]), a discrete approximation of the operator L (*i.e.* $\sum_{j=1}^N P_{i,j} u(x_j)$) can be thought of as an approximation of $Lu(x_i)$. Note the localization of P (near its diagonal) as shown in Figure 5g. Note also since $\phi_i(x) = \sum_{j=1}^N P_{i,j} \tau(x, x_j)$, ϕ_i is obtained by applying the discrete version of L against the fundamental solution of L and in that sense the elements ϕ_i or ϕ_i^{loc} (see Fig. 5a) are approximations of masses of Diracs.

10. ON NUMERICAL HOMOGENIZATION

10.1. Short overview

As mentioned in the introduction the field of numerical homogenization concerns the numerical approximation of the solution space of (1.1) with a finite-dimensional space. This problem is motivated by the fact that standard methods (such as finite-element method with piecewise linear elements [11]) can perform arbitrarily badly for PDEs with rough coefficients such as (1.1). Although some numerical homogenization methods (described below) are directly inspired from classical homogenization concepts such as periodic homogenization and scale separation [14], oscillating test functions, G or H -convergence and compensated compactness [37, 65, 80] and localized cell problems [73], one of the main objectives of numerical homogenization is to achieve a numerical approximation of the solution space of (1.1) with arbitrary rough coefficients (*i.e.*, in particular, without the assumptions found in classical homogenization, such as scale separation, ergodicity at fine scales and ϵ -sequences of operators). Numerical homogenization methods differ in (computational) cost and assumptions required for accuracy and by now, the field of numerical homogenization has become large enough that it is not possible to give a complete review in this short paper. For the sake of completeness, we will quote below the non exhaustive list of numerical homogenization methods discussed in [70]:

- The multi-scale finite element method [31, 44, 45, 87] can be seen as a numerical generalization of this idea of oscillating test functions found in H -convergence. A convergence analysis for periodic media revealed a resonance error introduced by the microscopic boundary condition [44, 45]. An over-sampling technique was proposed to reduce the resonance error [45].
- Harmonic coordinates play an important role in various homogenization approaches, both theoretical and numerical. These coordinates were introduced in [49] in the context of random homogenization. Next, harmonic coordinates have been used in one-dimensional and quasi-one-dimensional divergence form elliptic problems [8, 10], allowing for efficient finite dimensional approximations. These coordinates have been used in [13, 71, 72] to establish the anomalous diffusion of stochastic processes evolving in media characterized by an infinite number of scales with no scale separation and unbounded contrast and provide a proof of Davies’ conjecture [71] on the behavior of the heat kernel in periodic media. The connection of these coordinates with classical homogenization is made explicit in [3] in the context of multi-scale finite element methods. The idea of using particular solutions in numerical homogenization to approximate the solution space of (1.1) appears to have been first proposed in reservoir modeling in the 1980s [19, 86] (in which a global scale-up method was introduced based on generic flow solutions (*i.e.*, flows calculated from generic boundary conditions)). Its rigorous mathematical analysis has been done only recently [69] and is based on the fact that solutions are \mathcal{H}^2 -regular with respect to harmonic coordinates (recall that they are \mathcal{H}^1 -regular with respect to Euclidean coordinates). The main message here is that if the right hand side of (1.1) is in L^2 , then solutions can be approximated at small scales (in \mathcal{H}^1 -norm) by linear combinations of d (linearly independent) particular solutions (d being the dimension of the space). In that sense, harmonic coordinates are

- (simply) good candidates for being d linearly independent particular solutions. The idea of a global change of coordinates analogous to harmonic coordinates has been implemented numerically in order to up-scale porous media flows [19, 29, 30]. We refer, in particular, to a recent review article [19] for an overview of some main challenges in reservoir modeling and a description of global scale-up strategies based on generic flows.
- In [27, 33], the structure of the medium is numerically decomposed into a micro-scale and a macro-scale (meso-scale) and solutions of cell problems are computed on the micro-scale, providing local homogenized matrices that are transferred (up-scaled) to the macro-scale grid. See also [1, 2, 34]. We also refer to [38] for convergence results on the Heterogeneous Multiscale Method in the framework of G and Γ -convergence.
 - More recent work includes an adaptive projection based method [68], which is consistent with homogenization when there is scale separation, leading to adaptive algorithms for solving problems with no clear scale separation; finite difference approximations of fully nonlinear, uniformly elliptic PDEs with Lipschitz continuous viscosity solutions [21] and operator splitting methods [4, 5].
 - We refer to [16, 17] (and references therein) for recent results on homogenization of scalar divergence-form elliptic operators with stochastic coefficients. Here, the stochastic coefficients $a(x/\varepsilon, \omega)$ are obtained from stochastic deformations (using random diffeomorphisms) of the periodic and stationary ergodic setting.
 - We refer to [12] for a detailed study of the fluctuation error (error in approximating the limiting probability distribution of the random part of the difference between heterogeneous and homogenized solutions) of MsFEM and HMM in the context of stochastic homogenization and a rigorous assessment of the behavior of those multiscale algorithms in non-periodic situations. In particular, [12] shows that savings in computational cost based on scale separation may come at the price of amplifying the variances of these fluctuations.

10.2. On localization

The numerical complexity of numerical homogenization is related to the size of the support of the basis elements used to approximate the solution space of (1.1). The possibility to compute such bases on localized sub-domains of the global domain Ω without loss of accuracy is therefore a problem of practical importance. We refer to [9, 22, 28, 55, 70] for recent localization results for divergence-form elliptic PDEs. The strategy of [22] is to construct triangulations and finite element bases that are adapted to the shape of high conductivity inclusions *via* coefficient dependent boundary conditions for the subgrid problems (assuming a to be piecewise constant and the number of inclusions to be bounded). The strategy of [28] is to solve local eigenvalue problems, observing that only a few eigenvectors are sufficient to obtain a good pre-conditioner. Both [22, 28] require specific assumptions on the morphology and number of inclusions. The idea of the strategy is to observe that if a is piecewise constant and the number of inclusions is bounded, then u is locally \mathcal{H}^2 away from the interfaces of the inclusions. The inclusions can then be taken care of by adapting the mesh and the boundary values of localized problems or by observing that those inclusions will affect only a finite number of eigenvectors.

The strategy of [9] is to construct Generalized Finite Elements by partitioning the computational domain into a collection of preselected subsets and compute optimal local bases (using the concept of n -widths [74]) for the approximation of harmonic functions. Local bases are constructed by solving local eigenvalue problems (corresponding to computing eigenvectors of P^*P , where P is the restriction of a -harmonic functions from ω^* onto $\omega \subset \omega^*$, P^* is the adjoint of P , and ω is a sub-domain of Ω surrounded by a larger sub-domain ω^*). The method proposed in [9] achieves a near exponential convergence rate (in the number of pre-computed bases functions) for harmonic functions. Non-zero right hand sides (denoted by g) are then taken care of by finding (for each different g) particular solutions on preselected subsets with a constant Neumann boundary condition (determined according to the consistency condition). The near exponential rate of convergence observed in [9] is explained by the fact that the source space considered in [9] is more regular than L^2 (see [9]) and requires the computation of particular (local) solutions for each right hand side g and each non-zero boundary condition, the basis obtained in [9] is in fact adapted to a -harmonic functions away from the boundary).

The strategy of [70] is to use the transfer property of the flux-norm introduced in [15] (and the strong compactness of the solution space [70]) to identify the global basis and then (inspired by an idea stemming from classical homogenization [39, 40, 73, 88]) localize the computation of the basis to sub-domains of size

$\mathcal{O}(\sqrt{H} \ln(1/H))$ (without loss of accuracy) by replacing the operator $-\operatorname{div}(a\nabla\cdot)$ with the operator $\frac{1}{T}\cdot - \operatorname{div}(a\nabla\cdot)$ in the computation of this basis with a well chosen T balancing the created exponential decay in the Green's function with the deterioration of the transfer property. We also refer to [83] for a generalization of the transfer property to Hilbert triple settings and a different approach to localization for hyperbolic initial-boundary value problems based on the observation that weak solutions of symmetric hyperbolic systems propagate at finite speed.

In [41], it is shown that for scalar elliptic problems with L^∞ coefficients, there exists a local (generalized) finite element basis (AL basis) with $\mathcal{O}((\log \frac{1}{H})^{d+1})$ basis functions per nodal point, achieving the $\mathcal{O}(H)$ convergence rate. Although the construction provided in [41] involves solving global problems with specific right hand sides, its theoretical result supports the possibility of constructing localized bases.

The strategy of [55] is to introduce a modified Clément interpolation \mathcal{I}_H (in which the interpolation of u is not based on its nodal values but on volume averages around nodes), and writing V^f the kernel of \mathcal{I}_H (*i.e.*, the set of functions u such that $\mathcal{I}_H u = 0$), identify V_H^{ms} as the orthogonal complement of V^f with respect to the scalar product defined by $a(u, v) = \int_\Omega \nabla u a \nabla v$. The finite element basis ϕ_i is then identified by projecting nodal piecewise linear elements onto \mathcal{I}_H . The work [55] shows that the computation of ϕ_i can be localized to sub-domains of size $\mathcal{O}(H \ln \frac{1}{H})$ without loss of accuracy. The main differences between the strategy of [55] and ours are as follows: (1) The identification of the space V^f in [55] and the related projections lead to saddle point problems (even after localization) whereas the method proposed here leads to localized elliptic linear systems (which ensures the fact that the computational cost of our method remains minimal and plays an important role in its stability, which we observe numerically, in the high contrast regime) (2) The interpolation of solutions in [55] is not based on nodal values but on volume averages around nodes whereas our interpolation of solutions is based on nodal values (which allows for an accurate estimation of solutions from point measurements and the introduction of homogenized linear systems defined on nodal values) (3) The accuracy of our method does not depend on the aspect ratios of the triangles of the mesh formed by the interpolation points.

We refer to Section 10.3 for a description of the computational cost of our method.

10.3. Computational cost

Write d_{lf} the number nodes of the fine mesh contained in the local domain Ω_i (d_{lf} is the number of nodes/degrees of freedom of $\mathcal{T}_h \cap \Omega_i$). Write d_f the number of interior nodes of the fine mesh \mathcal{T}_h and d_c the number of interior nodes of the coarse mesh \mathcal{T}_H . Then the cost for solving local basis ϕ_i^{loc} , introduced in this paper, is $\mathcal{O}(d_{lf}d_c)$ (because each ϕ_i^{loc} is the solution of a banded/nearly diagonal linear system), and the cost for constructing the global stiffness matrix is $\mathcal{O}(d_{lf}(d_{lf}d_c/d_f)d_c) = \mathcal{O}(d_{lf}^2d_c^2/d_f)$. Notice that $d_{lf}d_c/d_f$ is the number of local domains which overlap with Ω_i . Therefore, the overall cost is $\mathcal{O}(d_{lf}d_c + d_{lf}^2d_c^2/d_f) = \mathcal{O}(d_{lf}d_c(1 + d_{lf}d_c/d_f))$, and the dominant part is $\mathcal{O}(d_{lf}^2d_c^2/d_f)$.

For quasi-uniform triangulations \mathcal{T}_h and \mathcal{T}_H , we have $d_f \sim h^{-d}$ and $d_c \sim H^{-d}$. For the method introduced in this paper, to achieve $\mathcal{O}(H)$ accuracy in \mathcal{H}^1 norm, we need $d_{lf} \sim (H \log(1/H))^{-d}$. Therefore the overall cost of (pre-)computing of all the localized basis elements ϕ_i^{loc} is $C_1 \sim (\frac{\log(1/H)}{h})^d$.

10.4. Connections with classical homogenization theory

We will now describe connections between classical homogenization theory [14, 46, 62, 65], our previous work [15, 70] and the present paper. Recall that in classical homogenization theory [65] one considers situations where the conductivity a depends on a (small) parameter ϵ (*i.e.*, $a := a(x, \epsilon)$) and one is interested in approximating the solution u_ϵ of (1.1) in the limit where ϵ converges towards zero by a function \hat{u}_ϵ that is easier to compute or simpler to represent.

In periodic homogenization [14, 46, 67], where the conductivity $a(x, \epsilon)$ is assumed to be of the form $a^{\text{per}}(\frac{x}{\epsilon})$ (and a^{per} is assumed to be periodic in its argument), this approximation can be achieved by observing that u_ϵ converges, as $\epsilon \downarrow 0$, towards the solution u_0 of $-\operatorname{div}(\hat{a}\nabla u_0) = g$ (with zero Dirichlet boundary condition) and where \hat{a} is a constant (effective, homogenized) conductivity determined by solving the so called (periodic) cell

problems $\operatorname{div}(a^{\text{per}}(e_i + \nabla\chi_i)) = 0$ over one period of a^{per} ($i \in \{1, \dots, d\}$, the vectors e_i form an orthonormal basis of \mathbb{R}^d and a^{per} is invariant under translation by e_i). u_0 is much simpler to compute than u_ϵ because its variations are at the $\mathcal{O}(1)$ -scale (referred to as the coarse scale) in contrast to the oscillations of u_ϵ that are at the ϵ -scale (referred to as the fine scale).

However the convergence of u_ϵ towards u_0 is limited to the L^2 -norm, which does not control the derivatives. Since in most applications (*e.g.*, computing stresses in elasticity and currents in electromagnetism) the approximation of derivatives is required, an important *corrector problem* arose: find an approximation of the form $\hat{u}_\epsilon(x) = u_0(x) + \epsilon u_1(x)$ that approximates the solution $u_\epsilon(x)$ of the original problem in \mathcal{H}^1 norm. Solving this problem was an essential step in the development of periodic homogenization. It turned out that the corrector term $\epsilon u_1(x)$ admits an explicit representation *via* the solution of the cell problems $\chi_i(\frac{x}{\epsilon})$ and the zero order approximation $u_0(x)$ of the form $\epsilon \sum_{i=1}^d \chi_i(\frac{x}{\epsilon}) \partial_i u_0(x)$. Thus the corrector term contains both fine and coarse scales. The concept of the cell problem played the key role in periodic homogenization since it is pivotal for both identifying the effective conductivity and an \mathcal{H}^1 -norm approximation of the solution of the original problem. Namely it allows to construct both the homogenized PDE and the corrector term. The advantage of computing \hat{u}_ϵ rather than directly computing u_ϵ comes from the fact that computing of $\chi_i(y)$, $i = 1, \dots, d$ and $u_0(x)$ is less expensive than computing u_ϵ .

The question of how far these ideas and concepts can be extended beyond periodic homogenization lead to development stochastic homogenization, *i.e.* the extension of the theory to coefficients of the form $a = \{a_{ij}(\frac{x}{\epsilon}, \omega)\}$, where ω is an atom of a probability space and these coefficients are assumed to be stationary and ergodic. While there is no periodic cell in this problem, the generalized cell problem was successfully introduced in [49, 73] and the homogenized problem was derived in terms of these problems analogously to the periodic case, that is the objective of identifying the homogenized PDE was achieved. However, the construction of the corrector terms turned out to be a much harder problem and by now it is only partially resolved [6, 12, 18, 47, 50]. It is also worth mentioning that stochastic homogenization, as it currently stands, is non-local in the sense that a deterministic and compactly supported perturbation of $a = \{a_{ij}(\frac{x}{\epsilon}, \omega)\}$ does not affect (in the limit where $\epsilon \downarrow 0$) the value of the effective conductivity nor the cell problems since these are defined taking averages (perturbations that leave the statistical properties of the random coefficient unchanged do not affect the homogenization result).

Many practical problems such as study of oil reservoirs, biological tissues, *etc.* contain multiple scales that are not necessarily well-separated. Again, one can ask to which extent the ideas and concepts of classical homogenization can be used in such problems. These questions, which one can address in the context of the model problem (1.1), are challenging because if coefficients $a(x)$ are just L^∞ functions, then they have “no structure” (no translational invariance as in the periodic and ergodic cases).

However the present work shows that, although there is no direct analog of the parameter ϵ , one can still develop a generalization of most of the concepts and objectives of classical homogenization. First observe that the linear space spanned by the basis elements ϕ_i (or ϕ_i^{loc} with support of size $\mathcal{O}(H \ln \frac{1}{H})$) constitutes a finite dimensional approximation of the solution space of (1.1) and this finite-dimensional approximation is made possible by the (strong) compactness of the solution space (in \mathcal{H}^1 -norm) when $g \in L^2(\Omega)$. Note that the requirement that g belongs to $L^2(\Omega)$ is also present in classical homogenization and the notion of compactness is also present in the (weaker) form of compactness by compensation [65]. Write S the stiffness matrix associated with the elements ϕ_i^{loc} defined by $S_{i,j} := \int_\Omega \nabla \phi_i^{\text{loc}} a \nabla \phi_j^{\text{loc}}$ and observe that if φ_i are piecewise linear nodal basis elements on a (coarse) mesh with vertices $(x_i)_{i \in \mathcal{N}}$ and if c^* the N -dimensional vector solution of the linear system $\sum_{j=1}^N S_{i,j} c_j = \int_\Omega \phi_i^{\text{loc}} g$ then the distance between u (the solution of (1.1)) and $\bar{u} := \sum_{i=1}^N c_i^* \varphi_i$ is at most $CH \|g\|_{L^2(\Omega)}$ in L^2 norm (this can be obtained in a straightforward manner by writing u^H the finite element solution of (1.1) over $(\phi_i^{\text{loc}})_{i \in \mathcal{N}}$, *i.e.* $u^H := \sum_{i=1}^N c_i^* \phi_i^{\text{loc}}$, applying the triangle inequality to $u - \bar{u} = u - u^H + u^H - \bar{u}$, and Thm. 1.1 of [66] to $u^H - \bar{u}$). In that sense, S can be viewed as a homogenized linear system, generalizing the concept of an homogenized PDE, and its entries can be seen as localized effective (edge) conductivities [23] generalizing the concept of effective conductivity. Furthermore noting that, although \bar{u} does not constitute a good approximation of the derivatives of u , the distance between u and u^H is at most $CH \|g\|_{L^2(\Omega)}$ in \mathcal{H}^1

norm, we deduce that the elements $\phi_i^{\text{loc}} - \varphi_i$ can be interpreted as generalized correctors. That is we define the generalize corrector as an \mathcal{H}^1 approximation of the exact solution which is constructed out of standard solutions that solve *generalized cell problems* and therefore does not depend on the RHS of the original equation. Several ways of introducing such problems were proposed in [15, 70] and the present paper. Of course in the present work the notion of corrector is superfluous since Rough Polyharmonic Splines directly lead to an \mathcal{H}^1 -accurate approximation of the solution space. It should also be noted that the elements $(\phi_i^{\text{loc}})_{i \in \mathcal{N}}$ provide an explicit representation allowing for the identification of the (local) effect of a local perturbation of a on the solution and on effective conductivities.

Acknowledgements. The work of H. Owhadi is partially supported by the National Science Foundation under Award Number CMMI-092600, the Department of Energy National Nuclear Security Administration under Award Number DE-FC52-08NA28613, the Air Force Office of Scientific Research under Award Number FA9550-12-1-0389 and a contract from the DOE Exascale Co-Design Center for Materials in Extreme Environments. The work of L. Zhang is supported by the Young Thousand Talents Program of China. The work of L. Berlyand is supported by DOE under Award Number DE-FG02-08ER25862. H. Owhadi thanks M. Desbrun and F. de Goes for stimulating discussions. H. Owhadi also thanks François Murat for helpful discussions on Lemma 3.1. L. Berlyand thanks M. Potomkin for useful comments and suggestions. We also thank an anonymous referee for carefully reading the manuscript and detailed comments and suggestions.

REFERENCES

- [1] A. Abdulle and M.J. Grote, Finite element heterogeneous multiscale method for the wave equation. *Multiscale Model. Simul.* **9** (2011) 766–792.
- [2] A. Abdulle and Ch. Schwab, Heterogeneous multiscale FEM for diffusion problems on rough surfaces. *Multiscale Model. Simul.* **3** (2004/05) 195–220.
- [3] G. Allaire and R. Brizzi, A multiscale finite element method for numerical homogenization. *Multiscale Model. Simul.* **4** (2005) 790–812.
- [4] T. Arbogast and K.J. Boyd, Subgrid upscaling and mixed multiscale finite elements. *SIAM J. Numer. Anal.* **44** (2006) 1150–1171.
- [5] T. Arbogast, C.-S. Huang and S.-M. Yang, Improved accuracy for alternating-direction methods for parabolic equations based on regular and mixed finite elements. *Math. Models Methods Appl. Sci.* **17** (2007) 1279–1305.
- [6] S.N. Armstrong and P.E. Souganidis, Stochastic homogenization of Hamilton-Jacobi and degenerate Bellman equations in unbounded environments. *J. Math. Pures Appl.* **97** (2012) 460–504.
- [7] M. Atteia, Fonctions spline et noyaux reproduisants d’Aronszajn-Bergman. *Rev. Française Informat. Recherche Opérationnelle* **4** (1970) 31–43.
- [8] I. Babuška, G. Caloz and J.E. Osborn, Special finite element methods for a class of second order elliptic problems with rough coefficients. *SIAM J. Numer. Anal.* **31** (1994) 945–981.
- [9] I. Babuška and R. Lipton, Optimal local approximation spaces for generalized finite element methods with application to multiscale problems. *Multiscale Model. Simul.* **9** (2011) 373–406.
- [10] I. Babuška and J.E. Osborn, Generalized finite element methods: their performance and their relation to mixed methods. *SIAM J. Numer. Anal.* **20** (1983) 510–536.
- [11] I. Babuška and J.E. Osborn, Can a finite element method perform arbitrarily badly? *Math. Comput.* **69** (2000) 443–462.
- [12] G. Bal and W. Jing, Corrector theory for MSFEM and HMM in random media. *Multiscale Model. Simul.* **9** (2011) 1549–1587.
- [13] G. Ben Arous and H. Owhadi, Multiscale homogenization with bounded ratios and anomalous slow diffusion. *Comm. Pure Appl. Math.* **56** (2003) 80–113.
- [14] A. Bensoussan, J.L. Lions and G. Papanicolaou, Asymptotic analysis for periodic structure. North Holland, Amsterdam (1978).
- [15] L. Berlyand and H. Owhadi, Flux norm approach to finite dimensional homogenization approximations with non-separated scales and high contrast. *Arch. Rational Mech. Anal.* **198** (2010) 677–721.
- [16] X. Blanc, C. Le Bris and P.-L. Lions, Une variante de la théorie de l’homogénéisation stochastique des opérateurs elliptiques. *C. R. Math. Acad. Sci. Paris* **343** (2006) 717–724.
- [17] X. Blanc, C. Le Bris and P.-L. Lions, Stochastic homogenization and random lattices. *J. Math. Pures Appl.* **88** (2007) 34–63.
- [18] A. Bourgeat and A. Piatnitski, Estimates in probability of the residual between the random and the homogenized solutions of one-dimensional second-order operator. *Asymptot. Anal.* **21** (1999) 303–315.
- [19] L.V. Branets, S.S. Ghai, L.L. and X.-H. Wu, Challenges and technologies in reservoir modeling. *Commun. Comput. Phys.* (2009) **6** 1–23.
- [20] R.A. Brownlee, *Error estimates for interpolation of rough and smooth functions using radial basis functions*. Ph.D. thesis. University of Leicester (2004).

- [21] L.A. Caffarelli and P.E. Souganidis, A rate of convergence for monotone finite difference approximations to fully nonlinear, uniformly elliptic PDEs. *Comm. Pure Appl. Math.* **61** (2008) 1–17.
- [22] C.-C. Chu, I.G. Graham and T.Y. Hou, A new multiscale finite element method for high-contrast elliptic interface problems. *Math. Comput.* **79** (2010) 1915–1955.
- [23] M. Desbrun, R. Donaldson and H. Owhadi. Modeling across scales: Discrete geometric structures in homogenization and inverse homogenization. *Reviews of Nonlinear Dynamics and Complexity. Special issue on Multiscale Analysis and Nonlinear Dynamics* (2012).
- [24] J. Duchon, Interpolation des fonctions de deux variables suivant le principe de la flexion des plaques minces. *Rev. Francaise Automat. Informat. Recherche Operationnelle Ser. RAIRO Anal. Numer.* **10** (1976) 5–12.
- [25] J. Duchon, Splines minimizing rotation-invariant semi-norms in Sobolev spaces, in Constructive theory of functions of several variables, *Proc. of Conf., Math. Res. Inst., Oberwolfach, 1976*, in vol. 571. of *Lect. Notes Math.* Springer, Berlin (1977) 85–100.
- [26] J. Duchon, Sur l'erreur d'interpolation des fonctions de plusieurs variables par les D^m -splines. *RAIRO Anal. Numér.* (1978) **12** 325–334.
- [27] W. E and B. Engquist, The heterogeneous multiscale methods. *Commun. Math. Sci.* **1** (2003) 87–132.
- [28] Y. Efendiev, J. Galvis and X. Wu, Multiscale finite element and domain decomposition methods for high-contrast problems using local spectral basis functions. *J. Comput. Phys.* **230** (2011) 937–955.
- [29] Y. Efendiev, V. Ginting, T. Hou and R. Ewing, Accurate multiscale finite element methods for two-phase flow simulations. *J. Comput. Phys.* **220** (2006) 155–174.
- [30] Y. Efendiev and T. Hou, Multiscale finite element methods for porous media flows and their applications. *Appl. Numer. Math.* **57** (2007) 577–596.
- [31] Y. Efendiev and T.Y. Hou, Multiscale finite element methods, Theory and applications, in vol. 4, *Surveys and Tutorials in the Applied Mathematical Sciences.* Springer, New York (2009).
- [32] I. Ekeland and R. Temam, Convex Analysis and Variational Problems, vol. 28 of *Classics in Appl. Math.* Society for Industrial and Applied Mathematics (1987).
- [33] B. Engquist and P.E. Souganidis, Asymptotic and numerical homogenization. *Acta Numerica* **17** (2008) 147–190.
- [34] B. Engquist, H. Holst and O. Runborg, Multi-scale methods for wave propagation in heterogeneous media. *Commun. Math. Sci.* **9** (2011) 33–56.
- [35] M. Giaquinta, Multiple integrals in the calculus of variations and nonlinear elliptic systems, vol. 105. *Ann. Math. Stud.* Princeton University Press, Princeton, NJ (1983).
- [36] D. Gilbarg and N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, 2nd edn. Springer-Verlag (1983).
- [37] E. De Giorgi, Sulla convergenza di alcune successioni di integrali del tipo dell'aera. *Rendi Conti di Mat.* **8** (1975) 277–294.
- [38] A. Gloria, Analytical framework for the numerical homogenization of elliptic monotone operators and quasiconvex energies. *SIAM MMS* **5** (2006) 996–1043.
- [39] A. Gloria, Reduction of the resonance error-Part 1: Approximation of homogenized coefficients. *Math. Models Methods Appl. Sci.* **21** (2011) 1601–1630.
- [40] A. Gloria and F. Otto, An optimal error estimate in stochastic homogenization of discrete elliptic equations. *Ann. Appl. Probab.* **22** (2012) 1–28.
- [41] L. Grasedyck, I. Greff and S. Sauter, The al basis for the solution of elliptic problems in heterogeneous media. *Multiscale Modeling and Simulation* **10** (2012) 245–258.
- [42] M. Grüter and K. Widman, The green function for uniformly elliptic equations. *Manuscripta Math.* **37** (1982) 303–342.
- [43] R.L. Harder and R.N. Desmarais, Interpolation using surface splines. *J. Aircr.* **9** (1972) 189–191.
- [44] T.Y. Hou, X.-H. Wu and Z. Cai, Convergence of a multiscale finite element method for elliptic problems with rapidly oscillating coefficients. *Math. Comput.* **68** (1999) 913–943.
- [45] T.Y. Hou and X.H. Wu, A multiscale finite element method for elliptic problems in composite materials and porous media. *J. Comput. Phys.* **134** (1997) 169–189.
- [46] V.V. Jikov, S.M. Kozlov and O.A. Oleinik, *Homogenization of Differential Operators and Integral Functionals.* Springer-Verlag (1991).
- [47] E. Kosygina, F. Rezakhanlou and S.R.S. Varadhan, Stochastic homogenization of Hamilton-Jacobi-Bellman equations. *Comm. Pure Appl. Math.* **59** (2006) 1489–1521.
- [48] O. Kounchev and H. Render, Polyharmonic splines on grids $\mathbb{Z} \times a\mathbb{Z}^n$ and their limits. *Math. Comput.* **74** (2005) 1831–1841.
- [49] S.M. Kozlov, The averaging of random operators. *Mat. Sb. (N.S.)* **109** (1979) 188–202, 327.
- [50] P.-L. Lions and P.E. Souganidis, Correctors for the homogenization of Hamilton-Jacobi equations in the stationary ergodic setting. *Comm. Pure Appl. Math.* **56** (2003) 1501–1524.
- [51] W.R. Madych and S.A. Nelson, Multivariate interpolation and conditionally positive definite functions. *Approx. Theory Appl.* **4** (1988) 77–89.
- [52] W.R. Madych and S.A. Nelson, Multivariate interpolation and conditionally positive definite functions. II. *Math. Comput.* **54** (1990) 211–230.
- [53] W.R. Madych and S.A. Nelson, Polyharmonic cardinal splines. *J. Approx. Theory* **60** (1990) 141–156.
- [54] W.R. Madych and S.A. Nelson, Polyharmonic cardinal splines: a minimization property. *J. Approx. Theory* **63** (1990) 303–320.
- [55] A. Malqvist and D. Peterseim, Localization of elliptic multiscale problems. Technical report [arXiv:1110.0692](https://arxiv.org/abs/1110.0692) (2012).

- [56] O.V. Matveev, Some methods for the reconstruction of functions of n variables defined on chaotic grids. *Dokl. Akad. Nauk* **326** (1992) 605–609.
- [57] O.V. Matveev, Spline interpolation of functions of several variables and bases in Sobolev spaces. *Trudy Mat. Inst. Steklov.* **198** (1992) 125–152.
- [58] O.V. Matveev, Interpolation of functions on chaotic grids. *Dokl. Akad. Nauk* **339** (199)4 594–597.
- [59] O.V. Matveev, Methods for the approximate recovery of functions defined on chaotic grids. *Izv. Ross. Akad. Nauk Ser. Mat.* **60** 111–156, 1996.
- [60] O.V. Matveev, On a method for the interpolation of functions on chaotic grids. *Mat. Zametki* **62** (1997) 404–417.
- [61] J.M. Melenk, On n -widths for elliptic problems. *J. Math. Anal. Appl.* **247** (2000) 272–289.
- [62] G.W. Milton, The theory of composites, vol. 6 of *Cambridge Monographs on Applied and Computational Mathematics*. Cambridge University Press, Cambridge (2002).
- [63] P. Ming and X. Yue, Numerical methods for multiscale elliptic problems. *J. Comput. Phys.* **214** (2006) 421–445.
- [64] R. Moser, Theory of partial differential equations. *MA6000A. Lect. Notes* (2012). Available on <http://people.bath.ac.uk/rm257/MA6000A/notes.pdf>.
- [65] F. Murat and L. Tartar, H-convergence. Séminaire d'Analyse Fonctionnelle et Numérique de l'Université d'Alger (1978).
- [66] F.J. Narcowich, J.D. Ward and H. Wendland, Sobolev bounds on functions with scattered zeros, with applications to radial basis function surface fitting. *Math. Comput.* **74** (2005) 743–763.
- [67] G. Nguetseng, A general convergence result for a functional related to the theory of homogenization. *SIAM J. Math. Anal.* **20** (1989) 608–623.
- [68] J. Nolen, G. Papanicolaou and O. Pironneau, A framework for adaptive multiscale methods for elliptic problems. *Multiscale Model. Simul.* **7** (2008) 171–196.
- [69] H. Owhadi and L. Zhang, Metric-based upscaling. *Comm. Pure Appl. Math.* **60** (2007) 675–723.
- [70] H. Owhadi and L. Zhang, Localized bases for finite dimensional homogenization approximations with non-separated scales and high-contrast. *SIAM Multiscale Model. Simul.* **9** (2011) 1373–1398. [arXiv:1011.0986](https://arxiv.org/abs/1011.0986).
- [71] H. Owhadi, Anomalous slow diffusion from perpetual homogenization. *Ann. Probab.* **31** (2003) 1935–1969.
- [72] H. Owhadi, Averaging versus chaos in turbulent transport? *Comm. Math. Phys.* **247** (2004) 553–599.
- [73] G.C. Papanicolaou and S.R.S. Varadhan, Boundary value problems with rapidly oscillating random coefficients, in *Random fields, Vol. I, II (Esztergom (1979))*, vol. 27. *Colloq. Math. Soc. János Bolyai*. North-Holland, Amsterdam (1981) 835–873.
- [74] A. Pinkus, *N-Widths in Approximation Theory*. Springer-Verlag (1985).
- [75] C. Rabut, *B-splines Polyharmoniques Cardinales: Interpolation, Quasi-interpolation, filtrage*. Thèse d'État. Université de Toulouse (1990).
- [76] Ch. Rabut, Elementary m -harmonic cardinal B -splines. *Numer. Algorithms* **2** (1992) 39–61.
- [77] Ch. Rabut, High level m -harmonic cardinal B -splines. *Numer. Algorithms* **2** (1992) 63–84.
- [78] M. Rossini, Detecting discontinuities in two-dimensional signals sampled on a grid. *JNAIAM J. Numer. Anal. Ind. Appl. Math.* **4** (2009) 203–215.
- [79] I.J. Schoenberg, Cardinal spline interpolation. *Conference Board of the Mathematical Sciences Regional Conf. Ser. Appl. Math.* No. 12. Society for Industrial and Applied Mathematics, Philadelphia, Pa., (1973).
- [80] S. Spagnolo, Sulla convergenza di soluzioni di equazioni paraboliche ed ellittiche. *Ann. Scuola Norm. Sup. Pisa* **22** (1968) 571–597; *errata*, S. Spagnolo, *Ann. Scuola Norm. Sup. Pisa* **22** (1968) 673.
- [81] G. Stampacchia, Le problème de dirichlet pour les équations elliptiques du second ordre à coefficients discontinus. *Ann. Inst. Fourier (Grenoble)* **15** (1965) 189–258.
- [82] G. Stampacchia, *Équations elliptiques du second ordre à coefficients discontinus*. Séminaire Jean Leray No. 3 (1963–1964). Numdam (1964).
- [83] William Symes. Transfer of approximation and numerical homogenization of hyperbolic boundary value problems with a continuum of scales. *TR12-20 Rice Tech Report* (2012).
- [84] J.L. Taylor, S. Kim and R.M. Brown, The green function for elliptic systems in two dimensions. [arXiv:1205.1089](https://arxiv.org/abs/1205.1089) (2012).
- [85] J. Vybiral, Widths of embeddings in function spaces. *J. Complexity* **24** (2008) 545–570.
- [86] C.D. White and R.N. Horne, Computing absolute transmissibility in the presence of finescale heterogeneity. *SPE Symposium on Reservoir Simulation* 16011 (1987).
- [87] X.H. Wu, Y. Efendiev and T.Y. Hou, Analysis of upscaling absolute permeability. *Discrete Contin. Dyn. Syst. Ser. B* **2** (2002) 185–204.
- [88] V.V. Yurinskii, Averaging of symmetric diffusion in a random medium. *Sibirsk. Mat. Zh.* **27** (1986) 167–180.