



Polymorphism analysis of the *hsp70* stress gene in Broiler chickens (*Gallus gallus*) of different breeds

Carmen Maria Mazzi¹, Jesus Aparecido Ferro¹, Maria Inês Tiraboschi Ferro¹, Vicente José Maria Savino², Antonio Augusto Domingos Coelho² and Marcos Macari³

¹Universidade Estadual Paulista, Faculdade de Ciências Agrárias e Veterinárias, Departamento de Tecnologia, Jaboticabal, SP, Brazil.

²Universidade de São Paulo, Escola Superior de Agricultura "Luiz de Queiroz", Departamento de Zootecnia, Piracicaba, SP, Brazil.

³Universidade Estadual Paulista, Faculdade de Ciências Agrárias e Veterinárias, Departamento de Morfologia e Fisiologia Animal, Jaboticabal, SP, Brazil.

Abstract

The promoter region and the beginning of the coding region of the *hsp70* stress gene were analysed in broiler chickens of a commercial breed (Hubbard-Pettersen), a breed selected for weight gain (PP1) and a non-selected breed (naked-neck Label Rouge). The naked neck gene (*Naked neck*, *Na*), which reduces feathering in birds and is thus related to heat resistance, was present in both PP1 and Label Rouge breeds. Genomic DNA was restricted with *Pst*I and Southern blotting analysis of the samples revealed the absence of polymorphic sites for that enzyme in the promoter region and beginning of the coding region of the *hsp70* gene of studied birds. PCR-SSCP analysis of these regions, however, indicated the presence of polymorphisms in the beginning of the coding region and the sequencing of the PCR products confirmed and identified two polymorphic sites in this region: a transition A → G in position +258 and a transversion C → G in position +276. Both mutations were considered to be silent, since they did not modify the aminoacid sequence of the protein Hsp70. The promoter region of the *hsp70* gene was identical in all studied birds, indicating that the regulation pattern of this gene must be the same in all birds at the promoter region. Three different alleles (*hsp70-1*, *hsp70-2* and *hsp70-3*) were identified for the *hsp70* gene from the observed mutations. The allele *hsp70-3* was detected in only two breeds, Hubbard-Pettersen and PP1, but at a low frequency (0,016 and 0,006, respectively).

Key words: broiler chickens, *hsp70* gene, *naked neck* (*Na*) gene, polymorphism, SSCP.

Received: November 29, 2002; Accepted: May 29, 2003.

Introduction

Heat stress in birds is one of the major concerns in the poultry industry, since it causes high mortality and/or low productivity, especially during the hot season. The introgression of the gene *naked neck* (*Na*) in bird breeds seems to improve the resistance of the birds to heat. The introduction of this gene in commercial breeds might contribute to the production of birds with high genetic potential that show better performance at high temperatures. The correlation between the presence of the *Na* gene and the resistance to heat stress is due to the fact that this gene reduces feathering by about 20% in heterozygote birds

(*Na/na*) and 40% in homozygote birds (*Na/Na*). Eberhart and Washburn (1993) stated that feathering reduction in naked neck birds probably caused their greater ability in dissipating heat in exposed areas, which was later confirmed by Yahav *et al.* (1998). Naked neck birds from a breed selected for weight gain (PP1) and heterozygote for the *Na* gene showed lower mortality and weight loss during severe gradual heat stress (28-42 °C) when compared to normally feathered birds (Mazzi, 1998).

Physiologists and geneticists have shown a great interest in a specific response seen in all live organisms when submitted to any kind of stress. The main characteristic during this specific response to stress is the increased expression of the so-called stress proteins (Hsps), especially Hsp70, which has been shown to be one of the most conserved stress proteins. Besides its role in cytoprotection, this protein is also considered a cellular thermometer (Craig and Gross, 1991).

Since birds are constantly challenged by heat stress in hot countries, many studies involving Hsp70 expression in broilers have been made. Gabriel (1996) showed that the expression of Hsp70 in the hepatocytes of broiler chickens submitted to heat stress is heat- and time-dependent. A positive correlation was seen between the change in cloacal temperature and Hsp70 levels in the liver of naked neck birds at five days of age (Dionello, 1998). Hsp70 expression was also shown to be tissue-dependent when the levels of this protein were studied in different organs of chicken embryos subjected to heat stress (Givisiez *et al.*, 2001). Naked neck birds, from a breed selected for weight gain (PP1), showed different Hsp70 levels in hepatic cells when submitted to increasing gradual heat stress (28–36 °C). Furthermore, the birds that were more resistant to heat (*Na/na*) had lower Hsp70 levels when compared to dominant homozygote PP1 birds (*Na/Na*), and also to normally feathered birds (*na/na*) from the same breed (Mazzi *et al.*, 2002). Mahmoud (2000) analysed the *hsp70* gene in birds submitted to heat stress and found polymorphic sites located upstream from the coding region. The birds that were more resistant to heat showed only one *Pst*I *hsp70* allele of 6.48 kb, whereas the other breeds showed two different alleles for that gene. According to these data, it was concluded that polymorphisms in the *hsp70* gene might be used by commercial breeders to produce birds that are more tolerant to heat.

The present study analysed the sequence of the promoter region and the beginning of the coding region of the *hsp70* gene in birds from a breed selected for weight gain (PP1) showing three genotypes for *Na* gene (*Na/Na*, *Na/na* and *na/na*), and one commercial normally-feathered breed (Hubbard-Pettersen) and one non-selected breed that was heterozygous for *Na* gene (non-selected Label Rouge) were also included. The objective was to evaluate the presence of polymorphism in these regions of the *hsp70* gene. The existence of polymorphisms opens up the possibility that one of them might be associated to the phenotype that shows more tolerance to heat.

Material and Methods

Birds

This study used thirty birds of the Hubbard-Pettersen commercial breed, thirty naked-neck birds (*Na/na*) of the non-selected Label Rouge breed and ninety birds of the experimental breed PP1, thirty of each genotype (*Na/Na*, *Na/na* and *na/na*).

Genomic DNA extraction

Blood samples (1 mL) were collected in syringes containing 100 µL EDTA (0.5 M, pH 8.0). Genomic DNA was obtained from 50 µL of blood using microcentrifuge tubes containing 800 µL of cold TKM buffer (1 M Tris-HCl; 1 M KCl; 1 M MgCl₂; 0.1 M EDTA; 1 M NaCl). Fifty

microlitres of 10% SDS pre-warmed to 37 °C were added and the samples were incubated in water bath at 55 °C for 30 min. After 300 µL of 6 M NaCl were added, the samples were gently homogenised and centrifuged at 14,000 rpm for 20 min, at room temperature. The supernatant from each microtube was transferred to test tubes containing 5 mL cold 100% ethanol. The tubes were gently inverted until the DNA could be visualised, which was then recovered with a Pasteur pipette, transferred to a fresh microtube and washed with cold ethanol at 70%. After centrifugation at 14,000 rpm for 4 min, the DNA was air-dried for 5 min and then dissolved in 1 mL TE (10 mM Tris-HCl pH 8.0; 1 mM EDTA). The absorbance of the samples was determined at 260 and 280 nm using a Hitachi spectrophotometer (Model U-2000) and the DNA concentration was calculated (Sambrook *et al.*, 1989).

Restriction with *Pst*I and Southern Blotting

Six pools of genomic DNA (five birds/pool) were prepared for each studied group. The pools contained 6 µg DNA and were digested with the restriction enzyme *Pst*I (Gibco-BRL) for 8 h at 37 °C. The samples were then separated by electrophoresis in a 1% agarose gel for 6 h, at 50 V. After electrophoresis, the gels were left in 0.2 N HCl for 10 min, washed rapidly using Milli-Q water and then immersed in a denaturing solution (0.5 N NaOH; 1.5 M NaCl) for 20 min. This procedure was done twice to assure DNA denaturation and the gels were washed once more with Milli-Q water and immersed in a neutralising solution (1 M Tris-HCl pH 7.5; 1.5 M NaCl) for 20 min. The DNA was transferred from the gel to nylon membranes (Immobilon-Ny+; SIGMA, N-8522) by Southern Blotting (Sambrook *et al.*, 1989). After transference, the membranes were washed for 5 min with a wash solution (0.1 X SSC; 0.1% SDS; 50 mM Tris-HCl pH 7.5) and, shortly after, crosslinked at 160,000 µJ/cm² to fix the DNA to the membrane. The membranes were placed in hybridisation bottles and 30 mL of pre-hybridisation solution (6X SSC; 10X Denhardt's reagent; 1% SDS; 250 µg denatured salmon sperm DNA/mL) was added. The bottles were placed in a hybridisation oven (HYBAID Instruments) at 65 °C for 2 h. The pre-hybridisation solution was discarded and 15 mL of pre-warmed hybridisation solution (6X SSC; 1% SDS, 65 °C) was added with 10 µL of probe labelled with ³²P-dCTP by random priming (Feinberg and Volgestein, 1983). The probe was a DNA fragment specific for the chicken *hsp70* gene, amplified using the primers Hsp70-F1 and Hsp70-R3 (Table 1). The fragment length was 554 bp and it comprised both the promoter and the beginning of the coding region of *hsp70* gene. After 16 h of hybridisation, the solution was changed to 50 mL of washing solution 1 (2X SSC, 0.1% SDS), pre-warmed to 65 °C. The membranes were washed twice for 20 min using washing solution 2 (0.1X SSC; 0.1% SDS, 65 °C). Sheets of X-Ray film

Table 1 - Primers used on PCR amplification of the chicken *hsp70* gene.

Primers	Sequence	Position on reference sequence
<i>Hsp70</i> -F1	5'GAGTGGCGCAGCGTAGAAAG 3'	18
<i>Hsp70</i> -F2	5'GATTGGTCCTTAGCGTTCTGGC 3'	208
<i>Hsp70</i> -F3	5'TCATCATGTCTGGCAAAGGG 3'	387
<i>Hsp70</i> -F4	5'AACCGCACACACCCAGCTATG 3'	497
<i>Hsp70</i> -F19	5'CAACAGAGATAGGGTGGGAG 3'	1993
<i>Hsp70</i> -R1	5'TTCCTTTGGTCAGTCAGCC 3'	382
<i>Hsp70</i> -R2	5'TGATCTCCACTTTGCCATGCTG 3'	479
<i>Hsp70</i> -R3	5'CACTTGGTTCTTGGCAGCATC 3'	571
<i>Hsp70</i> -R3b	5'CTGGGAGTCGTTGAAGTAAGCG 3'	856
<i>Hsp70</i> -R24	5'TGCCTTTATACACACCAACAG 3'	2426

The letters F and R in primer names refer to their orientation (forward and reverse, respectively).

were exposed to the membranes using adequate cassettes for 72 h at -70 °C and developed.

PCR amplification of the chicken *hsp70* gene

The chicken *hsp70* gene sequence deposited in GenBank under the accession number J02579 (Morimoto *et al.*, 1986) was called the reference sequence in the present study. The Gene Runner software (version 3.0) was used to design primers from this sequence (Table 1). The primers were used in PCR reactions to amplify specific regions of the *hsp70* gene, and reactions were assembled using the following combination of components: 1X Taq reaction buffer (Gibco-BRL), 0.2 pmol primer F, 0.2 pmol primer R, 0.1 mM dNTPs, 2 mM MgCl₂, 100 ng genomic DNA and 1 unit Taq DNA polymerase (Gibco-BRL) to a final volume of 30 µL.

The following regions of the *hsp70* gene were amplified: F1-R1 (364 bp), F2-R2 (271 bp), F3-R3 (184 bp), F4-R3b (359 bp) and F19-R24 (433 bp). The PCR program for F1-R1 was: initial denaturing at 95 °C for 2 min and 35 cycles of 95 °C for 1 min (denaturing), 60 °C for 1 min (primer annealing) and 72 °C for 1 min and 30 s (extension by Taq). The other reactions used similar PCR programs, but annealing temperatures were 65, 64.5, 64.5 and 61.5 °C, respectively. The positions of the primers in the *hsp70* gene sequence are shown in Figure 1.

SSCP analysis

After visualisation of the amplification products in agarose gel (1%), a 3-µL aliquot of each sample was diluted in 6 µL SSCP buffer (95% formamide; 20 mM EDTA; 0.05% bromophenol blue; 0.05% xylene-cyanol). The samples were then denatured for 10 min at 95 °C and incubated in ice for the same period before electrophoresis in a non-denaturing acrylamide:bis-acrylamide (49:1) gel, using the Bio-Rad mini-protean II system. The gels were composed of 600 µL 10X TBE (0.9 M Tris; 0.02 M EDTA; 0.9 M boric acid), acrylamide:bis (49:1) up to the desired concentra-




Figure 1 - Schematic representation of the position of the primers that were used in PCR-SSCP for detection of polymorphisms in the chicken *hsp70* gene.

tion, 60 µL 10% APS and 6 µL TEMED and Milli-Q water (qsp 6mL). Both the concentration of acrylamide and the running time were standardised for each analysed PCR product (15% for F1-R1, F2-R2 and F4-R3b; 20% for F3-R3 and 12% for F19-R24). The gels were silver-stained according to Bassan *et al.* (1991) so that the bands could be visualised.

Sequencing of the PCR products

In order to obtain the sequence of the promoter region and the beginning of the coding region of *hsp70* gene, the sequences F1-R2 (461bp) and F2-R3b (648 bp) were amplified for all birds using the primers shown in Table 1. The final volume of these reactions was 50 µL and the reagents were added as follows: 100 ng genomic DNA, 0.2 pmol primer F; 0.2 pmol primer R; 2 mM MgSO₄, 0.1 mM dNTP and one unit High Fidelity Taq DNA Polymerase (Invitrogen). The PCR program consisted of an initial denaturing (95 °C) for 2 min and 35 cycles of 95 °C for 1 min, 62.5 °C and 62 °C for 1 min (annealing of primer pairs F1-R2 and F2-R3b, respectively) and 68 °C for 1 min and 30 s (extension by Taq). Once the amplification was confirmed in 1% agarose gel, the PCR products were purified using the Concert[™] Rapid PCR Purification System Kit (Gibco-BRL). An aliquot of the purified product was used for quantification in 1% agarose gel and approximately 50-100 ng of the products were used for sequencing in a reaction with sequencing buffer (0.5X), 2 µL of Big Dye (version 3), 5 pmol of primer F or R and sterile water to a final

volume of 10 μ L. After 35 reaction cycles (95 °C for 1 min; 62.5 °C for 1 min and 72 °C for 1 min 30 s), the samples were washed with 80 μ L of 75% isopropanol for 15 min and centrifuged at 4,500 rpm for 25 min at 20 °C. The samples were washed using 200 μ L of 70% ethanol and centrifuged at 4,500 rpm for 15 min at 20 °C. The samples were then vacuum dried and loaded to a polyacrylamide gel in the ABI-377 DNA sequencer (Perkin Elmer).

Results

The results of Southern Blotting analysis of the total DNA digested with *Pst*I and hybridised with the probe specific for the chicken *hsp70* gene showed no sites for *Pst*I in the promoter region or the beginning of the coding region of the *hsp70* gene. One single 6 kb fragment hybridised with the probe in all animals (data not shown).

The PCR products F1-R1, F2-R2, F3-R3 and F19-R24 did not show any difference in their migration patterns in polyacrylamide gels when submitted to SSCP analysis. Nevertheless, different migration patterns in polyacrylamide gels were seen when F4-R3b fragments were analysed using this technique (Figure 2), indicating that this region (beginning of coding region) presents polymorphisms.

Data from sequencing of the promoter region and the beginning of the coding region of the *hsp70* gene were in accordance with the observations obtained by SSCP analysis of these regions. In all birds, the nucleotide sequence of the *hsp70* promoter region was identical. On the other hand,




Figure 2 - Migration pattern of F4-R3b *hsp70* fragment (359 bp) in acrylamide gel for some DNA chicken samples. H - Hubbard, R - PP1 *na/na* and h - PP1 *Na/na*.

at the beginning of the coding region, two polymorphic sites were found: one transition A \rightarrow G on position +258 and one transversion C \rightarrow G on position +276. Polymorphic sequences were submitted to BLASTx from NCBI and the two polymorphic sites were considered as silent mutation sites, since they did not cause a change in the aminoacid sequence of the *Hsp70* protein.

The polymorphisms found here and the combinations between them in the constitution of the genotype of the birds indicated the presence of three alleles for the chicken *hsp70* gene, which were called *hsp70-1*, *hsp70-2* and *hsp70-3*. The nucleotide sequences of the three alleles were deposited in the GenBank with accession numbers AY178441, AY178442 and AY178443 for *hsp70-1*, *hsp70-2* and *hsp70-3*, respectively.

One alignment of the three alleles using the software Clustal W is shown on Figure 3; differences among them only occur in positions +258 and +276. Figure 3 also shows the alignment between the three sequences and the reference sequence (Morimoto *et al.*, 1986). Some changes in the nucleotide sequences may be observed, specially in the




Figure 3 - Alignment (CLUSTALW) of alleles *hsp70-1*, *hsp70-2* and *hsp70-3* with the reference sequence (J02579 - Morimoto *et al.*, 1986). Underlined: conserved regions CAAT, HSEs 1 and 2, TATA box and initial ATG; in bold: transitions and transversions seen among the three alleles and the reference *hsp70* gene.

<i>hsp70-1</i>	GAAGGCTAAGCGGACTATAAAGAGGGCGCGAGCGGGCGCCGTAACGGCAGATCGCGCCGCA	
<i>hsp70-2</i>	GAAGGCTAAGCGGACTATAAAGAGGGCGCGAGCGGGCGCCGTAACGGCAGATCGCGCCGCA	
<i>hsp70-3</i>	GAAGGCTAAGCGGACTATAAAGAGGGCGCGAGCGGGCGCCGTAACGGCAGATCGCGCCGCA	
Ref. <i>hsp70</i>	GAAGGCTAAGCGGACTATAAAGAGGGCGCGA - CGGC - -CGTAACGGCAGATCGCGCCGCA	- 95

<i>hsp70-1</i>	GACAGCAGCGAGA - GCGGGCGGAGGAGACGTGACTGCGAGCGAGCAAGTACTGGCGGAG	
<i>hsp70-2</i>	GACAGCAGCGAGA - GCGGGCGGAGGAGACGTGACTGCGAGCGAGCAAGTACTGGCGGAG	
<i>hsp70-3</i>	GACAGCAGCGAGA - GCGGGCGGAGGAGACGTGACTGCGAGCGAGCAAGTACTGGCGGAG	
Ref. <i>hsp70</i>	GACAGCAGCGAGAAGCGGGCGGAGGAGACGTGACTGCGAGCGAGCAAGTACTGGCGGAG	- 35

<i>hsp70-1</i>	CGAGTGGCTGACTGACCA - GAGGAATCTATCATCATGTCTGGCAAAGGGCCGGCCATCGG	
<i>hsp70-2</i>	CGAGTGGCTGACTGACCA - GAGGAATCTATCATCATGTCTGGCAAAGGGCCGGCCATCGG	
<i>hsp70-3</i>	CGAGTGGCTGACTGACCA - GAGGAATCTATCATCATGTCTGGCAAAGGGCCGGCCATCGG	
Ref. <i>hsp70</i>	CGAGTGGCTGACTGACCAAGAGGAATCTATCATCATGTCTGGCAAAGGGCCGGCCATCGG	+26

+1		
<i>hsp70-1</i>	CATCGATCTGGGCACCACGTATTCTTGCGTGGGTGTCTTCCAGCATGGCAAAGTGGAGAT	
<i>hsp70-2</i>	CATCGATCTGGGCACCACGTATTCTTGCGTGGGTGTCTTCCAGCATGGCAAAGTGGAGAT	
<i>hsp70-3</i>	CATCGATCTGGGCACCACGTATTCTTGCGTGGGTGTCTTCCAGCATGGCAAAGTGGAGAT	
Ref. <i>hsp70</i>	CATCGATCTGGGCACCACGTATTCTTGCGTGGGTGTCTTCCAGCATGGCAAAGTGGAGAT	+86

<i>hsp70-1</i>	CATTGCCAACGACCAGGGGAACCGCACCACACCCAGCTATGTGGCCTTCACCGATACAGA	
<i>hsp70-2</i>	CATTGCCAACGACCAGGGGAACCGCACCACACCCAGCTATGTGGCCTTCACCGATACAGA	
<i>hsp70-3</i>	CATTGCCAACGACCAGGGGAACCGCACCACACCCAGCTATGTGGCCTTCACCGATACAGA	
Ref. <i>hsp70</i>	CATTGCCAACGACCAGGGGAACCGCACCACACCCAGCTATGTGGCCTTCACCGATACAGA	+146

<i>hsp70-1</i>	GCGCCTCATCGGGGATGCTGCCAAGAACCAAGTGGCAATGAACCCACCAACACCATCTT	
<i>hsp70-2</i>	GCGCCTCATCGGGGATGCTGCCAAGAACCAAGTGGCAATGAACCCACCAACACCATCTT	
<i>hsp70-3</i>	GCGCCTCATCGGGGATGCTGCCAAGAACCAAGTGGCAATGAACCCACCAACACCATCTT	
Ref. <i>hsp70</i>	GCGCCTCATCGGGGATGCTGCCAAGAACCAAGTGGCAATGAACCCACCAACACCATCTT	+206

<i>hsp70-1</i>	TGATGCCAAGCGTCTCATCGGCCGCAAGTATGATGACCCACAGTGCAGTCCGACATGAA	
<i>hsp70-2</i>	TGATGCCAAGCGTCTCATCGGCCGCAAGTATGATGACCCACAGTGCAGTCCGACATGAA	
<i>hsp70-3</i>	TGATGCCAAGCGTCTCATCGGCCGCAAGTATGATGACCCACAGTGCAGTCCGACATGAA	
Ref. <i>hsp70</i>	TGATGCCAAGCGTCTCATCGGCCGCAAGTATGATGACCCACAGTGCAGTCCGACATGAA	+266

<i>hsp70-1</i>	GCACTGGCCCTTCCGTGTGGTGAACGAGGGTGGCAAGCCCAAGGTGCAGGTGGAGTACAA	
<i>hsp70-2</i>	GCACTGGCCCTTCCGTGTGGTGAACGAGGGTGGCAAGCCCAAGGTGCAGGTGGAGTACAA	
<i>hsp70-3</i>	GCACTGGCCCTTCCGTGTGGTGAACGAGGGTGGCAAGCCCAAGGTGCAGGTGGAGTACAA	
Ref. <i>hsp70</i>	GCACTGGCCCTTCCGTGTGGTGAACGAGGGTGGCAAGCCCAAGGTGCAGGTGGAGTACAA	+326

<i>hsp70-1</i>	GGGTGAGATGAAGACCTTCTTCCAGAGGAGATCAGCTCTATGGTGCTACCAAGATGAA	
<i>hsp70-2</i>	GGGTGAGATGAAGACCTTCTTCCAGAGGAGATCAGCTCTATGGTGCTACCAAGATGAA	
<i>hsp70-3</i>	GGGTGAGATGAAGACCTTCTTCCAGAGGAGATCAGCTCTATGGTGCTACCAAGATGAA	
Ref. <i>hsp70</i>	GGGTGAGATGAAGACCTTCTTCCAGAGGAGATCAGCTCTATGGTGCTACCAAGATGAA	+386

Figure 3 (Cont.)

promoter region: a G deletion on position -218, a C deletion on position -214, two G insertions on positions -123 and -118, one C insertion on position -117 and two A deletions on positions -81 and -16. When the three alleles were compared to the reference sequence, it was observed that on positions +258 and +276, where the polymorphism occurs in

the studied population, the allele *hsp70-3* showed nucleotides that are identical to the reference sequence, *i.e.*, one nucleotide A (+258) and one C (+276).

The frequencies of each allele within the populations are presented in Table 2. The *hsp70-1* allele had the highest frequency in all groups of birds when compared to the other

Table 2 - Allelic and genotypic frequencies observed for each breed (n = 30 for Hubbard and Label Rouge and n = 90 for PP1).

Breed	Alleles				
	<i>hsp70-1</i>	<i>hsp70-2</i>	<i>hsp70-3</i>		
Hubbard	0.633	0.35	0.016		
PP1	0.82	0.15	0.006		
Label Rouge	0.683	0.316	-		
	Genotypes				
	<i>hsp70-1/hsp70-1</i>	<i>hsp70-2/hsp70-2</i>	<i>hsp70-1/hsp70-2</i>	<i>hsp70-1/hsp70-3</i>	<i>hsp70-2/hsp70-3</i>
Hubbard	0.33	0.03	0.60	-	0.03
PP1	0.67	0.08	0.20	0.03	0.01
Label Rouge	0.34	-	0.63	-	-

alleles. A low frequency of the *hsp70-3* allele was seen in populations PP1 (0.006) and Hubbard (0.016), while this allele was not seen in the non-selected Label Rouge population. Table 2 also shows the frequency of the five different genotypes observed in the populations. The highest frequency of the *hsp70-1/hsp70-1* genotype was seen in PP1 breed, whereas the commercial (Hubbard) and non-selected (Label Rouge) populations showed a higher frequency of the *hsp70-1/hsp70-2* genotype. The other genotypes had a low frequency, and the *hsp70-1/hsp70-3* genotype was seen only in the population PP1. Only the *hsp70-1/hsp70-1* and *hsp70-1/hsp70-2* genotypes were seen within the Label Rouge population, while the other groups presented at least three different genotypes. The values from the chi-square (χ^2) test for genotype frequencies found here indicated that all were in equilibrium (Hubbard = 0,3, $p > 0,05$; PP1 $Na/Na = 0,002$, $p > 0,05$; PP1 $Na/na = 0,33$ $p > 0,05$; PP1 $na/na = 0,022$, $p > 0,05$; Label Rouge = 0,093, $p > 0,05$).

Discussion

The absence of restriction sites for the *PstI* enzyme in the promoter region and the beginning of the coding region of the *hsp70* gene in the birds evaluated in this study is a different result from that reported by Mahmoud (2000), who found three different allelic fragments for the chicken *hsp70* gene when genomic DNA was restricted using *PstI*. The author also observed the presence of an *hsp70 PstI* allele of 6.48 kb in the birds that were more resistant to heat, and concluded that the polymorphisms in the *hsp70* gene exist in different bird populations and that it is possible that these polymorphisms might be used by commercial breeders to produce more tolerant birds.

Polymorphisms in the *hsp70* gene with *PstI* were also observed in other species, such as pigs (Ruohonen-Lehto *et al.*, 1993). Likewise, in patients with SLE (Systemic Lupus Erythematosus), a higher frequency was observed of the 8.5-kb *PstI* allele of the gene *hsp70-2*, which is heat-inducible, while normal individuals had a higher frequency

of the common 9-kb allele of *hsp70-2*. The presence of the polymorphic allele (8.5 kb) was associated with SLE susceptibility (Jarjour *et al.*, 1996). A frequency of 21.8% was observed for the *hsp70-2/PstI* allele in patients with MLV (Mediterranean Visceral Leishmaniosis), whereas normal individuals showed a 12.6% frequency of this allele (Meddeb-Garnaoui *et al.*, 2001). The same *PstI* allele of 8.5 kb was also associated with susceptibility to insulin-dependent diabetes mellitus (IDDM, Chuang *et al.*, 1996).

Different restriction patterns of the enzymes *HinfI* and *ScaI* were observed for the *hsp70* gene in *Cryptosporidium parvum* (Gobet and Toze, 2001); of *EcoRI*, *BamHI*, *DraI*, *SaII* and *HindIII* in nematodes (Hashmi *et al.*, 1997); of *BamHI* and *ClaI* in *Drosophyla* (Zatsepina *et al.*, 2001), of the *PvuII* enzyme in pigs (Ruohonen-Lehto *et al.*, 1993), and of the *BamHI* enzyme in humans susceptible to hypertension (Hamet *et al.*, 1994).

The absence of polymorphisms in the promoter region of the *hsp70* gene in the studied birds seems to indicate that this gene presents the same regulation pattern for this region in the different bird populations. Considering these observations, the differential expression of Hsp70 protein reported for PP1 birds submitted to graded heat stress of 28 to 36 °C (Mazzi *et al.*, 2002) does not seem to be related to the presence of polymorphisms in the promoter region of *hsp70* gene.

Zhang *et al.* (2002) detected, by PCR-SSCP and sequencing, polymorphism in regulatory and coding regions of the *hsp70* gene in chickens with different heat tolerance capability. Huang *et al.* (2002) detected single nucleotide polymorphisms (SNPs) in the 5' flanking region of the *hsp70-2* gene in boars. These SNPs have been associated with the quality of semen production during the summer and can, also, be used as markers for semen quality selection in these animals.

In our study, PCR-SSCP techniques were effective in revealing different genotypes of birds in the studied populations. Gasser *et al.* (2001) were also able to detect different *Cryptosporidium parvum* genotypes in diverse hosts using the same techniques.

The alignment of the three alleles (*hsp70-1*, *hsp70-2* and *hsp70-3*) detected in the present study and the *hsp70* gene sequence of Morimoto *et al.* (1986), considered as a reference sequence (Figure 3), revealed that among the three alleles, *hsp70-3* is the most similar to the reference gene, indicating that this allele must be the older within the population. The alterations seen in the promoter region of the alleles when compared to the reference sequence did not affect any critical regulatory regions such as the heat shock elements (HSE 1 and HSE 2), positioned at -170 to -158 and -180 to -166, respectively, where the transcription factors (HSFs) bind during the activation of the gene. The regions CAAT (-116 and -267) and TATA box (-139) were not affected by point mutations either.

Acknowledgments

The authors thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for financial support.

References

- Bassan BJ, Anollés GCE and Gresshoff PM (1991) Fast and sensitive silver staining of DNA in polyacrylamide gels. *Anal Biochem* 196:80-3.
- Chuang LM, Jou TS, Wu HP, Tai TY and Lin BJA (1996) Rapid method to study heat shock protein 70-2 gene polymorphism in insulin-dependent diabetes mellitus. *Pancreas* 13:268-272.
- Craig EA and Gross CA (1991) Is *hsp70* the cellular thermometer? *TIBS* 16:135-40.
- Dionello NJL (1998) Efeito do estresse calórico agudo em frangos de corte (2-5 dias de idade) sobre a síntese da proteína de choque térmico de 70 kD (*Hsp70*) e de seu RNA mensageiro em tecido hepático e cerebral. Thesis, Universidade Federal de Pelotas, Rio Grande do Sul.
- Eberhart DE and Washburn KW (1993) Assessing the effects of the naked neck gene on chronic heat stress resistance in two genetic populations. *Poult Sci* 72:1391-9.
- Feinberg AP and Volgestein BA (1983) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* 132:6-13.
- Gabriel JE, Ferro JA, Stefani RMP, Ferro MIT, Gomes SL and Macari M (1996) Effect of acute heat stress on the heat shock protein 70 messenger RNA on the heat shock protein expression in the liver of broilers. *Br Poult Sci* 37:443-9.
- Gasser RB, Zhu X, Cacio S, Chalmers R, Widmer G, Morgan UM, Andrew-Thompson RC, Pozio E and Bowring GF (2001) Genotyping *Cryptosporidium parvum* by single-strand conformation polymorphism analysis of ribosomal and heat shock gene regions. *Electrophoresis* 22:433-7.
- Givisiez, PEN, Silva MM, Mazzi CM, Ferro MIT, Ferro JA, Gonzalez E and Macari M (2001) Heat or cold chronic stress affects organ weights and *Hsp70* levels in chicken embryos. *Can J Anim Sci* 82:83-7.
- Gobet P and Toze S (2001) Sensitive Genotyping of *Cryptosporidium parvum* by PCR-RFLP Analysis of the 70-kilodalton Heat Shock Protein (*Hsp70*) Gene. *FEMS Microb Letters* 200:37-41.
- Hamet P, Sun YL, Malo D, Kong D, Kren V, Pravenec M, Kunes J, Dumas P, Richard L and Gagnon F (1994) Genes of stress in experimental hypertension. *Clin Exp Pharmacol Physiol* 21:907-11.
- Hashmi G, Hashmi S, Selvan S, Grewal P and Gaugler R (1997) Polymorphism in Heat Shock Protein Gene (*hsp70*) in Entomopathogenic Nematodes (Rhabditida). *J Therm Biol* 22:143-9.
- Huang SY, Chen MY, Lin EC, Tsou HL, Kuo YH, Ju CC and Lee WC (2002) Effects of single nucleotide polymorphisms in the 5'-flanking region of heat shock protein 70.2 gene on semen quality in boars. *Anim Reprod Sci* 70:99-109.
- Jarjour W, Reed AM, Gauthier J, Hunt SIII and Winfield JB (1996) The 8.5-kb *PstI* Allele of the Stress Protein Gene, *Hsp70-2*. *Human Immunology* 45:59-63.
- Mahmoud Kamel (2000) Genetic and environmental variations of chicken heat shock proteins. PhD Thesis, North Carolina State University, North Carolina.
- Mazzi CM (1998) Análise da expressão da proteína de estresse *Hsp70* em frangos de corte portadores do gene "naked neck" (pescoço pelado) submetidos a estresse térmico gradativo. Dissertação, Universidade Estadual Paulista, Jaboticabal, São Paulo.
- Mazzi CM, Ferro MIT, Coelho AAD, Savino VJM, Macari M, Ferro JA, Givisiez PEN, Giachetto PF, Silva MM and Dionello NJL (2002) Effect of heat exposure on the thermoregulatory responses of selected naked neck chickens. *Arq Bras Med Vet Zootec* 54:35-41.
- Meddeb-Garnaoui A, Gritli S, Garbouj S, Ben Fadhel M, El Kares R, Mansour L, Kaabi B, Chouchane L, Ben Salah A and Dellagi, K (2001) Association Analysis of HLA-Class II and Class III Gene Polymorphisms in the Susceptibility to Mediterranean Visceral Leishmaniasis. *Hum Immunol* 62:509-17.
- Morimoto RI, Hunt C, Huang S, Berg KL and Banerji SS (1986) Organization, Nucleotide Sequence, and Transcription of Chicken *HSP70* Gene. *J Biol Chem* 25:12692-99.
- Ruohonen-Lehto MK, Rothschild MF and Larson RG (1993) Restriction fragment length polymorphisms at the heat shock protein *Hsp70* gene (s) in pigs. *Anim Genet* 24:67-8.
- Sambrook J, Fritsch EF and Maniatis T (1989) *Molecular Cloning: a laboratory manual*. 2nd edition. Cold Spring Harbor Press, New York, 2:9.31-57.
- Yahav S, Luger D, Cahaner A, Dotan M, Rusal M and Hurwitz S (1998) Thermoregulation in naked neck chickens subjected to different ambient temperatures. *Br Poult Sci* 39:133-8.
- Zatsepina OG, Velikodvorskaia VV, Molodtsov VB, Garbuz D, Lermn DN, Bettencourt BR, Feder ME and Evgenev MB (2001) A *Drosophila melanogaster* Breed from Sub-Equatorial Africa has exceptional thermotolerance but decreased *Hsp70* expression. *J Exp Biol* 204:1869-81.
- Zhang X, Du H and Li J (2002) Single nucleotide polymorphism of chicken heat shock protein 70 gene. 7th World Congress on Genetics Applied to Livestock Production, Montpellier, France.