

Polymorphisms in *Pfmdr1*, *Pfcr1*, and *Pfnhe1* Genes Are Associated with Reduced *In Vitro* Activities of Quinine in *Plasmodium falciparum* Isolates from Western Kenya

Jelagat Cheruiyot,^{a,b} Luicer A. Ingasia,^a Angela A. Omondi,^{a,b} Dennis W. Juma,^a Benjamin H. Opot,^a Joseph M. Ndegwa,^{a,b} Joan Mativo,^a Agnes C. Cheruiyot,^{a,b} Redemptah Yeda,^a Charles Okudo,^a Peninah Muiruri,^a Ngalah S. Bidii,^a Lorna J. Chebon,^a Paul O. Angienda,^b Fredrick L. Eyase,^a Jacob D. Johnson,^a Wallace D. Bulimo,^a Ben Andagalu,^a Hoseah M. Akala,^a Edwin Kamau^a

Global Emerging Infections Surveillance Program, United States Army Medical Research Unit—Kenya, Kenya Medical Research Institute—Walter Reed Project, Kisumu and Nairobi, Kenya^a; School of Biological/Physical Sciences, Department of Zoology, Maseno University, Maseno, Kenya^b

In combination with antibiotics, quinine is recommended as the second-line treatment for uncomplicated malaria, an alternative first-line treatment for severe malaria, and for treatment of malaria in the first trimester of pregnancy. Quinine has been shown to have frequent clinical failures, and yet the mechanisms of action and resistance have not been fully elucidated. However, resistance is linked to polymorphisms in multiple genes, including multidrug resistance 1 (*Pfmdr1*), the chloroquine resistance transporter (*Pfcr1*), and the sodium/hydrogen exchanger gene (*Pfnhe1*). Here, we investigated the association between *in vitro* quinine susceptibility and genetic polymorphisms in *Pfmdr1* codons 86 and 184, *Pfcr1* codon 76, and *Pfnhe1* ms4760 in 88 field isolates from western Kenya. *In vitro* activity was assessed based on the drug concentration that inhibited 50% of parasite growth (the IC₅₀), and parasite genetic polymorphisms were determined from DNA sequencing. Data revealed there were significant associations between polymorphism in *Pfmdr1*-86Y, *Pfmdr1*-184F, or *Pfcr1*-76T and quinine susceptibility ($P < 0.0001$ for all three associations). Eighty-two percent of parasites resistant to quinine carried mutant alleles at these codons (*Pfmdr1*-86Y, *Pfmdr1*-184F, and *Pfcr1*-76T), whereas 74% of parasites susceptible to quinine carried the wild-type allele (*Pfmdr1*-N86, *Pfmdr1*-Y184, and *Pfcr1*-K76, respectively). In addition, quinine IC₅₀ values for parasites with *Pfnhe1* ms4760 3 DNNND repeats were significantly higher than for those with 1 or 2 repeats ($P = 0.033$ and $P = 0.0043$, respectively). Clinical efficacy studies are now required to confirm the validity of these markers and the importance of parasite genetic background.

Quinine (QN), a quinoline derivative, is used in many African countries as second-line treatment for uncomplicated malaria, as an alternate first-line treatment of severe malaria, and for treatment of malaria in the first trimester of pregnancy (1). World Health Organization (WHO) guidelines recommend a combination of QN plus doxycycline, tetracycline, or clindamycin for treatment of malaria (2). However, in most African countries, QN is used as a monotherapy (1, 2), probably due to the high cost of antibiotics (3). QN has been shown to have frequent clinical failures in Southeast Asia (4, 5), South America (5), and Africa (6–8). In a clinical trial conducted in Kenya concerning the treatment of severe *Plasmodium falciparum* malaria, QN was shown to have longer clearance times, longer fever clearance times, and higher recrudescence rates than malarone (9). The high rates of QN clinical failures can be explained by variations in pharmacokinetics, drug quality, and treatment compliance, but less so to resistance. Parasites with high-grade resistance to QN have not been documented (3). However, *in vitro* analysis has shown reduced sensitivity to QN in Asia (10) and South America (11), and much less so in Africa (12, 13). The high rates of reinfection and recrudescence can also be explained by the short half-life of QN (14).

Although the mechanisms of action and resistance to QN have not been fully elucidated, inhibition of heme detoxification in the parasite digestive vacuole has been implicated in QN antimalarial activity (15), and resistance is linked to polymorphisms in multiple genes, including the multidrug resistance 1 gene (*Pfmdr1*), chloroquine resistance transporter gene (*Pfcr1*), and the sodium/hydrogen exchanger gene (*Pfnhe1*) (16). In particular, mutations in *Pfmdr1* at codons 86, 184, 1042, and 1246 and in *Pfcr1* at codon

76 have been associated with reduced QN sensitivity (17–19). Interestingly, however, in some studies polymorphism in *Pfcr1*-76 has been shown not to have any effect on QN activity (20, 21), and the *Pfmdr1*-86 mutation only resulted in a decreased QN activity that did not reach statistical significance (20). Other novel mutations in the *Pfcr1* gene that have recently been shown to alter QN sensitivity are Q352K/R and C350R (20, 22, 23).

Pfnhe1 is a 5,760-bp gene that encodes 1,920 amino acids, with a microsatellite polymorphism consisting of variable DNNND repeat units, designated ms4760. Sequence polymorphisms in the *Pfnhe1* gene have been analyzed in laboratory strains and field isolates with varied susceptibilities to QN (16, 24–27). Several variants of ms4760 have been described in which ms4760–1, with 2 copies of the DNNND repeat unit, was significantly associated with reduced *in vitro* QN sensitivity in laboratory clones (16) and in field isolates (24). On the contrary, other studies have not found an association between polymorphisms in the *Pfnhe1* gene and

Received 6 February 2014 Returned for modification 23 March 2014

Accepted 11 April 2014

Published ahead of print 21 April 2014

Address correspondence to Edwin Kamau, Edwin.Kamau@us.army.mil.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.02472-14>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.02472-14

The authors have paid a fee to allow immediate free access to this article.

QN reduced susceptibility (26, 27). These conflicting results may indicate that the influence of *Pfnhe1* on QN susceptibility is strain dependent and can vary depending on the geographic origin of the parasites (28). Polymorphisms in the *Pfnhe1* ms4760 DDNHND NHHND and DDNNNDNHHND repeats have also been associated with varied *in vitro* QN susceptibilities (24–27, 29, 30).

Similar to most countries in Africa, QN is used as second-line treatment for uncomplicated malaria and treatment of severe malaria in Kenya (9, 12). A recent study analyzed the association of *in vitro* activities of QN with *Pfmdr1* codon 86, *Pfcr1* codon 76, and *Pfnhe1* polymorphism in isolates collected from Kilifi District, which is located on the Kenyan coast (24). These authors observed *Pfmdr1*-86 mutants, and 2 copies of DDNNND repeat units of *Pfnhe1* were associated with a decrease in QN susceptibility. In this study, we investigated the association between *in vitro* QN susceptibility and genetic polymorphisms in *Pfmdr1* codons 86 and 184, *Pfcr1* codon 76, and *Pfnhe1* ms4760 in field isolates from western Kenya, where malaria is holoendemic.

MATERIALS AND METHODS

Plasmodium falciparum parasites. *P. falciparum* field isolates used in this study were collected from patients with uncomplicated malaria, ages 6 months and older, attending outpatient clinics in Kisumu, Kisii, and Kericho District hospitals in western Kenya between January 2010 and December 2011. Details of this protocol have been described elsewhere (31). The research protocol was approved by the Ethical Review Committee of the Kenya Medical Research Institute (KEMRI number 1330) and the Walter Reed Army Institute of Research institutional review board (WRAIR number 1384). *P. falciparum* reference strains 3D7 (sensitive) and W2 (resistant) were used as controls. These clones were obtained from cryopreserved stocks and culture adapted for drug sensitivity assays.

In vitro drug sensitivity tests. QN was obtained from the Walter Reed Army Institute of Research (WRAIR), Silver Spring, MD. Parasites were maintained in continuous culture by using a method described elsewhere (32). A previously described SYBR green I-based drug sensitivity assay was used for *in vitro* drug sensitivity testing (33–35). *P. falciparum* parasites in continuous culture attaining 3 to 8% parasitemia (field isolates or reference strains) were adjusted to 2% hematocrit and 1% parasitemia. QN was prepared in 5 ml of 70% ethanol to attain 5 mg/ml, which was lowered to 2,554 nM in tissue culture medium. QN was then diluted across 10 concentrations ranging from 2,554 nM to 5 nM and predosed on 96-well microtiter plates as previously described (36). The assay was initiated by addition of 100 μ l reconstituted parasite components to each well of the drug plates and incubation at 37°C as previously described (35). The assay was terminated after 72 h, 100 μ l of lysis buffer containing SYBR green I (1 \times , final concentration) was added directly to the plates, and the mixtures were kept at room temperature in the dark for 24 h. Parasite replication inhibition was quantified by measuring the per well relative fluorescence units (RFU) of SYBR green I dye, using the Tecan Genios Plus system with excitation and emission wavelengths of 485 nm and 535 nm. The 50% inhibition concentration (IC₅₀) values for QN were calculated as described previously (31).

DNA extraction. Total genomic DNA of each *P. falciparum* isolate was extracted from filter paper blots (FTA; Whatman Inc., Bound Brook, NJ) by using a QIAamp DNA blood minikit (Qiagen, Valencia, CA), according to the manufacturer's instructions.

Genotypic characterization of *Pfmdr1* and *Pfcr1* genes. The 2^{- $\Delta\Delta$ CT} method (based on the threshold cycle [C_T]) of relative quantification was used to estimate copy number variation in *Pfmdr1*, as previously published (37). Polymorphisms in *Pfmdr1* codons 86 and 184 and *Pfcr1* codon 76 were determined by nested PCR and sequencing as previously described (31, 38).

***Pfnhe1* microsatellite genotyping.** Microsatellite repeats in *Pfnhe1* were characterized by nested PCR using 1.5 μ l of each primer. Primary

run primers were NHE-F (5'-AGTCGAAGGCGAATCAGATG-3') and NHE-R (5'-GATACTACGAACATGTTTCATG-3'). Secondary run primers NHE-F (5'-ATCCCTGTTGATATATCGAATG-3') and NHE-R (5'-TTGTCATTAGTACCCTTAGTTG-3'), as previously described (24). The 25- μ l reaction volume setup consisted of 10 \times PCR buffer to a final concentration of 1 \times , 25 mM MgCl₂ to a final concentration of 1.5 mM, 20 mM deoxynucleoside triphosphates mix to a final concentration of 2 mM, 10 μ M each of the primers to a final concentration of 100 nM, 5 U/ μ l AmpliTaq DNA polymerase to a final concentration of 1 U/reaction tube. Amplification of the ms4760 region was done under the following amplification conditions for the first run: initial hybridization at 94°C for 5 min, a subsequent 35 cycles of denaturation at 94°C for 30 s, annealing at 57.0°C for 30 s, and extension at 72°C for 30 s, and a final extension at 72°C for 7 min. Secondary run conditions were the following: initial hybridization at 94°C for 5 min, a subsequent 35 cycles of denaturation at 94°C for 30 s, annealing at 59.0°C for 30 s, and extension at 72°C for 30 s, and a final extension at 72°C for 7 min. The PCR products were resolved by electrophoresis on a 2% agarose gel, stained with 0.5 μ g/ml ethidium bromide, and visualized under UV light. Amplicons were purified and sequenced using ABI Prism BigDye Terminator v3.1 cycle sequencing ready reaction kits (Applied Biosystems, Foster City, CA), as directed by the manufacturer on a 3500XL genetic analyzer. Sequences were analyzed with the BioEdit sequence alignment editor (version 7.0.9.0). Sequences were analyzed for the number of DDNNND, DDNHNDNHHND, and DDNNND NHHND repeats in the *Pfnhe1* ms4760 microsatellite.

Statistical analysis. Continuous data in the form of IC₅₀ estimates were expressed as medians with interquartile ranges. Further, IC₅₀ data for isolates with various genetic polymorphism in *Pfmdr1*, *Pfcr1*, and *Pfnhe1* were compared by using nonparametric tests (one-way analysis of variance, Kruskal-Wallis test with Dunn's multiple comparison posttest, and the Mann-Whitney test). Associations between IC₅₀ estimates and genetic markers were considered significant when *P* was <0.05. Analyses were conducted using the Prism program (version 5.0.2; GraphPad Software, Inc., San Diego, CA). For analyses, all isolates with mixed genotypes were considered mutants.

Nucleotide sequence accession numbers. New *Pfnhe1* ms4760 microsatellite profiles were deposited in GenBank under the following accession numbers: KF719182, KF719183, KF719184, KF719185, and KF719186.

RESULTS

QN chemosensitivity. The QN IC₅₀ values for the two reference strains, 3D7 and W2, were established and used as internal controls in subsequent experiments. The median IC₅₀ values (in nM; interquartile ranges and *n* values in parentheses) were 27 (17–32; *n* = 4) for 3D7 and 362 (111–400; *n* = 4) for W2. A total of 88 culture-adapted field isolates were successfully analyzed, and QN IC₅₀s were determined. The median IC₅₀ (interquartile range) for the 88 isolates was 69.01 nM (19.05–336.0).

***Pfmdr1* and *Pfcr1* mutations.** Genotype analyses revealed 67% and 52% of the isolates carried the wild-type genotype at codons N86 and Y184 of the *Pfmdr1* gene, respectively, whereas 35% of the isolates carried the wild-type genotype at codon K76 of the *Pfcr1* gene. Notably, all the isolates had a single copy of the *Pfmdr1* gene.

Associations of QN *in vitro* sensitivity with *Pfmdr1* and *Pfcr1* polymorphisms. Figure 1 shows profiles for *in vitro* QN sensitivity per genetic polymorphism in the *Pfmdr1* and *Pfcr1* genes. The isolates revealed a significant association between the QN IC₅₀ and mutations in *Pfmdr1* codons 86 and 184 and *Pfcr1* codon 76 (*P* < 0.0001 for each association). The median IC₅₀ for parasites for *Pfmdr1* codons 86 and 184 were ~15-fold higher in the mutant genotypes compared to the wild-type, whereas with *Pfcr1* codon




FIG 1 *In vitro* QN sensitivity for polymorphisms at each genetic marker in *Pfmdr1*-86, *Pfmdr1*-184, and *Pfcrt*-76 codons. The analysis comparing alleles at each codon was done using the Mann-Whitney test. The horizontal (blue) bars indicate medians. The numbers of isolates analyzed are shown in brackets. There was a significant association between the QN IC₅₀ and mutation at each codon.

76, the median IC₅₀ was ~9-fold higher for the mutant genotype than for the wild type.

***Pfnhe1* ms4760 polymorphism.** The genetic polymorphisms of ms4760 in *Pfnhe1* gene was analyzed. Information on the QN IC₅₀ values and the genetic profiles is summarized in Table S1 of the supplemental material. These isolates contained 15 different genetic polymorphisms of ms4760 in the *Pfnhe1* gene. Five of the genetic profiles have not been previously described. The new profiles were deposited with GenBank and were assigned accession numbers [KF719182](#), [KF719183](#), [KF719184](#), [KF719185](#), and [KF719186](#). Previously described profiles that were present in our isolates included the following: ms4760-1, ms4760-2, ms4760-3, ms4760-4, ms4760-6, ms4760-8, ms4760-9, ms4760-16, and ms4760-53. The most common genetic polymorphisms were the following: ms4760-1, ms4760-3, and ms4760-9, in 22, 16, and 12 of the isolates analyzed, respectively. Combined, they represented 60% of the total isolates analyzed. The least common genetic polymorphisms were ms4760-16, [KF719184](#), and [KF719186](#), each of which was present in only one isolate (Fig. 2). Figure 3 shows alignments of the 15 sequences of *Pfnhe1* ms4760 identified in 88 western Kenya *P. falciparum* isolates.

Association between QN sensitivity and the number of DNNND repeats in *Pfnhe1*. There was a significant association between the QN IC₅₀ and the number of DNNND repeats (Fig. 4). The QN IC₅₀ for parasites with 3 DNNND repeats was significantly higher than for those with 1 or 2 repeats ($P = 0.033$ and $P = 0.0043$, respectively). There was no statistical difference in QN IC₅₀s between parasites with 1 or 2 repeats. Interestingly, there was no association between QN IC₅₀ and DDN HNDNHND repeats. The median IC₅₀ (interquartile range; n) for parasites with 1 DDN HNDNHND repeat was 80.72 nM (18.71–589.5; $n = 28$), for parasites with 2 repeats it was 57.79 nM (18.19–253.4; $n = 59$), and for those with 3 repeats it was 948.7 nM (118.4–1,779; $n = 2$).

Associations of QN sensitivity and genotype combination for *Pfmdr1*, *Pfcrt*, and *Pfnhe1*. Our data showed a significant as-




FIG 2 Prevalence rates of *Pfnhe1* ms4760 profiles among the 88 western Kenya isolates. Profiles ms-1 to ms-53 have been described previously, while the sequences for [KF719182](#) to [KF719186](#) were first described in this study.

sociation of genetic polymorphisms in the *Pfmdr1* gene at codons 86 and 184, the *Pfcrt* gene at codon 76, and the *Pfnhe1* gene with DNNND repeats. To further understand this observation, we analyzed the relationship between the QN IC₅₀ and the genotype combination of DNNND repeat number for the *Pfmdr1* and *Pfcrt* genes. In *Pfmdr1* codons 86Y (Fig. 5A) and 184F (Fig. 5B), data revealed parasites with 3 DNNND repeats had higher median IC₅₀s than those with 1 or 2 repeats. For codon 86Y, there was a statistically significant difference between parasites with 3 DNNND repeats versus 2 repeats ($P = 0.0302$). For codon 184F, there was a statistically significant difference between parasites with 3 DNNND repeats versus 1 or 2 repeats ($P = 0.0148$ and $P = 0.0031$, respectively). For *Pfcrt* codon 76T (Fig. 5C), data revealed parasites with 3 DNNND repeats had median IC₅₀s more than 5-fold higher than in parasites with 1 or 2 DNNND repeats, and there was a statistically significant difference between those with 3 DNNND repeats versus 1 or 2 repeats ($P = 0.0019$ and $P = 0.0041$, respectively).

Genetic profiles of parasites with a QN IC₅₀ outside the interquartile range. To further investigate the role of genetic polymorphisms in the *Pfmdr1*, *Pfcrt*, and *Pfnhe1* genes in determining parasite phenotypic characteristics in response to QN sensitivity, the genetic profiles of parasites with QN IC₅₀s below the lower interquartile range or above the upper interquartile range were analyzed. There were 23 parasites with a QN IC₅₀ of ≤ 19.05 and 22 parasites with a QN IC₅₀ of ≥ 336 . Table S2 in the supplemental material shows the genetic profiles of parasites with QN IC₅₀s below and above the interquartile ranges. For the parasites with values below the lower interquartile range, the predominant genotype present in 74% of the parasites was *Pfmdr1*-N86 *Pfmdr1*-Y184 *Pfcrt*-K76 (all codons carrying the wild-type genotype), whereas above the upper interquartile range, the predominant genotype present in 82% of the parasites was *Pfmdr1*-86Y *Pfmdr1*-184F *Pfcrt*-76T (all codons carrying mutant genotypes). Two genotype profiles, *Pfmdr1*-N86 *Pfmdr1*-Y184 *Pfcrt*-76T and *Pfmdr1*-N86 *Pfmdr1*-184F *Pfcrt*-76T were present in parasites both in the lower and above the interquartile ranges. In the parasites with values below the lower interquartile range, 52% of the parasites had 2 DNNND repeats, whereas for parasites above the upper interquartile range, 55% had 3 DNNND repeats.

Profile No	I	II	III	IV	V	VI	
MS_4760-1	KKKKKSGSNN-DNN	DNNND	DNNND	NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-2	KKKKKSGSNN-DNN	DNNND		DKNNKND-DDNNNDNH-KNDKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-3	KKKKKSGSNN-DNN	DNNND		DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-4	KKKKKSGSNN-DNN	DNNND	DNNND	NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-6	KKKKKSGSNN-DNN	DNNND		DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-8	KKKKKSGSNN-DNN	DNNND	DNNND	NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-9	KKKKKSGSNN-DNN	DNNND	DNNND	NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-16	KKKKKSGSNN-DNN	DNNND		NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND-DNHND	DNNSSH
MS_4760-20	KKKKKSGSNN-DNN	DNNND		DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
MS_4760-53	KKKKKSGSNN-DNN	DNNND		DKNNKND-DDNNNDNHKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
KF719182	KKKKKSGSNN-DNN	DNNND		NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
KF719183	KKKKKSGSNN-DNN	DNNND	DNNND	NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
KF719184	KKKKKSGSNN-DNN	DNNND	DNNND	NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
KF719185	KKKKKSGSNN-DNN	DNNND		NHND-DKNNKND	DDNNNDNHND	DDNNNDNHND	DNNSSH
KF719186	KKKKKSGSNN-DNN	DNNND		DKNNKND-DDNNNDNHND	DDNNNDNHND	DDNNNDNHND	DNNSSH

FIG 3 Alignment of 15 sequences of *Pfnhe1* ms4760 identified in 88 western Kenya *P. falciparum* isolates. Blocks I to VI have been described previously (29). The DNNND repeats are in block II, the DDNNNDNHND repeats are in block V, and the DDNNNDNHND repeats are in block VI. Profiles ms4760-1 to ms4760-53 have been described previously (25, 27, 32); sequences KF719182 to KF719186 are the described for the first time in this study. Other differences among sequences are highlighted.

DISCUSSION

In this study, we have clearly shown a significant association between polymorphisms in *Pfmdr1* codons 86 and 184 and *Pfcrt* codon 76 and QN susceptibility in *P. falciparum* parasites from western Kenya. Furthermore, the diversity of microsatellite repeats in *Pfnhe1* ms4760 within these isolates was underscored and also shown to be linked to QN susceptibility. Three repeats of the DNNND polymorphism in the *Pfnhe1* gene significantly reduced parasite susceptibility to QN. Most importantly, we have de-

scribed a predominant parasite genotype, *Pfmdr1*-86Y *Pfmdr1*-184F *Pfcrt*-76T, in parasites with high QN IC₅₀s.

The median IC₅₀ (interquartile range) for isolates from our study was 69.01 nM (19.05–336.0). The chemosensitivity threshold for QN has not been clearly defined. However, the historical WHO IC₅₀ against the W2 clone of *P. falciparum*, which is considered resistant, is 315 nM (39). Other different QN threshold cut-offs have been proposed: 300, 500, and 800 nM (40–42). In studies conducted in Senegal (43) and Kenya (24), only 1% and 7% (respectively) of the isolates tested against QN had IC₅₀s of >500 nM. In the study conducted in Kilifi, Kenya, none of the isolates had an IC₅₀ of >800 nM (24). In field isolates from the Republic of the Congo, 25.7% exceeded the 500 nM cutoff, whereas only 5.4% exceeded the 800 nM cutoff (29). In our study, 18.2% (16/88) of the isolates exceeded the 500 nM cutoff and 11.4% (10/88) exceeded the 800 nM cutoff. Although 70.4% (62/88) of the isolates had IC₅₀ estimates below 300 nM, and were therefore considered sensitive to QN, a relatively high number of samples, compared to those collected elsewhere, can be considered resistant to QN, with a threshold cutoff exceeding 800 nM. The *in vitro* data from our study were supported by the genotypic data. This was expected, since in western Kenya, the prevalence of the *Pfmdr1*-86Y *Pfmdr1*-184F *Pfcrt*-76T genotype, which is also responsible for chloroquine resistance, has remained persistently high (31). Also, *in vitro* data for field isolates from Uganda have shown a tight correlation between sensitivities to chloroquine and QN (44).

In a study that analyzed parasite isolates from Kilifi, Kenya, parasites carrying the *Pfmdr1*-86 mutation showed a trend toward decreased QN activity, but there was no significant association (24). Similarly, in a study that used field isolates from Uganda, polymorphisms in *Pfmdr1*-86 and *Pfmdr1*-184 were implicated in decreased sensitivity to QN but did not reach significance ($P = 0.22$ and $P = 0.34$, respectively) (44). However, the *Pfmdr1*-1246




FIG 4 Association between the QN IC₅₀ and the number of DNNND repeats. The analysis comparing polymorphisms in DNNND repeats was done using the Mann-Whitney test. The horizontal (blue) bars indicate the medians. The numbers of isolates analyzed are shown in brackets. *P* values comparing the repeats are indicated; 3 DNNND repeats had statistically higher QN IC₅₀ values, compared to isolates with 1 or 2 repeats.




FIG 5 Relationship between the QN IC₅₀ and the genotype combination of DNNND repeats for *Pfm $dr1$ -86* (A), *Pfm $dr1$ -184* (B), and *Pfc rt -76* (C). The analysis comparing the number of DNNND repeats with each allele was done using the Mann-Whitney test. The horizontal bars (blue) indicate the medians. The numbers of isolates analyzed are shown in brackets. With one exception, 3 DNNND repeats had statistically significantly higher QN IC₅₀ values than did isolates with 1 or 2 repeats for all mutant alleles analyzed. The association of *Pfm $dr1$ -86Y* with 1 versus 3 DNNND repeats was not statistically significant ($P = 0.21$).

mutation was shown to be statistically significant for modulating QN activity ($P = 0.029$). Transfection studies have also shown that *Pfm $dr1$ -1034*, *Pfm $dr1$ -1042*, and *Pfm $dr1$ -1246* mutations modulate resistance to QN (17). Here, we have shown statistically significant associations of mutations in *Pfm $dr1$* codons 86 and 184 with QN activity. The difference in these observations could be due to the genetic backgrounds of the parasites. QN resistance appears to be dependent on multiple genes, which indicates polymorphisms in *Pfm $dr1$* contribute to the overall phenotype of the parasite but in the backdrop of the genetic background of the parasite (17). It will be important to go back and analyze *Pfm $dr1$* codon 1246 in isolates from our study, as this mutation has been shown to be important in modulating QN activity both in transfection and among *in vitro* data of field isolates (16, 44).

Transfection studies have shown that K76T contributes to QN resistance, but the extent of its contribution differs between strains (16, 45). Similarly, our data clearly showed a significant association of K76T mutation with QN reduced sensitivity. This is contrary to what other studies have shown (24). However, similarly to the *Pfm $dr1$* gene, the genetic background seems to be important and must therefore ultimately determine parasite phenotype.

Studies of the association of polymorphisms in the *Pfn $he1$* ms4670 microsatellite with *in vitro* susceptibility to QN have had conflicting results (24, 27, 29, 46). In studies that showed a positive correlation, the number of repeats in block 2 (DNNND), block 5 (DDNHNDNHNNDD), and block 6 (DDNNNDNHND) have been associated with modulation of QN resistance (27, 46). In this study, we showed there was a significant association for parasites with 3 repeats of DNNND and QN susceptibility. Polymorphisms in DDNHNDNHNNDD or DDNNNDNHND repeats did not have an effect on QN susceptibility. Some studies have shown that an increase in DNNND repeats is associated with reduced susceptibility to QN (16, 46). Interestingly, the study that analyzed isolates from Kilifi, Kenya, showed 2 DNNND repeats, and not 3 DNNND repeats, were associated with reduced QN susceptibility (24). Further, that study showed that 3 DNNND repeats restored QN activity, contrary to our findings. Other studies did not find any association for polymorphisms in DNNND repeats with QN activity when evaluating isolates from African patient populations (27, 29). Interestingly, however, in one of the studies, there was a positive association for NHNDNHNNDD repeats with increased QN IC₅₀ ($P = 0.01$) among isolates from an African population (27). This is in line with findings from

studies conducted using isolates from Vietnam and the China-Myanmar border, where an increased number of DDNDNDND repeats was associated with high *in vitro* QN susceptibility (30, 43). The seemingly conflicting association of polymorphisms in the *Pfnhe1* ms4670 microsatellite with *in vitro* susceptibility to QN can also be explained by differences in parasite genetic backgrounds, which can be associated with geographic origins of the isolates.

Lastly, we established the contributions of *Pfmdr1*-86, *Pfmdr1*-184, and *Pfcrt*-76 in the context of DNNND polymorphisms to QN susceptibility. For all three genotypes (*Pfmdr1*-86Y, *Pfmdr1*-184F, and *Pfcrt*-76T), the cumulative effect was highest in the background of 3 DNNND repeats, reaching statistical significance in each one of the genotypes. When we analyzed the genotypes of samples with results above the upper interquartile range, comprising isolates with IC₅₀ estimates above the WHO cutoff for resistance, the majority of these isolates (82%) carried mutant alleles at the three genetic markers. When analyzed in combination with DNNND repeats, 55% of these isolates carried 3 repeats, as opposed to 32% with 2 repeats, and the remainder with 1 repeat. On the contrary, the majority of the isolates (74%) with values below the lower interquartile range had wild-type genotypes for all three codons analyzed, with most (52%) carrying 2 DNNND repeats.

Conclusions. In this study, we showed the associations of *Pfmdr1*-86, *Pfmdr1*-184, *Pfcrt*-76, and *Pfnhe1* ms4670 polymorphisms with parasite responses to QN. Further, we showed that these genetic markers are only relevant in the context of the genetic background of the isolates. Andriantsoanirina et al. (27) strongly suggested polymorphisms in *Pfnhe1* might not be valid molecular markers for *in vitro* susceptibility to QN in *P. falciparum* isolates from Africa. Those authors tested only 83 isolates from a few countries in Africa. Africa is a continent with extremely diverse geographical landscapes, different malaria ecologies, varied rates of malaria endemicity, and parasites with different genetic structures. The cumulative evidence is clearly showing that the genetic background of the parasite is critical in determining QN activity. This will have important implications, because parasites from each geographic region must therefore be analyzed to determine which markers confer reduced QN susceptibility. To further validate our observations and conclusions, it will be important that isolates from different malaria ecological zones and regions of malaria endemicity in Kenya are analyzed. Most importantly, more clinical efficacy studies must be conducted in Kenya and other African countries, given the importance of QN in the management of malaria.

ACKNOWLEDGMENTS

We thank the Director of KEMRI, USAMRU-K, and the entire Malaria Drug Research team for their support during the study period. We also thank John Okombo of KEMRI—Wellcome Trust Kilifi for sharing the *Pfnhe1* primers and guidance during data analyses.

This work was supported by the U.S. Department of Defense, Global Emerging Infections Surveillance and Response System (GEIS), a division at the Armed Forces Health Surveillance Center (AFHSC).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The opinions and assertions contained herein are the private opinions of the authors and are not to be construed as reflecting the views of the U.S. Army Medical Research Unit—Kenya or the U.S. Department of Defense.

We declare that there are no competing interests.

REFERENCES

1. World Health Organization. 2009. Global antimalarial drug policies database—WHO African region, September ed. World Health Organization, Geneva, Switzerland.
2. World Health Organization. 2010. Global report on antimalarial efficacy and drug resistance: 2000–2010. World Health Organization, Geneva, Switzerland.
3. Achan J, Talisuna AO, Erhart A, Yeka A, Tibenderana JK, Baliraine FN, Rosenthal PJ, D'Alessandro U. 2011. Quinine, an old anti-malarial drug in a modern world: role in the treatment of malaria. *Malar. J.* 10:144. <http://dx.doi.org/10.1186/1475-2875-10-144>.
4. Pukrittayakamee S, Supanaranond W, Looareesuwan S, Vanijanonta S, White NJ. 1994. Quinine in severe *falciparum* malaria: evidence of declining efficacy in Thailand. *Trans. R. Soc. Trop. Med. Hyg.* 88:324–327. [http://dx.doi.org/10.1016/0035-9203\(94\)90102-3](http://dx.doi.org/10.1016/0035-9203(94)90102-3).
5. Chongsuphajaisiddhi T, Sabchareon A, Attanath P. 1983. Treatment of quinine resistant *falciparum* malaria in Thai children. *Southeast Asian J. Trop. Med. Public Health* 14:357–362.
6. Kofoed PE, Mapaba E, Lopes F, Pussick F, Aaby P, Rombo L. 1997. Comparison of 3, 5 and 7 days' treatment with Quinimax for *falciparum* malaria in Guinea-Bissau. *Trans. R. Soc. Trop. Med. Hyg.* 91:462–464. [http://dx.doi.org/10.1016/S0035-9203\(97\)90286-8](http://dx.doi.org/10.1016/S0035-9203(97)90286-8).
7. Rogier C, Brau R, Tall A, Cisse B, Trape JF. 1996. Reducing the oral quinine-quinidine-cinchonin (Quinimax) treatment of uncomplicated malaria to three days does not increase the recurrence of attacks among children living in a highly endemic area of Senega. *Trans. R. Soc. Trop. Med. Hyg.* 90:175–178. [http://dx.doi.org/10.1016/S0035-9203\(96\)90128-5](http://dx.doi.org/10.1016/S0035-9203(96)90128-5).
8. Kremsner PG, Winkler S, Brandts C, Neifer S, Bienzle U, Graninger W. 1994. Clindamycin in combination with chloroquine or quinine is an effective therapy for uncomplicated *Plasmodium falciparum* malaria in children from Gabon. *J. Infect. Dis.* 169:467–470. <http://dx.doi.org/10.1093/infdis/169.2.467>.
9. Esamai F, Tenge CN, Ayuo PO, Ong'or WO, Obala A, Jakait B. 2005. A randomized open label clinical trial to compare the efficacy and safety of intravenous quinine followed by oral malarone vs. intravenous quinine followed by oral quinine in the treatment of severe malaria. *J. Trop. Pediatr.* 51:17–24. <http://dx.doi.org/10.1093/tropej/fmh069>.
10. Mayxay M, Barends M, Brockman A, Jaidee A, Nair S, Sudimack D, Pongvongsa T, Phompida S, Phetsouvanh R, Anderson T, White NJ, Newton PN. 2007. In vitro antimalarial drug susceptibility and *Pfcrt* mutation among fresh *Plasmodium falciparum* isolates from the Lao PDR (Laos) Am. J. Trop. Med. Hyg. 76:245–250.
11. Legrand E, Volney B, Meynard JB, Mercereau-Puijalon O, Esterre P. 2008. In vitro monitoring of *Plasmodium falciparum* drug resistance in French Guiana: a synopsis of continuous assessment from 1994 to 2005. *Antimicrob. Agents Chemother.* 52:288–298. <http://dx.doi.org/10.1128/AAC.00263-07>.
12. Toure AO, Kone LP, Jambou R, Konan TD, Demba S, Beugre GE, Kone M. 2008. In vitro susceptibility of *P. falciparum* isolates from Abidjan (Cote d'Ivoire) to quinine, artesunate and chloroquine. *Sante* 18:43–47. <http://dx.doi.org/10.1684/san.2008.0103>.
13. Pradines B, Mabika Mamfoumbi M, Parzy D, Owono Medang M, Lebeau C, Mourou Mbina JR, Doury JC, Kombila M. 1998. In vitro susceptibility of Gabonese wild isolates of *Plasmodium falciparum* to artemether, and comparison with chloroquine, quinine, halofantrine and amodiaquine. *Parasitology* 117:541–545. <http://dx.doi.org/10.1017/S0031182098003400>.
14. Shann F, Stace J, Edstein M. 1985. Pharmacokinetics of quinine in children. *J. Pediatr.* 106:506–510.
15. Hawley SR, Bray PG, Mungthin M, Atkinson JD, O'Neill PM, Ward SA. 1998. Relationship between antimalarial drug activity, accumulation, and inhibition of heme polymerization in *Plasmodium falciparum* *in vitro*. *Antimicrob. Agents Chemother.* 42:682–686.
16. Ferdig MT, Cooper RA, Mu J, Deng B, Joy DA, Su XZ, Wellem TE. 2004. Dissecting the loci of low-level quinine resistance in malaria parasites. *Mol. Microbiol.* 52:985–997. <http://dx.doi.org/10.1111/j.1365-2958.2004.04035.x>.
17. Reed MB, Saliba KJ, Caruana SR, Kirk K, Cowman AF. 2000. Pgh1 modulates sensitivity and resistance to multiple antimalarials in *Plasmodium falciparum*. *Nature* 403:906–909. <http://dx.doi.org/10.1038/35002615>.
18. Sidhu AB, Valderramos SG, Fidock DA. 2005. *Pfmdr1* mutations contribute to quinine resistance and enhance mefloquine and artemisinin

- sensitivity in *Plasmodium falciparum*. *Mol. Microbiol.* 57:913–926. <http://dx.doi.org/10.1111/j.1365-2958.2005.04729.x>.
19. Sidhu AB, Uhlemann AC, Valderramos SG, Valderramos JC, Krishna S, Fidock DA. 2006. Decreasing *Pfmdr1* copy number in *Plasmodium falciparum* malaria heightens susceptibility to mefloquine, lumefantrine, halofantrine, quinine, and artemisinin. *J. Infect. Dis.* 194:528–535. <http://dx.doi.org/10.1086/507115>.
 20. Valderramos SG, Valderramos JC, Musset L, Purcell LA, Mercereau-Puijalon O, Legrand E, Fidock DA. 2010. Identification of a mutant *Pfprt*-mediated chloroquine tolerance phenotype in *Plasmodium falciparum*. *PLoS Pathog.* 6:e1000887. <http://dx.doi.org/10.1371/journal.ppat.1000887>.
 21. Parola P, Pradines B, Simon F, Carlotti MP, Minodier P, Ranjeva MP, Badiaga S, Bertaux L, Delmont J, Morillon M, Silai R, Brouqui P, Parzy D. 2007. Antimalarial drug susceptibility and point mutations associated with drug resistance in 248 *Plasmodium falciparum* isolates imported from Comoros to Marseille, France in 2004–2006. *Am. J. Trop. Med. Hyg.* 77:431–437.
 22. Cooper RA, Ferdig MT, Su XZ, Ursos L, Mu J, Nomura T, Fujioka H, Fidock DA, Roepe PD, Wellems TE. 2002. Alternative mutation at position 76 of the vacuolar transmembrane protein *Pfprt* are associated with chloroquine resistance and unique stereospecific quinine and quinidine response in *Plasmodium falciparum*. *Mol. Pharmacol.* 61:35–42. <http://dx.doi.org/10.1124/mol.61.1.35>.
 23. Cooper RA, Lane KD, Deng B, Mu J, Patel JJ, Wellems TE, Su X, Ferdig MT. 2007. Mutations in transmembrane domains 1, 4 and 9 of the *Plasmodium falciparum* chloroquine resistance transporter alter susceptibility to chloroquine, quinine and quinidine. *Mol. Microbiol.* 63:270–282. <http://dx.doi.org/10.1111/j.1365-2958.2006.05511.x>.
 24. Okombo J, Kiara SM, Rono J, Mwai L, Pole L, Ohuma E, Borrmann S, Ochola LI, Nzila A. 2010. *In vitro* activities of quinine and other antimalarials and *pfnhe* polymorphisms in *Plasmodium* isolates from Kenya. *Antimicrob. Agents Chemother.* 54:3302–3307. <http://dx.doi.org/10.1128/AAC.00325-10>.
 25. Baliraine FN, Nsohya SL, Achan J, Tibenderana JK, Talisuna AO, Greenhouse B, Rosenthal PJ. 2011. Limited ability of *Plasmodium falciparum* *Pfprt*, *Pfmdr1*, and *Pfnhe1* polymorphisms to predict quinine *in vitro* sensitivity or clinical effectiveness in Uganda. *Antimicrob. Agents Chemother.* 55:615–622. <http://dx.doi.org/10.1128/AAC.00954-10>.
 26. Pelleau S, Bertaux L, Briolant S, Ferdig MT, Sinou V, Pradines B, Parzy D, Jambou R. 2011. Differential association of *Plasmodium falciparum* Na⁺/H⁺ exchanger polymorphism and quinine responses in field- and culture-adapted isolates of *Plasmodium falciparum*. *Antimicrob. Agents Chemother.* 55:5834–5841. <http://dx.doi.org/10.1128/AAC.00477-11>.
 27. Andriantsoanirina V, Menard D, Rabearimanana S, Hubert V, Bouchier C, Tichit M, Bras JL, Durand R. 2010. Association of microsatellite variations of *Plasmodium falciparum* Na⁺/H⁺ exchanger (*Pfnhe-1*) gene with reduced *in vitro* susceptibility to quinine: lack of confirmation in clinical isolates from Africa. *Am. J. Trop. Med. Hyg.* 82:782–787. <http://dx.doi.org/10.4269/ajtmh.2010.09-0327>.
 28. Nkrumah LJ, Riegelhaupt PM, Moura P, Johnson DJ, Patel J, Hayton K, Ferdig MT, Wellems TE, Akabas MH, Fidock DA. 2009. Probing the multifactorial basis of *Plasmodium falciparum* quinine resistance: evidence for a strain-specific contribution of the sodium-proton exchanger *PfNHE*. *Mol. Biochem. Parasitol.* 165:122–131. <http://dx.doi.org/10.1016/j.molbiopara.2009.01.011>.
 29. Briolant S, Pelleau S, Bogreau H, Hovette P, Zettor A, Castello J, Baret E, Amalvict R, Rogier C, Pradines B. 2011. *In vitro* susceptibility to quinine and microsatellite variations of the *Plasmodium falciparum* Na⁺/H⁺ exchanger (*Pfnhe-1*) gene: the absence of association in clinical isolates from the Republic of Congo. *Malar. J.* 10:37. <http://dx.doi.org/10.1186/1475-2875-10-37>.
 30. Sinou V, Quang le, Pelleau HS, Huong VN, Huong NT, Tai le M, Bertaux L, Desbordes M, Latour C, Long LQ, Thanh NX, Parzy D. 2011. Polymorphism of *Plasmodium falciparum* Na⁺/H⁺ exchanger is indicative of a low *in vitro* quinine susceptibility in isolates from Viet Nam. *Malar. J.* 10:164. <http://dx.doi.org/10.1186/1475-2875-10-164>.
 31. Eyase FL, Akala HM, Ingasia L, Cheruiyot A, Omondi A, Okudo C, Juma D, Yeda R, Andagalu B, Wanja E, Kamau E, Schnabel D, Bulimo W, Waters NC, Walsh DS, Johnson JD. 2013. The role of *Pfmdr1* and *Pfprt* in changing chloroquine, amodiaquine, mefloquine and lumefantrine susceptibility in western-Kenya *P. falciparum* samples during 2008–2011. *PLoS One* 8:e64299. <http://dx.doi.org/10.1371/journal.pone.0064299>.
 32. Trager W, Jensen JB. 1976. Human malaria parasites in continuous culture. *Science* 193:673–675. <http://dx.doi.org/10.1126/science.781840>.
 33. Rason MA, Randrianantoa T, Andrianantenaina H, Ratsimbaso A, Menard D. 2008. Performance and reliability of the SYBR Green I based assay for the routine monitoring of susceptibility of *Plasmodium falciparum* clinical isolates. *Trans. R. Soc. Trop. Med. Hyg.* 102:346–351. <http://dx.doi.org/10.1016/j.trstmh.2008.01.021>.
 34. Bacon DJ, Latour C, Lucas C, Colina O, Ringwald P, Picot S. 2007. Comparison of a SYBR green I-based assay with a histidine-rich protein II enzyme-linked immunosorbent assay for *in vitro* antimalarial drug efficacy testing and application to clinical isolates. *Antimicrob. Agents Chemother.* 51:1172–1178. <http://dx.doi.org/10.1128/AAC.01313-06>.
 35. Johnson JD, Denuall RA, Gerena L, Lopez-Sanchez M, Roncal NE, Waters NC. 2007. Assessment and continued validation of the malaria SYBR green I-based fluorescence assay for use in malaria drug screening. *Antimicrob. Agents Chemother.* 51:1926–1933. <http://dx.doi.org/10.1128/AAC.01607-06>.
 36. Tyner SD, Lon C, Se Y, Bethell D, Socheat D, Noeld H, Sea D, Satimai W, Schaecker K, Rutvisuttinun W, Fukuda MM, Chaorattanakawee S, Yingyuen K, Sundrakes S, Chaichana P, Saingam P, Buathong N, Sriwichai S, Chann S, Timmermans A, Saunders DL, Walsh DS. 2012. *Ex vivo* drug sensitivity profiles of *Plasmodium falciparum* field isolates from Cambodia and Thailand, 2005 to 2010, determined by a histidine-rich protein-2 assay. *Malar. J.* 11:198. <http://dx.doi.org/10.1186/1475-2875-11-198>.
 37. Price RN, Uhlemann AC, Brockman A, McGready R, Ashley E, Phaipun L, Patel R, Laing K, Looareesuwan S, White NJ, Nosten F, Krishna S. 2004. Mefloquine resistance in *Plasmodium falciparum* and increased *Pfmdr1* gene copy number. *Lancet* 364:438–447. [http://dx.doi.org/10.1016/S0140-6736\(04\)16767-6](http://dx.doi.org/10.1016/S0140-6736(04)16767-6).
 38. Vinayak S, Alam MT, Sem R, Shah NK, Susanti AI, Lim P, Muth S, Maguire JD, Rogers WO, Fandeur T, Barnwell JW, Escalante AA, Wongsrichanalai C, Arley F, Meshnick SR, Udhayakumar V. 2010. Multiple genetic backgrounds of the amplified *Plasmodium falciparum* multidrug resistance (*Pfmdr1*) gene and selective sweep of 184F mutation in Cambodia. *J. Infect. Dis.* 201:1551–1560.
 39. World Health Organization. 2001. Mark III *in vitro* micro-test for the assessment of the response of *Plasmodium falciparum* to chloroquine, mefloquine, quinine, amodiaquine, sulfadoxine/pyrimethamine, and artemisinin. CTD/MAL/9720, revision 2. World Health Organization, Geneva, Switzerland.
 40. Ralaimazava P, Durand R, Godineau N, Jezic Z, Pradines B, Bouchaud O, Bras JL. 2002. Profile and evolution of the chemosusceptibility of *falciparum* malaria imported into France in 2000. *Euro Surveill.* 7(7):pii=355. <http://europa.eu.int/ViewArticle.aspx?ArticleId=355>.
 41. Brasseur P, Kouamou J, Moyou-Somo R, Druilhe P. 1992. Multi-drug resistant *falciparum* malaria in Cameroon in 1987–1988. I. Stable figures of prevalence of chloroquine- and quinine-resistant isolates in the original foci. *Am. J. Trop. Med. Hyg.* 46:1–7.
 42. Pradines B, Rogier C, Fusai T, Tall A, Trape JF, Doury JC. 1996. *In vitro* sensitivity of 85 *Plasmodium falciparum* isolates in the Fatik region, Senegal. *Med. Trop. (Mars.)* 56:141–145. (In French.)
 43. Meng H, Zhang R, Yang H, Fan Q, Su X, Miao J, Cui L, Yang Z. 2010. *In vitro* sensitivity of *Plasmodium falciparum* clinical isolates from the China-Myanmar border area to quinine and association with polymorphism in the Na⁺/H⁺ exchanger. *Antimicrob. Agents Chemother.* 54:4306–4313. <http://dx.doi.org/10.1128/AAC.00321-10>.
 44. Nsohya SL, Kiggundu M, Nanyunja S, Joloba M, Greenhouse B, Rosenthal PJ. 2010. *In vitro* sensitivities of *Plasmodium falciparum* to different antimalarial drugs in Uganda. *Antimicrob. Agents Chemother.* 54:1200–1206. <http://dx.doi.org/10.1128/AAC.01412-09>.
 45. Sidhu AB, Verdier-Pinard D, Fidock DA. 2002. Chloroquine resistance in *Plasmodium falciparum* malaria parasites conferred by *Pfprt* mutations. *Science* 298:210–213. <http://dx.doi.org/10.1126/science.1074045>.
 46. Henry M, Briolant S, Zettor A, Pelleau S, Baragatti M, Baret E, Mosnier J, Amalvict R, Fusai T, Rogier C, Pradines B. 2009. *Plasmodium falciparum* Na⁺/H⁺ exchanger 1 transporter is involved in reduced susceptibility to quinine. *Antimicrob. Agents Chemother.* 53:1926–1930. <http://dx.doi.org/10.1128/AAC.01243-08>.