

Positive Solutions of a Nonlinear m -Point Integral Boundary Value Problem

Jessada Tariboon¹ and Weerawat Sudsutad²

Department of Mathematics, Faculty of Applied Science
King Mongkut's University of Technology North Bangkok
Bangkok 10800, Thailand

Centre of Excellence in Mathematics, CHE, Sri Ayutthaya Road,
Bangkok 10400, Thailand

¹jessadat@kmutnb.ac.th, ²wrw.sst@gmail.com

Abstract

We study sufficient conditions for the existence of positive solutions to the m -point integral boundary value problem

$$u'' + a(t)f(u) = 0, \quad t \in (0, 1),$$
$$u(0) = 0, \quad u(1) = \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} u(s) ds,$$

where $0 = \eta_0 < \eta_1 < \eta_2 < \dots < \eta_{m-2} < \eta_{m-1} = 1$, $0 < \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) < 2$, $\alpha_i \geq 0$ for $i \in \{1, \dots, m-3\} \cup \{m-1\}$ and $\alpha_{m-2} > 0$. $a \in C([0, 1], [0, \infty))$ and $f \in C([0, \infty), [0, \infty))$. We show the existence of at least one positive solution if f is either superlinear or sublinear by applying the fixed point theorem in cones.

Mathematics Subject Classification: 34B10; 34B15

Keywords: Positive solution; Fixed point theorem; Cone

1 Introduction

The study of multi-point boundary-value problems for linear second order ordinary differential equations was initiated by Il'in and Moiseev [1-2]. Gupta [3] studied three-point boundary-value problems for nonlinear second order ordinary differential equations. Since then, the existence of solutions for nonlinear three-point, m -point, integral and nonlocal boundary value problems has been studied by several authors by using the Leray-Schauder continuation theorem,

nonlinear alternatives of Leray-Schauder, coincidence degree theory and the fixed-point theorem in cones. We refer the reader to [4-21, 23-29] and the references therein.

However, most of those papers have studied the existence and multiplicity of solutions (or positive solutions) to boundary value problems with boundary conditions specified as solution values or the slope of solution values at given points or as more general nonlocal integral boundary conditions. There are also a few papers that consider nonlinear second order ordinary differential equations with boundary conditions specified as an area under the curve of solutions; see for example, [22].

In this paper, we consider the existence of at least one positive solution to the equation

$$u'' + a(t)f(u) = 0, \quad t \in (0, 1), \quad (1)$$

with the following boundary conditions that include an m -point integral condition

$$u(0) = 0, \quad u(1) = \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} u(s) ds, \quad (2)$$

where $0 = \eta_0 < \eta_1 < \eta_2 < \dots < \eta_{m-2} < \eta_{m-1} = 1$, $0 < \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) < 2$, $\alpha_i \geq 0$ for $i \in \{1, \dots, m-3\} \cup \{m-1\}$ and $\alpha_{m-2} > 0$. We note that the m -point boundary condition (2) are related to $m-1$ intervals of the area under the curve of solution $u(t)$ from $t = \eta_{i-1}$ to $t = \eta_i$ for $i = 1, \dots, m-1$.

By the positive solution of (1)-(2) we mean that a function $u(t)$ is positive on $0 < t < 1$ and satisfies the problem (1)-(2). Throughout this paper, we also assume the following.

(H1) $f \in C([0, \infty), [0, \infty))$ and the limits;

$$f_0 := \lim_{u \rightarrow 0^+} \frac{f(u)}{u}, \quad f_\infty := \lim_{u \rightarrow \infty} \frac{f(u)}{u},$$

exist. We denote that $f_0 = 0$ and $f_\infty = \infty$ correspond to the superlinear case, and $f_0 = \infty$ and $f_\infty = 0$ correspond to the sublinear case.

(H2) $a \in C([0, 1], [0, \infty))$, and there exists $t_0 \in [\eta_{m-2}, 1]$ such that $a(t_0) > 0$.

Moreover, we will work in the Banach space $C[0, 1]$, and only the sup norm is used. The proof of the main theorem is based upon an application of the following well-known Guo-Krasnoselskii fixed point theorem [30].

Theorem 1.1 *Let E be a Banach space, and let $K \subset E$ be a cone. Assume Ω_1, Ω_2 are open bounded subsets of E with $0 \in \Omega_1, \overline{\Omega_1} \subset \Omega_2$, and let*

$$A : K \cap (\overline{\Omega_2} \setminus \Omega_1) \longrightarrow K$$

be a completely continuous operator such that

(i) $\|Au\| \leq \|u\|$, $u \in K \cap \partial\Omega_1$, and $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_2$; or

(ii) $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_1$, and $\|Au\| \leq \|u\|$, $u \in K \cap \partial\Omega_2$.

Then A has a fixed point in $K \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

2 Preliminary Notes

In this section, we present some lemmas that will be used in the proof of our main results.

Lemma 2.1 *Let $\alpha_i \geq 0$ for $i = 1, 2, \dots, m-1$, and $\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2) \neq 2$. If $y \in C[0, 1]$, then the problem*

$$u''(t) + y(t) = 0, \quad 0 < t < 1, \tag{3}$$

$$u(0) = 0, \quad u(1) = \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} u(s) ds, \tag{4}$$

has a unique solution

$$\begin{aligned} u(t) = & - \int_0^t (t-s)y(s)ds + \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)y(s)ds \\ & - \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 y(s) ds \\ & + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 y(s) ds. \end{aligned} \tag{5}$$

Proof. From (3), we have

$$u''(t) = -y(t).$$

For $t \in [0, 1)$, integration from 0 to t , yields

$$u'(t) = - \int_0^t y(s) ds + u'(0).$$

For $t \in [0, 1]$, integration from 0 to t , gives

$$u(t) = - \int_0^t \left(\int_0^r y(s) ds \right) dr + u'(0)t,$$

i.e.,

$$u(t) = - \int_0^t (t-s)y(s)ds + u'(0)t. \quad (6)$$

So

$$u(1) = - \int_0^1 (1-s)y(s)ds + u'(0).$$

Integrating (6) from η_{i-1} to η_i for $0 \leq \eta_{i-1} < \eta_i \leq 1$, $i = 1, \dots, m-1$, and reversing the order of double integration, we get

$$\begin{aligned} \int_{\eta_{i-1}}^{\eta_i} u(s)ds &= - \int_{\eta_{i-1}}^{\eta_i} \left(\int_0^r (r-s)y(s)ds \right) dr + u'(0) \frac{\eta_i^2 - \eta_{i-1}^2}{2} \\ &= - \int_0^{\eta_i} \frac{(\eta_i - s)^2}{2} y(s)ds + \int_0^{\eta_{i-1}} \frac{(\eta_{i-1} - s)^2}{2} y(s)ds \\ &\quad + u'(0) \frac{\eta_i^2 - \eta_{i-1}^2}{2}, \quad i = 1, \dots, m-1. \end{aligned}$$

From condition (4), we have

$$\begin{aligned} u'(0) - \int_0^1 (1-s)y(s)ds &= - \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 y(s)ds \\ &\quad + \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 y(s)ds \\ &\quad + \frac{u'(0)}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2). \end{aligned}$$

Hence,

$$\begin{aligned} u'(0) &= \frac{2}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)y(s)ds \\ &\quad - \frac{1}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 y(s)ds \\ &\quad + \frac{1}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 y(s)ds. \end{aligned}$$

Therefore, (3)-(4) has a unique solution

$$\begin{aligned}
 u(t) = & - \int_0^t (t-s)y(s)ds + \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)y(s)ds \\
 & - \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 y(s)ds \\
 & + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 y(s)ds.
 \end{aligned}$$

□

Lemma 2.2 *Let $\alpha_i \geq 0$ for $i = 1, 2, \dots, m-1$, and $\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2) < 2$. If $y \in C([0, 1], [0, \infty))$, then the unique solution u of (3)-(4) satisfies $u(t) \geq 0$ for $t \in [0, 1]$.*

Proof. From the fact that $u''(t) = -y(t) \leq 0$, we know that the graph of $u(t)$ is concave down on $(0, 1)$. From this fact, we have that

$$\frac{u(\eta_1)}{\eta_1} \geq \frac{u(\eta_2)}{\eta_2} \geq \dots \geq \frac{u(\eta_{i-1})}{\eta_{i-1}} \geq \frac{u(\eta_i)}{\eta_i} \geq \dots \geq \frac{u(1)}{1}, \tag{7}$$

and

$$\int_{\eta_{i-1}}^{\eta_i} u(s)ds \geq \frac{1}{2}(\eta_i - \eta_{i-1})(u(\eta_i) + u(\eta_{i-1})), \tag{8}$$

where $\frac{1}{2}(\eta_i - \eta_{i-1})(u(\eta_i) + u(\eta_{i-1}))$ is the area of the trapezoid under the curve $u(t)$ from $t = \eta_{i-1}$ to $t = \eta_i$ for $i = 1, 2, \dots, m - 1$. Combining conditions (4), (7) and (8), we get

$$\begin{aligned}
 u(1) &= \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} u(s)ds \\
 &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i - \eta_{i-1})(u(\eta_i) + u(\eta_{i-1})) \\
 &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i - \eta_{i-1}) \left(u(\eta_i) + \frac{\eta_{i-1}}{\eta_i} u(\eta_i) \right) \\
 &= \frac{1}{2} \sum_{i=1}^{m-1} \frac{\alpha_i}{\eta_i} (\eta_i^2 - \eta_{i-1}^2) u(\eta_i). \tag{9}
 \end{aligned}$$

If $u(1) \geq 0$, then the concavity of u together with the boundary condition $u(0) = 0$ implies that $u(t) \geq 0$ for $t \in [0, 1]$.

If $u(1) < 0$, then conditions (4), (7) and (8) imply

$$\begin{aligned} u(1) &= \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} u(s) ds \\ &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i - \eta_{i-1}) (u(\eta_i) + u(\eta_{i-1})) \\ &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i - \eta_{i-1}) (\eta_i u(1) + \eta_{i-1} u(1)) \\ &= \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) u(1). \end{aligned}$$

This contradicts the fact that $\sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) < 2$. \square

Lemma 2.3 *Let $\alpha_i \geq 0$ for $i \in \{1, \dots, m-3\} \cup \{m-1\}$, $\alpha_{m-2} > 0$ and $\sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) > 2$. If $y \in C[0, 1]$ and $y(t) \geq 0$ for $t \in (0, 1)$, then (3)-(4) has no positive solution.*

Proof. Assume that (3)-(4) has a positive solution u , then $u(\eta_i) > 0$ for $i = 1, \dots, m-1$. By condition (9), we have

$$\begin{aligned} u(1) &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) \frac{u(\eta_i)}{\eta_i} \\ &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) \frac{u(\bar{\eta})}{\bar{\eta}} \\ &> \frac{u(\bar{\eta})}{\bar{\eta}}, \end{aligned}$$

where $\bar{\eta} \in \{\eta_1, \dots, \eta_{m-1}\}$ satisfies $\frac{u(\bar{\eta})}{\bar{\eta}} = \min\{\frac{u(\eta_i)}{\eta_i} \mid i = 1, \dots, m-1\}$. This contradicts the concavity of u .

If $u(1) = 0$, then applying $\alpha_{m-2} > 0$, we know that $\int_{\eta_{m-3}}^{\eta_{m-2}} u(s) ds = 0$ which implies $u(\eta_{m-2})u(\eta_{m-3}) \leq 0$. This contradicts the concavity of u . The proof is complete. \square

Lemma 2.4 *Let $\alpha_i \geq 0$ for $i \in \{1, 2, \dots, m-3\} \cup \{m-1\}$, $\alpha_{m-2} > 0$ and $\sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)^2 < 2$. If $y \in C[0, 1]$ and $y(t) \geq 0$ for $t \in (0, 1)$, then the unique solution u of problem (3)-(4) satisfies*

$$\inf_{t \in [\eta_{m-2}, 1]} u(t) \geq \gamma \|u\|,$$

where

$$\gamma = \min \left\{ \eta_{m-2}, \frac{1}{2} \sum_{i=1}^{m-2} \alpha_i (\eta_i^2 - \eta_{i-1}^2), \frac{\alpha_{m-2} (\eta_{m-2}^2 - \eta_{m-3}^2) (1 - \eta_{m-2})}{\eta_{m-2} (2 - \alpha_{m-2} (\eta_{m-2}^2 - \eta_{m-3}^2))}, \frac{1}{2} \sum_{i=1}^{m-2} \frac{\alpha_i}{\eta_i} (\eta_i^2 - \eta_{i-1}^2) (1 - \eta_i) \right\}. \tag{10}$$

Proof. Since $u''(t) = -y(t) \leq 0$, the graph of $u(t)$ is concave down on $(0, 1)$. If $u(t)$ is maximum at $t = \bar{t}$, then $\|u\| = u(\bar{t})$. We divide the proof into four cases.

Case I. If $0 < \eta_1 < \dots < \eta_{m-2} < \bar{t} \leq 1$ and $\inf_{t \in [\eta_{m-2}, 1]} u(t) = u(\eta_{m-2})$, then the concavity of u implies that

$$\frac{u(\eta_{m-2})}{\eta_{m-2}} \geq \frac{u(\bar{t})}{\bar{t}} \geq u(\bar{t}).$$

Therefore,

$$\inf_{t \in [\eta_{m-2}, 1]} u(t) \geq \eta_{m-2} \|u\|.$$

Case II. If $0 < \eta_1 < \dots < \eta_{m-2} \leq \bar{t} < 1$ and $\inf_{t \in [\eta_{m-2}, 1]} u(t) = u(1)$. Then (4), (8) and the concavity of u imply

$$\begin{aligned} u(1) &= \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} u(s) ds \\ &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i (\eta_i - \eta_{i-1}) (u(\eta_i) + u(\eta_{i-1})) \\ &\geq \frac{1}{2} \sum_{i=1}^{m-2} \alpha_i (\eta_i - \eta_{i-1}) (u(\eta_i) + u(\eta_{i-1})) \\ &= \frac{1}{2} \sum_{i=1}^{m-2} \alpha_i (\eta_i^2 - \eta_{i-1}^2) \frac{\left(\frac{u(\eta_i) + u(\eta_{i-1})}{2}\right)}{\left(\frac{\eta_i + \eta_{i-1}}{2}\right)} \\ &\geq \frac{1}{2} \sum_{i=1}^{m-2} \alpha_i (\eta_i^2 - \eta_{i-1}^2) \frac{u(\bar{t})}{\bar{t}} \\ &\geq \frac{1}{2} \sum_{i=1}^{m-2} \alpha_i (\eta_i^2 - \eta_{i-1}^2) u(\bar{t}). \end{aligned}$$

This implies

$$\inf_{t \in [\eta_{m-2}, 1]} u(t) \geq \frac{1}{2} \sum_{i=1}^{m-2} \alpha_i (\eta_i^2 - \eta_{i-1}^2) \|u\|.$$

Case III. If $0 < \eta_1 < \dots < \eta_i \leq \bar{t} < \eta_{i+1} < \dots < \eta_{m-2} < 1$ for some $i \in \{1, \dots, m-3\}$, and $\inf_{t \in [\eta_{m-2}, 1]} u(t) = u(1)$. From (9), we get that

$$u(1) \geq \frac{1}{2} \sum_{i=1}^{m-1} \frac{\alpha_i(\eta_i^2 - \eta_{i-1}^2)u(\eta_i)}{\eta_i} \geq \frac{\alpha_{m-2}}{2\eta_{m-2}} (\eta_{m-2}^2 - \eta_{m-3}^2)u(\eta_{m-2}). \tag{11}$$

From the concavity of u and (11), we get

$$\begin{aligned} u(\bar{t}) &\leq u(1) + \frac{u(1) - u(\eta_{m-2})}{1 - \eta_{m-2}}(0 - 1) \\ &= u(1) - \frac{u(1)}{1 - \eta_{m-2}} + \frac{u(\eta_{m-2})}{1 - \eta_{m-2}} \\ &\leq u(1) \left[1 - \frac{1}{1 - \eta_{m-2}} + \frac{2\eta_{m-2}}{\alpha_{m-2}(\eta_{m-2}^2 - \eta_{m-3}^2)(1 - \eta_{m-2})} \right] \\ &= \frac{\eta_{m-2}(2 - \alpha_{m-2}(\eta_{m-2}^2 - \eta_{m-3}^2))}{\alpha_{m-2}(\eta_{m-2}^2 - \eta_{m-3}^2)(1 - \eta_{m-2})} u(1). \end{aligned}$$

Thus,

$$\inf_{t \in [\eta_{m-2}, 1]} u(t) \geq \frac{\alpha_{m-2}(\eta_{m-2}^2 - \eta_{m-3}^2)(1 - \eta_{m-2})}{\eta_{m-2}(2 - \alpha_{m-2}(\eta_{m-2}^2 - \eta_{m-3}^2))} \|u\|.$$

Case IV. If $\bar{t} \leq \eta_1 < \eta_2 < \dots < \eta_{m-2} < 1$ and $\inf_{t \in [\eta_{m-2}, 1]} u(t) = u(1)$. It is easy to see from the concavity of u that

$$\frac{u(\eta_{m-2})}{1 - \eta_{m-2}} \geq \frac{u(\eta_{m-3})}{1 - \eta_{m-3}} \geq \dots \geq \frac{u(\eta_1)}{1 - \eta_1} \geq \frac{u(\bar{t})}{1 - \bar{t}} \geq u(\bar{t}).$$

This, together with (9), implies that

$$\begin{aligned} u(1) &\geq \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i \frac{(\eta_i^2 - \eta_{i-1}^2)}{\eta_i} (1 - \eta_i) u(\bar{t}) \\ &= \frac{1}{2} \sum_{i=1}^{m-2} \frac{\alpha_i}{\eta_i} (\eta_i^2 - \eta_{i-1}^2) (1 - \eta_i) u(\bar{t}). \end{aligned}$$

This implies that

$$\inf_{t \in [\eta_{m-2}, 1]} u(t) \geq \frac{1}{2} \sum_{i=1}^{m-2} \frac{\alpha_i}{\eta_i} (\eta_i^2 - \eta_{i-1}^2) (1 - \eta_i) \|u\|.$$

This completes the proof. □

3 Main Results

In this section we discuss the existence of at least one positive solution for the problem (1)-(2). We obtain the following existence results.

Theorem 3.1 *Assume (H1) and (H2) hold. Then the problem (1)-(2) has at least one positive solution in the case*
 (i) $f_0 = 0$ and $f_\infty = \infty$ (superlinear), or
 (ii) $f_0 = \infty$ and $f_\infty = 0$ (sublinear).

Proof. Superlinear case. Suppose then that $f_0 = 0$ and $f_\infty = \infty$. We wish to show the existence of a positive solution of (1)-(2). Now (1)-(2) has a solution $u = u(t)$ if and only if u solves the operator equation

$$\begin{aligned}
 u(t) = & - \int_0^t (t-s)a(s)f(u(s))ds \\
 & + \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)f(u(s))ds \\
 & - \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 a(s)f(u(s))ds \\
 & + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)f(u(s))ds \\
 := & Au(t).
 \end{aligned} \tag{12}$$

Denote

$$K = \left\{ u \mid u \in C[0, 1], u \geq 0, \min_{\eta_{m-2} \leq t \leq 1} u(t) \geq \gamma \|u\| \right\},$$

where γ is defined by (10). It is obvious that K is a cone in $C[0, 1]$. Moreover, by Lemma 2.4, $AK \subset K$. It is also easy to check that $A : K \rightarrow K$ is completely continuous.

Since $f_0 = 0$, we may choose $H_1 > 0$ so that $f(u) \leq \epsilon u$, for $0 < u < H_1$, where $\epsilon > 0$ satisfies

$$\left(\frac{2 \int_0^1 (1-s)a(s)ds + \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)ds}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \right) \epsilon \leq 1. \tag{13}$$

Thus, if $u \in K$ and $\|u\| = H_1$, then from (12) and (13), we get

$$\begin{aligned}
Au(t) &\leq \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)f(u(s))ds \\
&\quad + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)f(u(s))ds \\
&\leq \frac{2}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)\epsilon u(s)ds \\
&\quad + \frac{1}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)\epsilon u(s)ds \\
&\leq \frac{2\epsilon}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)ds \|u\| \\
&\quad + \frac{\epsilon}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)ds \|u\| \\
&\leq \left(\frac{2 \int_0^1 (1-s)a(s)ds + \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \right) \epsilon H_1. \quad (14)
\end{aligned}$$

Let $\Omega_1 = \{u \in C[0, 1] \mid \|u\| < H_1\}$, then (14) shows that $\|Au\| \leq \|u\|$, for $u \in K \cap \partial\Omega_1$.

Further, since $f_\infty = \infty$, there exists $\widehat{H}_2 > 0$ such that $f(u) \geq \rho u$, for $u \geq \widehat{H}_2$, where $\rho > 0$ is chosen so that

$$\rho \gamma \frac{\sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \int_{\eta_{m-2}}^1 (1-s)a(s)ds \geq 1. \quad (15)$$

Let $H_2 = \max\{2H_1, (\widehat{H}_2/\gamma)\}$ and $\Omega_2 = \{u \in C[0, 1] \mid \|u\| < H_2\}$, then $u \in K$ and $\|u\| = H_2$ implies

$$\min_{t \in [\eta_{m-2}, 1]} u(t) \geq \gamma \|u\| \geq \widehat{H}_2.$$

So,

$$\begin{aligned}
 Au(s) = & - \int_0^s (s-r)a(r)f(u(r))dr \\
 & + \frac{2s}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-r)a(r)f(u(r))dr \\
 & - \frac{s}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - r)^2 a(r)f(u(r))dr \\
 & + \frac{s}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - r)^2 a(r)f(u(r))dr.
 \end{aligned}$$

This implies

$$\begin{aligned}
 u(1) = & \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} u(s)ds \\
 = & \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} \left(- \int_0^s (s-r)a(r)f(u(r))dr \right. \\
 & + \frac{2s}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-r)a(r)f(u(r))dr \\
 & - \frac{s}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - r)^2 a(r)f(u(r))dr \\
 & \left. + \frac{s}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - r)^2 a(r)f(u(r))dr \right) ds \\
 = & - \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} \int_0^s (s-r)a(r)f(u(r))drds \\
 & + \frac{2 \int_0^1 (1-r)a(r)f(u(r))dr}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \left(\sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} sds \right)
 \end{aligned}$$

$$\begin{aligned}
& - \frac{\sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - r)^2 a(r) f(u(r)) dr}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \left(\sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} s ds \right) \\
& + \frac{\sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - r)^2 a(r) f(u(r)) dr}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \left(\sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} s ds \right) \\
& = \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s) f(u(s)) ds \\
& - \frac{1}{2} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 a(s) f(u(s)) ds \\
& + \frac{\int_0^1 (1-s) a(s) f(u(s)) ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \left(\sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) \right) \\
& - \frac{\sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 a(s) f(u(s)) ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \left(\sum_{i=1}^{m-1} \frac{\alpha_i}{2} (\eta_i^2 - \eta_{i-1}^2) \right) \\
& + \frac{\sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s) f(u(s)) ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \left(\sum_{i=1}^{m-1} \frac{\alpha_i}{2} (\eta_i^2 - \eta_{i-1}^2) \right) \\
& = \frac{\int_0^1 (1-s) a(s) f(u(s)) ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2) \\
& - \frac{\sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 a(s) f(u(s)) ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \\
& + \frac{\sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s) f(u(s)) ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \\
& = \frac{1}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \left[\int_0^{\eta_{i-1}} (\eta_i - \eta_{i-1}) \right. \\
& \quad \times (2 - (\eta_i + \eta_{i-1})) s a(s) f(u(s)) ds \\
& \quad + \int_{\eta_{i-1}}^{\eta_i} [(\eta_i^2 - \eta_{i-1}^2)(1-s) - (\eta_i - s)^2] a(s) f(u(s)) ds \\
& \quad \left. + \int_{\eta_i}^1 (\eta_i^2 - \eta_{i-1}^2)(1-s) a(s) f(u(s)) ds \right] \\
& \geq \frac{1}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \left[\int_{\eta_{i-1}}^{\eta_i} (\eta_i - \eta_{i-1}) \right. \\
& \quad \times [\eta_{i-1}(1-s) + (1-\eta_i)s] a(s) f(u(s)) ds
\end{aligned}$$

$$\begin{aligned}
 & \left. + (\eta_i^2 - \eta_{i-1}^2) \int_{\eta_i}^1 (1-s)a(s)f(u(s))ds \right] \\
 \geq & \frac{1}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2) \int_{\eta_i}^1 (1-s)a(s)f(u(s))ds \\
 \geq & \frac{\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_{\eta_{m-2}}^1 (1-s)a(s)f(u(s))ds. \tag{16}
 \end{aligned}$$

Hence, for $u \in K \cap \partial\Omega_2$,

$$\|Au\| \geq |u(1)| \geq \rho\gamma \frac{\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_{\eta_{m-2}}^1 (1-s)a(s)ds \|u\| \geq \|u\|.$$

By the first part of Theorem 1.1, it follows that A has a fixed point in $K \cap (\bar{\Omega}_2 \setminus \Omega_1)$, such that $H_1 \leq \|u\| \leq H_2$.

Sublinear case. Suppose that $f_0 = \infty$ and $f_\infty = 0$. We first choose $H_3 > 0$ such that $f(u) \geq Mu$ for $0 < u < H_3$, where

$$\gamma M \frac{\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_{\eta_{m-2}}^1 (1-s)a(s)ds \geq 1.$$

For $u \in K$ and $\|u\| = H_3$, by a similar method used to obtain (16), we can get that

$$\begin{aligned}
 Au(1) &= \sum_{i=1}^{m-1} \alpha_i \int_{\eta_{i-1}}^{\eta_i} Au(s)ds \\
 &\geq \frac{1}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2) \int_{\eta_i}^1 (1-s)a(s)f(u(s))ds \\
 &\geq \frac{\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_{\eta_{m-2}}^1 (1-s)a(s)f(u(s))ds \\
 &\geq \frac{\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_{\eta_{m-2}}^1 (1-s)a(s)Mu(s)ds \\
 &\geq \gamma M \frac{\sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_{\eta_{m-2}}^1 (1-s)a(s)ds \|u\| \\
 &\geq H_3. \tag{17}
 \end{aligned}$$

Let $\Omega_3 = \{u \in C[0, 1] \mid \|u\| < H_3\}$. Thus, (17) shows that $\|Au\| \geq \|u\|$, $u \in K \cap \partial\Omega_3$.

Next, in view of $f_\infty = 0$, there exists $\widehat{H}_4 > 0$ such that $f(u) \leq \lambda u$ for $u \geq \widehat{H}_4$, where $\lambda > 0$ satisfies

$$\left(\frac{2 \int_0^1 (1-s)a(s)ds + \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1}-s)^2 a(s)ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \right) \lambda \leq 1.$$

We consider the following two cases.

Case (i). Suppose f is bounded, say $f(u) \leq N$ for all $u \in [0, \infty)$. In this case, choose

$$H_4 = \max \left\{ 2H_3, \left(\frac{2 \int_0^1 (1-s)a(s)ds + \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1}-s)^2 a(s)ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \right) N \right\},$$

such that, for $u \in K$ with $\|u\| = H_4$, we have

$$\begin{aligned} Au(t) &= - \int_0^t (t-s)a(s)f(u(s))ds \\ &\quad + \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)f(u(s))ds \\ &\quad - \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 a(s)f(u(s))ds \\ &\quad + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)f(u(s))ds \\ &\leq \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)f(u(s))ds \\ &\quad + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)f(u(s))ds \\ &\leq \left(\frac{2 \int_0^1 (1-s)a(s)ds + \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1}-s)^2 a(s)ds}{2 - \sum_{i=1}^{m-1} \alpha_i (\eta_i^2 - \eta_{i-1}^2)} \right) N \\ &\leq H_4. \end{aligned}$$

Therefore, $\|Au\| \leq \|u\|$.

Case (ii). If f is unbounded, which implies that there is H_4 , $H_4 > \max\{2H_3, (\widehat{H}_4/\gamma)\}$, such that $f(u) \leq f(H_4)$ for $0 < u \leq H_4$. Then for $u \in K$ and $\|u\| = H_4$, we

have

$$\begin{aligned}
 Au(t) &= - \int_0^t (t-s)a(s)f(u(s))ds \\
 &\quad + \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)f(u(s))ds \\
 &\quad - \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_i} (\eta_i - s)^2 a(s)f(u(s))ds \\
 &\quad + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)f(u(s))ds \\
 &\leq \frac{2t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \int_0^1 (1-s)a(s)f(H_4)ds \\
 &\quad + \frac{t}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)f(H_4)ds \\
 &\leq \left(\frac{2 \int_0^1 (1-s)a(s)ds + \sum_{i=1}^{m-1} \alpha_i \int_0^{\eta_{i-1}} (\eta_{i-1} - s)^2 a(s)ds}{2 - \sum_{i=1}^{m-1} \alpha_i(\eta_i^2 - \eta_{i-1}^2)} \right) \lambda H_4 \\
 &\leq H_4.
 \end{aligned}$$

Hence, in either case, we may always let $\Omega_4 = \{u \in C[0, 1] \mid \|u\| < H_4\}$, and for $u \in K \cap \partial\Omega_4$. By the second part of Theorem 1.1, we can conclude that problem (1)-(2) has at least one positive solution. \square

4 Some examples

In this section, in order to illustrate our result, we consider some examples.

Example 4.1 Consider the eleven-point boundary value problem

$$u''(t) + e^{2t}(e^u u^e \sin u + u^2 \ln(1 + u)) = 0, \quad 0 < t < 1, \tag{18}$$

$$\begin{aligned}
 u(0) = 0, \quad u(1) &= 10e \int_0^{\frac{1}{10}} u(s)ds + \frac{4e}{5} \int_{\frac{1}{5}}^{\frac{3}{10}} u(s)ds + \frac{8e}{9} \int_{\frac{2}{5}}^{\frac{5}{10}} u(s)ds \\
 &\quad + \frac{12e}{13} \int_{\frac{3}{5}}^{\frac{7}{10}} u(s)ds + \frac{16e}{17} \int_{\frac{4}{5}}^{\frac{9}{10}} u(s)ds. \tag{19}
 \end{aligned}$$

Set $\eta_i = \frac{i}{10}$ for $i = 0, \dots, 10$, $a(t) = e^{2t}$ and $f(u) = e^u u^e \sin u + u^2 \ln(1 + u)$, $\alpha_{2j} = 0$ for $j = 1, \dots, 5$, and $\alpha_1 = 10e$, $\alpha_3 = \frac{4e}{5}$, $\alpha_5 = \frac{8e}{9}$, $\alpha_7 = \frac{12e}{13}$, $\alpha_9 = \frac{16e}{17}$.

We can show that

$$\sum_{i=1}^{10} \alpha_i (\eta_i^2 - \eta_{i-1}^2) = \frac{e}{2} < 2.$$

Through a simple calculation we can get $f_0 = 0$ and $f_\infty = \infty$. Thus, by the first part of Theorem 3.1, we can get that the problem (18)-(19) has at least one positive solution.

Example 4.2 Consider the eleven-point boundary value problem

$$u''(t) + \pi^{tt^\pi} f(u) = 0, \quad 0 < t < 1, \quad (20)$$

$$\begin{aligned} u(0) = 0, \quad u(1) = & 2\pi \int_{\frac{1}{10}}^{\frac{1}{5}} u(s) ds + \pi \int_{\frac{3}{10}}^{\frac{2}{5}} u(s) ds + \frac{10\pi}{11} \int_{\frac{5}{10}}^{\frac{3}{5}} u(s) ds \\ & + \frac{3\pi}{5} \int_{\frac{7}{10}}^{\frac{4}{5}} u(s) ds + \frac{8\pi}{17} \int_{\frac{4}{5}}^{\frac{9}{10}} u(s) ds + \frac{10\pi}{19} \int_{\frac{9}{10}}^1 u(s) ds, \end{aligned} \quad (21)$$

where

$$f(u) = \begin{cases} \frac{\pi \sin u + 2 \cos u}{u^2}, & 0 < u \leq \frac{\pi}{2} \\ \frac{\sqrt{8u+2}\sqrt{\pi} \sin^2 u}{\pi^{\frac{3}{2}}}, & u > \frac{\pi}{2}. \end{cases}$$

Set $\eta_i = \frac{i}{10}$ for $i = 0, \dots, 10$, $a(t) = \pi^{tt^\pi}$, $\alpha_{2j-1} = 0$ for $j = 1, \dots, 4$, and $\alpha_2 = 2\pi$, $\alpha_4 = \pi$, $\alpha_6 = \frac{10\pi}{11}$, $\alpha_8 = \frac{3\pi}{5}$, $\alpha_9 = \frac{8\pi}{17}$, $\alpha_{10} = \frac{10\pi}{19}$. We can show that

$$\sum_{i=1}^{10} \alpha_i (\eta_i^2 - \eta_{i-1}^2) = \frac{\pi}{2} < 2.$$

Through a simple calculation we can get $f_0 = \infty$ and $f_\infty = 0$. Thus, by the second part of Theorem 3.1, we can get that the problem (20)-(21) has at least one positive solution.

ACKNOWLEDGEMENT. This research was supported by King Mongkut's University of Technology North Bangkok, Thailand.

References

- [1] V.A. Il'in, E.I. Moiseev, Nonlocal boundary value problem of the first kind for a Sturm-Liouville operator in its differential and finite difference aspects, *Differential Equations*, vol. 23, no.7, 803-810, 1987.

- [2] V.A. Il'in, E.I. Moiseev, Nonlocal boundary value problem of the second kind for a Sturm-Liouville operator, *Differential Equations*, vol. 23, no.8, 979-987, 1987.
- [3] C.P. Gupta, Solvability of a three-point nonlinear boundary value problem for a second order ordinary differential equation, *Journal of Mathematical Analysis and Applications*, vol. 168, 540-551, 1992.
- [4] M. Benchohra, J. J. Nieto, A. Ouahab, Second-order boundary value problem with integral boundary conditions, *Boundary Value Problems*. **2011**, Article ID 260309, 9 pp., 2011.
- [5] A. Boucherif, Second-order boundary problems with integral boundary conditions, *Nonlinear Analysis. Theory, Method & Applications*. vol. 70, no. 1, 364-371, 2009.
- [6] M. Feng, W. Ge, Positive solutions for a class of m -point singular boundary value problems, *Mathematical and Computer Modelling*, vol. 46, no. 3-4, 375-383, 2007.
- [7] C.P. Gupta, S. K. Ntouyas, P. Ch. Tsamatos, Solvability of an m -point boundary-value problem for second-order ordinary differential equations, *Journal of Mathematical Analysis and Applications*, vol. 189, 575-584, 1995.
- [8] Y. Guo, W. Shan, W. Ge, Positive solutions for second-order m -point boundary value problems, *Journal of Computational and Applied Mathematics*, vol. 151, no. 2, 415-424, 2003.
- [9] J. Jiang, L. Liu, Positive solutions for nonlinear second-order m -point boundary-value problems, *Electronic Journal of Differential Equations*, vol. 2009, no. 110, 12 pp., 2009.
- [10] J. Jiang, L. Liu, Y. Wu, Second-order nonlinear singular Sturm-Liouville problems with integral boundary conditions, *Applied Mathematics and Computation*, vol. 215, no. 4, 1573-1582, 2009.
- [11] G. L. Karakostas, P. Ch. Tsamatos, Existence of multiple positive solutions for a nonlocal boundary value problem, *Topological Methods in Nonlinear Analysis*, vol. 19, 109-121, 2002.
- [12] G. L. Karakostas, P. Ch. Tsamatos, Multiple positive solutions of some Fredholm integral equations arisen from nonlocal boundary-value problems, *Electronic Journal of Differential Equations*, vol. 2002, no. 30, 17 pp., 2002.

- [13] L. Kong, Second order singular boundary value problems with integral boundary conditions, *Nonlinear Analysis. Theory, Method & Applications*, vol. 72, no.5, 2628-2638, 2010.
- [14] B. Liu, Positive solutions of a nonlinear three-point boundary value problem, *Applied Mathematics and Computation*, vol. 132, no. 1, 11-28, 2002.
- [15] B. Liu, Positive solutions of a nonlinear three-point boundary value problem, *Computers & Mathematics with Applications*, vol. 44, no. 1-2, 201-211, 2002.
- [16] B. Liu, Positive solutions of a nonlinear four-point boundary value problems, *Applied Mathematics and Computation*, vol. 155, no. 1, 179-203, 2004.
- [17] R. Ma, Positive solutions of a nonlinear three-point boundary-value problem, *Electronic Journal of Differential Equations*, vol. 1998, no. 34, 1-8, 1998.
- [18] R. Ma, Positive solutions of a nonlinear m-point boundary value problem, *Computers & Mathematics with Applications*, vol. 42, no. 6-7, 755-765, 2001.
- [19] R. Ma, H. B. Thompson, Positive solutions for nonlinear m-point eigenvalue problems, *Journal of Mathematical Analysis and Applications*, vol. 297, no. 1, 24-37, 2004.
- [20] R. Ma, A survey on nonlocal boundary value problems, *Applied Mathematics E-Notes*, vol. 7, 257-279, 2007.
- [21] Y. Sun, Positive solutions of nonlinear second-order m-point boundary value problem, *Nonlinear Analysis. Theory, Method & Applications*, vol. 61, no. 7, 1283-1294, 2005.
- [22] J. Tariboon, T. Sitthiwirattam, Positive solutions of a nonlinear three-point integral boundary value problem, *Boundary Value Problems*, vol. 2010, Article ID 519210, 11 pp., 2010.
- [23] J. R. L. Webb, A unified approach to nonlocal boundary value problems, *Dynamic systems and applications*, vol. 5, 510-515, Dynamic, Atlanta, GA, 2008.
- [24] J. R. L. Webb, Solutions of nonlinear equations in cones and positive linear operators, *Journal of London Mathematical Society*, vol. 82, no. 2, 420-436, 2010.

- [25] J. R. L. Webb, G. Infante, Positive solutions of nonlocal boundary value problems involving integral conditions, *NoDEA Nonlinear Differential Equations and Applications*, vol. 15, 45-67, 2008.
- [26] J. R. L. Webb, G. Infante, Positive solutions of nonlocal boundary value problems: a unified approach, *Journal of London Mathematical Society*, vol. 74, no.2, 673-693, 2006.
- [27] Z. Yang, Existence and nonexistence results for positive solutions of an integral boundary value problem, *Nonlinear Analysis. Theory, Method & Applications*, vol. 65, no. 8, 1489-1511, 2006.
- [28] Z. Yang, Existence of nontrivial solutions for a nonlinear Sturm-Liouville problem with integral boundary conditions, *Nonlinear Analysis. Theory, Method & Applications*, vol. 68, no. 1, 216-225, 2008.
- [29] G. Zhang, J. Sun, Positive solutions of m -point boundary value problems, *Journal of Mathematical Analysis and Applications*, vol. 291, no. 2, 406-418, 2004.
- [30] D. Guo, V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*. Academic Press, San Diego, CA, 1988.

Received: August, 2011