

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Mineral contributions to the diet and health of wild chimpanzees in three**
2 **East African forests, with notes on mechanisms and underlying mineral con-**
3 **centrations**

4
5 Running title: Mineral contributions to chimpanzee diet

6
7 Vernon Reynolds^{1,2}, Alejandra Pascual-Garrido³, Andrew W. Lloyd⁴, Peter Lyons⁴, Catherine
8 Hobaiter^{2,5}

9
10 ¹ School of Anthropology, Oxford University, Oxford, UK; ² Budongo Conservation Field Sta-
11 tion, Masindi, Uganda; ³ School of Archaeology, Oxford University, Oxford, UK; ⁴ School of
12 Environment & Technology, University of Brighton, Brighton, UK; ⁵ School of Psychology and
13 Neuroscience, University of St Andrews, St Marys College, South Street, St Andrews, Scot-
14 land

15
16

1
2
3 17 **Abstract**
4

5 18 We present new data on the ingestion of minerals from termite mound soil (TMS) by chim-
6
7 19 panzees (*Pan troglodytes schweinfurthii*) living in the Budongo Forest Reserve, Uganda.

8
9
10 20 Termite mound soil is here shown to be a rich source of minerals, containing high concen-
11
12 21 trations of iron and aluminium. Termite mound soil is not, however, a source of sodium. The
13
14 22 concentrations of iron and aluminium are the highest yet found in any of the mineral
15
16 23 sources consumed. Levels of manganese and copper, though not so high as for iron and al-
17
18 24 uminium, are also higher than in other dietary sources. We focus on the contribution of
19
20 25 termite mound soil to other known sources of mineral elements consumed by these apes,
21
22 26 and compare the mineral content of termite soil with that of control forest soil, decaying
23
24 27 wood, clay, and the normal plant-based chimpanzee diet at Budongo. Samples obtained
25
26 28 from Mahale Mountains National Park and Gombe National Park, both in Tanzania, show
27
28 29 similar mineral distribution across sources. Finally, we suggest three distinct but related
29
30 30 mechanisms by which minerals may come to be concentrated in the above-mentioned
31
32 31 sources.
33
34

35
36
37 32
38
39 33 **Keywords:** geophagy; *Pan troglodytes*; termite mound soil, minerals; diet; chimpanzees;
40
41 34 Uganda; Tanzania
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 35 **Introduction**
4

5
6 36 Some bird and mammalian species, including elephants, macaques, tamarins, gorillas, chim-
7
8 37 panzees and humans (Wilson 2003), consume soil of a variety of kinds, often in the form of
9
10 38 clay, for its mineralogical content. In doing so they obtaining both essential minerals, which
11
12 39 may have nutritional value, and non-nutritional minerals that can aid in the detoxification of
13
14 40 harmful compounds such as alkaloids in the diet (Klaus. Klaus-Hugi and Schmid, 1998), and
15
16 41 alleviate gastro-intestinal upsets (Mahaney et al., 1996). As pointed out by Wilson (2003), in
17
18 42 a review of the literature in this field, physiological tests are needed to determine bioavaila-
19
20 43 bility of mineral elements eaten in the course of geophagy. Probably no single characteristic
21
22 44 of soils eaten by animals and humans can account for their consumption (Abrahams, 1999;
23
24 45 Wilson, 2003), with mineral supplementation, medical, and detoxification functions all play-
25
26 46 ing a part (Aufreiter, Hancock, Mahaney, Strambolic-Robb and Sanmagudas, 1997; Aufreiter
27
28 47 et al., 2001; Ketch, Malloch, Mahaney and Huffman, 2001; Mahaney, 1993; Mahaney et al.,
29
30 48 1999; Vermeer and Ferrell, 1985; Wilson, 2003).
31
32
33
34
35

36
37
38 50 The normal diet of wild chimpanzees in the Budongo Forest, Uganda, is typical of East Afri-
39
40 51 can chimpanzee groups, and consists primarily of fruits and leaves, with additional flowers,
41
42 52 bark, and pith. Besides these plant-based items, meat and insects are eaten sporadically
43
44 53 when they become available. Both meat, obtained primarily by killing monkeys (Nishida,
45
46 54 Uehara and Nyundo, 1979; Goodall, 1986; Mitani and Watts, 2001; Newton-Fisher, Notman
47
48 55 and Reynolds, 2002) and insects, for example termites (O'Malley and Power 2014), are high-
49
50 56 ly nutritious sources of minerals as well as proteins, fats and other dietary requirements.
51
52
53 57 However the bulk of the food eaten by wild chimpanzees is plant-based and this constitutes
54
55 58 80% or more of the daily diet of most individuals. While high in some minerals e.g. potassi-
56
57
58
59
60

1
2
3 59 um and calcium, the Budongo chimpanzees' diet lacks (or has low quantities) of others e.g.
4
5 60 aluminium, copper, manganese, and sodium, and, as a result, they need to locate these
6
7 61 minerals from other sources (Reynolds, Lloyd, Babweteera and English, 2009). Earlier work
8
9 62 (Reynolds et al., 2009; Reynolds, Lloyd and English, 2012; Reynolds et al., 2015) explored a
10
11 63 number of dietary supplements for mineral acquisition, namely decaying pith of *Raphia fa-*
12
13 64 *rinifera* and the decaying wood of *Cleistopholis patens*, which provide appreciable amounts
14
15 65 of sodium (Reynolds et al., 2009, 2012), and clay, which provides substantial amounts of
16
17 66 iron (Reynolds et al., 2015). In this paper we show that termite mound soil is a further valu-
18
19 67 able source of minerals eaten by chimpanzees in the Budongo Forest Reserve, Uganda, by
20
21 68 the Kasekela group at Gombe National Park and by the M group at the Mahale Mountains
22
23 69 National Park (Aufreiter et al., 2001).
24
25
26
27
28 70 Some discussion revolves around the extent of bioavailability of the iron ingested in termite
29
30 71 mound soil (Aufreiter et al., 2001). In part this resolves itself into the question of whether
31
32 72 the iron is in ferric (Fe^{3+}) or ferrous (Fe^{2+}) form. If the former, it is not bioavailable; if the lat-
33
34 73 ter it is. Experimental work (Aufreiter et al., 2001) using a medium with low pH to simulate
35
36 74 digestive conditions suggests that most of the iron in soil is in ferric form and only a small
37
38 75 part is ferrous. This finding suggests that the nutritional value of ingested termite mound
39
40 76 soil may be limited. However we should note that in humans a ferric reductase enzyme, du-
41
42 77 odenal cytochrome B, reduces ferric Fe^{3+} to Fe^{2+} (McKie et al., 2001). This enzyme, if present
43
44 78 in chimpanzees, as seems likely, serves to increase the bioavailability of iron ingested in
45
46 79 termite mound soil. If present, ferrihydrite, a hydrous ferric oxide mineral, is likely to be sol-
47
48 80 ubilised (Wilson, 2003). Mahaney et al (1997) concluded that in geophagy soils eaten by
49
50 81 chimpanzees in the Kibale Forest, Uganda, 20% of ingested iron was bioavailable, sufficient
51
52 82 for nutritional significance. In a study of soils eaten by humans and sold in local markets in
53
54
55
56
57
58
59
60

1
2
3 83 Uganda, it was concluded that consumption of 5g of soil contributed 19-25% of daily needs
4
5 84 for iron (Abrahams and Parsons, 1997; Abrahams 1997). Geissler et al. (1998), by contrast,
6
7 85 found that despite consuming 30g daily of iron-rich termite mound soil, anaemia remained
8
9 86 prevalent in a human population in Kenya. Pregnant women were particularly prone to eat-
10
11 87 ing clays in Uganda and other tropical countries. Pregnant Chacma baboons (*Papio ursinus*)
12
13 88 spent more time consuming iron-rich clay at monitored geophagy sites in Western Cape,
14
15 89 South Africa than baboons of other age-sex classes (Pebsworth, Bardi and Huffman, 2011).
16
17
18
19
20

21 91 Whereas the majority of minerals discussed in this paper can be regarded as either major
22
23 92 minerals essential for life or minor minerals required only as trace elements, aluminium is
24
25 93 neither of these and is not essential for life. Its ingestion in termite mound soil, probably in
26
27 94 the form of kaolinite (Johns and Duquette, 1991; Mahaney et al., 1995) and in some cases
28
29 95 gibbsite (Bolton, Campbell and Burton, 1998), probably serves medicinal functions, by re-
30
31 96 ducing acidity in the gut and neutralising plant toxins such as condensed tannins (Hladik,
32
33 97 1977; Goodall, 1986). Condensed tannins are ingested by chimpanzees on a daily basis at
34
35 98 Budongo, being found at high concentrations in several species of figs (*Ficus sp*), particularly
36
37 99 in the seed component. One fig species with a high concentration of condensed tannins, *Fi-*
38
39 100 *cus sur*, is the second most frequently eaten food of the Budongo chimpanzees. Condensed
40
41
42
43
44 101 tannins thus appear to be well tolerated by chimpanzees (Reynolds, Plumtre, Greenham and
45
46 102 Harbone, 1998; Wrangham, 1993; Aufreiter et al., 2001).
47
48
49

50
51 104 Termite mound soil eating appears to be an opportunist and largely individual activity, oc-
52
53 105 ccurring when the animals pass by a termite mound in the forest, often moving from one
54
55 106 vegetative feeding site to another. Observations by researchers and field assistants indicate
56
57
58
59
60

1
2
3 107 that “Gombe chimpanzees eat termite mound soil, on average, once a day” (Wrangham,
4
5 108 1977) and the same may be true at Mahale and Budongo. Anecdotal reports suggest that at
6
7 109 all three sites termite mound soil eating is more frequent among females than males, but
8
9 110 quantitative data are lacking. Termite mounds present a hard surface (Figure 1) and chim-
11
12 111 panzees either bite off a piece with their teeth or break off a piece with their fingers (Figure
13
14 112 2). At Mahale, chimpanzees eat the soil of termite mounds through the year, possibly to as-
15
16 113 sess the reproductive condition of the termites (Uehara, 1982), as well as during times of
17
18 114 gastrointestinal distress. At Gombe, about once a day, as they pass termite mounds, chim-
19
20 115 panzees pick off and eat a “walnut” sized piece of termite mound soil (Goodall, 1986; Ma-
21
22 116 haney, Hancock, Aufreiter and Huffman, 1996; Huffman, 1997). Time spent feeding on ter-
23
24 117 mite mound soil is short: at Gombe, 32 bouts of geophagy were measured and the mean
25
26 118 duration was 1.7 min, range 1-8 min (data from Wrangham, 1977; quoted in Uehara,
27
28 119 1982:53). Co-feeding in large groups on termite mound soil, seen for example when feeding
29
30 120 on other soils such as clay, has not been observed. And, unlike clay, termite mound soil is
31
32 121 not eaten with leaves. At Budongo, if termites are present in termite mound soil, they are
33
34 122 also eaten (Newton-Fisher, 1999), but use of tools for termite fishing has not been observed
35
36 123 at Budongo, possibly because termite mounds of *Pseudacanthotermes* are less fishable, hav-
37
38 124 ing few or no external holes (Collins & McGrew, 1985), unlike those of *Macrotermes* species.
39
40 125 At Mahale, use of tools for termite fishing by the M group has only been seen occasionally
41
42 126 (Takahata, 1982); while at Gombe, chimpanzees termite fish year around, though concen-
43
44 127 trate this activity around the wet months (Goodall, 1986; Uehara, 1982). Goodall (1986:256)
45
46 128 also refers to Wrangham’s 1977 study at Gombe: “Analysis of samples of termite clay ... re-
47
48 129 vealed substantial quantities of potassium, magnesium and calcium and traces of copper,
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 130 manganese, zinc, and sodium ... feeding on termite clay may be to neutralise tannins and
4
5 131 other poisons present in plant foods (Hladik, 1977)".
6
7 132 In this paper we explore the concentrations of mineral elements in termite mound soil
8
9 133 across three sites, as compared to control soil samples and other dietary sources. We go on
10
11 134 to provide possible explanations for the mechanisms by which mineral elements are con-
12
13 135 centrated in different soil and plant-based sources.
14
15
16
17 136
18
19 137

21 138 **Methods**

23 139 **Subjects and sites**

24
25
26 140 Data were collected in the Budongo Forest Reserve, in north-western Uganda; and the
27
28 141 Gombe National Park and the Mahale Mountains National Park, both in western Tanzania.
29
30 142 Subjects at each of the three sites sampled were all wild East African chimpanzees (*Pan*
31
32 143 *trogodytes schweinfurthii*). Males and females of all age groups, except infants (aged 0-5
33
34 144 years old) were seen eating at the termite mounds from which samples were collected.
35
36 145 Samples described here were collected between July 2015 and October 2017. Termite spe-
37
38 146 cies are shown in Table 1.
39
40
41
42 147

44 148 **Soil sample collection**

45
46
47 149 Across sites, termite mound soil samples were collected by removing a 10-15g piece of
48
49 150 mound soil from a termite mound, using a sterile knife. Clean gloves were worn to prevent
50
51 151 contamination from human sweat. In addition, control samples were collected of forest soil.
52
53 152 At Budongo, control samples were taken from forest soil 1-3m laterally from the termite
54
55 153 mound and 15-20cm deep. At Gombe control samples were taken from forest soil 1m later-

1
2
3 154 ally from the termite mound and 15-20cm deep. Control samples were not collected at Ma-
4
5 155 hale. All samples were put into individual new plastic bags, marked with date, collector,
6
7 156 block number (an indication of location within the chimpanzee territory), and sample num-
8
9 157 ber, and taken back to base camp where they were dried at a temperature of 40° C until ful-
10
11 158 ly dry. Five grams of each dried sample was then transferred to new sterile plastic container
12
13
14 159 tubes for onward shipment to the UK.
15
16
17 160

19 161 **Laboratory Analysis of Soil Samples**

21 162 The soil samples were dried to constant weight in an oven at 105°C for 6 hours. The total
22
23 163 mass of the dried material was determined. Duplicate samples were prepared by taking 0.1g
24
25 164 of the material and 3ml of Aqua Regia in a 10ml centrifuge tube. The samples were digested
26
27 165 in a water bath at 85°C for 3 hours. 7ml of ultrapure Type 1 water was then added to each
28
29 166 sample and the samples mixed using a vortex mixer. A 1ml aliquot of each sample was dilut-
30
31 167 ed 10 fold with Type 1 water for analysis. The elemental content of each sample was then
32
33 168 determined using a Perkin Elmer Optima 2100 DV Inductively Coupled Plasma Optical Emis-
34
35 169 sion Spectrometer (ICP-OES). Standards and a blank were made up at 2, 4, 6, 8 and 10 ppm
36
37 170 concentrations with 3% HNO₃ and three replicates of each element were measured. Each
38
39 171 sample was analysed in triplicate and the average of the triplicate analysis taken for each
40
41 172 duplicate. The mean of the duplicate analyses of the individual soil samples was then taken
42
43 173 to be representative of that soil sample. The elemental content per kg of dried material was
44
45 174 calculated from the raw data.
46
47
48
49
50

51 175

54 176 **Statistical analyses**

55
56
57
58
59
60

1
2
3 177 The data for each variable were tested for normality of distributions and equality of error
4
5 178 variances. Where these assumptions were not upheld non-parametric tests were used. Re-
6
7 179 sults were considered significant at $\alpha=0.05$. All data were analysed using SPSS v24.
8
9

10 180

11
12 181

13 14 15 182 **Results**

16
17
18 183 Values are mg/kg except where otherwise stated. We found a wide variation in the concen-
19
20 184 tration of the mineral elements measured in termite mound and control soil samples (Table
21
22 185 2). Iron, aluminium, and potassium were the highest in both termite mound soil and control
23
24 186 samples across sites. Zinc, sodium and copper had the lowest concentrations in both soil
25
26 187 types (with the exception of Mahale where zinc was more abundant in termite mound soil,
27
28 188 see Table 2).
29

30
31
32 189

33 34 190 **Budongo**

35
36 191 Potassium, phosphorus, aluminium, and copper were all more concentrated in termite
37
38 192 mound soil than in control soil; no other minerals varied in their abundance between soil
39
40 193 types (Table 2). When compared with mineral concentration in the normal diet (data taken
41
42 194 from Reynolds et al., [2012](#), Table 3), potassium (Kruskal Wallis: $X^2= 0.95$ $p=0.329$) and phos-
43
44 195 phorus (Kruskal Wallis: $X^2= 0.80$ $p=0.373$) are found at similar concentrations in termite
45
46 196 mound soil. Concentrations of all other minerals measured differed. Termite mound soil had
47
48 197 concentrations of iron over 75 times higher (49.1 ± 19.6 g/kg, $n=39$) than found in the nor-
49
50 198 mal diet (649 ± 1309 mg/kg, $n=24$; Kruskal Wallis: $X^2= 44.1$ $p<0.001$); and a very large con-
51
52 199 centration of aluminium (termite mound soil $15,300 \pm 4690$ mg/kg, $n=39$), which is com-
53
54
55
56
57
58
59
60

1
2
3 200 pletely absent from the normal diet (n=24; Kruskal Wallis: $X^2= 46.4$ p<0.001). Of other min-
4
5 201 erals, calcium ($X^2= 9.09$ p=0.003), magnesium ($X^2= 5.13$ p=0.024) and sodium ($X^2= 44.1$
6
7 202 p<0.001) were higher in the normal diet, while manganese ($X^2= 43.9$ p<0.001) and copper
8
9 203 ($X^2= 18.6$ p<0.001) were higher in termite mound soil.
10
11
12 204

14 205 **Gombe**

16 206 As at Budongo, iron had the highest concentrations in both termite mound soil and control
17
18 207 samples from Gombe, followed by aluminium (see Table 2). Levels of magnesium were
19
20 208 higher across Gombe soil samples (n=19) than in Budongo soil samples (n=66; Mann-
21
22 209 Whitney: U=71, p<0.001); with concentrations in termite mound soil over 5 times higher in
23
24 210 Gombe (Table 2; Mann-Whitney: U=22, p<0.001). As at Budongo zinc, sodium and copper
25
26 211 had the lowest concentrations. Sodium was completely absent from termite mound soil at
27
28 212 Gombe, but was present in small amounts in control samples. So, as at Budongo, Gombe
29
30 213 termite mound soil provided the high concentrations of iron and aluminium, together with
31
32 214 some magnesium and other minerals, with the notable exception of sodium. Concentrations
33
34 215 of potassium, iron, aluminium, and copper were all higher in termite mound than in control
35
36 216 soil samples at Gombe; concentrations of sodium and sulphur were lower (Table 2).
37
38
39
40
41
42
43

44 218 **Mahale**

46 219 As at Budongo and Gombe, iron and aluminium were present in the highest concentrations,
47
48 220 although at Mahale aluminium, rather than iron, was highest; at almost double the concen-
49
50 221 trations present in Budongo or Gombe (Table 2; Kruskal-Wallis: $X^2= 25.13$; p<0.001). Also as
51
52 222 at Budongo and Gombe, sodium and copper had the lowest concentrations at Mahale. None
53
54
55
56
57
58
59
60

1
2
3 223 of the three sites compared had a consistently higher or lower overall concentration of min-
4
5 224 erals in any particular soil type.
6

7
8 225

9
10 226 **Comparisons between termite mound soil, clay, decaying wood, and the normal diet of**
11
12 227 **fruit and leaves at Budongo**
13

14 228 We compare the mineral content in termite mound soil with that present in clay (data from
15
16 229 Reynolds et al 2015, Table 3), decaying wood (*Raphia farinifera* and *Cleistopholis patens*)
17
18 230 (data from Reynolds et al., Tables 1 and 2 combined), and the normal diet of fruit and leaves
19
20 231 at Budongo (data from Reynolds et al., 2012, Table 3). The differences between means
21
22 232 shown in Table 3 are significant for all minerals shown.
23
24
25

26 233

27
28 234

29
30 235 **Discussion**
31

32
33 236 Given the distance between the three sites (Budongo to Gombe 740km, Gombe to Mahale
34
35 237 180 km) there is a high degree of similarity in the concentration of soil minerals between
36
37 238 them. Termite mound soil provides chimpanzees with iron (Fig 3) and aluminium (Fig 4) in
38
39 239 high concentrations at all three, with provision of other essential minerals at lower concen-
40
41 240 trations at all of them, and absence or near absence of sodium at them all. Thus, a clear pic-
42
43 241 ture emerges of the contribution of termite mound soil to the mineral intake of chimpan-
44
45 242 zees in East Africa and possibly elsewhere.
46
47
48

49 243

50
51 244 Similar differences between termite mound soil and control samples have been found by
52
53 245 Adams et al. (2017), Mahaney et al. (1996, 1999), Aufreiter et al. (2001), Sarcinelli et al.
54
55 246 (2009). This widespread difference indicates a process whereby some mineral elements be-
56
57
58
59
60

1
2
3 247 come concentrated in termite mound soil. What is the process? It could take place at the
4
5 248 stage of acquisition of soil by termites, which involves a prolonged process of embedding
6
7 249 grains of soil in ingested water and salivary secretions (Turner, 2005) after which they are
8
9
10 250 carried up into the mound to the building point. However, minerals that are relatively scarce
11
12 251 in control forest soil are also relatively scarce in termite mound soil. Sodium in particular,
13
14 252 scarce in forest soil, is very low or absent (i.e. below measurement detection limits) in ter-
15
16 253 mite mound soil (see also Tweheyo et al., 2006). The main process whereby minerals be-
17
18 254 come concentrated in termite mound soil is therefore unlikely to be selection by termites.
19
20

21 255

22
23 256 Low values (or absence) of sodium in termite mound soil were found in the initial samples of
24
25
26 257 termite mound soil collected as part of a study of minerals in clay (n=5; Reynolds et al.,
27
28 258 2015). This finding is now validated by a larger sample size across three different sites. The
29
30 259 complete absence of sodium from termite mound soil at Gombe, while present in control
31
32 260 samples, could indicate avoidance or rejection of sodium by termites or that they consume
33
34 261 sodium for their own requirements. The latter may be the correct explanation. Kaspari et al.
35
36 262 (2009, 2014) showed experimentally that numbers of termites in the soil and litter decom-
37
38 263 position rates were higher in Amazonian forest plots to which sodium had been applied than
39
40 264 in control plots. Whether sodium consumption is a common attribute of termites or can ex-
41
42 265 plain the relative lack of sodium in Gombe termite mound soil is not known (Scheffrahn
43
44 266 pers. comm.).
45
46
47
48

49 267

50
51 268 High values of aluminium and iron and low values of sodium were also found by Mahaney et
52
53 269 al. (1996, 1997, 1999) and Tweheyo et al. (2006) who emphasised the possible medicinal
54
55 270 use of aluminium in clay in the form of metahalloysite. Metahalloysite has the same formula
56
57
58
59
60

1
2
3 271 as kaolinite, $\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$ (Brindley, Robinson and MacEwan, 1946) and is used by humans
4
5 272 (commercially in the form of Kaopectate) to treat gastro-intestinal complaints (Hunter,
6
7 273 1973; Mahaney et al., 1997, 1999; Johns and Duquette, 1991; Wilson, 2003; Fairhead, 2016).
8
9
10 274 Smectite and gibbsite are further possible contributors to the efficacy of termite mound soil
11
12 275 (Wilson, 2003). Higher concentrations of mineral elements in termite mound soil than in
13
14 276 surrounding control soil were found by Aufreiter et al. (2001) and Adams et al. (2017) in a
15
16 277 study of arboreal termitaria in Peru.
17
18

19 278

21 279 **Mineral accretion**

22
23 280 It is of great interest that chimpanzees appear to have discovered these three “hidden”
24
25 281 sources of minerals: plant-based, soil-based, and animal-generated. In two of the three
26
27 282 (plant-based and animal-generated) mineral concentration comes about as a result of water
28
29 283 evaporation. In each case, water containing minerals is drawn up in decaying wood by capil-
30
31 284 lary action, in the case of termite mounds transported by termites. In the third case, clay,
32
33 285 low levels of minerals occur in the forest substrate and these are leached out of the soil by
34
35 286 rain-water that collects in holes under trees.
36
37
38

39 287

40
41
42 288 At *Raphia farinifera* and *Cleistopholis patens* sites, chimpanzees chew the fibrous, decaying
43
44 289 wood containing minerals left behind after evaporation, following which they spit out
45
46 290 ‘wedges’ of fibrous matter. At clay sites it appears that the minerals are ingested by chim-
47
48 291 panzees by chewing the clay when it is in semi-solid form, or extracting it from clay-water
49
50 292 with the use of leaf or moss sponges (Reynolds et al., 2015). At termite mound soil sites,
51
52 293 chimpanzees chew pieces of mound soil in a similar way to the way they chew clay.
53
54
55
56
57
58
59
60

1
2
3 294 In each of the above cases, a low level of minerals exists in the environment, too dispersed
4
5 295 and at concentrations too low for detection and acquisition by large mammals such as
6
7 296 chimpanzees. Concentration of minerals may come about in three ways:

9
10 297 (a) In the case of decaying *Raphia farinifera* palms, and *Cleistopholis patens* trees, these
11
12 298 are located in swamp forest which periodically floods, bringing in river water which
13
14 299 contains low levels of mineral elements leached from the soil and rocks along its
15
16 300 course. These elements are in low concentration (Reynolds et al., 2009, 2012, 2015).
17
18 301
19
20 302 swamp water upwards inside the tree's vertical, fibrous, pith-filled trunk. Because
21
22 303 the head of the *Raphia* palm has previously fallen off after the tree fruited, the top
23
24 304 of the trunk is now open and the whole trunk forms a cylinder filled with fibrous
25
26 305 pith. Water containing low levels of minerals can enter this cylinder from below and
27
28 306 rises up the fibres. As water evaporates from the top of the cylinder, it will leave its
29
30 307 mineral content behind. As a result we speculate that this becomes concentrated,
31
32 308 and it is this source that the chimpanzees have learned to access by making a hole in
33
34 309 the bark of the lower trunk (see Reynolds et al., 2009). In the case of *Cleistopholis*
35
36 310 *patens*, we believe minerals become concentrated in a similar way but without the
37
38 311 cylindrical process, merely by the adsorption by the decaying tree of mineral-
39
40 312 containing water, which evaporates upwards from the tree, leaving behind concen-
41
42 313 trated minerals, which are then accessed by chimpanzees chewing the decaying
43
44 314 wood.

45
46 315 (b) In the case, of clay, we don't believe evaporation plays a part. The action of rain wa-
47
48 316 ter and/or river water on forest soil, especially in hollows under trees, leads to disso-
49
50 317 lution and/or dispersion of minerals from the clay material which contains a high
51
52
53
54
55
56
57
58
59
60

1
2
3 318 level of aluminium and surrounding soil which has a high iron content (Eggeling
4 319 1947, Aufreiter 1997).
5
6
7 320 (c) In the case of termite mound soil, the actions of the termites themselves serve to concen-
8
9 321 trate the mineral elements in surrounding soil. The mechanisms by which this happens are
10
11 322 not clear and require further study. Studies by Sieber (1982) and Hesse (1955) focus on the
12
13 323 use of water by termites in processing surrounding soil before carrying it to the surface of
14
15 324 the mound. Turner (2005, 2011) describes, with associated videos, the process of drinking
16
17 325 and carrying soil by termites. In the case of forest termites, a further process may be im-
18
19 326 portant: the ingestion of organic matter in forest soil, thus having the incidental effect of in-
20
21 327 creasing the proportion of the mineral component. Further work is needed to elucidate the
22
23 328 causes of the differences between forest soil and termite mound soil.
24
25
26
27 329

330 **Summary and conclusions**

331 Termite mound soil provides the highest concentrations of aluminium and iron found in any
332 of the dietary items at the sites studied here. The normal diet of chimpanzees, while high in
333 calcium and moderately high in potassium and magnesium, lacks aluminium and copper and
334 is low in other essential minerals. Sodium, low in the normal diet, is absent or in low con-
335 centration in termite mound soil, which is thus not a dietary source of sodium for chimpan-
336 zees. This absence is in stark contrast to the high concentration of sodium found in decaying
337 wood, which is eaten (Fig 5, see also Reynolds et al. 2009). Thus, geophagy, meat eating,
338 and insectivory all add to the intake of essential minerals obtained by chimpanzees. In both
339 Budongo and Gombe, control forest soil taken from just a few meters away from the ter-
340 mite mounds contains substantially lower concentrations of potassium, aluminium and cop-
341 per. Thus we can see a concentrating effect in termite mound soil for some minerals, with
342 the notable exception of sodium. Termite mound soil at Mahale shows a similar pattern of

1
2
3 343 minerals to those at Budongo and Gombe, with high levels of iron and aluminium, and mod-
4
5 344 erate levels of potassium and magnesium. We suggest three possible mechanisms by which
6
7 345 minerals become concentrated: evaporation of water in decaying wood, concentration after
8
9 346 transport by termites, and dissolution or dispersion of mineral elements in clay after leach-
10
11 347 ing of soil by water. Chimpanzees have discovered these sources of minerals and now ex-
12
13 348 ploit them sporadically as an adjunct to their normal diet of fruits, leaves and other plant
14
15 349 parts.
16
17
18
19
20

21 351 **References**

- 22
23
24 352 Abrahams, P.W. (1997). Geophagy (soil consumption) and iron supplementation in Uganda.
25
26 353 *Tropical Medicine and International Health*, 2, 617-623.
27
28 354 Abrahams, P.W., & Parsons, J.A. (1997). Geophagy in the tropics: a literature review. *Journal*
29
30 355 *of Geography*, 162, 63-72.
31
32
33 356 Abrahams, P.W. (1999). The chemistry and mineralogy of three savanna lick soils. *Journal of*
34
35 357 *Chemical Ecology*, 25, 2215-2228.
36
37
38 358 Adams, D.B., Rehg, J.A., & Watsa, M. (2017). Observations of termitarium geophagy by
39
40 359 Rylands' bald-faced saki monkeys (*Pithecia rylandsi*) in Madre de Dios, Peru. *Primates*, 58(3),
41
42 360 449-459.
43
44
45 361 Aufreiter, S., Hancock, R.G.V., Mahaney, W.C., Strambolic-Robb, A., & Sanmagudas, K.
46
47 362 (1997). Geochemistry and mineralogy of soils eaten by humans. *International Journal of*
48
49 363 *Food Science and Nutrition*, 48, 293-305.
50
51
52 364 Aufreiter, S., Mahaney, W.C., Milner, M.W., Huffman, M.A., Hancock, R.G.V., Wink, M., &
53
54 365 Reich, M. (2001). Mineralogical and chemical interactions of soils eaten by chimpanzees of
55
56
57
58
59
60

1
2
3 366 the Mahale Mountains and Gombe Stream National Parks, Tanzania. *Journal of Chemical*
4
5 367 *Ecology*, 27, 285-311.
6
7 368 Bolton, K.A., Campbell, V.M., & Burton, F.D. (1998). Chemical analysis of soils of Kowloon
8
9 369 (Hong Kong) eaten by hybrid macaques. *Journal of Chemical Ecology*, 24, 195-205.
10
11
12 370 Brindley, G.W., Robinson, K., & MacEwan, D.M.C. (1946). The clay minerals halloysite and
13
14 371 meta-halloysite. *Nature*, 157, 225-226.
15
16 372 Collins, D.A., & McGrew W.C. (1985). Chimpanzees' (*Pan troglodytes*) choice of prey among
17
18 373 termites (Macrotermitinae) in Western Tanzania. *Primates*, 26 (4), 375-389.
19
20
21 374 Deblauwe, I., & Janssens, G.P.J. (2008). New insights in insect prey choice by chimpanzees
22
23 375 and gorillas in Southeast Cameroon: the role of nutritional value. *American Journal of Physi-*
24
25 376 *cal Anthropology*, 135, 42-55. Doi: 10.1002/ajpa.20703.
26
27
28 377 Eggeling W.J. (1947). Observations on the ecology of the Budongo Rain Forest, Uganda. *J.*
29
30 378 *Ecol.*, 34 (1): 20-87.
31
32 379 Fairhead, J.R. (2016). Termites, Mud Daubers and their Earths: A Multispecies Approach to
33
34 380 Fertility and Power in West Africa. *Conservation and Society*, 14, 359-367.
35
36 381 Geissler, P.W., Mwaniki, D.L., Thiong'o, F., Michaelsen, K.F., & Friis, H. (1998). Geophagy,
37
38 382 iron status and anaemia among primary school children in Western Kenya. *Tropical Medi-*
39
40 383 *cine and International Health*, 3, 529-534.
41
42
43 384 Goodall, J. (1986). *The Chimpanzees of Gombe: patterns of behaviour*. Cambridge, MA: Har-
44
45 385 vard University Press.
46
47
48 386 Hesse, P.R. (1955). A chemical and physical study of the soils of termite mounds in East Afri-
49
50 387 ca. *The Journal of Ecology*, 449-461.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 388 Hladik, C.M. (1977). Chimpanzees of Gombe and the chimpanzees of Gabon: some compara-
4
5 389 tive data on the diet. In: T.H. Clutton-Brock (Ed.) *Primate Ecology* (pp.481-501). New York:
6
7 390 Academic Press.
- 9
10 391 Hunter, J.M. (1973). Geophagy in Africa and in the United States: a culture-nutrition hy-
11
12 392 pothesis. *Geography Review*, 63, 170-195.
- 14
15 393 Huffman, M.A. (1997). Current evidence for self-medication in primates: A multidisciplinary
16
17 394 perspective. *Yearbook of Physical Anthropology*, 40, 171-200.
- 18
19 395 Johns, T. (1986). Detoxification function of geophagy and domestication of the potato. *Jour-*
20
21 396 *nal of Chemical Ecology*, 12, 635-646.
- 23
24 397 Johns, T., & Duquette, M. (1991). Detoxification and mineral supplementation as functions
25
26 398 of geophagy. *American Journal of Clinical Nutrition*, 53, 448-456. PMID: 1989412
- 28
29 399 Kaspari, M., Yanoviak, S.P., Dudley, R., Yuan, M., & Clay, N.A. (2009). Sodium shortage as a
30
31 400 constraint on the carbon cycle in an inland tropical rainforest. *Proceedings of the National*
32
33 401 *Academy of Sciences*, 106(46), 19405-19409. DOI: 10.1073/pnas.0906448106
- 35
36 402 Kaspari, M., Clay, N.A., Donoso, D.A., & Yanoviak, S.P. (2014). Sodium fertilization increases
37
38 403 termites and enhances decomposition in an Amazonian forest. *Ecology*, 95(4), 795-800.
- 39
40 404 Ketch, L.A., Malloch, D., Mahaney, W.C., & Huffman, M.A. (2001). Comparative microbial
41
42 405 analysis and clay mineralogy of soils eaten by chimpanzees (*Pan troglodytes schweinfurthii*)
43
44 406 in Tanzania. *Soil Biology and Biochemistry*, 33(2), 199-203.
- 46
47 407 Klaus, G., Klaus-Hugi, C., & Schmid, B. (1998). Geophagy by large mammals at natural licks in
48
49 408 the rain forest of Dzanga National Park, Central African Republic. *Journal of Tropical Ecology*,
50
51 409 14, 829-839.
- 53
54 410 Mahaney, W.C. (1993). Scanning electron microscopy of earth mined and eaten by moun-
55
56 411 tain gorillas in the Virunga Mountains, Rwanda. *Primates*, 34, 311-319.
- 57
58
59
60

- 1
2
3 412 Mahaney, W.C., Stambolic, A., Knesevich, M., Hancock, R.G.V., Aufreiter, S., Sanmugadis, K.,
4
5 413 Kessler, M.J., & Grynepas, M.D. (1995). Geophagy among rhesus macaques on Cayo Santiago,
6
7 414 Puerto Rico. *Primates*, 36, 323-333.
8
9
10 415 Mahaney, W.C., Hancock, R.G.V., Aufreiter, S., & Huffman, M.A. (1996). Geochemistry and
11
12 416 clay mineralogy of termite mound soil and the role of geophagy in chimpanzees of the Ma-
13
14 417 hale Mountains, Tanzania. *Primates*, 37, 121-134.
15
16
17 418 Mahaney, W.C., Milner, M.W., Sanmugadas, K., Hancock, R.G.V., Aufreiter, S., Wrangham,
18
19 419 R.W., & Pier, H.W. (1997). Analysis of geophagy soils in Kibale Forest, Uganda. *Primates*, 38,
20
21 420 159-176.
22
23
24 421 Mahaney, W.C., Zippin, J., Milner, M., Sanmugadas, K., Hancock, R.G.V., Aufreiter, S., Camp-
25
26 422 bell, S., Huffman, M.A., Wink, M., Malloch, D., & Kalm, V. (1999). Chemistry, mineralogy and
27
28 423 microbiology of termite mound soil eaten by the chimpanzees of the Mahale Mountains,
29
30 424 Western Tanzania. *Journal of Tropical Ecology*, 15, 565-588.
31
32
33 425 Mahaney, W.C., Milner, M.W., Aufreiter, S., Hancock, R.G.V., Wrangham, R.W. & Campbell,
34
35 426 S. (2005). Soils consumed by chimpanzees of the Kanyawara community in the Kibale Forest,
36
37 427 Uganda. *International Journal of Primatology*, 26, 1375-1398.
38
39
40 428 McKie, A.T., Barrow, D., Latunde-Dada, G.O., Rolfs, A., Sager, G., Mudaly, E., Mudaly, M.,
41
42 429 Richardson, C., Barlow, D., Bomford, A., Peters, T.J., Raja, K.B., Shirali, S., Hediger, M.A., Far-
43
44 430 zaneh, F., & Simpson, R.J. (2001). An iron-regulated ferric reductase associated with the ab-
45
46 431 sorption of dietary iron. *Science*, 291(5509), 1755-9.
47
48
49 432 Mitani, J.C., & Watts, D.P. (2001). Why do chimpanzees hunt and share meat? *Animal Be-*
50
51 433 *haviour*, 61, 915-924.
52
53
54
55
56
57
58
59
60

- 1
2
3 434 Newton-Fisher, N.E. (1999). Termite eating and food sharing by male chimpanzees in the
4
5 435 Budongo Forest, Uganda. *African Journal of Ecology*, 37, 369-371. DOI: 10.1046/j.1365-
6
7 436 2028.1999.00187.x
8
9
10 437 Newton-Fisher, N.E., Notman, H., & Reynolds, V. (2002). Hunting of mammalian prey by Bu-
11
12 438 dongo Forest chimpanzees. *Folia Primatologica*, 73(5), 281-3.
13
14
15 439 Nishida, T., Uehara, S., & Nyundo, R. (1979). Predatory behaviour among wild chimpanzees
16
17 440 of the Mahale Mountains. *Primates*, 20, 1-20.
18
19
20 441 O'Malley, R.C., & Power, M.L. (2014). The energetic and nutritional yields from insectivory
21
22 442 for Kasekela chimpanzees. *Journal of Human Evolution*, 71, 46-58. DOI:
23
24 443 10.1016/j.jhevol.2013.09.014
25
26
27 444 Pebsworth, P.A., Bardi, M., & Huffman, M.A. (2011). Geophagy in chacma baboons: patterns
28
29 445 of soil consumption by age class, sex, and reproductive state. *American Journal of Primatol-*
30
31 446 *ogy*, 74, 48-57.
32
33
34 447 Reynolds, V., Plumptre, A.J., Greenham, J., & Harborne, J. (1998). Condensed tannins and
35
36 448 sugars in the diet of chimpanzees (*Pan troglodytes schweinfurthii*) in the Budongo Forest,
37
38 449 Uganda. *Oecologia*, 115, 331-336.
39
40
41 450 Reynolds, V., Lloyd, A. W., Babweteera, F., & English, C.J. (2009). Decaying *Raphia farinifera*
42
43 451 palm trees provide a source of sodium for wild chimpanzees in the Budongo Forest, Uganda.
44
45 452 *PLoS ONE*, 4(7), e6194.
46
47
48 453 Reynolds, V., Lloyd, A.W., & English, C.J. (2012). Adaptation by Budongo Forest chimpanzees
49
50 454 (*Pan troglodytes schweinfurthii*) to loss of a primary source of dietary sodium. *African Pri-*
51
52 455 *mates*, 7(2), 156-162.
53
54
55
56
57
58
59
60

1
2
3 456 Reynolds, V., Lloyd, A.W., English, C.J., Lyons, P., Dodd, H., Hobaiter, C., Newton-Fisher, N.,
4
5 457 Mullins, C., Lamon, N., Schel, A-M., & Fallon, B. (2015). Mineral acquisition from clay by Bu-
6
7 458 dongo Forest chimpanzees. *PLoS ONE*, 10(7), e0134075. DOI: 10.1371
8
9
10 459 Sarcinelli, T.S., Schaefer, C.E.G., de Souza Lynch, L., Arato, H.D., Viana, J.H.M., de Alburquerque
11
12 460 Filho, M.R., & Gonçalves, T.T. (2009). Chemical, physical and micromorphological prop-
13
14 461 erties of termite mounds and adjacent soils along a toposequence in Zona da Mata, Minas
15
16 462 Gerais State, Brazil. *Catena*, 76(2), 107-113.
17
18
19 463 Takahata, Y. (1982). Termite fishing observed in the M group chimpanzees. Mahale Moun-
20
21 464 tains Chimpanzee Research Project, *Ecological Report No. 18*, 1-2.
22
23 465 Turner, J.S. (2005). Extended physiology of an insect-built structure. *American Entomologist*,
24
25 466 51(1), 36-38.
26
27
28 467 Turner, J.S. (2011). <http://www.esf.edu/efb/turner/media/termiteVideos.html>. See in par-
29
30 468 ticular videos entitled "Initiation of building", "Termite drinking", "A brief look at termite
31
32 469 repair".
33
34
35 470 Turner, J.S. (2001). On the mound of *Macrotermes michaelseni* as an organ of respiratory
36
37 471 gas exchange. *Physiological and Biochemical Zoology*, 74(6), 798-822.
38
39
40 472 Turner, J.S. (2000). Architecture and morphogenesis in the mound of *Macrotermes*
41
42 473 *michaelseni* (Sjöstedt) (Isoptera: Termitidae, Macrotermitinae) in northern Namibia.
43
44 474 *Cimbebasia*, 16, 143-175.
45
46
47 475 Tweheyo, M., Reynolds, V., Huffman, M.A., Pebsworth, P., Goto, S., Mahaney, W.C., Milner,
48
49 476 M.W., Waddell, A., Dirzowsky, R., & Hancock, R.G.V. (2006). Geophagy in chimpanzees (*Pan*
50
51 477 *trogodytes schweinfurthii*) of the Budongo forest reserve, Uganda. In *Primates of Western*
52
53 478 *Uganda* (pp. 135-152). Springer, New York, NY.
54
55
56 479 Uehara, S. (1982). Seasonal changes in the techniques employed by wild chimpanzees in the
57
58
59
60

1
2
3 480 Mahale Mountains, Tanzania, to feed on termites (*Pseudacanthotermes spiniger*). *Folia pri-*
4
5 481 *matol* 37, 44-76. DOI: 10.1159/000156020
6
7 482 Vermeer, D.E., & Ferrell, R.E. (1985). Nigerian geophagical clay: a traditional antidiarrheal
8
9 483 pharmaceutical. *Science*, 227, 634-636.
10
11
12 484 Wilson, M.J. (2003). Clay mineralogical and related characteristics of geophagic materials.
13
14 485 *Journal of Chemical Ecology*, 29(7), 1525-1547.
15
16
17 486 Wrangham, R.W. (1977). Feeding behaviour of chimpanzees in Gombe National Park, Tan-
18
19 487 zania. In T. Clutton-Brock (Ed.), *Primate ecology: Studies of feeding and ranging behaviour in*
20
21 488 *lemurs, monkeys and apes* (pp. 503-538). Academic Press, London.

22
23 489

24
25 490

26 27 28 491 **Acknowledgments**

29
30 492 At the Budongo Conservation Field Station we are indebted to Aliyo Jacob and Chandiya
31
32 493 Bosco, Geresomu Muhumuza, Monday Gideon, Geoffrey Muhanguzi, Caroline Asimwe, and
33
34 494 P. Nyeko for termite identification. For research permission and export permits in Uganda
35
36 495 we are grateful to the Uganda National Council for Science and Technology, the National
37
38 496 Forest Authority, and the Uganda Wildlife Authority. For securing the necessary permission
39
40 497 to import samples to UK for analysis we are grateful to Raymond Ward of the University of
41
42 498 Brighton. For help at Gombe National Park we are indebted to Anthony Collins, Deus Mjun-
43
44 499 gu, Nuhu Japhary Buke, Matendo Msafiri Katoto, and to Rudolf Scheffrahn, University of
45
46 500 Florida, USA, for termite identification. At Mahale Mountains National Park we are grateful
47
48 501 to Michio Nakamura, Kazuhiko Hosaka and Masaki Shimada. For research permission in
49
50 502 Gombe National Park and Mahale Mountains National Park we are indebted to the Tanzania
51
52 503 National Parks Service, the Tanzania Wildlife Research Institute, and the Tanzania Commis-
53
54
55
56
57
58
59
60

1
2
3 504 sion for Science and Technology. For logistical assistance we acknowledge The Jane Goodall
4
5 505 Institute at Gombe and the Mahale Wildlife Conservation Society at Mahale. We thank
6
7 506 Anne-Marijke Schel for permission to use the video material in Supporting Information, and
8
9 507 J.P.E.C. Darlington, Derek Pomeroy, S. Sutherland, and Andrew Smith for their comments
10
11 508 and advice. For financial support we acknowledge the Mohamed bin Zayed Species Conser-
12
13 509 vation Fund, the Royal Zoological Society of Scotland, the Leverhulme Trust, and the Boise
14
15 510 Fund.
16
17
18
19
20

21 512 **Supporting information**

22
23
24 513 **Video 1. Termite mound soil consumption.** Young adult male (Zig) in the Budongo Forest
25
26 514 Reserve, feeding on soil from a *Pseudacanthotermes spiniger* termite mound in 2011 (video
27
28 515 Anne-Marijke Schel, # 08-29-2011_123144.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 516 **Tables and Figures**
4

5
6 517 **Table 1. Termite species and sampling periods across sites.** TMS = termite mound soil, CTRL
7 518 = control soil. VR = V Reynolds, APG = A Pascual-Garrido, KH = K Hosaka, MS = M Shimada.
8

9 519

Site	Date(s) collected	Samples (N)	Termite species	Collectors
Budongo	July 2015 – Oct 2017	39 TMS, 27 CTRL	<i>Pseudacanthotermes spiniger</i> and <i>Cubitermes ugandensis</i>	VR
Gombe	Dec 2015	12 TMS, 7 CTRL	<i>Macrotermes bellicosus</i> , <i>Macrotermes michaelsoni</i> and <i>Macrotermes subhyalinus</i>	APG
Mahale	Aug – Sept 2015	11 TMS, 0 CTRL	Likely <i>Pseudacanthotermes spp.</i>	KH MS

23 520

24
25 521

26
27 522
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

523 **Table 2. Mineral element concentration in termite mound and control soil across sites.** All
 524 mineral concentrations reported in mean mg/kg \pm standard deviations; Significant differ-
 525 ences between termite mound and control soil are indicated in **bold**. Element key:
 526 Al=aluminium, Ca=calcium, Cu=copper, Fe=iron, K=potassium, Mg=magnesium,
 527 Mn=manganese, Na=sodium, P=phosphorus, S=sulphur, Zn=zinc

Mineral element	Budongo			Gombe			Mahale
	TMS (n=39)	CTRL (n=27)	Kruskal-Wallis	TMS (n=12)	CTRL (n=7)	Kruskal-Wallis	TMS (n=11)
Na	5 ± 15	14 ± 27	$X^2 = 1.43$; $p = 0.232$	0	47.1 ± 8	$X^2 = 16.84$; $p < 0.0001$	41.9 ± 43
K	1080 ± 395	685 ± 90	$X^2 = 25.5$; $p < 0.001$	1980 ± 724	1197 ± 291	$X^2 = 7.78$; $p = 0.005$	5140 ± 2659
S	237 ± 171	169 ± 188	$X^2 = 2.94$; $p = 0.86$	119 ± 50	339 ± 27	$X^2 = 12.60$; $p < 0.0001$	279 ± 133
P	694 ± 219	524 ± 109	$X^2 = 9.92$; $p = 0.002$	422 ± 115	329 ± 35	$X^2 = 2.86$; $p = 0.091$	264 ± 123
Ca	3270 ± 3179	2310 ± 1463	$X^2 = 0.83$; $p = 0.361$	1030 ± 939	466 ± 257	$X^2 = 3.46$; $p = 0.063$	1720 ± 648
Fe	49100 ± 19576	43657 ± 15489	$X^2 = 0.80$; $p = 0.372$	44500 ± 6380	28200 ± 4728	$X^2 = 12.00$; $p = 0.001$	32100 ± 3235
Zn	4.06 ± 15	0	$X^2 = 3.34$; $p = 0.068$	0	0	N/A	455 ± 293
Mn	1050 ± 421	1130 ± 418	$X^2 = 0.46$; $p = 0.498$	383 ± 244	357 ± 119	$X^2 = 0.00$; $p = 1.00$	585 ± 242
Al	18100 ± 4690	15300 ± 4182	$X^2 = 5.36$; $p = 0.021$	19400 ± 5428	11700 ± 2327	$X^2 = 7.76$; $p = 0.005$	32600 ± 8016
Cu	20.86 ± 27	1.41 ± 4.5	$X^2 = 12.62$; $p < 0.0001$	92.3 ± 62	18.8 ± 29	$X^2 = 7.39$; $p = 0.007$	10.2 ± 12
Mg	670 ± 294	604 ± 125	$X^2 = 0.12$; $p = 0.912$	3520 ± 2996	1600 ± 775	$X^2 = 2.06$; $p = 0.151$	5210 ± 2751

529

530

531

532

533

534

535

536

537

538

539 **Table 3. Mean quantities of minerals in termite mound soil, decaying wood, clay, and**
 540 **normal fruit + leaf diet (mg/kg) in Budongo samples.** All mineral concentrations reported in
 541 mean mg/kg \pm standard deviations; Significant differences between termite mound and
 542 other sources are indicated in **bold**. Element key: Al=aluminium, Ca=calcium, Cu=copper,
 543 Fe=iron, K=potassium, Mg=magnesium, Mn=manganese, Na=sodium, P=phosphorus. ¹Data
 544 taken from Reynolds et al., 2015; ²Data taken from Reynolds et al., 2012.

Mineral element	Termite mound soil (n=39)	Clay soil ¹ (n=10)	Decaying wood ^{1,2} (n=31)	Normal diet ² (n=24)	Kruskal-Wallis
Na	5 ± 15	234 ± 228	3032 ± 3826	293 ± 507	$X^2 = 84.33; p < 0.0001$
K	1080 ± 395	2528 ± 3613	9478 ± 14282	4074 ± 6485	$X^2 = 37.13; p < 0.0001$
P	694 ± 219	414 ± 534	1049 ± 2107	851 ± 964	$X^2 = 9.36; p < 0.025$
Ca	3270 ± 3179	2381 ± 3003	4221 ± 5675	13315 ± 30648	$X^2 = 17.75; p < 0.0001$
Fe	49100 ± 19576	8720 ± 3080	141 ± 152	649 ± 1310	$X^2 = 82.04; p < 0.0001$
Mn	1050 ± 421	306 ± 252	183 ± 369	66 ± 69	$X^2 = 67.67; p < 0.0001$
Al	18100 ± 4690	7885 ± 5245	0	0	$X^2 = 94.83; p < 0.0001$
Cu	20.9 ± 27	17 ± 13	0	0	$X^2 = 40.36; p < 0.0001$
Mg	670 ± 294	1012 ± 1165	2240 ± 2071	1557 ± 1272	$X^2 = 18.71; p < 0.0001$

546

547

548

549

550

551

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

552 **Figure 1. Termite mound (*Pseudacanthotermes spiniger*) in the Budongo Forest, Uganda.**



553
554
555
556
557

Figure 2. Site where chimpanzee has removed a piece of termite mound soil, Budongo Forest, Uganda.




558
559
560

561

562

563 Fig. 3 Iron content of termite mound soil at Budongo, Gombe, Mahale and in the normal diet of
564 chimpanzees at Budongo



565

566

567

568

569

570

571


572

573

574

575 Fig. 4 Aluminium content of termite mound soil at Budongo, Gombe, Mahale and in the normal diet
576 of chimpanzees at Budongo

577



578

579

580

581

582

583

584

585

586


587

588

589

590

591 Fig. 5 Presence of sodium in four sources of chimpanzee diet – Budongo data



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- Termite mound soil provides the highest concentrations of aluminium and iron found in any of the dietary items at the sites studied hitherto.
- We describe concentrating mechanisms in termite mound soil for some minerals.
- Chimpanzees have discovered these sources of minerals and how to exploit them.