

Possible Role of Bacterial Siderophores in Inflammation

Iron Bound to the *Pseudomonas* Siderophore Pyochelin Can Function as a Hydroxyl Radical Catalyst

Thomas J. Coffman,** Charles D. Cox,[‡] Brian L. Edeker,[‡] and Bradley E. Britigan**

*Department of Internal Medicine and Research Service, Veterans Administration Medical Center, Iowa City, Iowa 52246; and Departments of [‡]Internal Medicine and [§]Microbiology, University of Iowa College of Medicine, Iowa City, Iowa 52242

Abstract

Tissue injury has been linked to neutrophil associated hydroxyl radical ($\cdot\text{OH}$) generation, a process that requires an exogenous transition metal catalyst such as iron. In vivo most iron is bound in a noncatalytic form. To obtain iron required for growth, many bacteria secrete iron chelators (siderophores). Since *Pseudomonas aeruginosa* infections are associated with considerable tissue destruction, we examined whether iron bound to the *Pseudomonas* siderophores pyochelin (PCH) and pyoverdin (PVD) could act as $\cdot\text{OH}$ catalysts. Purified PCH and PVD were iron loaded (Fe-PCH, Fe-PVD) and added to a hypoxanthine/xanthine oxidase superoxide- ($\cdot\text{O}_2^-$) and hydrogen peroxide (H_2O_2)-generating system. Evidence for $\cdot\text{OH}$ generation was then sought using two different spin-trapping agents (5,5 dimethyl-pyrroline-1-oxide or N-t-butyl- α -phenyl-nitrone), as well as the deoxyribose oxidation assay. Regardless of methodology, $\cdot\text{OH}$ generation was detected in the presence of Fe-PCH but not Fe-PVD. Inhibition of the process by catalase and/or SOD suggested $\cdot\text{OH}$ formation with Fe-PCH occurred via the Haber-Weiss reaction. Similar results were obtained when stimulated neutrophils were used as the source of $\cdot\text{O}_2^-$ and H_2O_2 . Addition of Fe-PCH but not Fe-PVD to stimulated neutrophils yielded $\cdot\text{OH}$ as detected by the above assay systems. Since PCH and PVD bind ferric (Fe^{3+}) but not ferrous (Fe^{2+}) iron, $\cdot\text{OH}$ catalysis with Fe-PCH would likely involve $\cdot\text{O}_2^-$ -mediated reduction of Fe^{3+} to Fe^{2+} with subsequent release of "free" Fe^{2+} . This was confirmed by measuring formation of the Fe^{2+} -ferrozine complex after exposure of Fe-PCH, but not Fe-PVD, to enzymatically generated $\cdot\text{O}_2^-$. These data show that Fe-PCH, but not Fe-PVD, is capable of catalyzing generation of $\cdot\text{OH}$. Such a process could represent as yet another mechanism of tissue injury at sites of infection with *P. aeruginosa*. (*J. Clin. Invest.* 1990. 86:1030-1037.) Key words: pyoverdin • neutrophil • cystic fibrosis • lung injury • hydroxyl radical

This work was presented in part at the National and Midwest Sectional Meetings of the American Federation for Clinical Research, 28 April and 10 November 1989 and published in abstract form (*Clin. Res.* 1989 37:426a. [Abstr.] and *Clin. Res.* 1989. 37:908a. [Abstr.]).

Address reprint requests to Dr. Bradley E. Britigan, Department of Internal Medicine, University of Iowa, SW-54, GH, Iowa City, IA 52242.

Received for publication 26 December 1989 and in revised form 30 May 1990.

J. Clin. Invest.




© The American Society for Clinical Investigation, Inc.

0021-9738/90/10/1030/08 \$2.00

Volume 86, October 1990, 1030-1037

Introduction

Neutrophil-derived oxidants have been suggested as important contributors to host injury in a wide array of inflammatory states (1). In vitro, superoxide anion ($\cdot\text{O}_2^-$) and hydrogen peroxide (H_2O_2), generated by the neutrophil "respiratory burst" can react with an exogenous iron catalyst to form hydroxyl radical ($\cdot\text{OH}$) via the Haber-Weiss reaction shown below (2, 3).



Hydroxyl radical is a highly reactive oxidant. Several lines of evidence point to $\cdot\text{OH}$ as an important mediator of acute lung injury and other forms of phagocyte-associated tissue damage (1, 4, 5-7). Although it has been reported that neutrophils have the endogenous capacity for $\cdot\text{OH}$ formation (8-13), the experimental techniques used in these studies have been criticized for a lack of specificity (14). Recent studies using spin trapping and other techniques have strongly suggested that an exogenous transition metal catalyst must be present for neutrophil activation to result in $\cdot\text{OH}$ generation (3, 15-20).

In humans "free" iron is almost nonexistent, present at a level of $\sim 10^{-18}$ M (21). Most iron, whether intra- or extracellular, is bound to proteins or incorporated into other molecules. Iron bound to either of the two principal extracellular iron chelates, transferrin and lactoferrin, is not catalytic for the Haber-Weiss reaction (22-25). Although recent evidence suggests that iron bound to ferritin or hemoglobin may be able to act as a $\cdot\text{OH}$ catalyst (26-28), access of phagocyte-derived $\cdot\text{O}_2^-/\text{H}_2\text{O}_2$ to these intracellular iron complexes is limited by cellular antioxidant systems (1).

Iron is an essential nutrient for microbial growth and metabolism for which invading microorganisms must compete with host iron-binding proteins (29, 30). Host sequestration of iron has been suggested as an important means of defense from bacterial pathogens (21, 31, 32). Under iron-limited conditions many bacteria and fungi secrete low-molecular weight compounds with high iron-binding affinities known as siderophores (29, 30, 33, 34). These siderophores can abstract iron from some host sources, making it available for uptake and utilization by the microorganism (29, 30, 35, 36).

Pseudomonas aeruginosa, a bacterial pathogen associated with severe necrotizing pneumonia in compromised hosts as well as progressive pulmonary deterioration in cystic fibrosis

(37), secretes two siderophores; pyochelin (PCH)¹ and pyoverdin (PVD) (33, 34). Indirect evidence has been obtained that siderophore generation occurs *in vivo* at sites of *P. aeruginosa* infection (38) where neutrophil-derived $\cdot\text{O}_2^-$ and H_2O_2 would also be present. Accordingly, we assessed whether iron bound to PCH or PVD could catalyze $\cdot\text{OH}$ formation via the Haber-Weiss reaction since the generation of $\cdot\text{OH}$ from the interaction of stimulated phagocytes and siderophore-bound iron could contribute to the extensive tissue injury characteristic of pseudomonas infection.

Methods

Reagents. Diethylenetriaminepentaacetic acid (DTPA), SOD, PMA, DMSO, hypoxanthine, bovine serum albumin, dihydrocytochalasin B, zymosan A, catalase, *N*-*t*-butyl- α -phenyl-nitron (PBN), 3-(2-pyridyl)-5,6-bis (4-phenylsulfonic acid)-1,2,4 triazine (ferrozine), 2-deoxyribose, TCA, 2,9-dimethyl-1,10 phenanthroline (neocuproine) and thiobarbituric acid (TBA) were purchased from Sigma Chemical (St. Louis, MO). Xanthine oxidase was purchased from Sigma Chemical Co. or Boehringer Mannheim Biochemicals (Indianapolis, IN). Results were not altered by the source of xanthine oxidase. Ferrous ammonium sulfate, ferric chloride, and H_2O_2 were purchased from Fischer Scientific (Fairlawn, NJ). Zymosan was opsonized (OZ) by incubation in 100% normal pooled human serum (37°C for 30 min) as previously described (3).

Siderophore preparation. Pyochelin (PCH) and pyoverdin (PVD) were purified from *P. aeruginosa* broth culture as previously described (33, 34). Briefly, *P. aeruginosa* strain PA01 (ATCC 15692; American Type Culture Collection, Rockville, MD) was grown to log phase under iron-depleted conditions. PCH secreted into the media was extracted with dichloromethane and 1% acetic acid followed by purification by TLC. PVD was purified from a separate broth culture by a series of filtration/precipitation steps culminating in gel IEF. PCH was suspended in DMSO and PVD suspended in H_2O .

To iron load either PVD or PCH, sufficient FeCl_3 was added at pH 5.0 to achieve 50% saturation based on the known molar binding ratio of Fe/PCH (1:2) and Fe/PVD (1:1). 50% saturated PCH (Fe-PCH) and PVD (Fe-PVD) was chosen for study to eliminate the possible contribution of "free" iron resulting from inadvertent overloading of the molecules.

Neutrophil separation. Neutrophils were separated from venous blood of normal human volunteers using dextran sedimentation and a Ficoll-Hypaque gradient according to the method of Borregaard et al. (39). Neutrophils were then maintained on ice in HBSS without phenol red (University of Iowa Cell Culture Facility, Iowa City, IA) until usage.

Spin trapping. Electron spin resonance (ESR) detection of spin adducts was performed using a spectrometer (model E104 A ESR; Varian Associates, Palo Alto, CA). Desired reaction mixtures (0.5 ml) were prepared in glass tubes and transferred to a quartz ESR flat cell, which was in turn placed in the cavity of the ESR spectrometer. Sequential ESR scans were then obtained at 25°C. Unless otherwise noted ESR spectrometer settings were: microwave power, 20 mW; modulation frequency 100 kHz; modulation amplitude, 1.0 G; and response time, 1 s. Other settings are noted in the figure legends.

Deoxyribose oxidation. Deoxyribose oxidation was performed using a slight modification of the methods of Greenwald et al. (18). Briefly, 100 μM Fe-PVD or 50–200 μM Fe-PCH was added to buffer (H_2O or HBSS) containing deoxyribose (5 mM), and in some cases 5

$\times 10^6$ neutrophils or 200 μM hypoxanthine to a final volume of 1 ml. After the addition of 200 μM H_2O_2 , 100 ng/ml of the neutrophil stimulant PMA, or 0.06 U/ml xanthine oxidase, respectively, to initiate $\cdot\text{O}_2^-/\text{H}_2\text{O}_2$ generation reaction mixtures were incubated 15–30 min at 37°C. Reactions were terminated by addition of 1.0 ml TCA (6%) and 0.5 ml TBA (1% wt/vol in 0.5 M NaOH), after which cells (if present) were pelleted (12,400 g, 5 min). The supernatant was transferred to glass tubes, boiled for 15 min, and A_{532} of each mixture determined using a spectrophotometer (model DU-30; Beckman Instruments, Inc., Palo Alto, CA). 500 Units/ml catalase or 30 U/ml SOD were included in the original reaction mixture in some experiments.

Iron release from siderophores. Ferrozine/ Fe^{2+} complex formation as measured by A_{562} was used to assess $\cdot\text{O}_2^-$ -mediated reduction and subsequent release of siderophore bound Fe^{3+} (40). Mixtures containing 10 mM ferrozine, 200 μM hypoxanthine, 0.06 U/ml xanthine oxidase and siderophore (PVD, 100 μM , PCH, 50–100 μM) were assayed for an increase in A_{562} . 30 U/ml SOD was included in some experiments to confirm that iron release was dependent on the generation of $\cdot\text{O}_2^-$. Results were identical regardless of whether or not neocuproine was included in the reaction mixture to prevent formation of a copper-ferrozine complex. In some experiments 500 U/ml catalase was included to prevent the reoxidation of Fe^{2+} by H_2O_2 . Although this resulted in an increased rate of Fe^{2+} -ferrozine complex formation, it had no effect on total Fe^{2+} release observed.

Results

The reaction of xanthine oxidase with xanthine or hypoxanthine results in the generation of $\cdot\text{O}_2^-$ and H_2O_2 but not $\cdot\text{OH}$. Detection of $\cdot\text{OH}$ after the addition of an iron chelate to a mixture of (hypo)xanthine and xanthine oxidase has been routinely used to assess the capacity of that chelate to catalyze the Haber-Weiss reaction (22–24). Accordingly, 50% iron-saturated preparations of pyochelin (Fe-PCH) or pyoverdin (Fe-PVD) were added to a hypoxanthine/xanthine oxidase system and evidence for $\cdot\text{OH}$ formation sought using a previously described spin-trapping system consisting of 5,5, dimethyl-1-pyrroline-1-oxide (DMPO) and DMSO (3, 19, 20).

In the absence of DMSO, DMPO reacts with $\cdot\text{O}_2^-$ and $\cdot\text{OH}$ to yield 2,2, dimethyl-5-hydroperoxy-1-pyridinyloxy (DMPO/ $\cdot\text{OOH}$) and 2,2 dimethyl-5-hydroxy-1-pyridinyloxy (DMPO/ $\cdot\text{OH}$) (41, 42). However, DMPO/ $\cdot\text{OH}$ may also arise from the decomposition of DMPO/ $\cdot\text{OOH}$, rendering DMPO/ $\cdot\text{OH}$ detection unreliable as evidence for the presence of $\cdot\text{OH}$. DMSO reacts with $\cdot\text{OH}$ to yield methyl radical ($\cdot\text{CH}_3$), which can be spin trapped by DMPO as 2,2,5-trimethyl-1-pyridinyloxy (DMPO/ $\cdot\text{CH}_3$). When the concentration of DMSO exceeds DMPO, as routinely is the case in our system, the presence of $\cdot\text{OH}$ is manifested as DMPO/ $\cdot\text{CH}_3$ (3). Since DMPO/ $\cdot\text{CH}_3$ is not a direct decomposition product of DMPO/ $\cdot\text{OOH}$ its detection provides more specific spin-trap evidence of $\cdot\text{OH}$ generation (43).

Consistent with previous work (20), ESR spectra obtained during the reaction in H_2O of xanthine oxidase and hypoxanthine in the presence of DMPO, DMSO, and DTPA was composed primarily of DMPO/ $\cdot\text{OOH}$ and DMPO/ $\cdot\text{OH}$ (Figs. 1 A and 2 A). The small DMPO/ $\cdot\text{CH}_3$ peaks present were inhibited by the inclusion of SOD but not catalase (not shown), indicating they arose from spin trapping the small amount of $\cdot\text{OH}$ that may arise from DMPO/ $\cdot\text{OOH}$ breakdown (3, 43).

A marked increase in DMPO/ $\cdot\text{CH}_3$ was detected when Fe-PCH (Fig. 1 B), but not Fe-PVD (Fig. 2 B), was added to the hypoxanthine/xanthine oxidase system. The presence of

1. Abbreviations used in this paper: DMPO, 5,5-dimethyl-pyrroline-1-oxide; DTPA, diethylenetriaminepentaacetic acid; ESR, electronic spin resonance; OZ, opsonized zymosan; PBN, *N*-*t*- α -phenyl-nitron; PCH, pyochelin; PVD, pyoverdin; TBA, thiobarbituric acid.




Figure 1. Representative ESR spectra ($n = 3-5$) after the addition of 0.06 U/ml xanthine oxidase to a solution containing: (A) DMSO (140 mM), DMPO (100 mM), DTPA (0.1 mM), hypoxanthine (HX, 0.2 mM); (B) contents of A plus Fe-PCH (0.2 mM); (C) contents of A plus Fe-PCH (0.2 mM) and catalase (500 U/ml); and (D) contents of A plus apopyochelin. Location of high- and low-field peaks corresponding to $\text{DMPO}\cdot\text{CH}_3$, $\text{DMPO}\cdot\text{OH}$, and $\text{DMPO}\cdot\text{OOH}$ are indicated as CH_3 , OH , and OOH , respectively. Receiver gain was 3.2×10^4 and sweeprate 12.5 G/min.

catalase returned the $\text{DMPO}\cdot\text{CH}_3$ peak amplitude to that observed in the absence of Fe-PCH (Fig. 1 C) as would be expected if the $\text{DMPO}\cdot\text{CH}_3$ increase resulted from spin trapping of $\cdot\text{OH}$ formed via the Haber-Weiss reaction. No spectra were observed with the omission of xanthine oxidase (not shown). Substitution of apopyochelin for Fe-PCH (Fig. 1 D) yielded only background $\text{DMPO}\cdot\text{CH}_3$ peak amplitudes. These data suggest that Fe-PCH but not Fe-PVD is capable of catalyzing the Haber-Weiss reaction.

It has been reported that the stability of $\text{DMPO}\cdot\text{CH}_3$ is decreased in the presence of $\cdot\text{O}_2^-$ suggesting that failure to detect $\text{DMPO}\cdot\text{CH}_3$ may not be an absolute indicator of the lack of $\cdot\text{OH}$ formation (19, 44, 45). Recently we developed a new means of spin trapping $\cdot\text{OH}$ using DMSO and the spin trap PBN (46). In the presence of DMSO and PBN, the generation of $\cdot\text{OH}$ in aerated solutions yields a single stable nitroxide species, which we have assigned to $\text{PBN}\cdot\text{OCH}_3$ (46). The use of PBN in place of DMPO offers two advantages when investigating $\cdot\text{OH}$ generation from $\cdot\text{O}_2^-$ and H_2O_2 . First, unlike $\text{DMPO}\cdot\text{CH}_3$, $\text{PBN}\cdot\text{OCH}_3$ appears to be resistant to $\cdot\text{O}_2^-$ -




Figure 2. ESR spectra representative of 4 separate experiments obtained after the addition of xanthine oxidase (0.06 U/ml) to a solution containing: (A) DMSO (140 mM), DMPO (100 mM), DTPA (0.1 mM), and hypoxanthine (HX, 0.2 mM); and (B) contents of A plus Fe-PVD (0.1 mM). Location of high- and low-field peaks corresponding to $\text{DMPO}\cdot\text{CH}_3$, $\text{DMPO}\cdot\text{OH}$, and $\text{DMPO}\cdot\text{OOH}$ are indicated as CH_3 , OH , and OOH , respectively. ESR spectrometer settings were as in Fig. 1.

induced degradation (46). Second, reaction of $\cdot\text{O}_2^-$ with PBN does not yield a stable spin adduct.

Using this PBN/DMSO spin trapping system we reassessed the potential for Fe-PCH and Fe-PVD to act as $\cdot\text{OH}$ catalysts. The results are seen in Fig. 3. Consistent with the DMPO results, $\cdot\text{OH}$ production ($\text{PBN}\cdot\text{OCH}_3$) was detected in solutions containing hypoxanthine, xanthine oxidase, and Fe-PCH (Fig. 3 B) but not Fe-PVD (Fig. 3 D). Catalase inhibited $\text{PBN}\cdot\text{OCH}_3$ spin adduct formation (Fig. 3 C) consistent with Haber-Weiss mediated $\cdot\text{OH}$ formation.

These spin trapping data (Figs. 1-3) provided strong evidence for the ability of Fe-PCH but not Fe-PVD to act as a Haber-Weiss catalyst. However, to further confirm these results we used an alternative $\cdot\text{OH}$ detection system, the deoxyribose oxidation assay. In the presence of $\cdot\text{OH}$, 2-deoxyribose is oxidized to yield a compound which when exposed to thio-barbituric acid and boiled forms a chromogen with an absorbance maximum of 532 nm (A_{532}) (18, 47). The magnitude of




Figure 3. ESR spectra representative of three to five separate experiments obtained after the addition of xanthine oxidase (0.06 U/ml) to solutions containing: (A) DMSO (140 mM), PBN (10 mM), DTPA (0.1 mM), and hypoxanthine (HX, 0.2 mM); (B) contents of A plus Fe-PCH (0.2 mM); (C) contents of A plus Fe-PCH (0.2 mM) and catalase (500 U/ml); and (D) contents of A plus Fe-PVD (0.1 mM). The species detected in B is that of $\text{PBN}\cdot\text{OCH}_3$. Receiver gain was 5×10^4 and sweeprate 10 G/min.

TBA-reactive deoxyribose oxidation products formed correlates with the amount of $\cdot\text{OH}$ generated. Detection of such products appears to be a sensitive and relatively specific assay for the generation of $\cdot\text{OH}$. Analogous to results with the known $\cdot\text{OH}$ catalyst Fe-EDTA, addition of xanthine oxidase to a solution of hypoxanthine, 2-deoxyribose and Fe-PCH resulted in SOD- and catalase-suppressible chromogen formation (Table I). This was not observed with Fe-PVD (Table I) confirming our spin-trapping results. Variation in background chromogen formation observed in reactions containing hypoxanthine, 2-deoxyribose, and xanthine oxidase but no exogenous iron likely resulted from the presence of trace iron contamination of buffer and enzyme preparations (48).

It has been suggested that the reaction of H_2O_2 with some ferric iron chelates may yield a hydroxyl radical-like species in the absence of an exogenous source of $\cdot\text{O}_2^-$ (49–51). To determine whether Fe-PCH was capable of participating in such a reaction, formation of TBA-reactive deoxyribose oxidation products was determined after the addition of $200\ \mu\text{M}$ H_2O_2 to different iron chelates ($50\ \mu\text{M}$)-Fe-PCH, Fe^{3+} -EDTA, and Fe^{2+} -EDTA (Table I). In the absence of exogenous iron no evidence of $\cdot\text{OH}$ was detectable. Fe-PCH and Fe^{3+} -EDTA resulted in generation of a species that oxidized deoxyribose to a similar extent. The magnitude of this deoxyribose oxidation was considerably less than that observed with Fe^{2+} -EDTA.

To further confirm that chromagen formation in the deoxyribose assay resulted from the presence of $\cdot\text{OH}$ we attempted to assess the impact on chromagen formation of various scavengers that have been shown to have differing reaction rates for $\cdot\text{OH}$ vs. other oxidants (52). These studies were confounded by the fact that Fe-PCH was suspended in DMSO,

a potent $\cdot\text{OH}$ scavenger (52). Other solvents were not suitable, either because of volatility or inherent scavenger activity.

At sites of infection human phagocytes, particularly neutrophils, would be the likely source of $\cdot\text{O}_2^-$ and H_2O_2 that could interact with bacterial siderophores to form $\cdot\text{OH}$. In addition to inducing $\cdot\text{O}_2^-$ formation, neutrophil stimulation also results in extracellular release of a variety of enzymes and other compounds from cytoplasmic storage granules (53). Previous studies have demonstrated that the neutrophil granule components lactoferrin and myeloperoxidase diminish $\cdot\text{OH}$ formation by neutrophils supplemented with catalytic iron by chelating iron in a noncatalytic form and scavenging H_2O_2 , respectively (15, 20, 54). Among the enzymes secreted by stimulated neutrophils are a variety of proteases that, while unlikely to impact on $\cdot\text{OH}$ generation directly, could alter the structure and thereby the iron-binding characteristics of either PCH or PVD. Such changes could lead to either loss or gain, of Haber-Weiss catalytic properties. Thus the ability or inability of pseudomonas siderophores to catalyze formation of $\cdot\text{OH}$ by the hypoxanthine/xanthine oxidase system by no means assures similar results when stimulated neutrophils are the source of $\cdot\text{O}_2^-$ and H_2O_2 .

To determine what impact stimulated neutrophils had on the catalytic potential of Fe-PCH or Fe-PVD, spin-trap evidence of $\cdot\text{OH}$ was sought after PMA stimulation of human neutrophils in the presence of Fe-PCH or Fe-PVD. Consistent with earlier work, PMA stimulation of neutrophils in the presence of DMPO, DMSO, and DTPA, but without exogenous iron, yielded only $\cdot\text{O}_2^-$ -derived DMPO spin adducts (Fig. 4 A). Iron supplementation resulted in catalase suppressible DMPO/ $\cdot\text{CH}_3$, (Fig. 4, B and C). ESR spectra of solutions containing PMA stimulated neutrophils, DMPO, DTPA, DMSO, and Fe-PVD yielded no evidence of $\cdot\text{OH}$ (catalase inhibitable DMPO/ $\cdot\text{CH}_3$) formation (Fig. 4 D).

Unfortunately, when Fe-PCH was used as the iron source in buffers necessary to maintain neutrophil viability, nitroxide artifacts similar to those described in other systems (55) were encountered which prevented accurate interpretation of the results (not shown). Fortunately, substitution of PBN for DMPO eliminated the Fe-PCH induced artifacts. When neutrophils were stimulated with PMA in the presence of DTPA, DMSO, Fe-PCH and PBN, a nitroxide species consistent with PBN/ $\cdot\text{OCH}_3$ was detected, indicating $\cdot\text{OH}$ formation (Fig. 5 A). The omission of PMA (Fig. 5 B), or the inclusion of catalase (Fig. 5 C), prevented PBN/ $\cdot\text{OCH}_3$ formation. The substitution of Fe-PVD for Fe-PCH failed to promote PBN/ $\cdot\text{OCH}_3$ formation (Fig. 5 D).

PMA stimulation may lead to preferential secretion of secondary granule contents (56). To optimize exposure of Fe-PVD to primary granule proteases experiments were repeated in which neutrophils which had been pretreated with dihydrocytochalasin B to prevent phagosome closure (57) were stimulated with opsonized zymosan in the presence of Fe-PVD, DMSO, DTPA, and DMPO or PBN. Once again no spin-trap evidence of $\cdot\text{OH}$ formation was detected (not shown).

Similar to the approach using the hypoxanthine/xanthine oxidase system the above experiments were repeated using formation of TBA-reactive deoxyribose oxidation products as an indicator of $\cdot\text{OH}$ generation. Chromogen formation was seen when neutrophils were stimulated with PMA in the presence of Fe-PCH but not Fe-PVD (Table II).

Both PCH and PVD are unable to bind Fe^{2+} (33, 34, 58).

Table I. Ability of Pyochelin and Pyoverdin to Catalyze Hydroxyl Radical Formation by Hypoxanthine/Xanthine Oxidase or Hydrogen Peroxide

Ferripyochelin		Ferripyoverdin	
	A_{532}		A_{532}
HX/XO	0.068	HX/XO	0.000
HX/XO + Fe^{3+} -EDTA	0.390	HX/XO + Fe^{3+} -EDTA	0.600
HX/XO + Fe^{3+} -EDTA + catalase	0.052	HX/XO + Ferripyoverdin	0.005
HX/XO + Fe^{3+} -EDTA + SOD	0.092		
HX/XO + ferripyochelin	0.210		
HX/XO + ferripyochelin + catalase	0.055		
HX/XO + ferripyochelin + SOD	0.075		
HX/XO + apopyochelin	0.072		
H_2O_2	0.022		
H_2O_2 + Fe^{3+} -EDTA	0.174		
H_2O_2 + Fe^{2+} -EDTA	0.694		
H_2O_2 + ferripyochelin	0.165		

Formation of TBA-reactive deoxyribose oxidation products measured as A_{532} representative of 3–10 separate experiments after the addition of various iron chelates to H_2O_2 or the reaction of hypoxanthine (HX) and xanthine oxidase (XO). Background activity with H_2O_2 or HX/XO in the absence of exogenous iron is related to iron contaminants in buffer and/or enzyme preparations.




Figure 4. Representative ESR spectra ($n = 3-5$) of solutions containing: (A) neutrophils (5×10^6 /ml), DMSO (140 mM), DMPO (100 mM), DTPA (0.1 mM), and PMA (0.1 μ g/ml); (B) contents of A plus FeSO_4 (0.1 mM); (C) contents of A plus FeSO_4 (0.1 mM) and catalase (500 U/ml); and (D) contents of A plus Fe-PVD (0.1 mM). Location of high- and low-field peaks corresponding to $\text{DMPO} \cdot \text{CH}_3$, $\text{DMPO} \cdot \text{OH}$, and $\text{DMPO} \cdot \text{OOH}$ are indicated as CH_3 , OH , and OOH , respectively. ESR spectrometer settings were as in Fig. 1.

Therefore, $\cdot\text{O}_2^-$ mediated reduction of Fe^{3+} bound to either PCH or PVD should result in release of free Fe^{2+} that would then be available for oxidation by H_2O_2 . Documentation of such an event would suggest a possible means for interrupting redox cycling of iron bound to the siderophore, either by irreversibly binding the free Fe^{2+} or through the addition of competing Fe^{3+} chelators to limit reassociation of Fe^{3+} with the siderophore.

Ferrozine avidly binds Fe^{2+} forming a complex with a peak absorbance at 562 nm (40). The formation of this complex in a solution containing ferrozine, Fe-PCH or Fe-PVD, and a source of $\cdot\text{O}_2^-$ would reflect the ability of $\cdot\text{O}_2^-$ to reduce, and cause the release of, siderophore bound Fe^{3+} . Addition of xanthine oxidase to a solution of hypoxanthine, ferrozine, and Fe-PCH resulted in a gradual increase in A_{562} (Fig. 6). No increase in A_{562} was seen with Fe-PVD, apopyochelin, or the omission of xanthine oxidase. These data confirm that $\cdot\text{O}_2^-$ can reduce and release iron bound to Fe-PCH but not Fe-PVD.

Discussion

Infection with *Pseudomonas aeruginosa* is associated with local accumulation of leukocytes and leads to both acute and chronic damage to surrounding tissues (37, 59, 60). In a variety of inflammatory conditions, local tissue injury may be related to generation of the highly reactive oxidant $\cdot\text{OH}$ (1, 4-7), formed through the reaction of neutrophil-derived $\cdot\text{O}_2^-$ and




Figure 5. Representative ESR spectra ($n = 5$) of solutions containing: (A) neutrophils (5×10^6 /ml), DMSO (140 mM), PBN (10 mM), DTPA (0.1 mM), and Fe-PCH (0.2 mM); (B) contents of A plus PMA (0.1 μ g/ml); (C) contents of A plus PMA (0.1 μ g/ml) and catalase (500 U/ml); (D) contents of B except Fe-PVD (0.1 mM) was substituted for Fe-PCH. The species observed in B is that of $\text{PBN} \cdot \text{OCH}_3$. ESR spectrometer settings were as in Fig. 2.

H_2O_2 with an iron catalyst (2). Because of its possible implications for *Pseudomonas*-associated tissue damage we investigated whether the *Pseudomonas* siderophores PCH and PVD bind iron in a manner that promotes $\cdot\text{OH}$ formation when $\cdot\text{O}_2^-$ and H_2O_2 are present.

Three separate approaches were used to measure $\cdot\text{OH}$ production, two spin-trapping systems and the deoxyribose oxidation assay. Regardless of methodology, when Fe-PCH was added to an enzymatic $\cdot\text{O}_2^-$ - and H_2O_2 -generating system evidence of $\cdot\text{OH}$ production was detected. Iron complexed to PVD by contrast did not appear to promote $\cdot\text{OH}$ formation. Inhibition of Fe-PCH catalyzed $\cdot\text{OH}$ generation by inclusion of catalase or SOD in the system is consistent with formation of this species by a Haber-Weiss mechanism. Although previously hypothesized (61), to our knowledge this is the first report of the ability of iron bound to a bacterial siderophore to be capable of catalyzing the Haber-Weiss reaction.

Central to the hypothesis that siderophore bound iron can participate in Haber-Weiss catalysis is that the ferric iron can be reduced to ferrous iron by $\cdot\text{O}_2^-$ with subsequent reoxidation to the ferric state by H_2O_2 . Consistent with previous work suggesting that PCH could not bind Fe^{2+} (33, 34, 58, 62) inclusion of ferrozine and Fe-PCH in an enzymatic $\cdot\text{O}_2^-$ -gener-

Table II. Ability of Pyochelin and Pyoverdin to Catalyze Hydroxyl Radical Formation by Stimulated Neutrophils (PMN)

	A_{532}
PMN + Fe-EDTA	0.532
PMN + Fe-EDTA + catalase	0.018
PMN + Fe-EDTA + SOD	0.008
PMN + ferripyochelin	0.120
PMN + ferripyochelin + catalase	0.001
PMN + ferripyochelin + SOD	0.019
PMN + ferripyoverdin	0.003

Formation of TBA-reactive deoxyribose oxidation products measured as A_{532} representative of seven separate experiments resulting from the stimulation of human neutrophils (PMN) by PMA in the presence of the iron chelates noted.




Figure 6. Increase in A_{562} over time reflecting formation of Fe^{2+} -ferrozine complex after addition of xanthine oxidase to a mixture of hypoxanthine and ferrozine alone (labeled XO) and in the presence of Fe-PCH (Fe-PCH + XO) or Fe-PVD (Fe-PVD + XO). Also shown are results with the addition of SOD to the Fe-PCH/xanthine oxidase mixture (Fe-PCH + XO + SOD) and with a mixture of hypoxanthine, ferrozine, and Fe-PCH to which xanthine oxidase was not added (Fe-PCH). Results are representative of three to four experiments.

ating system resulted in formation of the Fe^{2+} -ferrozine complex due to $\cdot O_2^-$ -mediated release of PCH-bound iron. Thus Fe-PCH-induced $\cdot OH$ generation likely involves the interaction of free Fe^{2+} with H_2O_2 to yield $\cdot OH$ and Fe^{3+} which then can either reassociate with the siderophore or be reduced again by $\cdot O_2^-$. Alternatively, we found that the reaction of H_2O_2 directly with Fe-PCH may also yield a species resembling $\cdot OH$. Similar results have been reported with other ferric iron chelates (49–51) but the mechanism of this reaction remains in doubt.

It has been suggested that reaction of H_2O_2 with Fe^{2+} under some circumstances may yield an oxidant species which is not $\cdot OH$ but rather an Fe^{2+} - H_2O_2 complex (ferryl species) (52, 63, 64). We are unable to eliminate the possibility that the species catalyzed by Fe-PCH is an $\cdot OH$ -like species rather than $\cdot OH$ itself. However since each of these oxidants is highly reactive, from a biologic standpoint it may not be a critical point.

In vivo the most important source of $\cdot O_2^-$ and H_2O_2 would be stimulated phagocytes, particularly neutrophils. In addition to $\cdot O_2^-$ and H_2O_2 release, a wide array of enzymes and proteins are also released during neutrophil stimulation (53). Previous reports have found that at least two of the granular components affect the potential for $\cdot OH$ formation in association with the neutrophil respiratory burst. Lactoferrin and myeloperoxidase inhibit $\cdot OH$ formation by iron-supplemented neutrophils by sequestering iron in a noncatalytic form and consuming H_2O_2 , respectively (15, 20, 54). In contrast, it seemed possible that neutrophil proteases through their action on the peptide PVD could alter the potential of iron bound to PVD to participate in the Haber-Weiss reaction.

However, in spite of these theoretical considerations using the same assays employed with the hypoxanthine/xanthine oxidase system, we again found that only Fe-PCH would catalyze $\cdot OH$ production by stimulated PMN. By inference, neu-

trophil proteases do not endow Fe-PVD with catalytic properties and lactoferrin and myeloperoxidase release do not eliminate Fe-PCH-catalyzed $\cdot OH$ generation. An alternative explanation for the apparent lack of $\cdot OH$ formation with the coincubation of Fe-PVD and stimulated neutrophils would be if Fe-PVD inhibited the neutrophil respiratory burst. However, as assessed by either oxygen consumption or ferricytochrome *c* reduction no such inhibition was detected (results not shown).

Although our data clearly suggest that Fe-PCH is capable of catalyzing formation of $\cdot OH$ in the presence of neutrophil or enzymatic sources of $\cdot O_2^-$ and H_2O_2 it remains unclear as to the relevance of such an observation to in vivo conditions. Indirect evidence has been presented that *Pseudomonas* secrete PVD and PCH in vivo (38). However, no data are available as to what the concentration of either siderophore may be at sites of *Pseudomonas* infection. We are currently developing assay systems to quantitate levels of PCH and PVD in biologic fluids (e.g. bronchoalveolar lavage samples). Nevertheless, the concentration of siderophores used in this study were the same or less than those which accumulated routinely, in in vitro broth culture of *P. aeruginosa* (33, 34), providing some evidence of biologic relevance.

Assuming that concentrations of Fe-PCH sufficient to generate biologically relevant amounts of $\cdot OH$ are present at sites commonly involved in pseudomonas infection (e.g., lung) the potential of such $\cdot OH$ to damage local tissue is unclear. $\cdot OH$ is an extremely reactive oxidant. If formed in vivo it would likely diffuse only a few angstroms before encountering an oxidizable biomolecule. Consequently to be involved in injury to host cells, formation of $\cdot OH$ by the above mechanism would likely need to occur in close proximity to host cell membrane. PCH is very lipophilic (33), and we have obtained preliminary evidence that Fe-PCH readily becomes associated with eukaryotic cell membranes (Coffman, T. J., and B. E. Britigan, unpublished). Such targeting of catalytic iron to host membrane could markedly enhance Fe-PCH toxicity. We are currently examining whether the presence of Fe-PCH enhances the susceptibility of relevant eukaryotic cells (e.g., pulmonary epithelial cells) to $\cdot O_2^-/H_2O_2$ -mediated cytotoxicity.

A variety of extracellular secretory products of *P. aeruginosa* have been incriminated in the tissue destruction observed in *Pseudomonas* infection (37, 59, 60). Our finding that the pseudomonas siderophore Fe-PCH catalyzes $\cdot OH$ formation suggests another possible and novel mechanism for inflammatory damage may be present. Some 50 other siderophores produced by an assortment of bacteria and fungi have been identified (29). Ability to form siderophores has been suggested as a virulence factor for some bacterial pathogens other than *P. aeruginosa* (65, 66). Evaluation of other microbial siderophores may result in the identification of as yet other such compounds capable of serving as Haber-Weiss catalysts.

It has previously been reported that *Staphylococcus aureus* grown in vitro so as to enhance its intracellular iron stores is more susceptible to killing by H_2O_2 and human monocytes, but not neutrophils (67–69). These data have been interpreted as evidence for the involvement of $\cdot OH$ catalyzed by bacteria-associated iron in phagocyte microbicidal activity and inflammatory tissue injury. In recent work we have been unable to document formation of $\cdot OH$ using spin trapping techniques following incubation of similarly prepared iron-rich *Staphylococci* with the hypoxanthine/xanthine oxidase sys-

tem, monocytes, or neutrophils (Cohen, S. M., B. E. Britigan, J. S. Chai, T. L. Roeder, and G. M. Rosen, manuscript submitted for publication). The work reported in the present communication may also have somewhat greater relevance to in vivo conditions than that with iron-rich organisms. In general, sites of bacterial infection are felt to be low-iron microenvironments from the standpoint of the microorganism (21, 31, 32, 38). Thus, in vivo, iron-rich organisms would be unlikely to occur while this environment would induce bacterial production and secretion of siderophores such as PCH (29, 30, 33, 34, 36). In addition, with regard to inflammatory tissue injury, catalytic iron associated with bacteria would be expected to result in generation of $\cdot\text{OH}$ in the immediate proximity of the organism. Given the limited diffusibility of $\cdot\text{OH}$ this would make it less likely to damage surrounding tissue. Siderophores on the other hand are freely diffusible, allowing them potentiality to target catalytic iron to host cells.

In summary, we have obtained experimental evidence that iron bound to one of two pseudomonas siderophores, PCH, is capable of in vitro catalysis of the Haber-Weiss reaction. Such a process in vivo could contribute to tissue injury associated with *P. aeruginosa* infection. Further work supportive of such a hypothesis could suggest new means to limit tissue injury associated with *P. aeruginosa* and possibly other bacterial infections.

Acknowledgments

We acknowledge the technical assistance of Angela Christoffersen and thank Naomi Erickson for help in preparation of the manuscript.

This work was supported in part by U. S. Public Health Service grants HL-44275, AI-28412, and AI-13120, the Veterans Administration Research Service, the Cystic Fibrosis Foundation and the Sandoz Foundation for Gerontologic Research. This work was performed during the tenure of Pfizer Scholar Award and Veterans Administration Research Associate career awards to Bradley E. Britigan.

References

- Weiss, S. J. 1986. Oxygen, ischemia and inflammation. *Acta Physiol. Scand. Suppl.* 548:9-37.
- Haber, F., and J. Weiss. 1934. The catalytic decomposition of hydrogen peroxide by iron salts. *Proc. R. Soc. Lond. A. Math Phys. Soc.* 147:332-351.
- Britigan, B. E., G. M. Rosen, Y. Chai, and M. S. Cohen. 1986. Do human neutrophils make hydroxyl radical? Determination of free radicals generated by human neutrophils activated with a soluble or particulate stimulus using electron paramagnetic resonance spectroscopy. *J. Biol. Chem.* 261:4426-4431.
- Ward, P. A., G. O. Till, R. Kunkel, and C. Beauchamp. 1983. Evidence for the role of hydroxyl radical in complement and neutrophil-dependent tissue injury. *J. Clin. Invest.* 72:789-801.
- Kuroda, M., K. Murakami, and Y. Ishikawa. 1987. Role of hydroxyl radicals derived from granulocytes in lung injury induced by phorbol myristate acetate. *Am. Rev. Respir. Dis.* 135:1435-1444.
- Till, G. O., J. R. Hatherill, W. W. Tourtellotte, M. J. Lutz, and P. A. Ward. 1985. Lipid peroxidation and acute lung injury after thermal trauma to skin. *Am. J. Pathol.* 119:376-384.
- Fox, R. B. 1984. Prevention of granulocyte mediated oxidant lung injury in rats by a hydroxyl radical scavenger, dimethylthiourea. *J. Clin. Invest.* 74:1456-1464.
- Tauber, A. I., and B. M. Babior. 1977. Evidence for hydroxyl radical production by human neutrophils. *J. Clin. Invest.* 60:374-379.
- Weiss, S. J., P. K. Rustagi, and A. F. Lebuglio. 1978. Human granulocyte generation of hydroxyl radical. *J. Exp. Med.* 147:316-323.
- Rosen, H., and S. J. Klebanoff. 1979. Hydroxyl radical generation by polymorphonuclear leukocytes measured by electron spin resonance spectroscopy. *J. Clin. Invest.* 64:1725-1729.
- Green, M. R., H. A. Q. Hill, M. J. Okolow-Zubkowska, and A. W. Segal. 1979. The production of hydroxyl and superoxide radicals by stimulated human neutrophils, measurement by EPR spectroscopy. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 100:23-26.
- Repine, J. E., J. W. Eaton, M. W. Anders, J. R. Hoidal, and R. B. Fox. 1979. Generation of hydroxyl radical by enzymes, chemicals, and human phagocytes in vitro. Detection with three anti-inflammatory agent, dimethyl sulfoxide. *J. Clin. Invest.* 64:1642-1651.
- Sagone, A. L., Jr., and R. M. Husney. 1987. Oxidation of salicylates by stimulated granulocytes: evidence that these drugs act as free radical scavengers in biological systems. *J. Immunol.* 138:2177-2183.
- Cohen, M. S., B. E. Britigan, D. J. Hassett, and G. M. Rosen. 1988. Do human neutrophils form hydroxyl radical? Evaluation of an unresolved controversy. *Free Rad. Biol. Med.* 5:81-88.
- Winterbourn, C. C. 1986. Myeloperoxidase is an effective inhibitor of hydroxyl radical production: implications for the oxidative reactions of neutrophils. *J. Clin. Invest.* 78:545-550.
- Thomas, M. J., P. S. Shirley, C. C. Hedrick, and L. R. Dechatalet. 1986. Role of free radical processes in stimulated human polymorphonuclear leukocytes. *Biochemistry.* 25:8042-8048.
- Kaur, H., Z. Fagerheim, M. Grooveld, A. Puppo, and B. Halliwell. 1988. Aromatic hydroxylation of phenylalanine as an assay for hydroxyl radicals: application to activated neutrophils and heme protein leghemoglobin. *Anal. Biochem.* 172:360-367.
- Greenwald, R. A., S. W. Rush, S. A. Mark, and Z. Weitz. 1989. Conversion of superoxide generated by polymorphonuclear leukocytes to hydroxyl radical: a direct spectrophotometric detection system based on degradation of deoxyribose. *Free Rad. Biol. Med.* 6:385-392.
- Pou, S., M. S. Cohen, B. E. Britigan, and G. M. Rosen. 1989. Spin trapping and human neutrophils: limits of detection of hydroxyl radical. *J. Biol. Chem.* 264:11299-12302.
- Britigan, B. E., G. M. Rosen, B. Y. Thompson, Y. Chai, and M. S. Cohen. 1986. Stimulated human neutrophils limit iron-catalyzed hydroxyl radical formation as detected by spin trapping techniques. *J. Biol. Chem.* 261:17026-17032.
- Bullen, J. J., H. J. Rogers, and E. Griffiths. 1978. Role of iron in bacterial infection. *Curr. Top. Microbiol. Immunol.* 80:1-35.
- Aruoma, O. I., and B. Halliwell. 1987. Superoxide-dependant and ascorbate-dependant formation of hydroxyl radicals from hydrogen peroxide in the presence of iron. Are lactoferrin and transferrin promoters of hydroxyl radical generation? *Biochem. J.* 241:273-278.
- Winterbourn, C. C. 1983. Lactoferrin-catalyzed hydroxyl radical production: Additional requirements for a chelating agent. *Biochem. J.* 210:15-19.
- Baldwin, D. A., E. R. Jenny, and P. Aisen. 1984. The effect of human transferrin and milk lactoferrin on hydroxyl radical formation from superoxide and hydrogen peroxide. *J. Biol. Chem.* 259:13391-13394.
- Buettner, G. R. 1987. The reaction of superoxide, formate radical, and hydrated electron with transferrin and its model compound, Fe(III)-ethylenediamine-*N,N'*-bis [2-(2-hydroxyphenyl) acetic acid] as studied by pulse radiolysis. *J. Biol. Chem.* 262:11995-11998.
- Biamond, P., H. G. van Eijk, A. J. G. Swaak, and J. F. Koster. 1984. Iron mobilization from ferritin by superoxide derived from stimulated polymorphonuclear leukocytes. Possible mechanism in inflammation diseases. *J. Clin. Invest.* 73:1576-1579.
- Puppo, A., and B. Halliwell. 1988. Formation of hydroxyl radicals from hydrogen peroxide in the presence of iron. Is hemoglobin a biological Fenton reagent? *Biochem. J.* 249:185-190.
- Sadrzadeh, S. M. H., E. Graf, S. S. Panter, P. E. Hallaway, and J. W. Eaton. 1984. Hemoglobin: a biologic Fenton reagent. *J. Biol. Chem.* 259:11354-11356.
- Neilands, J. B. 1981. Microbial iron compounds. *Annu. Rev. Biochem.* 50:715-31.

30. Raymond, K. N., G. Muller, and B. F. Matzanke. 1984. Complexation of iron by siderophores: A review of their solution and structural chemistry and biological function. *Top. Curr. Chem.* 123:50-101.
31. Kluger, M. J., and B. A. Rothenburg. 1979. Fever and reduced iron: their interaction as a host defense response to bacterial infection. *Science (Wash. DC)*. 203:374-376.
32. Finkelstein, R. A., C. V. Sciortino, and M. A. McIntosh. 1983. Role of iron in microbe-host interactions. *Rev. Inf. Dis.* 5:5759-5777.
33. Cox, C. D., K. L. Rinehart, Jr., M. L. Moore, and C. J. Cook, Jr. 1981. Pyochelin: novel structure of an iron-chelating growth promoter for *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA.* 78:4256-4260.
34. Cox, C. D., and P. Adams. 1985. Siderophore activity of pyoverdinin for *Pseudomonas aeruginosa*. *Infect. Immun.* 48:130-138.
35. Doring, G., M. Pfestorf, K. Botzenhart, and M. A. Abdallah. 1988. Impact of proteases on iron uptake of *Pseudomonas aeruginosa* pyoverdinin from transferrin and lactoferrin. *Infect. Immun.* 56:291-293.
36. Sriyosachati, S., and C. D. Cox. 1986. Siderophore-mediated iron acquisition from transferrin by *Pseudomonas aeruginosa*. *Infect. Immun.* 52:885-891.
37. Fick, R. B., and S. J. Hata. 1989. Pathogenetic mechanisms in lung diseases caused by *Pseudomonas aeruginosa*. *Chest.* 95:2065-2135.
38. Brown, M. R. W., H. Anwar, and P. A. Lambert. 1984. Evidence that mucoid *Pseudomonas aeruginosa* in the cystic fibrosis lung grows under iron restricted conditions. *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Lett.* 21:113-117.
39. Borregaard, N., J. M. Heiple, E. R. Simons, and R. A. Clark. 1983. Subcellular localization of the b cytochrome component of the human microbicidal oxidase: translocation during activation. *J. Cell. Biol.* 97:52-61.
40. Boyer, R. F., and C. J. McCleary. 1987. Superoxide ion as a primary reductant in ascorbate-mediated ferritin iron release. *Free Rad. Biol. Med.* 3:389-395.
41. Finkelstein, E., G. M. Rosen, and E. J. Rauckman. 1980. Spin trapping of superoxide and hydroxyl radical: practical aspects. *Arch. Biochem. Biophys.* 200:1-16.
42. Finkelstein, E., G. M. Rosen, and E. J. Rauckman. 1982. Production of hydroxyl radical by decomposition of superoxide spin trapped adducts. *Mol. Pharmacol.* 21:262-265.
43. Britigan, B. E., M. S. Cohen, and G. M. Rosen. 1987. Detection of the production of oxygen-centered free radicals by human neutrophils using spin trapping techniques: a critical perspective. *J. Leukocyte Biol.* 41:349-362.
44. Samuni, A., C. D. V. Black, C. M. Krishna, H. L. Malech, E. F. Bernstein, and A. Russo. 1988. Hydroxyl radical production by stimulated neutrophils reappraised. *J. Biol. Chem.* 263:13797-14801.
45. Samuni, A., C. M. Krishna, P. Riesz, E. Finkelstein, and A. Russo. 1989. Superoxide reaction with nitroxide spin adducts. *Free Rad. Biol. Med.* 6:141-148.
46. Britigan, B. E., T. J. Coffman, and G. R. Buettner. 1990. Spin trapping evidence for the lack of significant hydroxyl radical production during the respiration burst of human phagocytes using a spin adduct resistant to superoxide mediated destruction. *J. Biol. Chem.* 265:2650-2656.
47. Halliwell, B., and J. M. C. Gutteridge. 1981. Formation of a thiobarbituric acid-reactive substance from deoxyribose in the presence of iron salts. The role of superoxide and hydroxyl radicals. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 128:347-351.
48. Vile, G. F., and C. C. Winterbourn. 1986. High affinity iron binding by xanthine oxidase. *J. Free Rad. Biol. Med.* 2:393-396.
49. Inoue, S., and S. Kawanishi. 1987. Hydroxyl radical production and human DNA damage induced by ferric nitrilotriacetate and hydrogen peroxide. *Cancer Res.* 47:6522-6527.
50. Walling, C., R. E. Partch, and T. Weil. 1975. Kinetics of the decomposition of hydrogen peroxide catalyzed by ferric ethylenediaminetetraacetate complex. *Proc. Natl. Acad. Sci. USA.* 72:140-142.
51. Aruoma, O. I., B. Halliwell, E. Gajewski, and M. Dizdaroglu. 1989. Damage to the bases in DNA induced by hydrogen peroxide and ferric iron chelates. *J. Biol. Chem.* 264:20509-20512.
52. Winterbourn, C. C. 1987. The ability of scavengers to distinguish $\cdot\text{OH}$ production in the iron catalyzed Haber-Weiss reaction: comparison of four assays for $\cdot\text{OH}$. *Free Rad. Biol. Med.* 3:33-39.
53. Henson, P. M., J. E. Henson, C. Fittschen, G. Kamini, D. L. Bratton, and D. W. H. Riches. 1988. Phagocytic cells: degranulation and secretion. In *Inflammation: Basic Principles and Clinical Correlates*. J. I. Gallin, I. M. Goldstein, and R. Snyderman, editors. Raven Press, New York. 363-390.
54. Britigan, B. E., D. J. Hassett, G. M. Rosen, D. R. Hamill, and M. S. Cohen. 1989. Neutrophil degranulation inhibits potential hydroxyl radical formation: differential impact of myeloperoxidase and lactoferrin release on hydroxyl radical production by iron supplemented neutrophils assessed by spin trapping. *Biochem. J.* 264:447-455.
55. Tero-Kubota, S., Y. Ikegami, T. Kurokawa, R. Sasaki, K. Sugioka, and M. Nakano. 1982. Generation of free radicals and initiation of radical reactions in nitron- Fe^{2+} -phosphate buffer systems. *Biochem. Biophys. Res. Commun.* 108:1025-1031.
56. Wang-Iverson, P., K. B. Pryzwansky, J. K. Spitznagel, and M. H. Cooney. 1978. Bactericidal activity of phorbol myristate acetate treated human polymorphonuclear leukocytes. *Infect. Immun.* 22:945-955.
57. Root, R. K., and J. A. Metcalf. 1977. H_2O_2 release from human granulocytes during phagocytosis: Relationship to superoxide anion formation and cellular metabolism of H_2O_2 . Studies with normal and cytochalasin B treated cells. *J. Clin. Invest.* 60:1266-1279.
58. Beier, R. C., and R. D. Stipanovich. 1989. Fast atom bombardment of metal-pyochelin complexes: metastable analysis at constant B/E of zinc-pyochelin. *Biomed. Environ. Mass. Spectr.* 18:185-191.
59. Pier, G. B. 1985. Pulmonary disease associated with *Pseudomonas aeruginosa* in cystic fibrosis: current status of the host bacterium interaction. *J. Infect. Dis.* 151:515-580.
60. Suter, S., O. B. Schaad, L. Roux, U. E. Nydegger, and F. A. Waldvogel. 1984. Granulocyte neutral proteases and *Pseudomonas* elastase as possible causes of airway damage in patients with cystic fibrosis. *J. Infect. Dis.* 149:523-531.
61. Rosen, G. M., and E. Finkelstein. 1985. Use of spin traps in biological systems. *Adv. Free Rad. Biol. Med.* 1:345-375.
62. Cox, C. D. 1980. Iron reductases from *Pseudomonas aeruginosa*. *J. Bacteriol.* 141:199-204.
63. Koppenol, W. H. 1986. The reaction of ferrous EDTA with hydrogen peroxide: Evidence against hydroxyl radical formation. *J. Free Rad. Biol. Med.* 1:281-285.
64. Rush, J. D., and W. H. Koppenol. 1986. Oxidizing intermediates in the reaction of ferrous EDTA with hydrogen peroxide. *J. Biol. Chem.* 261:6730-6733.
65. Williams, P. H., and N. H. Carbonetti. 1986. Iron, siderophores, and the pursuit of virulence: independence of the aerobactin and enterochelin iron uptake systems in *Escherichia coli*. *Infect. Immun.* 51:942-947.
66. Carbonetti, N. H., P. S. H. Boonchais, V. Vaisanen-Rhen, T. K. Korhonen, and P. H. Williams. 1986. Aerobactin-mediated iron uptake by *Escherichia coli* isolates from human extra-intestinal infections. *Infect. Immun.* 51:966-8.
67. Repine, J. E., R. B. Fox, and E. M. Berger. 1981. Hydrogen peroxide kills *Staphylococcus aureus* by reacting with staphylococcal iron to form hydroxyl radical. *J. Biol. Chem.* 256:7094-7096.
68. Repine, J. E., R. B. Fox, E. M. Berger, and R. N. Harada. 1981. Effect of staphylococcal iron content on the killing of *Staphylococcus aureus* by polymorphonuclear leukocytes. *Infect. Immun.* 32:407-410.
69. Hoepelman, I. M., W. A. Bezemer, C. M. J. E. Vandembroucke-Grauls, J. J. M. Marx, and J. Verhoef. 1990. Bacterial iron enhances oxygen radical-mediated killing of *Staphylococcus aureus* by phagocytes. *Infect. Immun.* 58:26-31.