

Postpartum program actions in primary health care: an integrative review

Tatiane Baratieri (<https://orcid.org/0000-0002-0270-6395>)¹

Sonia Natal (<https://orcid.org/0000-0001-6155-4785>)¹

Abstract *Puerperium is a period of significant morbimortality for women, and Primary Health Care (PHC) is important in developing actions to meet women's health needs. This study aimed to systematize the knowledge produced on postpartum care programs actions within PHC at both national and international levels. This is an integrative review of the literature in databases LILACS (Latin American and Caribbean Health Sciences Literature), BDENF (Nursing Database), SciELO (Scientific Electronic Library Online) and PubMed (US National Library of Medicine). Search was performed in the period April-May 2017. Forty-three papers met the selection criteria. Results indicate that PHC has the physical structure to provide puerperae with care, but has a shortage of human and material resources; there is low postpartum consultation coverage and home visits; there is a good evaluation of the incentive for breastfeeding, but focused on the child; international screening of Postpartum Depression through the Edinburgh Postnatal Depression Scale and care shortage for this condition in Brazil. Postpartum care still focuses on care for the newborn and is mostly restricted to the immediate and late puerperium.*

Key words *Primary health care, Postpartum, Women's health*

¹ Universidade Federal de Santa Catarina. R. Eng. Agrônomo Andrei Cristian Ferreira s/n, Trindade. 88040-900 Florianópolis SC Brasil. baratieri.tatiane@gmail.com

Introduction

Although life expectancy is higher for women than for men in most countries, a number of social and health factors combine to have a lower quality of life for women¹. Gender inequality leads to many risks to women's health, including physical and sexual violence, sexually transmitted infections, malaria, chronic obstructive pulmonary disease and maternal morbidity and mortality¹.

Mortality rates during pregnancy and childbirth remain high in developing countries, despite their worldwide decline in recent decades, with 99% of maternal deaths occurring in these countries¹. Avoidable deaths continue to occur in alarming proportions, mainly due to hypertensive diseases, hemorrhages, sepsis/infections and abortion-related complications^{1,2}.

Maternal health is sensitive to Primary Health Care (PHC)^{2,3}. Thus, increasing the quality of this point of care is key to reducing the mortality rates of this population. Deaths and morbidities during pregnancy, childbirth and puerperium are avoidable by implementing integrated and universal access actions through light technologies and primary care^{2,3}.

It should be noted that most maternal deaths are concentrated in the immediate puerperium (1-10 days postpartum), besides being a significant morbidity stage^{4,5} that extends to late puerperium (from the 11th to the 45th day) and remote puerperium (after 45 days)⁶, with studies that indicate persistent conditions for more than 4 years postpartum^{7,8}. PHC is primarily responsible for postpartum women care through integration of technical knowledge and an ability to receive, support and detect early physical and emotional changes, performing prevention, treatment and follow-up on women referring to others services when needed^{9,10}.

Considering the significant morbimortality of women in the puerperal period, and the fact that PHC is the caregiver responsible for solving most health problems, it is important to develop efficient and effective actions to meet the needs of women's health in the puerperium.

To that end, health policies at national and international levels work together and establish guidelines, actions and strategies for puerperal care, especially in primary care, understanding that adequate resources are essential requirements for comprehensive care^{1,6}. Despite these initiatives, there is evidence that postpartum care in PHC requires adjustments, with improvements in physical and material structure, management

and care in health services, professional qualification, women-centered care, overcoming technician care, thus contributing to improved women's health^{7,10-12}.

Knowing the activities developed and results achieved by puerperal care programs is important for the planning and improvement of PHC care for postpartum women's health. Based on the above, we enquire as to the characteristics and results of the studies that address postpartum care in the context of PHC. Thus, this study aimed to systematize knowledge produced on postpartum care in PHC both at national and international levels.

Methods

This is an integrative review of the literature, which is characterized by a method that allows the search, critical evaluation and synthesis of available evidence about the subject investigated, whose final product consists of the current state of knowledge, implementation of interventions and identification of gaps that guide the development of other studies¹³.

The operational stages of the integrative review are identification of the theme and selection of the hypothesis or research question; establishment of inclusion/exclusion criteria of studies/sampling or search in literature; definition of the information to be extracted from the selected studies/categorization of studies; evaluation of studies included in the integrative review; interpretation of results; and presentation of knowledge review/synthesis¹³.

The following guiding question was formulated in the first step: What are the characteristics – type of study, criteria, parameters, indicators - and results of the studies that address postpartum care in PHC?

In the second step, we searched for references through the Virtual Health Library (BVS), with simultaneous search in the main national and international databases, and we selected LILACS (Latin American and Caribbean Health Sciences Literature) and BDENF (Nursing Database). We conducted advanced search in SciELO (Scientific Electronic Library Online) and PubMed (National Library of Medicine of the United States).

We established the following inclusion criteria for the selection of papers: original work, answering the guiding question; published in Portuguese, English or Spanish; published in a scientific journal with at least Qualis B2 for public health; and when Qualis evaluation was unavailable, we veri-

fied the Journals Impact Factors (JCR), including papers with a minimum rating of 1.0. Duplicated studies in one or more databases were excluded.

The search for references occurred between April and May 2017 using the following descriptors indexed in Health Sciences Descriptors (DeCS), all of which were grouped using Boolean expressions AND and OR: *período pós-parto* OR *postpartum period* OR *período pós-parto* OR *saúde materna* OR *maternal health* OR *salud materna* OR *saúde da mulher* OR *women's health* OR *salud de la mujer* OR *saúde materno-infantil* OR *maternal and child health* OR *salud materno infantil* AND *Atenção Primária à Saúde* OR *Primary Health Care* OR *Atención Primaria de Salud* OR *Saúde da Família* OR *Family Health* OR *Salud de la familia*. In PubMed, we used the following *Mesh Terms*: *postpartum period* OR *maternal health* OR *women's health* AND *primary health care* OR *family health*.

The wide range of descriptors used is justified since the postpartum subject is handled in conjunction with others related to women's health, and restriction to descriptor "*postpartum period*" would limit the possibility of finding studies that approach the subject. Given the relevance of some studies, we selected references that considered the inclusion criteria of the study cited by the selected articles.

In order to evaluate the methodological quality of studies found, we used specific tools for each type of study. Evaluative studies were analyzed through by adapting the meta-evaluation criteria established by Stufflebeam¹⁴, excluding studies classified as "weak" and "reasonable", that is, with more than 50% of the evaluation questions answered negatively. The other studies were evaluated using the Critical Appraisal Skills Program (CASP)¹⁵, which consists of specific checklists for each type of study, and does not suggest a scoring system, so the authors chose to use the same classification of Stufflebeam¹⁴, excluding reasonable and poor studies. These studies, except evaluative ones, were also classified according to their level of evidence, according to the criteria of Stetler et al.¹⁶.

For the accomplishment of the third stage (definition of the information to be extracted from the selected studies), the tool elaborated by Ursi¹⁷ was used, which addresses, among other information, the study identification, methodological characteristics, evaluation of methodological rigor, interventions and results found.

The results of the selected studies were systematized according to similarities of data analyzed in each research. Data synthesis and inter-

pretation were based on the results of the critical evaluation of selected studies. A comparison was made with theoretical knowledge, identification of conclusions and implications resulting from the integrative review.

In order to minimize bias, two reviewers performed the search, evaluation and selection of studies, and then a consensus discussion was held on papers to be included in the review.

Results

After completing cross-referencing between the DeCS / Mesh Terms, 1,974 references were found, and at the end of the material selection strategies, 43 papers were shortlisted for analysis and discussion, as shown in Figure 1.

Year of publication of papers ranged from 1995 to 2016, with 69.8% (30) in the last five years, with 20.9% (9) published in 2013. Regarding the study site, 60.5% (26) were performed in the American continent, mainly Brazil, with 51.2% (22). Other studies were held in Europe, 6 (14%), Oceania, 6 (13.9%), Asia, 2 (4.7%), Africa, 2 (4.7%) and one in three countries, Bangladesh and Nepal (Asia) and Malawi (Africa). The main three languages were English, 51.2% (22), followed by Portuguese, 44.2% (19), and 4.7% (2) Spanish.

Most studies were quantitative (76.7%) (33), and 37.2% (16) are evaluative, 20.9% (9) qualitative, 13.9% (6) cross-sectional, 11.6% (5) clinical trial, 7% (3) cohort and the same quantity of quasi-experimental, and one was descriptive. Regarding evaluation of methodological quality, 12.5% (2) of the 16 evaluative papers were rated excellent, 56.25% (9) very good and 31.25% (5) good, with two value judgments^{18,19}. The others were classified by level of evidence, predominantly at level IV, 59.3% (16), followed by level III, 22.2% (6) and level II, 18.5% (5); and regarding CASP, most were classified as very good, 51.8% (14), 25.9% (7) as excellent and 22.2% (6) as good.

The main results of the analyzed papers are shown below, grouped by similarities for a better understanding:

Evaluation of the structure of puerperal care health facilities

Of the evaluative papers, three^{11,20,21} addressed the evaluation of the structure for puerperal care, all performed in Brazil. Puerperal care is mainly provided by doctors and nurses^{11,20}, with problems mainly in fixing and qualifying doctors in




Figure 1. Flowchart of the paper selection process as per PRISMA/2015.

PHC¹¹. In the Santos (SP) study, the introduction of nutritionists was evaluated; this initiative is insufficient, professionals are found in about 65% of facilities for one or two periods of the week²¹.

Regarding the evaluation of the physical and material structure, the studies found that most of the health establishments had a minimum infrastructure for puerperae care, such as single room and availability of materials, vaccines and medicines. However, one of the limitations of care is the availability and maintenance of the functionality of materials^{11,20,21}. The studies also pointed out poor health education actions and inadequate or non-existent physical space for this purpose^{11,20,21}.

Postpartum consultation: demand/supply and reasons for non-adherence

Studies evaluating prenatal care, as recommended by the Pre-Natal and Birth Humanization Program (PHPN) in Brazil, have shown, since implementation to date, that the recommendation to have six or more prenatal consultations, all basic exams, HIV testing, the 2nd dose, booster or immunizing tetanus vaccine and puerperium consultation is poorly complied with^{20,22,23}.

Regarding Brazilian studies that evaluated the indicator of puerperium consultation up to 42 days postpartum, a low achievement was observed, varying from 16.8% to 58%²⁴⁻²⁶. A lack or little flexibility of prior scheduling of puerperal consultation^{11,18} and hosting the First Comprehensive Health Week^{18,21} were identified. Studies indicate that active search for defaulting women is performed in more than half of the studies, but it is not a systematic practice for all puerperae^{11,20,21,25}.

A study carried out in nine poorer regions of Peru showed that 58% of women reported postpartum control within 42 days²⁷. A survey conducted in the UK in 1995 found that 91% of women attended the puerperal consultation six weeks into postpartum²⁸.

International and national literature shows the reasons for not adhering to the puerperal consultation, and they are: professionals' lack of information on postpartum actions^{4,5}; lack of medicines and equipment; lack of professional qualification; cultural barriers⁴; and perception that postpartum care is intended for the child^{4,5,7}. In addition, puerperae educational actions are still traditional, which also discourages their participation²⁹.

In Australia, it has been shown that problems related to the occurrence of postpartum care in an organized and higher quality way is due to the lack of consistent and cohesive guidelines for community care of new mothers and their babies⁵. Two Australian studies^{5,7} and one from the UK²⁸ identified that puerperae value and are open to primary care^{5,7,28}, provided that consultation is a time of understanding, support and reassurance and are more likely to talk about their needs when asked about them^{5,7}.

Addressing women in puerperal consultation

Regarding actions performed by professionals during puerperium consultation, a survey in the UK indicates that 93% of the women had abdominal examination and 70% vaginal examination. However, many women are submitted to exams with no obvious reason (such as unnecessary vaginal examination), while other tests that may be useful for some conditions (such as hemogram for fatigued women) are used infrequently²⁸.

Brazilian studies indicate that the most common activities carried out in the puerperium by health professionals, besides encouraging breastfeeding, are guidelines on contraception methods^{18,25}, food and nutrition^{18,21} and use of ferrous sulfate¹⁸. Less frequent actions are cytopathology of the uterine cervix²⁴, clinical breast examination^{18,24}, investigation of women's emotional state, orientation regarding return to sexual intercourse, evaluation of lochia and examination of the genital region¹⁸. A national study that evaluated puerperal care among adolescents classified it as adequate for 43.2% of the users, and the other ones received intermediate or inadequate care, considering in this evaluation the consultation of puerperal review, home visit in the first week after childbirth, clinical and gynecological evaluation, guidance on family planning, and guidance on breastfeeding²⁵. In Peru, it was found that only 38.5% of the women used ferrous sulfate and 12.6% vitamin A 30 days after delivery, and 31% had chosen a contraceptive method²⁷.

During the first 3 months postpartum, Australian studies indicated that 89% of women had sexual health problems, and although most women had contact with PHC professionals during that time, only 24% remembered being asked about the theme by general practitioners and 14% by nurses^{30,31}. Women expressed dissatisfaction with the quality of postnatal primary

health care, reporting a strong shift from maternal health to baby health after birth³⁰.

Home visit as a postpartum care tool

Of the 42 studies selected for the review, nine addressed the topic of home visits (HV) in the postpartum period, four of them national^{18,21,25,32} and five international³³⁻³⁷. Four studies indicate a low level of HV^{18,21,25,33}, and the same occurs more frequently in Brazil when it comes to family health facilities, performed mainly by Community Health Workers^{18,25}. HV are more likely when the team monitors women during the prenatal period^{18,33}.

As to the positive impact of HV programs during the gestational-puerperal cycle, international studies have pointed to reduced tobacco use among puerperae³⁴, improved care and development of the child^{36,37}, greater social and psychological support to women and households³⁷, and three international³⁴⁻³⁶ and two national²¹⁻³² studies reported increased exclusive breastfeeding was identified up to six months. A multiprofessional team is required to perform postpartum HV, which influences the improvement of the quality of care and nears ties between the team and women/households^{21,37,38}.

Breastfeeding

From the reviewed studies, breastfeeding appeared as the main action addressed by professionals in Brazil, and is a subject poorly investigated internationally when addressing actions regarding puerperae in PHC. Seven Brazilian studies have evaluated the impact of its prevalence before and after breastfeeding promotion initiatives (such as "breastfeeding-friendly basic facility" or "Brazil breastfeeding network"), identifying that such actions contribute to an increased rate of breastfeeding, both exclusive up to six months and subsequent to this period^{19,32,39-41}. Two studies did not find statistical significance in their results, although they also showed superiority in the percentage of breastfeeding after the implementation of the initiative^{42,43}.

A study carried out in Francisco Morato/SP, which evaluated the knowledge of health professionals about breastfeeding showed that they recognize the importance of breastfeeding for the mother-child binomial, but more easily mention the benefits to the child. In addition, although women receive guidance, they have limited knowledge on the subject⁴⁴.

Postpartum depression (PPD): non-specialized management in PHC

The international literature shows that PPD screening programs in PHC can improve women's mental health outcomes in the postpartum period⁴⁵⁻⁴⁹, even in low-income regions with low-cost technologies, as pointed out by Chilean⁴⁷ and Australian⁵⁰ studies. The Edinburgh Postnatal Depression Scale (EPDS) validated in several countries is recognized as the main PPD tracking tool by non-specialized professionals in PHC^{45-49,51}.

It is emphasized that public health nurses are well positioned to identify and treat depressed women and refer them when needed. A small investment in the training of nurses to identify and treat PPD may have a positive long-term effect^{45,48}.

According to an Australian study⁵⁰, a health professional's investigation about the past or current mental health of women is associated with the search for help throughout the perinatal period. In Denmark, it was found that puerperae who developed a psychiatric episode had higher rates of consultation by health professionals before, during and after pregnancy⁵².

Barriers to treatment of PPD were evidenced by a study in Chile, such as lack of user knowledge about the disease, negative conceptualization and rejection of the available treatment options; poor network support and long waiting times and a lack of coordination between clinical and administrative decisions⁵³.

Among studies that addressed PPD, three were national and identified insufficient knowledge by PHC professionals to prevent and treat PPD, lack of a systematic tool for PPD screening and that work focus during gestation and postpartum is still biological⁵⁴⁻⁵⁶.

Discussion

The bibliographic search in this study started with a large number of references (1,974), and 43 were selected for analysis according to inclusion and exclusion criteria. Most of the references that made up the initial sample of the search addressed child health or women's health at other periods of life unrelated to puerperium.

We can affirm that literature that addresses women's care in the puerperium in the context of PHC is scarce, since most studies selected did not provide a comprehensive approach to puerperal care, so evidence was limited to one-off issues of specific actions, and evaluative studies were re-

stricted to process evaluations only. Of the studies that evaluated the puerperium process, only Silva et al.¹⁸ conducted a more extensive investigation of the actions carried out in puerperium, while others, mainly Brazilians, mostly limited themselves to investigating the number of postpartum consultations and impact of breastfeeding promotion programs.

An increased national and international scientific production was noted in the last 5 years, indicating a rising concern for women's health in the puerperium, possibly by the promotion of strategies aimed at reducing maternal and child morbimortality, for example, proposed by the Sustainable Development Goals, and adopted by several countries. Despite worldwide decline, in the last decades, maternal and infant mortality remains a public health problem, especially in developing countries, and developed countries identified the need to qualify care for women in the postpartum period to reduce maternal morbidity, which may also have fueled increased studies in recent years.

According to the studies analyzed, puerperal care provided in PHC predominates in postpartum consultation and breastfeeding indicators, in national papers and home visits and PPD programs conducted mainly at the international level. Other aspects of care, such as assessment of physical structure and materials, health professionals and other activities performed in the puerperium (general guidelines, physical assessment and social vulnerability assessment) are poorly investigated.

In Brazil, investigations are restricted to the immediate and late puerperium, addressing remote puerperium only when it comes to breastfeeding, and the latter focusing on the child. International studies dealt with remote puerperium in a maximum of two and a half years postpartum^{34,36}, with a predominance of PDD investigation^{46-48,50-52}. While many women traverse puerperium without interurrences, others experience significant issues that may persist for weeks, months or even more than 4 years after giving birth^{7,8}, indicating these women's need for care over time.

Considering that maternal health is PHC-sensitive^{2,3} and that this point of care should provide care for women longitudinally, there is a need to qualify such actions in order to follow-up on women in all stages of puerperium, including the remote stage, neglected not only by public policies and clinical guidelines^{3,6,57}, but also in scientific studies. It is noteworthy that most of PHC

care is low cost and has significant results in reducing maternal morbimortality with integrated and universal access actions^{2,3}.

Results showed that, in developing countries (Brazil and Peru), the puerperal consultation still does not occur with a satisfactory frequency, and maternal mortality rates remain high in such countries. There is a need for change in PHC access policies for women, their partners and/or family members during this period, which could affect the improvement of maternal and child morbimortality rates. Most studies approach the topic of puerperal consultation only as to its frequency rather than investigating their quality.

The highest risk of mortality is during the immediate and late postpartum period, with a higher morbimortality rate in the first postpartum week, which makes it a critical time for women and their child. Postpartum consultation is a primary intervention to reduce maternal morbimortality through prevention, early detection and treatment of complications, and advice on contraception^{4,5}. However, this action does not seem to provide an opportunity for women to have their health needs met, either because of poor quality of care²⁸ or because of the low frequency of these consultations²³.

In addition to the low frequency of puerperal consultation, results of the studies on the topic “women approach in puerperium consultation” indicate the incipient and poor care provided in PHC, with fragmented, biologicist care, childcare priority, focus on the “mother” role and not in women and their needs, in both national and international studies.

International clinical guidelines^{3,9,57,58} guide the delivery of care to women in the puerperium, primarily by PHC professionals, through comprehensive and coordinated care, valuing women’s health needs, stimulating autonomy, self-care and care for their child. These guidelines are based on a high degree of scientific evidence and stand out for the quality of care delivery, which must be better investigated in the studies that address puerperal consultation and better implemented in the practice of providing care to women in the puerperium within PHC in different Brazilian and international settings.

HV is an important moment for the PHC team to detect early physical and emotional changes⁹, develop educational actions, identify household risks and vulnerabilities, establish a professional-user bond and improve the protection of women and children health in the postpartum period¹⁸.

Regarding the mentioned health care strategy, international studies have analyzed HV programs and their impacts, showing their importance for puerperal care. Brazilian studies are limited to investigating the frequency of visits and, incipiently, actions performed in HV, especially during the first week postpartum, and the main professional to perform this function is the community health worker, who also focuses on child and biologicist care.

With respect to breastfeeding, although it is an indicator of women’s health in the postpartum period, studies still address the issue with greater emphasis on the child. This topic and the indicator number of postpartum consultations were the ones with most emphasis in Brazilian studies, with predominantly quantitative analysis, indicating a gap in investigations of the quality of care provided in PHC.

Breastfeeding benefits to women’s health are countless¹⁹ and the main difficulties in its maintenance occur in the first months postpartum, and PHC is an incentive environment for this practice⁴¹, considering its proximity to women and their priority role in the provision of puerperal care⁹.

PPD care identified in the papers analyzed was extensively investigated, especially in international studies. EPDS has emerged as an important tool for PPD screening by PHC professionals in developing countries to ensure early detection and appropriate therapeutic intervention in cases of PPD. National studies point to “empirical” PPD care, with professionals unprepared for screening, prevention and treatment of this condition⁵⁴⁻⁵⁶.

PPD is an international phenomenon and the most common complication in the postpartum period. Viewed as a public health problem whose prevalence ranges from 10% to 15% in developed countries, reaching up to 40% in developing countries^{45,53,56}, this condition affects women’s quality of life and link with the newborn and its development^{53,54,56}.

PHC health professionals are well positioned to provide psychosocial support to all women in the perinatal period and to perform preventive approaches among women who may be more likely to develop PPD or related disorders, particularly in the immediate postpartum period^{9,58}. These professionals should play an active role in screening and selecting PPD treatment, considering contextual factors, such as culture, socioeconomic condition, among others^{9,58}.

The limitations found for the accomplishment of this review referred to the wide range of

descriptors used, resulting in a large number of papers to perform the reading of titles and abstracts. However, this strategy was necessary, due to the scarcity of studies that use the postpartum period as a descriptor. This limitation was overcome by a thorough reading of titles and abstracts, ensuring selection of the studies that addressed the guiding question. Furthermore, a variety of research methods used to address the actions of puerperal programs was observed, which prevents a more accurate comparative analysis of the results achieved. As any type of study, this integrative review has advantages and limitations of the method itself, and care should be taken with regard to conclusions related to the findings. However, authors highlight the potential of the method used, since it has validity among experts in the review area.

Final considerations

Results indicate that there is a range of actions to be performed by PHC to assist women in the puerperium, which can be carried out using light and low-cost technologies, and that this point of care is paramount in helping to reduce of maternal morbimortality through counseling and support for recovery from pregnancy and birth, early identification and appropriate management of physical and emotional health needs.

Despite the recognized importance of puerperal care, Brazilian studies are mostly confined to investigating the number of postpartum visits, the number of home visits and the effect of breastfeeding programs, neglecting the comprehensive nature of postpartum care to women in PHC.

National and international studies indicate that postpartum care still focuses on newborn

care and are mostly restricted to the immediate and late puerperium, and there is a need to improve the implementation of primary health care for women, especially in relation to the coordination and longitudinality of care, which evidences the need for better quality of care for puerperae in PHC.

It is noteworthy that PPD has been the target of countless studies in the international literature, given its high prevalence and public health issue, and evidenced the high potential of PHC professionals to act in the prevention, screening and treatment of said disease. In Brazil, this is a poorly studied topic worthy of attention of postpartum public policies, with results that point to the lack of qualification of PHC professionals for PPD care.

This review evidenced knowledge gaps regarding the subject matter, with the need for more robust studies that address comprehensive women care, as well as aspects of the quality of care provided in PHC. In addition, there is a need for evaluative research with a qualitative and quantitative approach in order to understand the implementation of puerperal care in PHC and the relationship of results found with the puerperal care program.

It is important to emphasize that the puerperium investigation can identify a range of activities to be developed and considered in PHC, both physical, material and human structure, as well as professional qualification, program planning and management, besides external factors cultural aspects of the woman receiving care. Thus, this study serves as a subsidy both to guide the elaboration and improvement of actions to the benefit of women in the puerperium in PHC by professionals and managers and to direct the elaboration of empirical studies that can investigate the integrality of puerperal care.

Collaborations

T Baratieri worked on the manuscript design, methodology, drafting and approval of the final version to be published. S Natal worked on manuscript design, methodology, critical review and approval of the final version.

References

- World Health Organization (WHO). Global Health Observatory (GHO) data. *Women and health* [Internet] 2017 [cited 2017 Jun 10]. Available from: <http://www.who.int/gho/en/>
- Bhutta ZA, Cabral S, Chan C-w, Keenan WJ. Reducing maternal, newborn, and infant mortality globally: An integrated action agenda. *Obstet Gynecol Int J* 2012; 119(Supl. 1):13-17.
- World Health Organization (WHO). *WHO recommendations on Postnatal care of the mother and newborn*. Geneva: WHO Library Cataloguing-in-Publication Data; 2013.
- Nabukera SK, Witte K, Muchunguzi C, Bajunirwe F, Batwala VK, Mulogo EM, Farr C, Barry S, Salihu HM. Use of postpartum health services in rural Uganda: knowledge, attitudes, and barriers. *J Community Health* 2006; 31(2):84-93.
- Brodribb W, Zadoroznyj M, Dane A. The views of mothers and GPs about postpartum care in Australian general practice. *BMC Fam Pract* 2013; 14(1):1-9.
- Brasil. Ministério da Saúde (MS), Instituto Sírio-Libanês de Ensino e Pesquisa. *Protocolos da Atenção Básica: Saúde das Mulheres*. Brasília: MS; 2016.
- Hartley S, Sutherland G, Brown S, Yelland J. 'You're more likely to tell the GP if you're asked': women's views of care from general practitioners in the first postpartum year. *Aust J Prim Health* 2012; 18(4):308-312.
- Gartland D, MacArthur C, Woolhouse H, McDonald E, Brown S. Frequency, severity and risk factors for urinary and faecal incontinence at 4 years postpartum: a prospective cohort. *Int J Obstet Gynecol* 2016; 123(7):1203-1211.
- National Institute for Health and Care Excellence (NICE). Antenatal and postnatal mental health: clinical management and service guidance. London: NICE; 2015.
- Correa MS, Feliciano KV, Pedrosa EN, Souza AI. Women's perception concerning health care in the post-partum period: a meta-synthesis. *Open J Obstet Gynecol* 2014; 4:416-426.
- Oliveira DC, Teixeira RC, Tomiyoshi JT, Correa ACdP, Mandu ENT. Estrutura organizacional da atenção pós-parto na estratégia saúde da família. *Esc Anna Nery Rev Enferm* 2013; 17(3):446-554.
- Corrêa MSM, Feliciano KVO, Pedrosa EN, Souza AI. Acolhimento no cuidado à saúde da mulher no puerpério. *Cad Saude Publica* 2017; 33(3):1-12.
- Mendes KDS, Silveira RCCP, Galvão CM. Revisão integrativa: método de pesquisa para a incorporação de evidências na saúde e na enfermagem. *Texto & contexto enferm* 2008; 17(4):758-764.
- Stufflebeam DL. Program Evaluation Models Metaevaluation Checklist (Based on The Program Evaluation Standards). *Evaluation Checklists Project* [Internet] 1999 [cited 2017 May 15]. Available from: <http://dmeforpeace.org/learn/program-evaluations-models-metaevaluation-checklist>
- Critical Appraisal Skills Programme (Casp). *Programme AS. Making sense of evidence. CASP Checklists* [Internet] 2017 [cited 2017 May 15]. Available from: <http://www.casp-uk.net/casp-tools-checklists>
- Stetler CBMD, Rucki S, Broughton S, Corrigan B, Fitzgerald J, Giuliano K, Havener P, Sheridan A. Utilization-focused integrative reviews in a nursing service. *Appl Nurs Res* 1998; 11(4):195-206.
- Ursi E. *Prevenção de lesões de pele no perioperatório: revisão integrativa da literatura* [dissertação]. Ribeirão Preto: Universidade de São Paulo; 2005.
- Silva LLB, Feliciano KVO, Oliveira LNFP, Pedrosa EN, Corrêa MSM, Souza AI. Cuidados prestados à mulher na visita domiciliar da "Primeira Semana de Saúde Integral". *Rev Gaucha Enferm* 2016; 37(3):1-8.
- Rito RV, Oliveira MI, Brito Ados S. Degree of compliance with the ten steps of the Breastfeeding Friendly Primary Care Initiative and its association with the prevalence of exclusive breastfeeding. *J Pediatr (Rio J)* 2013; 89(5):477-484.
- Parada CMG. Avaliação da assistência pré-natal e puerperal desenvolvidas em região do interior do Estado de São Paulo em 2005. *Rev. bras. saúde mater. infant* 2008; 8(1):113-124.
- Laporte-Pinfildi ASC, Zangirolani LTO, Spina N, Martins PA, Medeiros MAT. Atenção nutricional no pré-natal e no puerpério: percepção dos gestores da Atenção Básica à Saúde. *Rev nutr* 2016; 29(1):109-123.
- Serruya SJ, Lago T, Cecatti JG. Avaliação preliminar do programa de humanização no pré-natal e nascimento no Brasil. *Rev bras ginecol obstet* 2004; 26(7):517-525.
- Corrêa MD, Tsunehiro MA, Lima MOP, Bonadio IC. Avaliação da assistência pré-natal em unidade com estratégia saúde da família. *Rev Esc Enferm USP* 2014; 48(n. esp.):23-31.
- Santos SR, Cunha AJLA, Gamba CM, Machado FG, Leal Filho JMM, Moreira NLM. Avaliação da assistência à saúde da mulher e da criança em localidade urbana da região Sudeste do Brasil. *Rev Saude Publica* 2000; 34(3):266-271.
- Vilarinho LM, Nogueira LT, Nagahama EEI. Avaliação da qualidade da atenção à saúde de adolescentes no pré-natal e puerpério. *Esc Anna Nery Rev Enferm* 2012; 16(2):312-319.
- Hass CN, Teixeira LB, Beghetto MG. Adequabilidade da assistência pré-natal em uma estratégia de saúde da família de Porto Alegre-RS. *Rev Gaucha Enferm* 2013; 34(3):22-30.
- Velásquez Hurtado JE, Solís Alcedo L, Vigo Valdez WE, Rosas Aguirre AM, Giusti Hundskopf P, Alfaro Fernandez P, Cabrera Arredondo H. Evaluación de las prácticas de cuidado materno infantil en áreas con pobreza extrema del Perú, 2012. *Rev Peru Med Exp Salud Publica* 2014; 31(2):243-253.
- Bick DE, MacArthur C. Attendance, content and relevance of the six week postnatal examination. *Midwifery* 1995; 11(2):69-73.

29. Guerreiro EM, Rodrigues DP, Queiroz ABA, Ferreira MA. Educação em saúde no ciclo gravídico-puerperal: sentidos atribuídos por puérperas. *Rev Bras Enferm* 2014; 67(1):13-21.
30. Woolhouse H, McDonald E, Brown SJ. Changes to sexual and intimate relationships in the postnatal period: women's experiences with health professionals. *Aust J Prim Health* 2014; 20(3):298-304.
31. McDonald E, Woolhouse H, Brown SJ. Consultation about sexual health issues in the year after childbirth: a cohort study. *Birth* 2015; 42(4):354-361.
32. Graça LCC, Figueiredo MCB, Conceição MTCC. Contributions of the nursing intervention in primary healthcare for the promotion of breastfeeding. *Rev Lat Am Enfermagem* 2011; 19(2):429-436.
33. Sitrin D, Guenther T, Murray J, Pilgrim N, Rubayet S, Ligowe R, Pun B, Malla H, Moran A. Reaching mothers and babies with early postnatal home visits: the implementation realities of achieving high coverage in large-scale programs. *PLoS One* 2013; 8(7):e68930.
34. Mejdoubi J, van den Heijkant SC, van Leerdam FJ, Crone M, Crijnen A, HiraSing RA. Effects of nurse home visitation on cigarette smoking, pregnancy outcomes and breastfeeding: a randomized controlled trial. *Midwifery* 2014; 30(6):688-695.
35. Bashour HN, Kharouf MH, AbdulSalam AA, El Asmar K, Tabbaa MA, Cheikha SA. Effect of postnatal home visits on maternal/infant outcomes in Syria: a randomized controlled trial. *Public Health Nurs* 2008; 25(2):115-125.
36. Robling M, Bekkers M-J, Bell K, Butler CC, Cannings-John R, Channon S, Martin BC, Gregory JW, Hood K, Kemp A, Kenkre J, Montgomery AA, Moody G, Owen-Jones E, Pickett K, Richardson G, Roberts ZE, Ronaldson S, Sanders J, Stamuli E, Torgerson D. Effectiveness of a nurse-led intensive home-visitation programme for first-time teenage mothers (Building Blocks): a pragmatic randomised controlled trial. *Lancet* 2016; 387(10014):146-155.
37. Saïas T, Lerner E, Greacen T, Simon-Vernier E, Emer A, Pintaux E, Guédeney A, Dugravier R, Tereno S, Falissard B, Tubach F; CAPEDP Study Group, Revah-Levy A. Evaluating fidelity in home-visiting programs a qualitative analysis of 1058 home visit case notes from 105 families. *PLoS One* 2012; 7(5):e36915.
38. Laporte-Pinfildi ASC, Medeiros MAT. Nutritional care during prenatal and postpartum periods: A report of experiences in a city on São Paulo's coast. *Rev nutr* 2016; 29(6):947-961.
39. Cardoso LO, Vicente AS, Damião JJ, Rito RV. The impact of implementation of the Breastfeeding Friendly Primary Care Initiative on the prevalence rates of breastfeeding and causes of consultations at a basic healthcare center. *J Pediatr* 2008; 84(2):147-153.
40. Alves ALN, Oliveira MIC, Moraes JR. Iniciativa Unidade Básica Amiga da Amamentação e sua relação com o aleitamento materno exclusivo. *Rev Saude Publica* 2013; 47(6):1130-1140.
41. Passanha A, Benicio MHD, Venancio SI, Reis MCG. Implantação da Rede Amamenta Brasil e prevalência de aleitamento materno exclusivo. *Rev Saude Publica* 2013; 47(6):1141-1148.
42. Almeida CC, Scochi MJ, Souza RKT, Carvalho WO. Prevalência de aleitamento materno antes e após a implantação de um programa de redução de morbimortalidade infantil, no município de Campo Mourão (PR). *Cien Saude Colet* 2010; 15(2):575-583.
43. Brandao DS, Venâncio SI, Giugliani ER. Association between the Brazilian Breastfeeding Network implementation and breastfeeding indicators. *J Pediatr* 2015; 91(2):143-151.
44. Ciconi RCV, Venancio SI, Escuder MML. Avaliação dos conhecimentos de equipes do Programa de Saúde da Família sobre o manejo do aleitamento materno em um município da região metropolitana de São Paulo. *Rev bras saúde mater infant* 2004; 4(2):193-202.
45. Leung SS, Leung C, Lam T, Hung S, Chan R, Yeung T, Miao M, Cheng S, Leung SH, Lau A, Lee DT. Outcome of a postnatal depression screening programme using the Edinburgh Postnatal Depression Scale: a randomized controlled trial. *J Public Health* 2010; 33(2):292-301.
46. Milgrom J, Erickson J, Negri L, Gemmill AW. Screening for postnatal depression in routine primary care: properties of the Edinburgh Postnatal Depression Scale in an Australian sample. *Aust N Z J Psychiatry* 2005; 39(9):833-839.
47. Rojas G, Fritsch R, Solis J, Jadresic E, Castillo C, González M, Guajardo V, Lewis G, Peters TJ, Araya R. Treatment of postnatal depression in low-income mothers in primary-care clinics in Santiago, Chile: a randomised controlled trial. *Lancet* 2007; 370(9599):1629-1637.
48. Glavin K, Smith L, Sørsum R, Ellefsen B. Redesigned community postpartum care to prevent and treat postpartum depression in women—a one-year follow-up study. *J Clin Nurs* 2010; 19(21-22):3051-3062.
49. Segre LS, O'Hara MW, Brock RL, Taylor D. Depression screening of perinatal women by the Des Moines Healthy Start Project: program description and evaluation. *Psychiatr Serv* 2012; 63(3):250-255.
50. Reilly N, Harris S, Loxton D, Chojenta C, Forder P, Austin MP. The impact of routine assessment of past or current mental health on help-seeking in the perinatal period. *Women Birth* 2014; 27(4):e20-e7.
51. Abiodun O. Postnatal depression in primary care populations in Nigeria. *Gen Hosp Psychiatry* 2006; 28(2):133-136.
52. Munk-Olsen T, Pedersen HS, Laursen TM, Fenger-Grøn M, Vedsted P, Vestergaard M. Use of primary health care prior to a postpartum psychiatric episode. *Scand J Prim Health Care* 2015; 33(2):127-133.
53. Rojas. Barreras de acceso a tratamiento de la depresión posparto en centros de atención primaria de la Región Metropolitana: un estudio cualitativo. *Rev Med Chile* 2015; 143(4):424-432.

54. Meira B, Pereira P, Silveira M, Gualda D, Santos Jr H. Desafios para profissionais da atenção primária no cuidado à mulher com depressão pós-parto. *Texto & Contexto Enferm* 2015; 24(3):706-712.
55. Felix TA, Siqueira DA, Ximenes Neto FRG, Nascimento KV, Mira QLM. Actuación de enfermería frente a la depresión postparto en las consultas de puericultura. *Enferm Glob* 2013; 12(29):404-419.
56. Santos Junior HPO, Gualda DMS, Silveira MFA, Hall WA. Postpartum depression: the (in) experience of Brazilian primary healthcare professionals. *J Adv Nurs* 2013; 69(6):1248-1258.
57. National Institute for Health and Care Excellence (Nice). *Postnatal care up to 8 weeks after birth* [Internet] 2015 [cited 2017 Oct 27]. Available from: nice.org.uk/guidance/cg37
58. Austin MP, Highet, Guidelines Expert Advisory Committee. *Clinical practice guidelines for depression and related disorders: anxiety, bipolar disorder and puerperal psychosis, in the perinatal period. A guideline for primary care health professionals*. Melbourne: Beyond-blue; 2011.

Article submitted 14/08/2017

Approved 11/04/2018

Final version submitted 13/04/2018