

# Postpartum Stress: Current Concepts and the Possible Protective Role of Breastfeeding

Maureen Wimberly Groer, RN, PhD, FAAN,

Mitzi Wilkinson Davis, RN, PhD, Jean Hemphill, RN, MSN

**Objective:** To review stress during the postpartum and the research supporting that a unique, protective biology exists in breastfeeding mothers that may reduce reactivity to stress.

**Data Sources:** Publications from nursing and biomedical literature.

**Study Selection:** Studies reviewed were those that have contributed to concurrent conceptualizations of postpartum stress. Additionally, studies with sufficient participants were analyzed for common findings. Animal literature was reviewed for studies on the stress response in lactating and nonlactating animal models.

**Data Extraction:** Stress during the postpartum may be conceptualized as physical, intrapersonal, and interpersonal. Animal data and a few recent human studies suggest that the neuroendocrinology of the lactating mother may down-regulate the magnitude of the stress response.

**Data Synthesis:** A diminished stress response may serve to protect the breastfeeding maternal-infant dyad from environmental stimuli and to direct the physiology of the mother toward milk production, energy conservation, and nurturance.

**Conclusion:** Nurses understand the benefits of breastfeeding for optimal infant health, but new research suggests that maternal health may also be benefited in a biologic and as yet unexplored way. *JOGNN*, 31, 411-417; 2002.

**Keywords:** Breastfeeding—Neuroendocrinology—Physiology—Postpartum—Stress

Accepted: November 2001

The purpose of this review is to examine the stressors that occur in the postpartum period and to explore a possible stress-protecting role of breastfeeding. Research suggests that lactational physiology may produce a diminution of the biologic stress response and even the psychologic perceptions of stress.

Although for most parents there is joy and satisfaction in the birth of a new baby, many situations or events occur that are stressors, with the potential for taxing parents' ability to function. Smith (1989) compared primiparae to multiparae in terms of postpartum adjustments that may lead to stress. For primiparae, the most frequent reported stressors were baby feeding, fatigue, breast soreness, return of the figure, limiting visitors, regulating demands, growth and development, interpreting behavior, physical care of the baby, and labor and delivery. Multiparae reported major stressors of fatigue, regulating demands, emotional tension, sibling jealousy, and the labor and delivery experience.

## Stressors

Horowitz and Damato (1999) conducted a triangulated study on postpartum stress. Ninety-five women were studied at Week 6 postpartum. Mothers described experiences that they felt were stressful, which the researchers found fell into four categories (roles, task, resources, and relationships). The following numbers of stressors were reported: 86 reports in the roles category, 60 in tasks (such as child care, housework), 41 in resources, and 13 in

relationships. This study and others cited below support that the specific stressors associated with the postpartum can be divided into somewhat overlapping categories of physical, intrapersonal, and interpersonal phenomena.

### *Physical Stressors*

A variety of physical symptoms occurs in the postpartum, as illustrated by participants in a study by Gjerdingen, Froberg, Chaloner, and McGovern (1993). Those participants reported breast symptoms, fatigue, hemorrhoids, poor appetite, constipation, increased sweating, acne, hand numbness/tingling, dizziness, hot flashes, illness days, respiratory symptoms, sexual concerns, and hair loss. Some physical conditions such as backache (Breen, Ransil, Groves, & Oriol, 1994; MacEvelly & Buggy, 1996), urinary incontinence (Wilson, Herbison, & Herbison, 1996), and sexual dysfunction (Glazener, 1997) have been studied because of their duration, frequency, and/or adverse impact. Physical symptoms may be stressors in themselves or may be due to stress as demonstrated by Lenz, Parks, Jenkins, and Jarrett's (1986) finding that negative life events stress was a predictor of postpartum illness in mothers of 6-month-old infants. Of the physical symptoms reported by mothers, fatigue is consistently identified as a major concern (Brown & Lumley, 1998; Chapman, Macey, Borum, & Bennett, 1985; Gardner, 1991; Gjerdingen et al., 1993; Graef et al., 1988; Gruis, 1977; Ruchala & Halstead, 1994). It has been found to correlate positively with perceived stress in breastfeeding women at 3 days and 3, 6, and 9 weeks postpartum (Wambach, 1998). Pugh and Milligan (1993) proposed that maternal fatigue may alter self-care, mothering behaviors, and infant caregiving. On average, the amount of sleep in a 24-hour period during the 4th week postpartum is 7.53 hours, but only 6.15 hours occur during the night. Furthermore, the sleep pattern is discontinuous (Quillin, 1997). Sleep difficulties continue for some women for many months, and the effects of sleep deprivation can be acute and cumulative (Tulman, Fawcett, Groblewski, & Silverman, 1990). Fatigue is related to both quantity and quality of sleep and can alter, through neuroendocrine pathways, the stress response, immune function, and well-being.

### *Intrapersonal Stress*

A common intrapersonal stress is concern about weight. Stein and Fairburn (1996) found that eating disorder symptoms in a sample of 98 primagravidas increased in the first 3 months postpartum and then plateaued over the next 6 months. Walker's (1998) analysis of 227 new mothers' written descriptions of feelings about weight concluded that 8% had weight-related distress.

The development of maternal identity (Rubin, 1984) and maternal role attainment, defined by Mercer (1986, p. 198) as a process in which "the mother achieves com-

petence in the role and integrates the mothering behaviors into her established role set, so that she is comfortable with her identity as a mother," are clearly intrapersonal challenges in the postpartum period. Maternal stress impedes sensitivity to the infant. Ragozin, Basham, Crnic, Greenberg, and Robinson (1982) found a relationship between high levels of maternal stress at 1 month postpartum and decreased maternal sensitivity to cues at 4

**T**he postpartum is a time of joy and satisfaction and also a time of vulnerability or even of crisis.

months postpartum. In a study of role adaptation (Dormire, Strauss, & Clarke, 1989), adolescent primiparae, when highly stressed, demonstrated less sensitivity to the messages conveyed by their infants' behaviors and less effective maternal behaviors.

A dramatic intrapersonal consequence of stress is postpartum depression. With an incidence consistently estimated at more than 10%, postpartum depression has a deep and persistent negative impact on mothers, infants, partners, and other children. Hall, Kotch, Browne, and Rayens (1996) found the effects of common, everyday stressors on depressive symptoms in 738 women 1 to 2 months postpartum were both direct and mediated by self-esteem. Everyday stressors were more strongly associated with depressive symptoms than were major life events. Beck's (1996) meta-analysis of 44 studies of depression and various predictor variables found that child care stress and life stress were major predictors of depression and only prenatal depression had a greater effect size.

### *Interpersonal Stress*

Pregnancy and childbirth are periods of stressful transition with respect to women's social roles. Gruis (1977) found that at 1 month postpartum almost all mothers were concerned about their abilities to meet the needs of family members, take effective and safe care of their infants, and find time for themselves. This finding was replicated by Harrison and Hicks (1983) in mothers 3 to 8 weeks postpartum. In a study of work and postpartum health, McGovern et al. (1997) found that 90% of women reported role function limitations and that time off from work of more than 20 weeks had a positive effect on maternal mental health and role function.

Longer term effects on functional status (the performance of behaviors associated with roles) is indicated by the finding that even at 6 months postpartum, 20% of women had not resumed their full level of household

activities and 30% had not fully resumed their social and community activities (Tulman et al., 1990). The multiplicity and overlapping of roles often provokes conflict and stress (Spurlock, 1995). Thomas (1997) suggested that stress associated with women's role commitments may contribute to immunosuppression and disease vulnerability. This would support Gjerdingen et al.'s (1993) suggestion that increased gynecologic and breast symptoms among working women at 9 to 12 months postpartum may represent a delayed response to the stress of combining multiple roles of employment, new parenthood, and marriage.


A decrease in satisfaction with or strain in the marital relationship following birth is common (Lewis, 1988), and Affonso and Mayberry (1990) found mate/spouse interactions to be stressors of high intensity. Mercer (1986) found that maternal perception of overall family functioning (among and between individuals, dyads, and family-social units such as work and school) was adversely affected by life events stress, which was measured in women at both high and low risk at 8 months postpartum. Unrealistic cultural assumptions about motherhood result in work overload, difficulty with transition to motherhood, stress, guilt, and sometimes depression (Lee, 1997).

### Biologic Factors: Lactation and Stress Responses

The postpartum is associated with significant stressors, and excessive stress poses a threat to maternal, child, and family well-being. It is important to determine if postpartum biology plays a role in the mother's stress response and in stress's potentially deleterious effects on her health. Stress must be understood in terms of both the biologic processes that mediate response and the psychological processes that affect perception, appraisal, and coping.

This article focuses on the biologic processes mediating stress reactivity and alterations or adaptations in those processes occurring during lactation. Basics of the physiologic response are illustrated in Figure 1. In general, a stressor, which may be physiologic or psychosocial or both, simultaneously activates the sympathetic adrenomedullary system leading to release of the catecholamines epinephrine and norepinephrine and the hypothalamic-pituitary-adrenal system (HPA axis) leading to release of corticotropin, cortisol, and endogenous opioid peptides. Specifically, the hypothalamus secretes corticotropin releasing hormone (CRH), which stimulates the anterior pituitary hormone to secrete adrenocorticotrophic hormone (ACTH). This ACTH stimulates the adrenal cortex to synthesize and secrete glucocorticoids. All of these phenomena are influenced by the neuroendocrine and immune physiology of the postpartum woman.

It is interesting to note how few studies of postpartum stress and adaptation or even postpartum physiology dis-



**FIGURE 1**  
The physiologic response to stress.

*Note.* SNS = Sympathetic nervous system. CRH = corticotropin releasing hormone. ACTH = adrenocorticotrophic hormone.

tinguish lactating from nonlactating women. All postreproduction animal studies are necessarily of lactating animals, whereas in humans breastfeeding is a choice. Clearly, the neuroendocrinology of a lactating mother is profoundly different from that of a mother who is not lactating. From a biobehavioral perspective, it is reasonable to propose that the postpartum stress response might differ in breastfeeding and bottle-feeding mothers. Available literature supports this premise.

Lactating animals have a well-described diminished physiologic reactivity to stressors. Human studies, although few in number, support the existence of the same response in breastfeeding mothers. Although mothers may be subjected to many stressors in the postpartum, nature has apparently designed a way to decrease the magnitude of their physiologic response to stress if they breastfeed. The unique endocrine and immune state of the lactating mother is directed toward protection of the maternal-infant dyad, reducing maternal reactivity to the environment, promoting calmness and nurturing behavior, maintaining quantity and quality of milk, enhancing

immune function, and opposing glucocorticoid-mediated catabolism (Altemus, Deuster, Galliven, Carter, & Gold, 1995). It is characterized by down-regulated stress responses in both adrenergic and HPA reactivity, which return to normal very quickly when lactation ceases (Windle, Brady, et al., 1997) as do blood levels of cortisol, ACTH, and prolactin (Fischer, Patchev, Hellbach, Hassan, & Almeida, 1995).

**S**tressors during the postpartum may be divided into somewhat overlapping categories of physical, intrapersonal, and interpersonal phenomena.

The decrease in stress reactivity appears to originate in late pregnancy, when there is a surge in cortisol, due in part to placental CRH. This CRH from the placenta stimulates ACTH and glucocorticoid secretion. At the end of pregnancy, cortisol levels are very high and may approach levels seen in clinical depression or anorexia nervosa or in athletes who exercise strenuously (Magiakou, Mastorakos, Webster, & Chrousos, 1997).

Cortisol facilitates many physiologic and behavioral adaptations of late pregnancy, including catabolic mobilization of maternal energy stores to meet fetal needs and preparation of the breasts for later lactation (Atkinson & Waddell, 1995). In humans, CRH, which has risen throughout pregnancy, peaks at labor and falls dramatically at 1 day postpartum (Smith, Chan, Bowman, Harewood, & Phippard, 1993).

During pregnancy, there is a central suppression of CRH, as increasing placental CRH appears to suppress CRH release from the hypothalamus. This suppression and the gradual return to normal as the HPA axis “resets” itself may be implicated in the genesis of postpartum dysphoria. Lower levels of afternoon cortisol in the early postpartum have been associated with postpartum depression (Harris et al., 1996), as has the longer and more severe postpartum blunting of ACTH response to exogenous CRH administration (Magiakou et al., 1997).

The blunting of the HPA axis may be due to neuroendocrine mechanisms. Levels of estrogen and progesterone that were high during pregnancy drop dramatically at birth and remain low if a female lactates. Prolactin and oxytocin levels rise after birth and remain high if the mother lactates. Estrogen is an important factor in the secretion of ACTH, as it directly stimulates CRH production (Magiakou et al., 1997). The low levels of estrogen during lactation may therefore suppress CRH release by the hypothalamus. Plasma catecholamines also appear to

be reduced in response to stressors in lactating rats. Higuchi, Negoro, and Arita (1989) found that immobilization stress produced a smaller elevation in both plasma epinephrine and norepinephrine in lactating compared with nonlactating rats.

In addition, oxytocin and prolactin, which are high during lactation, may inhibit the HPA axis. Cook (1997) demonstrated this effect when he found that cortisol release decreased in both lactating and nonlactating sheep after infusion with oxytocin and prolactin and exposure to the stress of a barking dog. The stress increased prolactin in the nonlactators but not in lactating rats. Lightman and Young (1989) showed a similar effect on oxytocin release.

The antistress nature of lactation may be governed, in part, by the hormone oxytocin, which is maintained at high levels throughout lactation. Oxytocin is also released upon HPA axis activation, particularly in response to emotional stress. In stressed male rats, the oxytocin level rises remarkably within the brain, even though peripheral levels may not change (Engelmann, Ebner, Landgraf, Holsboer, & Wotjakl, 1999). Oxytocin infusion in both male and female rats produces sedation, lower blood pressure, decreased corticosteroid levels, and evidence of increased vagal activity. A variety of stimuli produce an elevation in oxytocin release in nonlactating animals, which may account for certain behavioral phenomena such as positive mood and prosocial behavior (Uvnas-Moberg, 1997). Infusion of oxytocin in 182 healthy men produced attenuated arousal and anger (Pietrowsky, Krug, Fehm, & Born, 1992). In human mothers, this hormone has been found to be amnesic and to reduce anxiety and produce sedation. The possibility of oxytocin-induced pain reduction is suggested by animal studies (Carter & Altemus, 1997) and supported by our knowledge of a mother with a chronic pain syndrome who reported that the only time she was free of debilitating pain was when she was breastfeeding.

A difference in behavioral response to stress has been clearly demonstrated in lactating and nonlactating animals exposed to a mild stress (10 minutes of white noise). Virgin rats showed HPA activation and increased activity levels, displacement grooming behaviors, and rearing, whereas lactating rats showed an entirely different repertoire of behaviors that were directed toward their pups. They did not show activation of the HPA response in response to the stress (Windle, Woods, et al., 1997). During lactation, there may also be increased central nervous system gamma-aminobutyric acid, a major inhibitory neurotransmitter, which may inhibit the affective state and behavior of the lactating animal (Qureshi, Hansen, & Soderstein, 1987).

The small amount of research conducted on the reactivity to stressors in lactating humans supports the premise that the breastfeeding mother has a diminished



response to stressors. Weisenfeld, Malatesta, Whitman, Grannose, and Vile (1985) found that compared to bottle-feeding mothers, breastfeeding mothers showed reduced galvanic skin response and heart rate changes while listening to a tape of a crying infant. Lactating women were found to have lower plasma levels of ACTH, cortisol, glucose, and norepinephrine than did nonlactating women following 20 minutes of graded treadmill exercise as a stressor (Altemus et al., 1995).

Mezzacappa, Kelsey, and Katlin (1999) studied blood pressure and continuous impedance cardiography in breastfeeding and bottle-feeding mothers and found a greater cardiovascular response to psychologic stressors in the mothers who bottle-fed their infants. They also reported lower incidences of psychiatric and cardiovascular disorders in the breastfeeding group. Light et al. (2000) found that breastfeeding mothers had lower blood pressure compared with bottle-feeding mothers, both during a laboratory stress experiment and later at home. As predicted from animal studies, they found that oxytocin release during stress was higher in lactating mothers and this hormone suppressed stress responsivity.

## Summary

This review has provided evidence for consideration of the postpartum as potentially stressful for a variety of physical, intrapersonal, and interpersonal factors. Nevertheless, postpartum women may be protected from the deleterious effects of stress through biologic mechanisms that have been extensively studied in animals but not in humans.

Lactating animals have a diminished physiologic reactivity to stressors, and a few human studies confirm this for breastfeeding mothers.

This protection appears to be governed by the neuroendocrinology associated with lactation, which is the natural, biologic normative behavior of all postpartum animal mothers. This protection makes evolutionary sense in that the maternal-infant dyad is protected and supported by these responses. Little attention has been paid to this obvious biologic difference in studies of human mothers. Lactational state has rarely been a variable in studies of the human postpartum, although only in humans is lactation a choice. We encourage maternal-child researchers to analyze their data with this perspective in mind.

Practice implications of these concepts include the possibility that the bottle-feeding mother might be physiolog-

ically more fragile and more susceptible to postpartum illness. Further research is required to provide sufficient evidence for this effect. Of course, nurses need to support all postpartum mother-infant dyads and to promote nurturant behaviors, including feeding, whether it is by breast or bottle.

## REFERENCES

- Affonso, D. D., & Mayberry, L. J. (1990). Common stressors reported by a group of childbearing American women. *Health Care for Women International, 11*(3), 331-345.
- Altemus, M., Deuster, P., Galliven, E., Carter, C., & Gold, P. (1995). Suppression of hypothalamic-pituitary-adrenal axis responses to stress in lactating women. *Journal of Clinical Endocrinology and Metabolism, 80*, 2954-2959.
- Atkinson, H., & Waddell, B. (1995). The hypothalamic-pituitary-adrenal axis in rat pregnancy and lactation: Circadian variation and interrelationship of plasma adrenocorticotropin and corticosterone. *Endocrinology, 136*, 512-520.
- Beck, C. T. (1996). A meta-analysis of the relationship between postpartum depression and infant temperament. *Nursing Research, 45*(4), 225-230.
- Breen, T. W., Ransil, B. J., Groves, P. A., & Oriol, N. E. (1994). Factors associated with back pain after childbirth. *Anesthesiology, 81*(1), 29-34.
- Brown, S., & Lumley, J. (1998). Maternal health after childbirth: Results of an Australian population based survey. *British Journal of Obstetrics and Gynaecology, 105*, 156-161.
- Carter, C. S., & Altemus, M. (1997). Integrative functions of lactational hormones in social behavior and stress management. *Annals of the New York Academy of Science, 87*, 164-174.
- Chapman, J., Macey, M., Borum, P., & Bennett, S. (1985). Concerns of breastfeeding mothers from birth to four months. *Nursing Research, 34*, 374-377.
- Cook, C. (1997). Oxytocin and prolactin suppress cortisol response to acute stress in both lactating and non-lactating sheep. *Journal of Dairy Research, 64*, 327-339.
- Dormire, S. L., Strauss, S. S., & Clarke, B. A. (1989). Social support and adaptation to the parent role in first-time adolescent mothers. *Journal of Obstetric, Gynecologic, and Neonatal Nursing, 18*, 327-337.
- Engelmann, M., Ebner, K., Landgraf, R., Holsboer, F., & Wotjak, C. T. (1999). Emotional stress triggers intrahypothalamic but not peripheral release of oxytocin in male rats. *Journal of Neuroendocrinology, 11*(11), 867-872.
- Fischer, D., Patchev, V., Hellbach, S., Hassan, S., & Almeida, O. (1995). Lactation as a model of naturally reversible hypercorticalism plasticity in the mechanisms governing hypothalamo-pituitary-adrenocortical activity in rats. *Journal of Clinical Investigation, 96*, 1208-1215.
- Gardner, D. (1991). Fatigue in postpartum women. *Applied Nursing Research, 4*, 57-62.
- Gjerdingen, D. K., Froberg, D. G., Chaloner, K. M., & McGovern, P. M. (1993). Changes in women's physical health during the first postpartum year. *Archives of Family Medicine, 2*(3), 277-283.

- Glazener, C.M.A. (1997). Sexual function after childbirth: Women's experiences, persistent morbidity and lack of professional recognition. *British Journal of Obstetrics and Gynaecology*, 104, 330-335.
- Graef, P., McGhee, K., Rozycki, J., Fescina-Jones, D., Clark, J. A., Thompson, J., & Brooten, D. (1988). Postpartum concerns of breastfeeding mothers. *Journal of Nurse-Midwifery*, 33(2), 62-66.
- Gruis, M. L. (1977). Beyond maternity: Postpartum concerns of mothers. *MCN: American Journal of Maternal Child Nursing*, 2, 182-188.
- Hall, L. A., Kotch, J. B., Browne, D., & Rayens, M. K. (1996). Self-esteem as a mediator of the effects of stressors and social resources on depressive symptoms in postpartum mothers. *Nursing Research*, 45(4), 231-238.
- Harris, B., Lovett, L., Smith, J., Read, G., Walker, R., & Newcombe, R. (1996). Cardiff puerperal mood and hormone study: III. Postnatal depression at 5 to 6 weeks postpartum, and its hormonal correlates across the peripartum period. *British Journal of Psychiatry*, 168, 739-744.
- Harrison, M., & Hicks, S. (1983). Postpartum concerns of mothers and their sources of help. *Canadian Journal of Public Health*, 74, 325-328.
- Higuchi, T., Negoro, H., & Arita, J. (1989). Reduced responses of prolactin and catecholamine to stress in the lactating rat. *Journal of Endocrinology*, 122, 495-498.
- Horowitz, J., & Damato, E. (1999). Mother's perceptions of postpartum stress and satisfaction. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 28, 595-605.
- Lee, C. (1997). Social context, depression, and the transition to motherhood. *British Journal of Health Psychology*, 2, 93-108.
- Lenz, E. R., Parks, P. L., Jenkins, L. S., & Jarrett, G. E. (1986). Life change and instrumental support as predictors of illness in mothers of 6-month-olds. *Research in Nursing and Health*, 9, 17-24.
- Lewis, J. M. (1988). The transition to parenthood: II. Stability and change in marital structure. *Family Process*, 27, 273-283.
- Light, K., Smith, T., Johns, J., Brownley, K., Hofheimer, J., & Amico, J. (2000). Oxytocin responsivity in mothers of infants: A preliminary study of relationships with blood pressure during laboratory stress and normal ambulatory activity. *Health Psychology*, 19, 560-567.
- Lightman, S., & Young, W. (1989). Lactation inhibits stress-mediated secretion of corticosterone and oxytocin and hypothalamic accumulation of corticotropin-releasing factor and enkephalin messenger ribonucleic acid. *Endocrinology*, 124, 2358-2364.
- MacEvilly, M., & Buggy, D. (1996). Back pain and pregnancy: A review. *Pain*, 64, 405-414.
- Magiakou, M., Mastorakos, G., Webster, E., & Chrousos, G. (1997). The hypothalamic-pituitary-adrenal axis and the female reproductive system. *Annals of the New York Academy of Sciences*, 816, 42-56.
- McGovern, P., Dowd, B., Gjerdingen, D., Moscovice, I., Kochevar, L., & Lohman, W. (1997). Time off work and the postpartum health of employed women. *Medical Care*, 35, 507-521.
- Mercer, R. T. (1986). *First-time motherhood*. New York: Springer.
- Mezzacappa, E., Kelsey, R., & Katlin, E. (1999, March). A preliminary study of maternal cardiovascular function and breast-feeding in the first year. Poster presentation at Annual Meeting, American Psychosomatic Society, Vancouver, BC, Canada.
- Pietrowsky, R., Krug, R., Fehm, H., & Born, J. (1992). The effects of "stress hormones" on emotional sensitivity. *Zeitschrift für Experimentelle Angewandte Psychologie*, 39(2), 278-292.
- Pugh, L., & Milligan, R. (1993). A framework for the study of childbearing fatigue. *Advances in Nursing Science*, 15, 60-70.
- Quillin, S. (1997). Infant and mothers sleep patterns during the 4th postpartum week. *Issues in Comprehensive Pediatric Nursing*, 20(2), 115-123.
- Qureshi, G. A., Hansen, S., & Soderstein, P. (1987). Offspring control of cerebrospinal fluid GABA concentrations in lactating rats. *Neuroscience Letters*, 75, 85-88.
- Ragozin, A. S., Basham, R. B., Crnic, K. A., Greenberg, M. T., & Robinson, N. M. (1982). Effects of maternal age on parenting. *Developmental Psychology*, 18, 627-634.
- Rubin, R. (1984). *Maternal identity and the maternal experience*. New York: Springer.
- Ruchala, P. L., & Halstead, L. (1994). The postpartum experience of low-risk women: A time of adjustment and change. *Maternal-Child Nursing Journal*, 22(3), 83-89.
- Smith, M. P. (1989). Postnatal concerns of mothers: An update. *Midwifery*, 5(4), 182-188.
- Smith, R., Chan, E., Bowman, M., Harewood, W., & Phippard, A. (1993). Corticotropin-releasing hormone in baboon pregnancy. *Journal of Clinical Endocrinology and Metabolism*, 76, 1063-1068.
- Spurlock, J. (1995). Multiple roles of women and role strains. *Health Care for Women International*, 16(6), 501-508.
- Stein, A., & Fairburn, C. G. (1996). Eating habits and attitudes in the postpartum period. *Psychosomatic Medicine*, 58(4), 321-325.
- Thomas, S. P. (1997). Distressing aspects of women's roles, vicarious stress, and health consequences. *Issues in Mental Health Nursing*, 18(6), 539-557.
- Tulman, L., Fawcett, J., Groblewski, L., & Silverman, L. (1990). Changes in functional status after childbirth. *Nursing Research*, 39, 70-75.
- Uvnas-Moberg, K. (1997). Oxytocin-linked antistress effects—The relaxation growth response. *Acta Physiologica Scandinavica*, 640, 38-42.
- Walker, L. O. (1998). Weight-related distress in the early months after childbirth. *Western Journal of Nursing Research*, 20, 30-44.
- Wambach, K. (1998). Maternal fatigue in breastfeeding primiparae during the first nine weeks postpartum. *Journal of Human Lactation*, 14(3), 219-229.
- Weisenfeld, A., Malatesta, C., Whitman, P., Grannose, C., & Vile, R. (1985). Psychophysiological response of breast and bottle-feeding mothers to their infants' signals. *Psychophysiology*, 22, 79-86.
- Wilson, P. D., Herbison, R. M., & Herbison, G. P. (1996). Obstetric practice and the prevalence of urinary incontinence three months after delivery. *British Journal of Obstetrics and Gynaecology*, 103(2), 154-161.

- Windle, R., Brady, M., Kunanandam, T., Da Costa, A., Wilson, B., Harbuz, M., Lightman, S., & Ingram, C. (1997). Reduced response of the hypothalamo-pituitary-adrenal axis to alpha-1-agonist stimulation during lactation. *Endocrinology*, 138, 3741-3748.
- Windle, R. J., Woods, S., Shanks, N., Perks, P., Conde, G. L., Da Costa, A., Ingram, C. D., & Lightman, S. L. (1997). Endocrine and behavioral responses to noise stress: Comparison of virgin and lactating female rats during non-disrupted maternal activity. *Journal of Neuroendocrinology*, 9(6), 407-414.

*Maureen Wimberly Groer is a professor and associate dean for research and evaluation, The University of Tennessee, College of Nursing, Knoxville.*

*Mitzi Wilkinson Davis is an associate professor, The University of Tennessee, College of Nursing, Knoxville.*

*Jean Hemphill is a doctoral student, The University of Tennessee, College of Nursing, Knoxville.*

*Address for correspondence: Maureen Wimberly Groer, RN, PhD, FAAN, The University of Tennessee, College of Nursing, 1200 Volunteer Blvd., Knoxville, TN 37996-4180. E-mail: mgroer@utk.edu.*