

Research Article

Potential antiviral agents of *Rosmarinus officinalis* extract against herpes viruses 1 and 2

Wafa A. AL-Megrin¹, Norah A. AISadhan²,  Dina M. Metwally^{3,4}, Razan A. Al-Talhi³,  Manal F. El-Khadragy^{1,5} and Lina J. M. Abdel-Hafez⁶

¹Biology Department, Faculty of Science, Princess Nourah Bint Abdulrahman University, Riyadh, Kingdom of Saudi Arabia; ²College of Medicine, Almaarefa University, Riyadh, Kingdom of Saudi Arabia; ³Department of Zoology, Faculty of Science, Faculty of Science, King Saud University, Riyadh, Kingdom of Saudi Arabia; ⁴Parasitology Department, Faculty of Veterinary Medicine, Zagazig University, Zagazig, Egypt; ⁵Department of Zoology and Entomology, Faculty of Science, Helwan University, Cairo, Egypt; ⁶Department of Microbiology and Immunology, Faculty of Pharmacy, October 6 University, Giza, Egypt

Correspondence: Manal F. El-Khadragy (manalelkhadrage@yahoo.com)



Herpes simplex viruses 1 and 2 (HSV-1 and HSV-2) belong to the herpesviridae family and cause neurological disorders by infecting the nervous system. The present study aimed to investigate the effects of *Rosmarinus officinalis* L. (rosemary) extract against HSV-1 and HSV-2 *in vitro*. The antioxidant activity of this extract was investigated by superoxide anion and 2,2-diphenyl-1-picrylhydrazyl (DPPH) free-radical assays. Rosemary extract was evaluated by an HSV-1 antiviral assay, in which viral replication in Vero cells was determined and quantified using a cytopathic effect assay. The present study showed that rosemary extract at 30 µg/ml caused 55% inhibition of HSV-1 plaques, whereas 40 µg/ml rosemary extract caused 65% inhibition of HSV-2 plaques. The extracts completely inhibited HSV-1 and HSV-2 plaque formation at 50 µg/ml. Scavenging activity of the superoxide anion radical was observed at 65.74 mg/ml, whereas 50% scavenging activity of the DPPH radical was observed at 67.34 mg/ml. These data suggest that rosemary extract may be suitable as a topical prophylactic or therapeutic agent for herpes viral infections. However, further research is required to elucidate the plant's active constituents, which may be useful in drug development.

Introduction

Herpes simplex viruses 1 and 2 (HSV-1 and HSV-2) belong to the herpesviridae family, primarily causing oral herpes lesions with HSV-1 and genital lesions with HSV-2 [1]. During the primary infection, HSV causes damage to the central nervous system, and can even result in encephalitis and meningitis [2]. Infections in immune-compromised patients are most severe and have been reported to be lethal [3]. Treatment of herpes infections is thus a major cause of concern owing to the difficulty in eliminating it from the ganglion, the high cost of treatment, and the increasing drug resistance.

At present, the standard therapy for the management of HSV infections is based on inhibition of the viral DNA polymerase by nucleoside analogs [4], including acyclovir, penciclovir, and their derivatives [4,5]. The resistance of HSV to acyclovir has become an important clinical problem, especially in immune-compromised patients undergoing long-term therapy [5]. Antiviral drugs may not be a perfect choice in many cases due to drug resistance [6]. Hence, it is essential to develop new antiviral agents, which can act against a broad range of viruses.

Rosemary (*Rosmarinus officinalis*, L.) is an herb commonly used as a spice and flavoring agent in food processing. The leaves are a very good source of carnosol, carnosic acid, rosmanol, 7-methyl-epirosmanol,

Received: 01 April 2020
Revised: 15 May 2020
Accepted: 29 May 2020

Accepted Manuscript online:
29 May 2020
Version of Record published:
10 June 2020

isorosmanol, rosmadial, and caffeic acid, which have substantial *in vitro* antioxidant activities [7]. Antibacterial effects of rosemary have also been reported [8–10] and linked to their polyphenolic composition. Furthermore, rosemary oil may be used in drug-resistant infections [11]. Rosemary leaves possess a variety of biological activities, including anti-cancer and anti-inflammatory effects [12].

Traditional drugs have been used to prevent or treat HSV infections for many years. Presently, new antiviral agents with efficacy and without severe undesirable effects are being studied. Previous studies have suggested that some essential oils are effective against HSV-1 [13,14]. Therefore, the present work was undertaken to develop new anti-herpes drugs.

Materials and methods

Viruses and cell lines

Vero cells were grown in Dulbecco's modified Eagle's medium (DMEM; Gibco, Brazil) supplemented with 10% fetal bovine serum (FBS; Gibco, Brazil) and gentamycin (80 µg/ml). The cells were maintained at 37°C in a humidified atmosphere with 5% CO₂. The herpes simplex virus types 1 and 2 were propagated in Vero cells and titrated based on plaque-forming units using plaque assay counts, as previously described [4], and stored at –80°C until the experiments were performed.

Plant extractions

Dried rosemary (*R. officinalis*, L.) fine powder (100 g) was extracted in water at a 50× (w/v) concentration for 24 h at room temperature (30 ± 2°C). The extract was filtered, lyophilized, and stored at –20°C.

Determination of flavonoid and phenol content

The aluminum chloride colorimetric method was used for total flavonoid content determination in the sample following the previously described method [15]. The total polyphenolic content was determined using a Folin-Ciocalteu reagent, which measures the oxidation of polyphenols to a blue-colored complex with a maximum absorbance at 750 nm [16].

Determination of DPPH radical scavenging activity

The free radical scavenging capacity was evaluated by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay [17], where the percent of DPPH decolorization of the sample was calculated by the following equation:

$$\text{DPPH (\% scavenging activity)} = \left(\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \right) \times 100$$

where A_{control} is the absorbance of DPPH solution without the sample and A_{sample} is the absorbance of DPPH solution with the sample. Superoxide anion scavenging activity was measured by determining the amount of reduced phenazine methosulfate [17]. Vitamin C was used as a positive control.

Superoxide radical scavenging assay

This activity was estimated by the reduction of nitro blue tetrazolium (NBT) based on the previously described method [18]. The non-enzymatic phenazine methosulfate-nicotinamide adenine dinucleotide (PMS/NADH) system forms superoxide anion (O₂^{•-}) radicals, which reduce NBT to a purple formazan dyes. Briefly, to a one 1 ml reaction solution contained 20 mM phosphate buffer (pH 7.4), 73 µM NADH, 50 µM NBT, 15 µM PMS, and different concentrations (20–100 mg/ml) of extract solution. After incubation at room temperature for 5 min, the color intensity was measured at 562 nm against the blank to determine the amount of formed day. Vitamin C was used as positive control.

Reducing power assay

The Fe³⁺ reducing power of the extract was determined according to a method by Abdel Moneim [18]. Briefly, 50, 100, 150, and 200 µl of the samples were mixed with 1.9 ml 0.2 M phosphate buffer, pH 6.6, and 2 ml of 1% potassium ferricyanide. The mixture was then incubated at 50°C for 20 min. Afterward, the mixture was stopped by adding 2 ml of 10% trichloroacetic acid and then centrifuging at 3000 rpm for 10 min. The upper layer of supernatant (2 ml) was mixed with distilled water (2 ml) and 0.1% FeCl₃ solution (0.5 ml). The absorbance was measured at 700 nm against a blank with a spectrophotometer and vitamin C was used as a standard. Reaction mixtures with a higher absorbance indicated a greater reducing power.

Table 1 Total phenolics and flavonoids contents of rosemary extracts

Conditions	Total phenolics*	Total flavonoids†
Rosemary extract	35.6 ± 0.8	22.3 ± 1.1

*Total phenolics are expressed as µg/mg gallic acid equivalent of polyphenols/mg extract.

†Flavonoids are expressed as µg/mg quercetin equivalents of flavonoids/mg extract.

Data are represented as mean ± SEM of three independent experiments each performed in duplicate.




Figure 1. Determination of DPPH radical scavenging activity

The antiradical activities of rosemary extract and ascorbic acid against the DPPH radical, as measured by changes in absorbance at 517 nm.

Cytotoxicity assay

Vero cells were exposed to different rosemary concentrations (1–5000 µg/ml) for 72 h and an MTT assay (3-(4,5-dimethylthiazol-2,5-diphenyl tetrazolium bromide) was performed to assess metabolic activity and therefore, cytotoxicity [4].

Antiviral activity assay

The antiviral activity assay was performed as previously described by Schnitzler [14]. Acyclovir (Sigma[®], St. Louis, MO, U.S.A.) at 0.05–2 µg/ml served as the positive control.

Viral plaque number reduction assay

A previously described procedure was followed for this assay [4], with minor modifications. Approximately 100 plaque-forming units of HSV-1 and HSV-2 were adsorbed for 1 h at 37°C on confluent cells and overlaid with MEM plus 1.5% carboxymethyl cellulose (CMC, Sigma[®], St. Louis, MO, U.S.A.), either in the presence or absence of different concentrations of orange peel. After 72 h, the cells were fixed and stained with naphthol blue black (Sigma) and plaques were counted. The 50% inhibitory concentration (IC₅₀) was defined as the concentration that inhibits the formation of 50% of the viral plaques when compared with untreated controls [19].

Statistical analysis

Data are presented as the mean ± the standard error of the mean ($x \pm \text{SEM}$) of at least three separate experiments.

Results and discussion

The total polyphenolic content in aqueous rosemary extract was 35.6 µg/mg gallic acid equivalents of polyphenols/mg extract, whereas the flavonoid content was 22.3 µg/mg quercetin equivalents of flavonoids/mg extract (Table 1). Similarly, Kim et al. [20] reported that a hot water extract of rosemary had 42.35 mg total phenolics/g and 26.98 µg/mg quercetin equivalents of flavonoids/mg extract. However, Couto et al. [21] obtained low polyphenol and flavonoid concentrations. The radical scavenging activity of the extract was 67.34 mg/ml, indicating that rosemary is a good antioxidant (Figure 1). The correlation between antioxidant activity and presence of polyphenols has been widely




Figure 2. Determination of superoxide anion scavenging activity

The antiradical activities of rosemary extract and ascorbic acid against O₂⁻ radicals, as measured by changes in absorbance at 560 nm.

Table 2 The antiviral activity of rosemary extract against HSV-1 and HSV-2

Rosemary	Cell cytopathic effect	
	HSV-1	HSV-2
10 µg/ml	+	+
15 µg/ml	+	+
20 µg/ml	+	+
25 µg/ml	+	+
30 µg/ml	-	+
40 µg/ml	-	+
50 µg/ml	-	-

studied in herbs [20,22] and reported by Estevez et al. [23], who isolated carnosic acid from rosemary.

Figure 2 shows the dose-response curves for the superoxide anion (O₂⁻) radical scavenging activity of the rosemary extract. The results indicate that the O₂⁻ radical scavenging activity was 65.74 mg/ml, suggesting that rosemary extract possesses strong antioxidant effects due to superoxide anion radical scavenging property.

Oxidation by NADPH oxidase leads to the generation of O₂⁻, which can be converted to H₂O₂ by superoxide dismutase (SOD) or reacts with nitric oxide (NO) to form peroxynitrite. Hydrogen peroxide can be further converted to oxygen and water by catalase and glutathione peroxidase. Superoxide radicals have been observed to inactivate enzymes, degrade DNA, kill cells, and damage cell membranes [24]. The antioxidant activity of rosemary extract was stronger than the superoxide anion scavenging activity.

The reducing power of a compound may serve as a significant indicator of its potential for antioxidant activity. During the reducing power assay, the presence of reductants (antioxidants) in the tested samples resulted in the reduction of the ferricyanide (Fe³⁺) complex to the ferrous form (Fe²⁺). Figure 3 shows that Fe²⁺ was monitored by measurement of absorbance at 700 nm, which increased linearly with increasing rosemary extract concentrations in the reaction mixture. It has been reported that a substances reducing power may likely be due to its hydrogen-donating ability [25].

The MTT assay results (Figure 4) show that the rosemary extract cytotoxicity in cultured Vero cells was up to 977 µg/ml, indicating that rosemary extract had low toxic effects. This assay was previously used to discriminate between antiviral and cytotoxicity concentrations of various essential oils [26].

To study the antiviral activity of rosemary extract against HSV-1 and HSV-2, the cytopathic inhibitory assay was performed to determine the cytopathic effects (CPEs) of the viruses and the data are shown in Table 2. A confluent monolayer of Vero cells was infected with HSV-1 and HSV-2 viruses. A high concentration (50 µg/ml) of rosemary extract showed moderate antiviral activity against the viruses, with a CPE seen at 35 µg/ml.

As shown in Figure 5, rosemary extract at 30 µg/ml caused 55% inhibition of HSV-1 plaques and 100% inhibition against HSV-1 at 50 µg/ml. The 50% inhabitation of acyclovir against HSV-1 was less than 0.9 µg/ml. A plaque inhibition assay was conducted to determine the IC₅₀.




Figure 3. Reducing power of rosemary extract

Reducing power of rosemary extract and ascorbic acid against Fe^{3+} ions, as measured by changes in absorbance at 700 nm.




Figure 4. Cytotoxicity assay

The cytotoxic effects of rosemary extract evaluated by MTT in Vero cells.

Figure 6 shows the *in vitro* activity of rosemary extract against anti-HSV-2. The HSV-2 strain was more sensitive, with 40 µg/ml of rosemary extract causing 65% inhibition of HSV-2 plaques and showing 100% inhibition at 50 µg/ml. The 50% inhibition of acyclovir against HSV-2 was less than 0.8 µg/ml. The anti-HSV-1 and 2 effects may be due to enhanced absorption of the extracts into the Vero cells. It may be that a phytochemical is participating in




Figure 5. The anti-herpes virus 1 activity of rosemary extract and acyclovir as a standard antiviral agent
In vitro activity of rosemary extract and acyclovir against anti-HSV-1. IC₅₀ is the concentration where a 50% cytotoxic effect is observed. Data are represented as mean ± SEM of two independent experiments, each performed in triplicate.




Figure 6. The anti-herpes virus 2 activity of rosemary extract and acyclovir as a standard antiviral agent
In vitro activity of rosemary extract and acyclovir against anti-HSV-2. IC₅₀ is the concentration where a 50% cytotoxic effect is observed. Data are represented as mean ± SEM of two independent experiments, each performed in triplicate.

phenolic binding with the protein coat of the virus, blocking ligands from the viral surface, thus inactivating the virus. However, further studies are needed to confirm these observations.

Several studies showed that polyphenols, triterpenes, anthraquinones, saponins, and polysaccharides that were isolated from natural plants can inhibit the replication of herpes viruses [27]. A large number of plant-derived and synthetic anti-HSV agents have also been described [4,28].

Conclusions

The present *in vitro* studies revealed that rosemary extract exhibited antiviral activity against HSV-1 and HSV-2, along with antioxidant effects. Further studies are necessary to identify the mechanism responsible for this activity.

Data Availability

All data generated or analyzed during this study are included in this article.

Competing Interests

The authors declare that there are no competing interests associated with the manuscript.

Author Contribution

M.F.E. and L.J.M.A. conceived and designed the experiments; W.A. and L.J.M.A. performed the experiments; N.A., R.A., L.J.M.A., and D.M. analyzed the data, contributed reagents/materials/analysis tools; and M.F.E. and L.J.M.A. wrote the paper.

Acknowledgements

This study was funded by the Deanship of Scientific Research at Princess Nourah bint Abdulrahman University through the Fast-Track research funding program.

Abbreviations

CPE, cytopathic effect; DPPH, 2,2-diphenyl-1-picrylhydrazyl; HSV, herpes simplex virus; NADH, nicotinamide adenine dinucleotide; NBT, nitro blue tetrazolium; PMS, phenazine methosulfate.

References

- Hong, W., Li, T., Song, Y., Zhang, R., Zeng, Z., Han, S. et al. (2014) Inhibitory activity and mechanism of two scorpion venom peptides against herpes simplex virus type 1. *Antiviral Res.* **102**, 1–10, <https://doi.org/10.1016/j.antiviral.2013.11.013>
- Swanson, 2nd, P.A. and McGavern, D.B. (2015) Viral diseases of the central nervous system. *Curr. Opin. Virol.* **11**, 44–54, <https://doi.org/10.1016/j.coviro.2014.12.009>
- Jiang, Y.C., Feng, H., Lin, Y.C. and Guo, X.R. (2016) New strategies against drug resistance to herpes simplex virus. *Int. J. Oral. Sci.* **8**, 1–6, <https://doi.org/10.1038/ijos.2016.3>
- Alvarez, A.L., Habtemariam, S., Abdel Moneim, A.E., Melon, S., Dalton, K.P. and Parra, F. (2015) A spiroketal-enol ether derivative from *Tanacetum vulgare* selectively inhibits HSV-1 and HSV-2 glycoprotein accumulation in Vero cells. *Antiviral Res.* **119**, 8–18, <https://doi.org/10.1016/j.antiviral.2015.04.004>
- Shabani, M., Nasr Esfahani, B., Sadeq Ehdaei, B., Moghim, S., Mirzaei, A., Sharifi, M. et al. (2019) Inhibition of herpes simplex virus type 1 replication by novel hsa-miR-7704 in vitro. *Res. Pharm. Sci.* **14**, 167–174
- Kondel, R., Shafiq, N., Kaur, I.P., Singh, M.P., Pandey, A.K., Ratho, R.K. et al. (2019) Effect of acyclovir solid lipid nanoparticles for treatment of Herpes Simplex Virus (HSV) infection in an animal model of HSV-1 infection. *Pharm. Nanotechnol.*, <https://doi.org/10.2174/2211738507666190829161737>
- Raskovic, A., Milanovic, I., Pavlovic, N., Milijasevic, B., Ubavic, M. and Mikov, M. (2015) Analgesic effects of rosemary essential oil and its interactions with codeine and paracetamol in mice. *Eur. Rev. Med. Pharmacol. Sci.* **19**, 165–172
- Fu, Y., Zu, Y., Chen, L., Efferth, T., Liang, H., Liu, Z. et al. (2007) Investigation of antibacterial activity of rosemary essential oil against *Propionibacterium acnes* with atomic force microscopy. *Planta Med.* **73**, 1275–1280, <https://doi.org/10.1055/s-2007-981614>
- Moreno, S., Scheyer, T., Romano, C.S. and Vojnov, A.A. (2006) Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. *Free Radic. Res.* **40**, 223–231, <https://doi.org/10.1080/10715760500473834>
- Sacco, C., Bellumori, M., Santomauro, F., Donato, R., Capei, R., Innocenti, M. et al. (2015) An in vitro evaluation of the antibacterial activity of the non-volatile phenolic fraction from rosemary leaves. *Nat. Prod. Res.* **29**, 1537–1544, <https://doi.org/10.1080/14786419.2014.986728>
- Raskovic, A., Milanovic, I., Pavlovic, N., Cebovic, T., Vukmirovic, S. and Mikov, M. (2014) Antioxidant activity of rosemary (*Rosmarinus officinalis* L.) essential oil and its hepatoprotective potential. *BMC Complement. Altern. Med.* **14**, 225, <https://doi.org/10.1186/1472-6882-14-225>
- Perez-Sanchez, A., Barrajon-Catalan, E., Caturla, N., Castillo, J., Benavente-Garcia, O., Alcaraz, M. et al. (2014) Protective effects of citrus and rosemary extracts on UV-induced damage in skin cell model and human volunteers. *J. Photochem. Photobiol. B* **136**, 12–18, <https://doi.org/10.1016/j.jphotobiol.2014.04.007>
- Civitelli, L., Panella, S., Marcocci, M.E., De Petris, A., Garzoli, S., Pepi, F. et al. (2014) In vitro inhibition of herpes simplex virus type 1 replication by *Mentha suaveolens* essential oil and its main component piperitenone oxide. *Phytomedicine: Int. J. Phytotherapy Phytopharmacol.* **21**, 857–865, <https://doi.org/10.1016/j.phymed.2014.01.013>
- Schnitzler, P. (2019) Essential oils for the treatment of Herpes Simplex Virus infections. *Chemotherapy* **64**, 1–7, <https://doi.org/10.1159/000501062>
- Mahboubi, M., Kazempour, N. and Boland Nazar, A.R. (2013) Total phenolic, total flavonoids, antioxidant and antimicrobial activities of scrophularia striata boiss extracts. *Jundishapur J. Nat. Pharmaceut. Prod.* **8**, 15–19, <https://doi.org/10.17795/jjnpp-7621>
- Chandra, S., Khan, S., Avula, B., Lata, H., Yang, M.H., Elsohly, M.A. et al. (2014) Assessment of total phenolic and flavonoid content, antioxidant properties, and yield of aeroponically and conventionally grown leafy vegetables and fruit crops: a comparative study. *Evid. Based Complement. Alternat. Med.* **2014**, 253875, <https://doi.org/10.1155/2014/253875>
- Musa, K.H., Abdullah, A. and Al-Haiqi, A. (2016) Determination of DPPH free radical scavenging activity: application of artificial neural networks. *Food Chem.* **194**, 705–711, <https://doi.org/10.1016/j.foodchem.2015.08.038>
- Abdel Moneim, A.E. (2013) The neuroprotective effects of purslane (*Portulaca oleracea*) on rotenone-induced biochemical changes and apoptosis in brain of rat. *CNS Neurol. Disord. Drug Targets* **12**, 830–841
- Akahoshi, Y., Kanda, J., Ohno, A., Komiya, Y., Gomyo, A., Hayakawa, J. et al. (2017) Acyclovir-resistant herpes simplex virus 1 infection early after allogeneic hematopoietic stem cell transplantation with T-cell depletion. *J. Infect. Chemother.* **23**, 485–487, <https://doi.org/10.1016/j.jiac.2017.02.001>
- Kim, I.-S., Yang, M.-R., Lee, O.-H. and Kang, S.-N. (2011) Antioxidant activities of hot water extracts from various spices. *Int. J. Mol. Sci.* **12**, 4120–4131, <https://doi.org/10.3390/ijms12064120>
- Couto, R.O., Conceição, E.C., Chaul, L.T., Oliveira, E.M.S., Martins, F.S., Bara, M.T.F. et al. (2012) Spray-dried rosemary extracts: physicochemical and antioxidant properties. *Food Chem.* **131**, 99–105, <https://doi.org/10.1016/j.foodchem.2011.08.036>
- Yashin, A., Yashin, Y., Xia, X. and Nemzer, B. (2017) Antioxidant activity of spices and their impact on human health: a review. *Antioxidants (Basel)* **6**, 70
- Estevez, M., Ventanas, S., Ramirez, R. and Cava, R. (2005) Influence of the addition of rosemary essential oil on the volatiles pattern of porcine frankfurters. *J. Agric. Food Chem.* **53**, 8317–8324, <https://doi.org/10.1021/jf051025q>

- 24 Kang, S.N. (2016) Ethanol extracts from mistletoe (*Viscum album* L.) act as natural antioxidants and antimicrobial agents in uncooked pork patties during refrigerated storage. *Asian-Australas J. Anim. Sci.* **29**, 109–118, <https://doi.org/10.5713/ajas.15.0253>
- 25 Al-Olayan, E., Elkhadragey, M.F., Othman, M.S., Aref, A., Kassab, R. and Abdel Moneim, A.E. (2014) The potential protective effect of *Physalis peruviana* L. against carbon tetrachloride-induced hepatotoxicity in rats is mediated by suppression of oxidative stress and downregulation of MMP-9 expression. *Oxid. Med. Cell. Longev.* **2014**, <https://doi.org/10.1155/2014/381413>
- 26 Ansari, M., Sharififar, F., Arabzadeh, A.M., Mehni, F., Mirtadzadini, M., Iranmanesh, Z. et al. (2014) In vitro evaluation of anti-herpes simplex-1 activity of three standardized medicinal plants from Lamiaceae. *Anc. Sci. Life* **34**, 33–38, <https://doi.org/10.4103/0257-7941.150777>
- 27 Frederico, E., Cardoso, A., Moreira-Marconi, E., de Sa-Caputo, D.D.C., Guimaraes, C.A.S., Dionello, C.D.F. et al. (2017) Anti-viral effects of medicinal plants in the management of dengue: a systematic review. *Afr. J. Tradit. Complement. Altern. Med.* **14**, 33–40, <https://doi.org/10.21010/ajtcam.v14i4S.5>
- 28 Dkhil, M.A., Al-Quraishy, S. and Delic, D. (2015) The antioxidant and anti-herpes simplex viruses activity of *Morus alba* leaves extract. *Pakistan J. Zool.* **47**, 1563–1569