

RESEARCH ARTICLE

Potential of exogenously sourced kinetin in protecting *Solanum lycopersicum* from NaCl-induced oxidative stress through up-regulation of the antioxidant system, ascorbate-glutathione cycle and glyoxalase system

Mohammad Abass Ahanger¹, Mohammed Nasser Alyemeni², Leonard Wijaya², Saud A. Alamri², Pravej Alam³, Muhammad Ashraf^{2,4}, Parvaiz Ahmad^{2,5*}

1 Department of Botany, Govt Post-Graduation College Rajouri, Jammu and Kashmir, India, **2** Botany and Microbiology Department, College of Science, King Saud University, Riyadh, Saudi Arabia, **3** Biology Department, College of Science and Humanities, Prince Sattam bin Abdulaziz University, Alkharj, Kingdom of Saudi Arabia (KSA), **4** International Centre for Chemical and Biological Sciences, University of Karachi, Pakistan, **5** Department of Botany, S.P. College, Srinagar, Jammu and Kashmir, India

* parvaizbot@yahoo.com



OPEN ACCESS

Citation: Ahanger MA, Alyemeni MN, Wijaya L, Alamri SA, Alam P, Ashraf M, et al. (2018) Potential of exogenously sourced kinetin in protecting *Solanum lycopersicum* from NaCl-induced oxidative stress through up-regulation of the antioxidant system, ascorbate-glutathione cycle and glyoxalase system. PLoS ONE 13(9): e0202175. <https://doi.org/10.1371/journal.pone.0202175>

Editor: Yuan Huang, Huazhong Agriculture University, CHINA

Received: April 2, 2018

Accepted: July 30, 2018

Published: September 4, 2018

Copyright: © 2018 Ahanger et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Abstract

The protective role of exogenously applied kinetin (10 μM KN, a cytokinin) against the adverse effects caused by NaCl-induced (150 mM) stress in *Solanum lycopersicum* was investigated. Application of KN significantly enhanced growth and biomass production of normally grown plants (non-stressed) and also mitigated the adverse effect of NaCl on stressed plants to a considerable extent. Among the examined parameters, chlorophyll and carotenoid contents, photosynthetic parameters, components of the antioxidant system (both enzymatic and non-enzymatic), osmotica accumulation, and mineral uptake exhibited a significant increase following the application of KN. Furthermore, KN application reduced the generation of reactive free radical hydrogen peroxide, coupled with a significant reduction in lipid peroxidation and an increase in membrane stability. The activities of antioxidant enzymes, and glyoxylase system were found to be promoted in plants exposed to NaCl, and the activities were further promoted by KN application, thereby protecting *S. lycopersicum* plants against NaCl-induced oxidative damage. Further strengthening of the antioxidant system in KN supplied plants was ascribed to regulation of ascorbate-glutathione cycle, phenols and flavonoids in them. The levels of proline and glycine betaine increased considerably in KN-treated plants, thereby maintaining relative water content. Moreover, exogenous KN application reduced the inhibitory effects of NaCl on K⁺ and Ca²⁺ uptake, which resulted in a considerable reduction in tissue Na⁺/K⁺ ratio.

Introduction

Salinity-induced decline in growth and development of plants occurs due to many factors, resulting in cumulative effects on key physiological processes, such as disruption in ion homeostasis leading to altered water balance, and inhibition of photosynthesis and other biochemical processes, including enzymes' activity [1–3]. However, to counteract the negative effects of environmental extremes and maintain efficient metabolism, plants employ several tolerance mechanisms, including: (a) accumulation of compatible organic osmolytes, so that water uptake and the associated physiological pathways are least affected; (b) efficient partitioning and compartmentalization of toxic levels of ions into less sensitive tissues or organelles, such as the vacuole; and (c) up-regulation of the antioxidant system for rapid neutralization of stress-generated toxic ROS [3–6]. High salt (NaCl) concentrations are believed to induce both osmotic and ionic stresses. Among the osmolytes, low molecular weight solutes such as glycine betaine (GB), proline, amino acids, and soluble sugars have been reported to contribute effectively to osmotic adjustment and protection against osmotic stress [7,8]. The roles of these osmolytes in cellular osmotic adjustment, stabilization of membrane integrity, and the detoxification of toxic ions during salt stress have been well reported [9]. Accumulation of osmolytes has been observed to protect polypeptide dissociation and preserve the photosynthetic oxygen-evolving complex [10] as well as the activity of ribulose-1, 5-bisphosphate carboxylase (Rubisco) during NaCl-induced stress [1]. Among the antioxidants (both enzymatic and non-enzymatic), superoxide dismutase (SOD), catalase (CAT), peroxidases and reductases [including ascorbate peroxidase (APX) and glutathione reductase (GR)], ascorbic acid (AsA), and glutathione (GSH and GSSG) are the key candidates which contribute to stress tolerance by mediating protection of major physiological and biochemical processes [5]. Antioxidants mediate protection through their involvement in maintaining redox buffering for the optimal functioning of major pathways, including photosynthetic electron transport and enzyme activity [2,3,5,11]. However, for the elimination of toxic methylglyoxal glyoxylase system constituted of two key enzymatic components is up-regulated [12]. Thus, each antioxidant plays its specific role in counteracting a specific type of stress generated ROS. The biggest challenge before the scientists is to enhance the crop production under stressful environment. To achieve this goal. External supplementation of phytohormones could be a sustainable approach under for crop production under salinity.

Plant growth hormones have also been reported to play key roles in mediating several responses during stressful cues for growth and developmental regulation [2]. Recently, KN has been reported to enhance salt tolerance in soybean through its interaction with other growth hormones [13]. Exogenous application of plant growth hormones, such as cytokinins, has been shown to enhance the yield potential of agricultural crops under field conditions [14–16]. Cytokinins are believed to stimulate processes such as water uptake and cell division and enhance the synthesis of chlorophylls and organ development, leading to rapid regeneration and proliferation of shoot tissues [17,18]. KN is a synthesized form of artificial cytokinins and has been reported to ameliorate growth inhibition in several crop plants exposed to stress, such as waterlogging [19], salinity [20], and metal stress [21]. Exogenous application of KN can alleviate toxic effects on growth and photosynthesis by modulating the endogenous profile of other growth hormones, including salicylic acid, gibberellic acid, and jasmonic acid, whereas it acts antagonistically against others such as abscisic acid [13]. However, the KN-induced stress tolerance mechanism has received little attention, including KN-mediated salt exclusion and the restricted absorption from soil solution and subsequent distribution within the tissues or compartmentalization into vacuoles [22].

Tomato (*Solanum lycopersicum* L.) is one of the largest commercially consumed vegetables. It has high contents of bioactive compounds, such as β -carotene, lycopene, flavonoids, and ascorbic acid, and has accordingly been considered as an effective anti-oxidative and anti-cancerous fruit [23]. In the present study, we aimed to evaluate whether or not application of KN enhances antioxidant metabolism and osmoregulation in *S. lycopersicum*, plants and whether this can reverse the adverse effects of salt stress on growth, pigment synthesis, and ion accumulation.

Material and methods

Tomato seeds (var. Pusa Rohini) were obtained from Indian Agricultural Research Institute, Pusa, New Delhi, India. The seeds were surface sterilized using NaOCl (5%) and allowed to germinate in Petri plates lined with blotting paper wetted with distilled water. After germination, healthy seedlings were transplanted into plastic pots of 20-cm diameter and filled with reconstituted sand supplemented with vermicompost and were supplemented with full-strength Hoagland's nutrient solution (200 mL) on every alternate day for 10 days. For inducing salinity stress modified Hoagland's nutrient solution supplemented with NaCl (150 mM) was used for another 20 days and 10 μ M KN (20 mL per pot) was sprayed foliarly using Teepol (0.1%) as surfactant. During the entire growth period pots were kept under green house with natural climatic conditions having day/night temperatures of 26 /16°C, relative humidity of 70–75% and an average photoperiod of 18 hrs light/6 hrs night. After 30 days of growth plants were uprooted carefully and analyzed for different parameters following standard protocols described hereunder.

Plant length and dry weight

Plant length was measured after uprooting using a manual scale. Dry weight was recorded after oven-drying the tissue at 70°C for 24 h.

Estimation of photosynthetic pigments

Fresh leaf tissues were extracted in acetone (80%) using pestle and mortar. Optical density of the supernatant was measured spectrophotometrically at 480, 645, and 663 nm [24].

Estimation of chlorophyll fluorescence and gas exchange parameters

Chlorophyll fluorescence parameters were measured in fully expanded leaves using PAM chlorophyll fluorimeter (H. Walz, Effeltrich, Germany) [25]. Measurement of CO₂ assimilation rate (*A*), stomatal conductance (*g_s*) and transpiration rate (*E*) was carried in uppermost fully expanded leaves using infrared gas analyzer (LCA-4 model, Analytical Development Company, Hoddesdon, England).

Determination of leaf relative water content (RWC)

Method described by Smart and Bingham [26] was employed for the determination of RWC and calculation was done using following formula:

$$\text{RWC} = \frac{\text{Fresh weight} - \text{Dry weight}}{\text{Turgid weight} - \text{Dry weight}} \times 100 \quad (1)$$

Estimation of hydrogen peroxide (H₂O₂), lipid peroxidation (MDA) and electrolyte leakage

For estimation of content of hydrogen peroxide (H₂O₂) spectrophotometric method described by Velikova et al. [27] was followed and absorbance was measured at 390 nm. Standard curve of H₂O₂ was used for calculation and expressed as nM g⁻¹ FW.

Lipid peroxidation was estimated in terms of formation of malondialdehyde content and extinction coefficient of 155 mM⁻¹cm⁻¹ was used for calculation [28].

Electrolyte leakage was determined using the method of Dionisio-Sese and Tobita [29] and formula employed for computation was:

$$\text{Electrolyte leakage (\%)} = (EC_1 - EC_0) / (EC_2 - EC_0) \times 100 \quad (2)$$

Determination of antioxidant enzyme activities and ascorbate-glutathione cycle

For extraction of antioxidant enzymes 0.5 g of fresh leaf tissue was macerated in ice-cold potassium phosphate buffer (100 mM, pH 7.0) containing PVP (1%) in a pre-chilled mortar and pestle. The homogenate was centrifuged at 12000 × g for 15 min at 4°C and supernatant was used as an enzyme source for assay of SOD, CAT, APX, and GR.

Superoxide dismutase (SOD, EC1.15.1.1) activity was estimated by measuring its ability to inhibit the photochemical reduction of nitroblue tetrazolium (NBT) at 560 nm. Assay mixture containing 100 mM phosphate buffer (pH 7.4), methionine (10 mM), EDTA (1.0 mM), riboflavin (50 μM), NBT (75 μM) and 100 μL enzyme extract was incubated under fluorescent light for 15 min [30]. Activity was expressed as U mg⁻¹ protein.

For assaying activity of catalase (CAT, EC1.11.1.6) decomposition of H₂O₂ was monitored at 240 nm for 2 min. Reaction mixture (2 mL) contained 50 mM phosphate buffer (pH 6.0), 0.1 mM EDTA, 20 mM H₂O₂, and 100 μL enzyme extract. Extinction coefficient of 39.4 mM⁻¹cm⁻¹ was used and activity was expressed as U mg⁻¹ protein [31].

Glutathione S-transferase (GST, EC: 2.5.1.18) was estimated according to the procedure of Hasanuzzaman and Fujita [32] and change in absorbance was read spectrophotometrically at 340 nm for 2 min. The activity of GST was calculated using the extinction coefficient of 9.6 mM⁻¹ cm⁻¹.

For determination of APX (APX, EC1.11.1.1) activity, 2 mL reaction mixture containing 50 mM phosphate buffer (pH 7.5), 0.1 mM EDTA, 0.3 mM ascorbate, 100 μL enzyme extract and 100 μL H₂O₂ was monitored for 2 min at 290 nm [33]. For calculation molar extinction coefficient of 2.8 mM⁻¹ cm⁻¹ was used and activity was expressed as U mg⁻¹ protein.

Glutathione reductase (GR, EC1.6.4.2) was assayed by measuring the change in absorbance at 340 nm for 3 min. Reaction mixture contained potassium phosphate buffer (100 mM; pH 7.0), EDTA (1.0 mM), NADPH (50 μM), oxidized glutathione (100 μM) and 100 μL enzyme in a final volume of 3 mL. Activity was expressed as U mg⁻¹ protein [34].

Methods of Miyake and Asada [35] and Nakano and Asada [33] were employed for the estimation of monodehydroascorbate reductase (MDHAR, EC1.6.5.4) and dehydroascorbate reductase (EC:1.8.5.1) activities. The MDHAR activity was expressed as nmol NADPH oxidized / (EUmg⁻¹ protein). The DHAR activity was determined by monitoring change in absorbance at 265 nm for 1 min with a spectrophotometer (Beckman 640D, USA) using an extinction coefficient of 14 mM⁻¹ cm⁻¹.

Ascorbate was estimated in fresh leaves after homogenizing in ice-cold meta-phosphoric acid (5%) containing EDTA (1 mM). Homogenate was centrifuged at 10,000 rpm for 20 min and the supernatant was used for the estimation of ascorbate [36].

The glutathione pool was determined using the method of Yu et al. [37] with slight modifications [38]. Standard curves with known concentrations of GSH and GSSG were used for calculations.

Estimation of glyoxalase system

For determination of glyoxalase I (EC: 4.4.1.5) and glyoxylase II (EC: 3.1.2.6) method described by Hasanuzzaman and Fujita [32] was followed. Assay mixture for glyoxylase I contained K-phosphate buffer (100 mM, pH 7.0), magnesium sulphate (15 mM), GSH (1.7 mM) and methylglyoxal (3.5 mM) in a final volume of 900 μ L. The change in absorbance was recorded at 240 nm after adding methylglyoxal. For glyoxylase II, the reaction mixture (1.5 mL) contained 100 mM Tris-HCl buffer (pH 7.2), DTNB (0.2 mM) and 1 mM S-D-lactoylglycathione. Extinction coefficients of $3.37 \text{ mM}^{-1} \text{ cm}^{-1}$ and $13.6 \text{ mM}^{-1} \text{ cm}^{-1}$ were used for the calculation of glyoxylase I and II, respectively.

Total phenols and flavonoids

Plant samples were extracted in methanol and centrifuged at 10,000 rpm for 10 min. Total phenolic content was determined by the Folin-Ciocalteu reagent using the method of Chun et al. [39] and expressed as mg gallic acid equivalent (GAE) g^{-1} of extract (mg g^{-1}). For estimation of flavonoid content method of Zhishen et al. [40] was followed using catechin as standard. Absorbance was recorded at 510 nm using a spectrophotometer flavonoid content was expressed as mg catechin equivalents g^{-1} of extract (mg g^{-1}).

Estimation of proline and glycine betaine

Proline was extracted using toluene and the absorbance was read at 520 nm [41]. Standard curve of proline was used for calculation and content was expressed as $\mu\text{mol proline g}^{-1}$ fresh weight.

The method described by Grieve and Grattan [42] was adopted for the estimation of glycine betaine, and formation of periodide crystals was read at 365 nm after reacting the mixture with cold KI-I₂ reagent under acidic conditions.

Determination of tissue Na, K, and Ca

Dry plant samples (0.5 g) were acid-digested using sulfuric acid and nitric acid (1:5, v/v) at 60°C. The digested samples were cooled and read in a flame photometer for the estimation of Na, K, and Ca.

Statistical analysis

Data were presented as the means of three replicates with \pm SE calculated. The data were subjected to a statistical analysis using analysis of variance for a completely randomized design, followed by the Duncan's multiple range test, and least significant difference (LSD) calculated at 0.05%.

Results

Growth and biomass yield

The tomato plants treated with NaCl (150 mM) exhibited a significant reduction in all growth parameters measured. For example, shoot and root lengths decreased by 63.11% and 61.64%, respectively, and shoot and root dry weights by 62.50% and 57.14%, respectively, compared

with the control. Exogenous application of KN (10 μ M) to tomato plants enhanced shoot length by 45.91% and root length by 52.83% compared with those of salt stressed plants alone. However, KN application to NaCl-treated plants (150 mM NaCl + KN) improved the shoot and root dry weights by 60.00% and 66.66%, respectively over NaCl treated plants alone (Fig 1A and 1B).

Pigment content

Relative to the control, NaCl treatment reduced the synthesis of chlorophyll *a*, chlorophyll *b*, total chlorophyll, and carotenoids by 52.12%, 39.58%, 47.88%, and 40.00%, respectively over control. However, exogenously applied KN mitigated the negative effects of NaCl by enhancing the synthesis of chlorophyll *a* by 22.22%, chlorophyll *b* by 31.03%, total chlorophyll by 25.67%, and carotenoids by 9.80% compared with those of the NaCl-stressed plants alone (Table 1).

Chlorophyll fluorescence

Table 2 represents the effect of NaCl and KN on chlorophyll fluorescence. NaCl stress declines the F_v/F_m by 18.82%, ϕ PSII by 13.84%, ϕ_{exc} by 28.81% and qP by 27.47% but it enhanced the NPQ by 20.68% compared to control plants. External supplementation of KN restored the chlorophyll fluorescence by enhancing F_v/F_m , ϕ PSII, ϕ_{exc} and qP by 42.02%, 57.14%, 23.19% and 24.24% respectively over the plants treated with NaCl alone. NPQ showed decline in KN supplied NaCl stressed plants by 26.31% over NaCl treated plants alone.

Gas exchange parameters

The NaCl treated plants showed decrease in P_n , A , g_s and E by 27.17%, 44.95%, 77.07% and 68.39% respectively compared to control plants. However, application of KN to NaCl stressed plants enhanced the P_n by 82.88%, A by 52.97%, g_s by 147.11% and E by 74.54% over NaCl-alone treated plants (Fig 2A–2D).




Fig 1. Ameliorating role of kinetin (Kn) on length and dry weight of shoot and root under NaCl toxicity in tomato. Data presented are the means \pm SE (n = 3). Different letters indicate significant difference at $P \leq 0.05$.

<https://doi.org/10.1371/journal.pone.0202175.g001>

Table 1. Exogenous kinetin (Kn) enhanced the pigment (mg g⁻¹ FW) content in tomato under NaCl toxicity. Data presented are the means ± SE (n = 3). Different letters indicate significant difference at P ≤ 0.05.

Treatments	Chl a	Chl b	Total Chl	Carotenoids
0	0.94±0.021b	0.48±0.006a	1.42±0.091b	0.85±0.066b
0+ Kn	0.99±0.026a	0.49±0.006a	1.47±0.097a	0.98±0.075a
150 mM NaCl	0.45±0.005d	0.29±0.002c	0.74±0.056d	0.51±0.031d
150 mM +Kn	0.55±0.008c	0.38±0.004b	0.93±0.072c	0.56±0.045c

<https://doi.org/10.1371/journal.pone.0202175.t001>

Relative water content

NaCl application resulted in a 35.91% reduction in RWC, however, supplementation of KN (NaCl + KN) enhanced the RWC by 17.48% over NaCl treated plants alone (Fig 2E).

H₂O₂, MDA and EL

Salt (NaCl) stress caused an increment in the production of hydrogen peroxide (H₂O₂), lipid peroxidation (measured in terms of malondialdehyde (MDA) formation), and electrolyte leakage by 2.74 fold, 2.66 fold and 6.37 fold respectively over control plants. However, application of KN to NaCl treated plants decreased the H₂O₂ by 1.53 fold, MDA by 1.42 fold and EL by 1.19 fold over NaCl treated plants alone (Fig 3A–3C).

Antioxidant enzymes and ascorbate-glutathione cycle

Relative to the control, NaCl-stressed plants exhibited an increase of 61.43% in SOD, 74.34% in CAT, and 77.79% in GST activities, which were further enhanced by 16.99%, 29.60%, and 25.29% respectively, by the application of KN compared with those treated with NaCl-alone.

The effect of NaCl and KN on ascorbate-glutathione cycle is presented in Fig 4A–4C. The content of ascorbic acid was observed to be decreased by 41.86% due to NaCl stress, however, KN application mitigated this reduction by causing a slight increase of 28.00% compared with that in the NaCl-treated plants alone (Fig 5A). The NaCl enhanced the APX and GR activity by 86.05% and 28.74% respectively over control, however. External supplementation of KN, further enhanced the APX activity by 22.89% and Gr activity by 6.93% compared to NaCl treated plants alone (Fig 5B and 5C). Although NaCl reduced the activity of MDHAR (38.30%) and DHAR (32.31%) over control, KN (NaCl + KN) was observed to reinforce their activity by 60.56% and 25.74% respectively over NaCl stressed plants (Fig 5D and 5E). Relative to the control, NaCl treated plants exhibited an increase of 50.00% and 87.07% in GSH and GSSG, respectively. Compared with the NaCl-stressed plants, NaCl + KN-treated plants exhibited an increase of 17.42% in GSH and 56.04% in GSSG content (Fig 5F and 5G).

Table 2. Effect of kinetin (Kn) on chlorophyll fluorescence in tomato under NaCl toxicity. Data presented are the means ± SE (n = 3). Different letters indicate significant difference at P ≤ 0.05.

Treatments	Efficiency of PSII (Fv/Fm)	Quantum yield of PSII (φPSII)	Capture efficiency of PSII (φ _{exc})	Photochemical quenching (qP)	Non-photochemical quenching (NPQ)
0	0.85±0.06b	0.65±0.03c	0.59±0.007b	0.91±0.08a	0.58±0.007c
0+ Kn	0.95±0.09a	0.72±0.05b	0.63±0.02a	0.94±0.09a	0.40±0.002d
150 mM NaCl	0.69±0.02c	0.56±0.005d	0.42±0.002d	0.66±0.01c	0.95±0.08a
150 mM +Kn	0.98±0.12a	0.88±0.08a	0.53±0.004c	0.82±0.05b	0.70±0.05b

<https://doi.org/10.1371/journal.pone.0202175.t002>




Fig 2. Effect of kinetin (Kn) on gas exchange parameters and relative water content in tomato under NaCl toxicity. Data presented are the means \pm SE (n = 3). Different letters indicate significant difference at $P \leq 0.05$.

<https://doi.org/10.1371/journal.pone.0202175.g002>

Glyoxalase system

NaCl stress increased the activity of glyoxylase I (20.40%) while it decreased that of glyoxylase II (25.64%) activity over control. Application of KN (10 μ M) to NaCl stressed plants further stimulated the activity by 14.28% (glyoxylase I) and 107% (glyoxylase II) over those of NaCl stressed counterparts (Fig 6A).




Fig 3. Effect of kinetin (KN) on H₂O₂, MDA accumulation and EL in tomato under NaCl toxicity. Data presented are the means \pm SE (n = 3). Different letters indicate significant difference at $P \leq 0.05$.

<https://doi.org/10.1371/journal.pone.0202175.g003>




Fig 4. Effect of kinetin (KN) on activity of SOD, CAT and GST in tomato under NaCl toxicity. Data presented are the means ± SE (n = 3). Different letters indicate significant difference at P ≤ 0.05.

<https://doi.org/10.1371/journal.pone.0202175.g004>

Total phenols and flavonoids

NaCl treatment increased total phenols (78.61%) and decreased flavonoids (64.99%) over control plants. Contrary to this, application of KN to NaCl stressed plants further increased the contents of total phenols by 26.18% and ameliorated the decline in flavonoids by 51.38% over NaCl stressed plants alone (S1 Table).

Proline and glycine betaine

In NaCl-treated plants, free proline and glycine betaine were observed to increase by 3.81 fold and 6.50 fold, respectively, compared with the control; however, exogenous application of KN (NaCl + KN) further increased the contents of free proline and glycine betaine by 1.12 fold and 1.32 fold, respectively, compared with salt stressed plants alone (Fig 6B and 6C).




Fig 5. Effect of kinetin (KN) on ascorbate-glutathione cycle in tomato under NaCl toxicity. Data presented are the means \pm SE (n = 3). Different letters indicate significant difference at $P \leq 0.05$.

<https://doi.org/10.1371/journal.pone.0202175.g005>

Tissue Na, K and Ca

Salt treatment caused a 71.80% and 44.54% decline in the uptake of K^+ and Ca^{2+} , respectively. KN-supplemented salt-treated plants exhibited a reduction of 48.81% in Na^+ accumulation, which was reflected in a significant decline in the Na^+/K^+ ratio compared to that in the NaCl-treated plants (Table 3).




Fig 6. Effect of kinetin (Kn) on Gly I and Gly II, proline and glycine betaine accumulation in tomato under NaCl toxicity. Data presented are the means \pm SE (n = 3). Different letters indicate significant difference at $P \leq 0.05$.

<https://doi.org/10.1371/journal.pone.0202175.g006>

Table 3. Effect of kinetin (Kn) on Na, K and Ca (mg g⁻¹ DW) uptake in tomato under NaCl toxicity. Data presented are the means ± SE (n = 3). Different letters indicate significant difference at P ≤ 0.05.

Treatments	Na ⁺	K ⁺	Na/K ratio	Ca ²⁺
0	15.27±0.81c	31.11±1.39b	0.49±0.009c	5.32±0.032b
0+ Kn	14.19±0.76d	34.12±1.42a	0.47±0.008c	6.11±0.035a
150 mM NaCl	37.61±1.44a	8.77±0.66d	4.28±0.022a	2.95±0.019d
150 mM +Kn	19.25±0.96b	15.32±0.82c	1.25±0.011b	4.77±0.028c

<https://doi.org/10.1371/journal.pone.0202175.t003>

Discussion

Salt stress reduced the length and biomass accumulation of *S. lycopersicum*, whereas KN maintained and stimulated its growth under both normal and stress conditions. Salinity stress has been reported to hamper the export of cytokinins such as KN from the root to the shoot [43], thereby decreasing the shoot KN concentration required for normal growth and development [44]. In the present study, exogenously applied KN may have maintained cell division at an optimal level. Our results of amelioration of the salinity-induced decline in length and biomass by KN, are consistent with the findings of Iqbal and Ashraf [45] and Kaya et al. [46] who also reported cytokinin-induced growth promotion in wheat and maize respectively. In *Solanum melongena*, Singh and Prasad [21] have also demonstrated enhanced growth and mitigation of metal stress due to exogenous application of KN. The salinity induced reduction in growth was correlated with a significant reduction in the synthesis of chlorophyll pigments and decrease in membrane damage. Earlier, Dhindsa et al. [47] and Chen et al. [48], have demonstrated a significant effect of stress-triggered membrane leakage on chlorophyll biosynthesis. The results for salt-induced reduced chlorophyll contents corroborate with those of Rasool et al. [49] in *Cicer arietinum* and Ahanger and Agarwal [3] in *Triticum aestivum*. Higher concentrations of NaCl have been reported to reduce pigment synthesis and pigment protein complex functioning [50] in addition to reducing the *de novo* synthesis of proteins and associated chlorophyll components [51]. For example, Das et al. [52] have reported improved yield in mulberry after KN mediated enhancement in chlorophyll synthesis. Enough cytokinin concentration triggers etioplast-chloroplast transitions accompanied by the greater appearance of the metabolites implicated in the chlorophyll tetrapyrrole biosynthesis pathway [53], which probably is regulated by the effect of cytokinin on the expression of key chlorophyll biosynthesis genes including *HEMA1*, *GUN4*, *GUN5*, and *CHLM* [54].

Photosynthetic efficiency was reduced by NaCl-mediated decline in the chlorophyll fluorescence parameters, and KN proved ameliorative by increasing *Fv/Fm*, ϕ PSII, ϕ_{exc} , *qP* and decreasing NPQ. Stresses are believed to damage the antenna pigments thereby restricting the transfer of electrons from PSII to PSI [55]. Our results suggest that KN-mediated improved efficiency may largely depend on the protected functioning of the photosynthetic light reactions synchronized functionally with the enzymes of PSII and PSI. Moreover, functioning of the photosystems is coupled to generation of trans-thylakoid proton gradient creating pH difference for regulating transfer of electrons to PSI, and this transfer has been reported under the direct control of proton gradient regulation-5 (PGR5) protein [56]. It can be inferred that greater PSII activity in KN supplemented plants may have prevented the formation of singlet oxygen thereby protecting the chloroplast structure from the oxidative damage. Application of KN was reported to protect the pigment protein complex from high temperatures in *Zea mays* [57]. KN also nullified the effect of waterlogging [58] on the Rubisco and Hills activity, and of salinity [16] on leaf area, Hill's reaction and chloroplast structure resulting in a significant improvement in net carboxylation efficiency. KN application to *Solanum lycopersicum* was found to be effective in

maintaining and protecting CO_2 assimilation rate, g_s and E under normal as well as NaCl treated conditions, thereby leading to better growth performance. Application of phytohormones has been observed to assuage the damaging effects of NaCl on the photosynthetic efficiency significantly [59], and thus future research in this direction can be worthwhile. It has been opined that phytohormones can improve the synthesis of proteins and polysaccharides involved in photosynthetic regulation [60]. However, in the present study, KN-mediated synthesis of redox components may have contributed to photoprotection. In another study, KN was reported to trigger the synthesis of carotenoids, which protected the photosynthetic apparatus from excess ROS by up-regulating the activities of pigment-synthesizing enzymes coupled with reduction in the degrading enzymes, the activity of which is otherwise reported to be increased due to stress [61].

Reduced H_2O_2 accumulation, lipid peroxidation and electrolyte leakage in KN treated plants under saline stress were observed. Such a protective role of KN has been reported earlier also [62]. Siddiqui et al. [63] reported the amelioration of storage-induced degradation of cauliflower due to treatment with 6-benzylaminopurine. Salinity triggered membrane leakage and hence lipid peroxidation can be mitigated by KN supply [8,49,64]. However, KN mediated enhanced membrane integrity can be ascribed to its involvement in maintenance of tissue water content, the antioxidant system and peroxidation rates, which are greatly affected by stresses.

Application of KN promoted the activities of antioxidant enzymes and enhanced the contents of AsA and glutathione (GSH and GSSG), thereby imparting protection to *S. lycopersicum* plants against NaCl-induced oxidative stress. Cytokinin-induced up-regulation of antioxidant system has been reported by Wu et al. [20], Wang et al. [65]; and Siddiqui et al. [63]. During antioxidant dismutation of ROS, SOD eliminates superoxide radicals with CAT and APX carrying the further dismutation. KN-induced up-regulation of SOD may cause modulation of the substrate superoxide and H_2O_2 , thereby reducing the formation of more toxic hydroxyl (OH^\cdot) radicals [21]. Wang et al. [65] have demonstrated cytokinin-induced temperature stress tolerance in bentgrass through induction of the activities of antioxidant enzymes. The KN-induced accumulation of AsA, GSH and GSSG can protect NaCl-stressed tomato plants from ROS-induced damage. APX, GR, DHAR, MDHAR, AsA, and GSH are the components of the ROS scavenging pathway (the ascorbate-glutathione cycle) and KN-induced up-regulation of these components strengthens the tolerance strategies against any possible oxidative damage. For example, salt stressed *Solanum melongena* treated with exogenous cytokinin exhibited reduced ROS accumulation and greater protection to photosynthetic pathways leading to better yield productivity. Under high salinity, Mittal et al. [66] have reported enhanced growth in *Brassica juncea* accompanied by increased CAT activity. H_2O_2 produced by the SOD activity can be eliminated by either CAT in the cytoplasm or APX in the ascorbate–glutathione cycle, which comprises a series of redox reactions, involving the active engagement of ascorbate, glutathione, and NADPH [67]. APX plays an important role in scavenging H_2O_2 in chloroplasts and the cytosol thereby preventing its diffusion to other organelles to avert the damage. Optimal functioning of the AsA-GSH pathway due to exogenous KN maintained the redox components including AsA and glutathione, thus reducing the oxidative stress effects of salinity. The increased activity of antioxidants is correlated with the enhanced tolerance of plants to stress [3,68,69]. Up-regulation of the activities of DHAR and MDHAR due to KN application led to the optimization of AsA and glutathione integrating the functioning of other enzymatic components including APX and GR with the non-enzymatic components, therefore causing an overall effect on the H_2O_2 neutralization and the availability of NADP for greater protection of electron transport chain [70]. In the present study, protection to the photosynthetic electron transport chain by KN could have been due to precise regulation of $\text{NADP}^+/\text{NADPH}$ ratio, which prevents the flow of electrons to molecular oxygen

subsequently restricting the generation of superoxide radical [20]. Cortleven et al. [71] have demonstrated that *Arabidopsis* mutants deficient in cytokinin production display greater intensity of stress-induced photodamage by exhibiting a declined D1-repair, which was found correlated with the reduced synthesis of AsA and glutathione. The present investigation also infers the protective role of KN for photosynthetic system through the up-regulation of antioxidant system. However, further studies on investigating the expression patterns of antioxidant molecules as affected by KN application are required.

Higher concentrations of GSH in KN treated plants may have contributed significantly to the maintenance of the glyoxylase system for elimination of methylglyoxal, which may decline the chances of any serious genotoxic effect [72]. Glyoxylase I and II form the key enzymatic components of the glyoxylase system, and KN-induced up-regulation in their activities, accompanied by the increased GSH, may have led to exploit the beneficial effects of methylglyoxal like crosstalk with important signaling molecules like Ca, ROS and ABA [73]. Reports discussing the involvement of cytokinins in optimization of methylglyoxal are very scanty, and thus further analysis at proteomic or genomic levels shall be handy in further consolidations of any possible role in methylglyoxal metabolism. Glyoxylase I uses GSH as a cofactor in converting methylglyoxal to S-D-lactoylglutathione, and glyoxylase II generates GSH thereby contributing to redox homeostasis and protection against toxic species [74]. Transgenic plants exhibiting increased expression of glyoxylase I exhibit apparent improvement in stress tolerance via enhanced production of GSH [75]. Therefore, it could be inferred that enhancing endogenous GSH by the application of KN and the functioning of methylglyoxal scavenging system can improve tolerance potential of plants. For example, in salt stressed rice, Rahman et al. [12] have also demonstrated up-regulation of the glyoxylase system. Enhanced activity of the glyoxylase system due to exogenous application of KN may have protected the electron transport system by preventing damage to chloroplast and mitochondrial ultrastructures [72].

Antioxidant system was reinforced by the greater synthesis of phenols and flavonoids in the KN-treated plants (S1 Table). It has been reported that greater production of phenolics regulate plant developmental processes like lignin and pigment biosynthesis, thereby providing structural integrity and scaffolding support to the plants [76]. KN-induced improvement in accumulation of phenols enhances the possibility of crosstalk with the signaling molecules like nitric oxide for improving the efficiency of assimilatory pathways like nitrogen metabolism [77]. Ahanger and Agarwal [3] demonstrated that greater accumulation of phenolic compounds imparts better ROS scavenging potential, resulting in greater protection of membranes and nucleic acids, and at times particularly under stress can contribute to osmoregulation [69]. In the present study KN-mediated accumulation of phenols may have contributed to strengthening of cell wall structures and prevention of oxidative damage to membrane lipids and proteins by modulating their peroxidation kinetics [78]. For example, presence of cytokinins greatly affected the synthesis of phenols in *Tulbaghia simmleri* [79].

In our investigation, salt treatment triggered the accumulation of proline and glycine betaine, and further enhancement resulted due to exogenous application of KN. In salt-stressed *Morus alba*, Harinasut et al. [80] have also reported a manifold increase in the accumulation of proline. Accumulation of proline is attributable to the up-regulation of proline synthesizing enzymes and subsequent down-regulation of the catabolizing enzymes, or alternatively can be due to its decreased incorporation into proteins [8,81,82]. KN-induced enhancement in the accumulation of proline and glycine betaine to some extent might have helped tomato plants to avert the salinity-induced decline in tissue RWC. Enhanced RWC due to exogenous application of cytokinins in copper and zinc treated *Lupinus termis* [83] and Pb-stressed safflower [62] plants has been reported, which may be attributed to increased hydraulic conductivity which is however significantly affected by stresses [84]. Proline and glycine betaine maintain

water balance of plants thereby minimizing the deleterious effects of stresses on metabolism [6] particularly by protecting protein turnover, enzyme activities and the expression of stress-protective proteins [3,5,69,85]. KN mediated increased proline and glycine betaine synthesis under normal and NaCl-stressed conditions is an indicative of its protective role for metabolically important pathways by bringing osmoregulation, though the role of exogenously applied cytokinin in regulating osmotic balance is yet inconclusive (S1 Table). Therefore, KN-mediated increase in tissue water content resulted in maintained cell wall extensibility and cell division, which cause increased growth and biomass accumulation.

Exposure of *S. lycopersicum* plants to NaCl stress impeded the uptake of essential mineral elements such as K and Ca. Excess uptake of Na⁺ intensifies its antagonistic relationship with important ions, including K⁺. The salinity-induced reduction in ion uptake is in agreement with Garg and Manchanda [86] in *Cajanus cajan*, Kohler et al. [87] in lettuce and Ahmad et al. [88] in *Brassica juncea*. Exogenous application of KN not only prevented Na toxicity by reducing its uptake, but it also caused a significant increase in the uptake of K⁺ reflecting in declined Na⁺/K⁺ ratio, an important attribute for regulating major cellular pathways and protecting them from Na toxicity [5,88]. Reduced Na⁺/K⁺ ratio results from the efficient exclusion of toxic ions leads to maintenance of osmotic potential and hence stress tolerance [89], and tissue Ca²⁺ concentration facilitates K⁺/Na⁺ selectivity [90]. During NaCl stress, cytosolic Ca²⁺ is maintained at high concentrations, as a consequence of transport from the apoplast and the intracellular compartments [91]. Such transient increases in cytosolic Ca²⁺ due to KN may contribute to initiate the signal transduction pathways that result in enhanced salt tolerance. Sodium chloride (NaCl) competes with Ca²⁺ at membrane binding sites [92]. Foliar application of KN to tomato plants might have mitigated the reduction in K and Ca uptake caused by salinity, resulting in increased K⁺/Na⁺ and Ca²⁺/Na⁺ ratios and plant performance by regulating the enzyme activity, membrane stability and stomatal functioning[2].

Conclusions

On the basis of the results of this study, it can be concluded that exogenously applied KN resulted in enhanced growth performance of *S. lycopersicum* plants under NaCl stress by up-regulating antioxidant metabolism and osmolyte accumulation. Reduced ROS production in parallel with reduced lipid peroxidation in KN-supplemented plants resulted in the maintenance of cellular functioning, greater photoprotection and mineral uptake, which caused enhanced protection of photosynthetic pigments. KN-mediated growth promotion under stressed conditions may be ascribed to strengthening of the ROS and methylglyoxal detoxifying systems by way of integration between different signaling components. Overall, these observations indicate the suitability of foliar application of KN for exploiting the genetic potential of *S. lycopersicum* under NaCl stress.

Supporting information

S1 Table. Effect of kinetin (KN) on total phenol and flavonoids in tomato under NaCl toxicity. Data presented are the means \pm SE (n = 3). Different letters indicate significant difference at $P \leq 0.05$.
(DOCX)

S1 Dataset. individual level data. Effect of Kn on growth, biomass yield, biochemical attributes, lipid peroxidation, antioxidants and secondary metabolites in tomato Under NaCl stress.
(XLSX)

Acknowledgments

The authors would like to extend their sincere appreciation to the Deanship of Scientific Research at King Saud University for funding this Research group NO.(RGP-199).

Author Contributions

Data curation: Mohammed Nasser Alyemini, Leonard Wijaya.

Formal analysis: Mohammed Nasser Alyemini, Leonard Wijaya, Pravej Alam.

Funding acquisition: Pravej Alam.

Investigation: Mohammad Abass Ahanger.

Methodology: Mohammad Abass Ahanger.

Software: Mohammed Nasser Alyemini, Leonard Wijaya, Pravej Alam.

Supervision: Parvaiz Ahmad.

Validation: Leonard Wijaya, Muhammad Ashraf, Parvaiz Ahmad.

Writing – original draft: Mohammad Abass Ahanger, Parvaiz Ahmad.

Writing – review & editing: Saud A. Alamri, Muhammad Ashraf.

References

1. Khan MIR, Asgher M, Khan NA (2014). Alleviation of salt-induced photosynthesis and growth inhibition by salicylic acid involves glycinebetaine and ethylene in mungbean (*Vigna radiata* L.). *Plant Physiol Biochem.* 80: 67–74. <https://doi.org/10.1016/j.plaphy.2014.03.026> PMID: 24727790
2. Ahmad P, Abdel Latef AA, Hashem A, Abd_Allah EF, Guzel S, Tran LSP (2016). Nitric oxide mitigates salt stress by regulating levels of osmolytes and antioxidant enzymes in chickpea. *Front Plant Sci.* 7: 347. <https://doi.org/10.3389/fpls.2016.00347> PMID: 27066020
3. Ahanger MA, Agarwal RM (2017). Salinity stress induced alterations in antioxidant metabolism and nitrogen assimilation in wheat (*Triticum aestivum* L) as influenced by potassium supplementation. *Plant Physiol Biochem.* 115: 449–460. <https://doi.org/10.1016/j.plaphy.2017.04.017> PMID: 28478373
4. Ahanger MA, Agarwal RM (2016). Potassium up-regulates antioxidant metabolism and alleviates growth inhibition under water and osmotic stress in wheat (*Triticum aestivum* L). *Protoplasma* 254: 1471–1486. <https://doi.org/10.1007/s00709-016-1037-0>
5. Ahanger MA, Agarwal RM, Tomar NS, Shrivastava M (2015). Potassium induces positive changes in nitrogen metabolism and antioxidant system of oat (*Avena sativa* L. cultivar Kent). *J Plant Interact.* 10: 211–223. <https://doi.org/10.1080/17429145.2015.1056260>
6. Ahanger MA, Tyagi SR, Wani MR, Ahmad P (2014). Drought Tolerance: Role of Organic Osmolytes, Growth Regulators, and Mineral Nutrients. In: Ahmad P, Wani MR, editors. *Physiological Mechanisms and Adaptation Strategies in Plants Under Changing Environment*: Springer New York. pp. 25–55.
7. Munns R, Tester M (2008). Mechanisms of Salinity Tolerance. *Ann Rev Plant Biol.* 59: 651–681. <https://doi.org/10.1146/annurev.arplant.59.032607.092911> PMID: 18444910
8. Ahmad P (2010). Growth and antioxidant responses in mustard (*Brassica juncea* L.) plants subjected to combined effect of gibberellic acid and salinity. *Arch Agron Soil Sci.* 56: 575–588. <https://doi.org/10.1080/03650340903164231>
9. Ashraf M, Foolad MR (2007). Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environ Exp Bot.* 59: 206–216. <https://doi.org/10.1016/j.envexpbot.2005.12.006>
10. Murata N, Mohanty PS, Hayashi H, Papageorgiou GC (1992). Glycinebetaine stabilizes the association of extrinsic proteins with the photosynthetic oxygen-evolving complex. *FEBS Lett.* 296: 187–189. [https://doi.org/10.1016/0014-5793\(92\)80376-r](https://doi.org/10.1016/0014-5793(92)80376-r) PMID: 1733776
11. Khan MIR, Nazir F, Asgher M, Per TS, Khan NA (2015). Selenium and sulfur influence ethylene formation and alleviate cadmium-induced oxidative stress by improving proline and glutathione production in wheat. *J Plant Physiol.* 173: 9–18. <https://doi.org/10.1016/j.jplph.2014.09.011> PMID: 25462073

12. Rahman A, Nahar K, Hasanuzzaman M, Fujita M (2016). Calcium supplementation improves Na⁺/K⁺ ratio, antioxidant defense and glyoxalase systems in salt-stressed rice seedlings. *Front Plant Sci.* 7: 609. <https://doi.org/10.3389/fpls.2016.00609> PMID: 27242816
13. Hamayun M, Hussain A, Khan SA, Irshad M, Khan AL, et al. (2015). Kinetin modulates physio-hormonal attributes and isoflavone contents of soybean grown under salinity stress. *Front Plant Sci.* 6: 377. <https://doi.org/10.3389/fpls.2015.00377> PMID: 26082785
14. Al-Hakimi AMA (2007) Modification of cadmium toxicity in pea seedlings by kinetin. *Plant Soil Environ.* 53: 129–135. <https://doi.org/10.17221/2228-pse>
15. Gangwar S, Singh VP, Prasad SM, Maurya JN (2010). Modulation of manganese toxicity in *Pisum sativum* L. seedlings by kinetin. *Sci Horticult.* 126: 467–474. <https://doi.org/10.1016/j.scienta.2010.08.013>
16. Aldesuquy H, Baka Z, Mickky B (2014) Kinetin and spermine mediated induction of salt tolerance in wheat plants: Leaf area, photosynthesis and chloroplast ultrastructure of flag leaf at ear emergence. *Egypt J Basic Appl Sci.* 1: 77–87. <https://doi.org/10.1016/j.ejbas.2014.03.002>
17. Letham DS, Palni LMS (1983). The biosynthesis and metabolism of cytokinins. *Ann. Rev Plant Physiol.* 34: 163–197. <https://doi.org/10.1146/annurev.pp.34.060183.001115>
18. Zwack PJ, Rashotte AM (2015). Interactions between cytokinin signalling and abiotic stress responses. *J Exp Bot.* 66: 4863–4871. <https://doi.org/10.1093/jxb/erv172> PMID: 25911740
19. Younis ME, El-Shahaby OA, Nemat Alla MN, El-Bastawisy ZM (2003). Kinetin alleviates the influence of waterlogging and salinity on growth and affects the production of plant growth regulators in *Vigna sinensis* and *Zea mays*. *Agronomie* 23: 277–285. <https://doi.org/10.1051/agro:2003010>
20. Wu X, Zhu Z, Li X, Zha D (2012). Effects of cytokinin on photosynthetic gas exchange, chlorophyll fluorescence parameters and antioxidative system in seedlings of eggplant (*Solanum melongena* L.) under salinity stress. *Acta Physiol Plant.* 34: 2105–2114. <https://doi.org/10.1007/s11738-012-1010-2>
21. Singh S, Prasad SM (2014). Growth, photosynthesis and oxidative responses of *Solanum melongena* L. seedlings to cadmium stress: Mechanism of toxicity amelioration by kinetin. *Sci Horticult.* 176: 1–10. <https://doi.org/10.1016/j.scienta.2014.06.022>
22. Shabala S, Pang J, Zhou M, Shabala L, Cui TA, Nick P, et al. (2009). Electrical signalling and cytokinins mediate effects of light and root cutting on ion uptake in intact plants. *PlantCell Environ.* 32: 194–207. <https://doi.org/10.1111/j.1365-3040.2008.01914.x> PMID: 19021884
23. Gerszberg A, Hnatuszko-Konka K, Kowalczyk T, Kononowicz AK (2015). Tomato (*Solanum lycopersicum* L.) in the service of biotechnology. *Plant Cell Tissue Org Cult.* 120: 881–902. <https://doi.org/10.1007/s11240-014-0664-4>
24. Arnon DI (1949) Copper enzymes in isolated chloroplasts, polyphenoxidase in *Beta vulgaris*. *Plant Physiol.* 24: 1–15. <https://doi.org/10.1104/pp.24.1.1> PMID: 16654194
25. Li PM, Cai RG, Gao HY, Peng T, Wang ZL (2007). Partitioning of excitation energy in two wheat cultivars with different grain protein contents grown under three nitrogen applications in the field. *Physiol Plant.* 129: 822–829. <https://doi.org/10.1111/j.1399-3054.2007.00880.x>
26. Smart RE, Bingham GE (1974). Rapid estimates of relative water content. *Plant Physiol.* 53: 258–260. <https://doi.org/10.1104/pp.53.2.258> PMID: 16658686
27. Velikova V, Yordanov I, Edreva A (2000). Oxidative stress and some antioxidant systems in acid rain-treated bean plants. *Plant Sci.* 151: 59–66. [https://doi.org/10.1016/s0168-9452\(99\)00197-1](https://doi.org/10.1016/s0168-9452(99)00197-1)
28. Madhava Rao KV, Sresty TVS (2000). Antioxidative parameters in the seedlings of pigeonpea (*Cajanus cajan* (L.) Millspaugh) in response to Zn and Ni stresses. *Plant Sci.* 157: 113–128. [https://doi.org/10.1016/s0168-9452\(00\)00273-9](https://doi.org/10.1016/s0168-9452(00)00273-9) PMID: 10940475
29. Dionisio-Sese ML, Tobita S (1998). Antioxidant responses of rice seedlings to salinity stress. *Plant Sci.* 135: 1–9. [https://doi.org/10.1016/s0168-9452\(98\)00025-9](https://doi.org/10.1016/s0168-9452(98)00025-9)
30. Dhindsa RS, Matowe W (1981). Drought tolerance in two mosses: Correlated with enzymatic defence against lipid peroxidation. *J Exp Bot.* 32: 79–91. <https://doi.org/10.1093/jxb/32.1.79>
31. Aebi H (1984) Catalase in vitro. *Methods in enzymology*: Elsevier. pp. 121–126.
32. Hasanuzzaman M, Fujita M (2013). Exogenous sodium nitroprusside alleviates arsenic-induced oxidative stress in wheat (*Triticum aestivum* L.) seedlings by enhancing antioxidant defense and glyoxalase system. *Ecotoxicol.* 22: 584–596. <https://doi.org/10.1007/s10646-013-1050-4> PMID: 23430410
33. Nakano Y, Asada K (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.* 22: 867–880. <https://doi.org/10.1093/oxfordjournals.pcp.a076232>
34. Foster JG, Hess JL (1980). Responses of superoxide dismutase and glutathione reductase activities in cotton leaf tissue exposed to an atmosphere enriched in oxygen. *Plant Physiol.* 66: 482–487. <https://doi.org/10.1104/pp.66.3.482> PMID: 16661460

35. Miyake C, Asada K (1992). Thylakoid-bound ascorbate peroxidase in spinach chloroplasts and photoreduction of its primary oxidation product monodehydroascorbate radicals in thylakoids. *Plant Cell Physiol.* 33: 541–553. doi.org/10.1093/oxfordjournals.pcp.a078288
36. Huang C, He W, Guo J, Chang X, Su P, Zhang L (2005). Increased sensitivity to salt stress in an ascorbate-deficient Arabidopsis mutant. *J Exp Bot.* 56: 3041–3049. <https://doi.org/10.1093/jxb/eri301> PMID: [16263910](https://pubmed.ncbi.nlm.nih.gov/16263910/)
37. Yu CW, Murphy TM, Lin CH (2003). Hydrogen peroxide-induced chilling tolerance in mung beans mediated through ABA-independent glutathione accumulation. *Funct.Plant Biol.* 30: 955. <https://doi.org/10.1071/fp03091>
38. Paradiso A, Berardino R, de Pinto MC, Sanita di Toppi L, Storelli MM, Tommasi F, et al. (2008). Increase in ascorbate-glutathione metabolism as local and precocious systemic responses induced by cadmium in durum wheat plants. *Plant Cell Physiol.* 49: 362–374. <https://doi.org/10.1093/pcp/pcn013> PMID: [18234716](https://pubmed.ncbi.nlm.nih.gov/18234716/)
39. Chun OK, Kim DO, Lee CY (2003). Superoxide radical scavenging activity of the major polyphenols in fresh plums. *J Agricul Food Chem.* 51: 8067–8072. <https://doi.org/10.1021/jf034740d> PMID: [14690398](https://pubmed.ncbi.nlm.nih.gov/14690398/)
40. Zhishen J, Mengcheng T, Jianming W (1999). The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 64: 555–559. [https://doi.org/10.1016/S0308-8146\(98\)00102-2](https://doi.org/10.1016/S0308-8146(98)00102-2)
41. Bates LS, Waldren RP, Teare ID (1973). Rapid determination of free proline for water-stress studies. *Plant Soil* 39: 205–207. <https://doi.org/10.1007/bf00018060>
42. Grieve CM, Grattan SR (1983). Rapid assay for determination of water soluble quaternary ammonium compounds. *Plant Soil* 70: 303–307. <https://doi.org/10.1007/bf02374789>
43. Kuiper D, Schuit J, Kuiper PJC (1990). Actual cytokinin concentrations in plant tissue as an indicator for salt resistance in cereals. *Plant Soil* 123: 243–250. <https://doi.org/10.1007/bf00011276>
44. Brault M, Maldiney R (1999). Mechanisms of cytokinin action. *Plant Physiol. Biochem.* 37: 403–412. [https://doi.org/10.1016/S0981-9428\(99\)80046-1](https://doi.org/10.1016/S0981-9428(99)80046-1)
45. Iqbal M, Ashraf M (2005). Presowing seed treatment with cytokinins and its effect on growth, photosynthetic rate, ionic levels and yield of two wheat cultivars differing in salt tolerance. *J Integ Plant Biol.* 47: 1315–1325. <https://doi.org/10.1111/j.1744-7909.2005.00163.x>
46. Kaya C, Tuna AL, Okant AM (2010). Effect of foliar applied kinetin and indole acetic acid on maize plants grown under saline conditions. *Turk J Agricul Forest.* 34: 529–538. <https://doi.org/10.3906/tar-0906-173>
47. Dhindsa RS, Plumb-Dhindsa P, Thorpe TA (1981). Leaf Senescence: Correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J Exp Bot.* 32: 93–101. <https://doi.org/10.1093/jxb/32.1.93>
48. Chen CT, Li CC, Kao CH (1991). Senescence of rice leaves XXXI. Changes of chlorophyll, protein, and polyamine contents and ethylene production during senescence of a chlorophyll-deficient mutant. *J Plant Growth Regul.* 10: 201–205. <https://doi.org/10.1007/bf02279335>
49. Rasool S, Ahmad A, Siddiqi TO, Ahmad P (2013). Changes in growth, lipid peroxidation and some key antioxidant enzymes in chickpea genotypes under salt stress. *Acta Physiol Plant.* 35: 1039–1050. <https://doi.org/10.1007/s11738-012-1142-4>
50. El-Tayeb MA (2005). Response of barley grains to the interactive effect of salinity and salicylic acid. *Plant Growth Regul.* 45: 215–224. <https://doi.org/10.1007/s10725-005-4928-1>
51. Ahanger MA, Morad-Talab N, Abd-Allah EF, Ahmad P, Hajiboland R (2016). Plant growth under drought stress. *Water Stress and Crop Plants: John Wiley & Sons, Ltd.* pp. 649–668.
52. Das C, Sengupta T, Chattopadhyay S, Setua M, Das NK, Saratchandra B (2002). Involvement of kinetin and spermidine in controlling salinity stress in mulberry (*Morus alba* L. cv. S1). *Acta Physiol Plant.* 24: 53–57. <https://doi.org/10.1007/s11738-002-0021-9>
53. Yaronskaya E, Vershilovskaya I, Poers Y, Alawady AE, Averina N, Grimm B (2006). Cytokinin effects on tetrapyrrole biosynthesis and photosynthetic activity in barley seedlings. *Planta* 224: 700–709. <https://doi.org/10.1007/s00425-006-0249-5> PMID: [16506064](https://pubmed.ncbi.nlm.nih.gov/16506064/)
54. Cortleven A, Marg I, Yamburenko MV, Schlicke H, Hill K, Grimm B et al. (2016). Cytokinin regulates the etioplast-chloroplast transition through the two-component signaling system and activation of chloroplast-related genes. *Plant Physiol.* 172: 464–478. <https://doi.org/10.1104/pp.16.00640> PMID: [27388681](https://pubmed.ncbi.nlm.nih.gov/27388681/)
55. Roach T, Krieger-Liszky A (2014). Regulation of photosynthetic electron transport and photoinhibition. *Curr Protein Peptide Sci.* 15: 351–362. <https://doi.org/10.2174/1389203715666140327105143> PMID: [24678670](https://pubmed.ncbi.nlm.nih.gov/24678670/)

56. Tikkanen M, Rantala S, Aro E-M (2015). Electron flow from PSII to PSI under high light is controlled by PGR5 but not by PSBS. *Front Plant Sci.* 6: 521. <https://doi.org/10.3389/fpls.2015.00521> PMID: [26217370](https://pubmed.ncbi.nlm.nih.gov/26217370/)
57. Bako SP (2006). Effects of plant age, ascorbate and kinetin applications on integrity of the photosynthetic pigment complex in maize (*Zea mays* L.) plants grown under heat stress. *Asian J Plant Sci.* 5: 357–362. <https://doi.org/10.3923/ajps.2006.357.362>
58. El-Shihaby OA, Younis ME, El-Bastawisy ZM, Nemat Alla MM (2002). Effect of kinetin on photosynthetic activity and carbohydrate content in waterlogged or seawater-treated *Vigna sinensis* and *Zea mays* plants. *Plant Biosyst.* 136: 277–290. <https://doi.org/10.1080/11263500212331351189>
59. Fatma M, Masood A, Per TS, Khan NA (2016). Nitric oxide alleviates salt stress inhibited photosynthetic performance by interacting with sulfur assimilation in mustard. *Front Plant Sci.* 7: 521. <https://doi.org/10.3389/fpls.2016.00521> PMID: [27200007](https://pubmed.ncbi.nlm.nih.gov/27200007/)
60. Aldesuquy H (2015). Synergistic effect of phytohormones on pigment and fine structure of chloroplasts in flag leaf of wheat plants irrigated by seawater. *Egypt J Basic Appl Sci.* 2: 310–317. <https://doi.org/10.1016/j.ejbas.2015.07.002>
61. Behera RK, Mishra PC, Choudhury NK (2002). High irradiance and water stress induce alterations in pigment composition and chloroplast activities of primary wheat leaves. *J Plant Physiol.* 159: 967–973. <https://doi.org/10.1078/0176-1617-00823>
62. Sayed SA (1999). Effects of lead and kinetin on the growth, and some physiological components of safflower. *Plant Growth Regul.* 29: 167–174. <https://doi.org/10.1023/A:1006216630915>
63. Siddiqui MW, Singh JP, Nayyar MA, Barman K, Ahmad MS, Kumar V (2015). 6-Benzylaminopurine affects lipid peroxidation and membrane permeability and thereby preserves curd quality and antioxidants during storage of cauliflower. *Acta Physiol Plant.* 37: 96. <https://doi.org/10.1007/s11738-015-1848-1>
64. Tuna AL, Kaya C, Dikilitas M, Higgs D (2008). The combined effects of gibberellic acid and salinity on some antioxidant enzyme activities, plant growth parameters and nutritional status in maize plants. *Environ Exp Bot.* 62: 1–9. <https://doi.org/10.1016/j.envexpbot.2007.06.007>
65. Wang Z, Pote J, Huang B (2003). Responses of cytokinins, antioxidant enzymes, and lipid peroxidation in shoots of creeping bentgrass to high root-zone temperatures. *J Amer Soc Horticult Sci.* 128: 648–655.
66. Mittal S, Kumari N, Sharma V (2012). Differential response of salt stress on *Brassica juncea*: Photosynthetic performance, pigment, proline, D1 and antioxidant enzymes. *Plant Physiol Biochem.* 54: 17–26. <https://doi.org/10.1016/j.plaphy.2012.02.003> PMID: [22369937](https://pubmed.ncbi.nlm.nih.gov/22369937/)
67. Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trend Plant Sci.* 7: 405–410. [https://doi.org/10.1016/s1360-1385\(02\)02312-9](https://doi.org/10.1016/s1360-1385(02)02312-9)
68. Ahmad P, Nabi G, Ashraf M (2011). Cadmium-induced oxidative damage in mustard [*Brassica juncea* (L.) Czern. & Coss.] plants can be alleviated by salicylic acid. *South Afr J Bot.* 77: 36–44. <https://doi.org/10.1016/j.sajb.2010.05.003>
69. Ahanger MA, Tomar NS, Tittal M, Argal S, Agarwal RM (2017). Plant growth under water/salt stress: ROS production; antioxidants and significance of added potassium under such conditions. *Physiol Mol Biol Plant* 23: 731–744. <https://doi.org/10.1007/s12298-017-0462-7> PMID: [29158624](https://pubmed.ncbi.nlm.nih.gov/29158624/)
70. Foyer CH, Noctor G (2011). Ascorbate and glutathione: The heart of the redox hub. *Plant Physiol.* 155: 2–18. <https://doi.org/10.1104/pp.110.167569> PMID: [21205630](https://pubmed.ncbi.nlm.nih.gov/21205630/)
71. Cortleven A, Nitschke S, Klaumunzer M, AbdElgawad H, Asard H, Grimm B, et al. (2014). A novel protective function for cytokinin in the light stress response is mediated by the ARABIDOPSIS HISTIDINE KINASE2 and ARABIDOPSIS HISTIDINE KINASE3 receptors. *Plant Physiol.* 164: 1470–1483. <https://doi.org/10.1104/pp.113.224667> PMID: [24424319](https://pubmed.ncbi.nlm.nih.gov/24424319/)
72. Gupta BK, Sahoo KK, Ghosh A, Tripathi AK, Anwar K, Das P, et al. (2017). Manipulation of glyoxalase pathway confers tolerance to multiple stresses in rice. *Plant Cell Environ.* 41: 1186–1200. <https://doi.org/10.1111/pce.12968> PMID: [28425127](https://pubmed.ncbi.nlm.nih.gov/28425127/)
73. Li ZG (2016). Methylglyoxal and glyoxalase system in plants: old players, new concepts. *The Bot Rev.* 82: 183–203. <https://doi.org/10.1007/s12229-016-9167-9>
74. Yadav SK, Singla-Pareek SL, Reddy M, Sopory SK (2005). Methylglyoxal detoxification by glyoxalase system: A survival strategy during environmental stresses. *Physiol Mol Biol Plant.* 11: 1.
75. Yadav SK, Singla-Pareek SL, Ray M, Reddy MK, Sopory SK (2005). Methylglyoxal levels in plants under salinity stress are dependent on glyoxalase I and glutathione. *Biochem Biophys Res Commun* 337: 61–67. <https://doi.org/10.1016/j.bbrc.2005.08.263> PMID: [16176800](https://pubmed.ncbi.nlm.nih.gov/16176800/)
76. Bhattacharya A, Sood P, Citovsky V (2010) The roles of plant phenolics in defence and communication during *Agrobacterium* and *Rhizobium* infection. *Mol Plant Pathol.* 11: 705–719. <https://doi.org/10.1111/j.1364-3703.2010.00625.x> PMID: [20696007](https://pubmed.ncbi.nlm.nih.gov/20696007/)

77. Klein A, Keyster M, Ludidi N (2015). Response of soybean nodules to exogenously applied caffeic acid during NaCl-induced salinity. *South Afr J Bot.* 96: 13–18. <https://doi.org/10.1016/j.sajb.2014.10.016>
78. Oteiza PI, Erlejman AG, Verstraeten SV, Keen CL, Fraga CG (2005). Flavonoid-membrane Interactions: A protective role of flavonoids at the membrane surface? *Clin Develop Immunol* 12: 19–25. <https://doi.org/10.1080/10446670410001722168> PMID: 15712595
79. Aremu AO, Plačková L, Gruz J, Bíba O, Šubrtová M, Novak O, et al. (2015). Accumulation pattern of endogenous cytokinins and phenolics in different organs of 1-year-old cytokinin pre-incubated plants: implications for conservation. *Plant Biol.* 17: 1146–1155. <https://doi.org/10.1111/plb.12367> PMID: 26177040
80. Harinasut P, Srisunak S, Pitukchaisopol S, Charoensataporn R (2000). Mechanisms of adaptation to increasing salinity of mulberry: Proline content and ascorbate peroxidase activity in leaves of multiple shoots. *Sci Asia* 26: 207–211.
81. Delauney AJ, Verma DPS (1993). Proline biosynthesis and osmoregulation in plants. *Plant J* 4: 215–223. <https://doi.org/10.1046/j.1365-313x.1993.04020215.x>
82. Iqbal N, Umar S, Khan NA (2015). Nitrogen availability regulates proline and ethylene production and alleviates salinity stress in mustard (*Brassica juncea*). *J Plant Physiol.* 178: 84–91. <https://doi.org/10.1016/j.jplph.2015.02.006> PMID: 25800225
83. Gadallah M, El-Enany A (1999) Role of kinetin in alleviation of copper and zinc toxicity in *Lupinus termis* plants. *Plant Growth Regul.* 29: 151–160. <https://doi.org/10.1023/A:1006245628188>
84. Ehlert C, Maurel C, Tardieu F, Simonneau T (2009). Aquaporin-mediated reduction in maize root hydraulic conductivity impacts cell turgor and leaf elongation even without changing transpiration. *Plant Physiol.* 150: 1093–1104. <https://doi.org/10.1104/pp.108.131458> PMID: 19369594
85. Thakur M, Sharma AD (2005). Salt-stress-induced proline accumulation in germinating embryos: Evidence suggesting a role of proline in seed germination. *J Arid Environ.* 62: 517–523. <https://doi.org/10.1016/j.jaridenv.2005.01.005>
86. Garg N, Manchanda G (2009). Role of arbuscular mycorrhizae in the alleviation of ionic, osmotic and oxidative stresses induced by salinity in *Cajanus cajan* (L.) Millsp. (pigeonpea). *J Agron Crop Sci.* 195: 110–123. <https://doi.org/10.1111/j.1439-037x.2008.00349.x>
87. Kohler J, Hernández JA, Caravaca F, Roldán A (2009). Induction of antioxidant enzymes is involved in the greater effectiveness of a PGPR versus AM fungi with respect to increasing the tolerance of lettuce to severe salt stress. *Environ Exp Bot.* 65: 245–252. <https://doi.org/10.1016/j.envexpbot.2008.09.008>
88. Ahmad P, Hashem A, Abd-Allah EF, Alqarawi AA, John R, Egamberdieva D, et al. (2015). Role of *Trichoderma harzianum* in mitigating NaCl stress in Indian mustard (*Brassica juncea* L.) through antioxidative defense system. *Frontiers in Plant Science* 6: 868. <https://doi.org/10.3389/fpls.2015.00868> PMID: 26528324
89. Wu QS, Zou YN, Liu W, Ye XF, Zai HF, Zhao LJ (2010). Alleviation of salt stress in citrus seedlings inoculated with mycorrhiza: changes in leaf antioxidant defense systems. *Plant Soil Environ.* 56: 470–475. <https://doi.org/10.17221/54/2010-pse>
90. Liu J, Zhu J (1998). A calcium sensor homolog required for plant salt tolerance. *Science* 280: 1943–1945. <https://doi.org/10.1126/science.280.5371.1943> PMID: 9632394
91. Knight H, Trewavas AJ, Knight MR (1997). Calcium signalling in *Arabidopsis thaliana* responding to drought and salinity. *Plant J.* 12: 1067–1078. <https://doi.org/10.1046/j.1365-313x.1997.12051067.x> PMID: 9418048
92. Davenport RJ, Reid RJ, Smith FA (1997). Sodium-calcium interactions in two wheat species differing in salinity tolerance. *Physiol Plant.* 99: 323–327. <https://doi.org/10.1034/j.1399-3054.1997.990216.x>