



The Open Neurology Journal

Content list available at: www.benthamopen.com/TONEUJ/

DOI: 10.2174/1874205X01610010042



RESEARCH ARTICLE

Potential Role of Caffeine in the Treatment of Parkinson's Disease

Mohsin H.K. Roshan^{*}, Amos Tambo and Nikolai P. Pace*Department of Physiology and Biochemistry, Faculty of Medicine and Surgery, University of Malta- Msida, Malta*

Received: February 24, 2016

Revised: April 30, 2016

Accepted: May 2, 2016

Abstract: Parkinson's disease [PD] is the second most common neurodegenerative disorder after Alzheimer's disease, affecting 1% of the population over the age of 55. The underlying neuropathology seen in PD is characterised by progressive loss of dopaminergic neurons in the substantia nigra pars compacta with the presence of Lewy bodies. The Lewy bodies are composed of aggregates of α -synuclein. The motor manifestations of PD include a resting tremor, bradykinesia, and muscle rigidity. Currently there is no cure for PD and motor symptoms are treated with a number of drugs including levodopa [L-dopa]. These drugs do not delay progression of the disease and often provide only temporary relief. Their use is often accompanied by severe adverse effects. Emerging evidence from both *in vivo* and *in vitro* studies suggests that caffeine may reduce parkinsonian motor symptoms by antagonising the adenosine A_{2A} receptor, which is predominately expressed in the basal ganglia. It is hypothesised that caffeine may increase the excitatory activity in local areas by inhibiting the astrocytic inflammatory processes but evidence remains inconclusive. In addition, the co-administration of caffeine with currently available PD drugs helps to reduce drug tolerance, suggesting that caffeine may be used as an adjuvant in treating PD. In conclusion, caffeine may have a wide range of therapeutic effects which are yet to be explored, and therefore warrants further investigation in randomized clinical trials.

Keywords: Adenosine A_{2A} receptors, Basal ganglia, Caffeine, Dopamine D_2 receptor, Parkinson's disease, α -synuclein.

1. INTRODUCTION

Parkinson's disease [PD] was named after James Parkinson, an English physician who described it as *Shaking Palsy* in 1817 [1]. It is the second most common neurodegenerative disorder after Alzheimer's Disease [AD], affecting 1% of population over the age of 55 [2, 3]. Clinical manifestations of PD includes a triad of motor symptoms (bradykinesia, rigidity, and resting tremor) and a range of neuropsychiatric and autonomic manifestations [4]. The precise aetiology of PD is unknown but studies have shown it to be multifactorial with an underlying genetic and environmental component. A number of single gene defects have been identified and these account for only 10% of PD cases; with sporadic forms accounting for the remaining cases [5]. The underlying neuropathology seen in PD is characterised by progressive loss of dopaminergic neurons in the substantia nigra pars compacta [SN_c] with the presence of Lewy bodies [LBs] and Lewy neurites [LNs] which are primarily composed of aggregates of alpha-synuclein [α -synuclein] [6].

Emerging data from epidemiological studies suggests that PD is less common in caffeine users compared to the general population [7, 8]. Furthermore, both *in vitro* and animal studies indicate that caffeine in low to moderate doses has neuroprotective effects [9]. Data suggests that caffeine antagonises adenosine A_{2A} receptors as well as prevents aggregation of α -synuclein [10]. This literature review aims to explore the neurophysiological effects of caffeine and its potential role as an adjuvant in the pharmacotherapy of PD.

1.1. Risk Factors

Although the aetiology of PD remains unknown, there are a number of important risk factors that play an important role in the development of PD.

* Address correspondence to this author at the Department of Physiology and Biochemistry, Faculty of Medicine and Surgery, University of Malta- Msida, Malta; Tel: +35699358550; E-mail: mohsin.roshan.13@um.edu.mt

1.1.1. Age and Gender

PD mostly affects individuals over the age of 50, although rare exceptions in younger individuals are reported. The risk of developing PD increases with age due to changes in neuronal circuitry and an increase in inflammatory response in neurons. Age-related cognitive decline and olfactory dysfunction are changes observed in approximately 80% of individuals in the early stages of the disease [11 - 13]. Such changes are associated with decreased neuronal density and cholinergic and serotonergic neurotransmitter concentrations in areas such as the locus coeruleus which may attribute to loss of olfaction [14]. In addition, PD is more prevalent in males due to influences of genetic, environmental, and physiological factors. For instance, since the farming industry is male-dominated, a higher proportion of male farmers are likely to be exposed to high levels of pesticides, which may trigger PD-inducing neurotoxicity.

Genomic imprinting inactivates one of the two X-chromosomes in females through methylation. This process may silence a number of genes involved in the pathogenesis of PD on the X-chromosome, halving the effect of any risk alleles and thus reducing susceptibility to PD [15, 16]. In addition, higher concentrations of endogenous oestrogen in females may have neuroprotective effects since endogenous oestrogen prevents neurotoxin build-up and loss of dopaminergic neurons in the SNC. This effect is not observed in males since oestrogen levels in healthy males are relatively lower than in females [17]. Nevertheless, the protective effect of oestrogen diminishes in postmenopausal women, and their risk of developing PD is then equivalent to males [18].

1.1.2. Family History

Monogenic forms account for less than 5% of cases of PD. Genomic variants in at least seven genes are associated with both autosomal dominant and autosomal recessive forms of PD. These genes (*SNCA*, *LRRK2*, *EIF4G1*, *VPS35*, *parkin*, *PARK2*, *PINK1* and *DJ1/PARK7*) are reviewed extensively elsewhere [5, 19].

For instance, α -synuclein (*SNCA*) is central to both familial and sporadic PD. A number of missense mutations in *SNCA* (A30P, E46K, and A53T) increase the tendency of α -synuclein protein to form β -sheets which aggregate and form LBs [20]. Multiple copies of this mutated gene have been reported in a number of familial cases of PD [21].

1.1.3. Occupational Risks and Head Trauma

Farmers are more likely to be exposed to pesticides, which are linked to neurodegenerative disorders. A number of pesticides contain neurotoxins such as dipyridylum herbicides that are structurally similar to 1, methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine [MPTP], a fast acting neurotoxin. Such herbicides can induce parkinsonian symptoms and predispose individuals to the disease by generating reactive oxygen species which destroy dopaminergic neurons in the SN_c [22, 23]. Moreover, some pesticides are known to inhibit neuronal aldehyde dehydrogenase [ALDH] activity which increases individual susceptibility to PD, particularly in individuals carrying genetic polymorphisms in *ALDH2* [24].

Head injuries can cause cerebral haemorrhages. Chronic head injuries involving concussion are reported to predispose individuals to neurodegenerative disorders, including PD [25, 26]. Concussion often leads to local inflammation where pro-inflammatory cytokines cause neuronal cell death and remodelling of the brain parenchyma, which is often witnessed in autopsies of Parkinson diseased brains [27]. Moreover, brain injury triggers astrocyte activation which leads to neuronal injury and scarring since astrocytes are no longer able to maintain the integrity of the blood-brain barrier [BBB], thereby allowing toxic molecules to reach the CNS [28].

1.1.4. Lower Levels of Folic Acid

Low levels of folic acid are commonly associated with birth defects. Emerging research on rodents suggests that low levels of folic acid in diet may increase the risk of developing PD. Evidence shows that folic acid normally reduces mitochondrial dysfunction and promotes the generation of new mitochondria that may help protect neuronal cell death. Folate deficiency may lead to greater susceptibility to oxidative damage in mitochondria and apoptosis of dopaminergic neurons in PD [29]. However, at least one prospective clinical trial failed to identify any link between high folic acid intake and a reduced risk of development of PD [30].

1.2. Neurophysiology of Basal Ganglia

The basal ganglia [BG] are comprised of a number of nuclei in the subcortical region. They play important roles in the initiation of muscular movements, learning, cognition, and help to coordinate changes in posture. They are comprised of a number of structures, which includes the substantia nigra - a midbrain structure consisting of the pars

compacta [SN_c] and pars reticulata [SN_r]. The SN_c plays an important role within the basal ganglia in synthesising and secreting dopamine which binds to Dopamine-1 receptors [D₁R] in the BG [31]. This subsequently activates GABAergic neurons favouring direct pathways, thus facilitating movement [32]. When dopamine binds to Dopamine-2 receptors [D₂R], indirect pathways are activated which subsequently decreases thalamo-cortical firing and inhibits movement as illustrated in Fig (1). The striatum also projects into a number of areas which includes the GP_i (contains GABAergic output neurons), the thalamic nucleus, reticular formation (affects muscle tone), and superior colliculus (affects eye movements). Alteration in these systems can be observed in PD patients, along with the bradykinesia which is a result of depletion of dopaminergic neurons in these pathways [32].




Fig. (1). Direct and indirect pathways of Basal Ganglia in initiating motor activity. In the direct pathway, cerebral cortical input to the striatum causes activation of inhibitory neurons in the striatum which then causes an increased inhibitory output to the globus pallidus internal [GP_i]. There is a decreased inhibitory output from GP_i to ventral anterior [VA] and ventral lateral [VL] nuclei of the thalamus which then projects *via* excitatory pathways into the premotor cortex. The direct pathway is involved in regulating tonic excitation in the premotor cortex which is an area involved in planning and initiating movement. The indirect pathway is inhibitory to movement when excitatory projection from cerebral cortex facilitates inhibitory projection neurons in globus pallidus external [GPe]. These then inhibit tonic inhibitory output neurons which decreases tonic inhibition of subthalamic nucleus [STN] resulting in increased excitatory output to GP_i. Excitatory input to GP_i increases inhibitory output from GP_i to thalamus which then decreases excitatory feedback to cerebral cortex leading to inhibition of motor activity. Dopamine promotes action of direct pathway while suppressing the activity of indirect pathway.

1.3. Pathophysiology of PD

PD occurs when the activity of neurons in GP_e is reduced. This results in an increased discharge of STN neurons. The degeneration of dopaminergic SNC neurons leads to increased activity of the indirect pathway which causes an oscillation in electrical discharge to neighbouring neurons affecting the overall activity throughout the BG [34]. Pathological changes in basal ganglia results in the development of a resting tremor, loss of coordination, bradykinesia and postural changes. Rigidity is a product of dopaminergic neuron destruction in SN_c which leads the striatum to over stimulate the corticospinal motor system [33, 34].

1.3.1. Molecular Histopathology of PD

The accumulation of LBs and LNs in neurons is a characteristic histological finding in PD. LBs and LNs are aggregates of α -synuclein, parkin, and synphilin-1 proteins. α -synuclein is found predominantly in the brain and expressed in mitochondria of the striatal and thalamic neurons [35, 36]. The precise role of α -synuclein is not known

but a number of studies suggest that α -synuclein helps to regulate dopamine release and supply synaptic vesicles to presynaptic terminals [37]. A number of missense mutations in the α -synuclein gene (A53T, A30P, and E46K) allow the α -synuclein protein to form aggregates of β -sheets which then form toxic oligomers commonly observed in hereditary forms of PD [38, 39]. Mutated forms of α -synuclein may have prion protein-like characteristics since it can induce β -confirmation in normal α -synuclein proteins. It does so by acting as a template to promote misfolding and aggregation of normal α -synuclein which can then spread on to other regions of the brain [40]. In addition, LBs and LNs have also been reported in striatal and nigral neurons of individuals who have undergone fetal-nigral mesencephalic cell transplantation [FNMCT], further supporting the hypothesis of prion protein-like characteristic of mutated α -synuclein [40]. Long-term accumulation of mutant α -synuclein protein also leads to dementia and behavioural as well as cognitive impairment. These are characteristic features of late stages in PD [41]. In addition to α -synuclein protein, defects in the parkin gene which produces mutated ubiquitin ligase prevents removal of α -synuclein aggregates, and allows LBs and LNs to form [42]. Recently, studies reported observing ferric deposition in parkinsonian brains. It was postulated that high level of iron present in SN_c may induced oxidative stress which then leads to loss of dopaminergic neurons [43].

2. NEUROPROTECTIVE AGENTS

Neuroprotection refers to the preservation of neuronal structure and function. Neuronal integrity can be compromised in diseases such as PD, and the aim of neuroprotective agents is to protect or slow down the progression of disease by delaying the process of neuronal loss. There are a number of candidates which help to reduce neurotoxicity by inhibiting neurotoxins, ROS, preserving mitochondrial function and preventing protein aggregations [44]. Neuroprotective agents include the following:

2.1. Polyphenols

Polyphenols are naturally occurring compounds found in plants that are responsible for protecting them against UVB radiation. Polyphenols have antioxidant and anti-carcinogenic effects which are yet to be explored and could potentially be used to treat a number of degenerative diseases including PD. In addition, polyphenols found in black tea are reported to inhibit α -synuclein aggregation and degeneration of dopaminergic neurons through an unknown mechanism [45].

2.2. Nicotine

Nicotine is a constituent of cigarettes and is a known potent parasympathomimetic alkaloid that acts on nicotinic acetylcholine receptors [nAChR] found in brain. Nicotine can protect nigrostriatal damage and help to improve cognitive functions according to a number of randomized controlled trials [46]. The exact mechanism of action of nicotine in nigrostriatal pathways remains unknown but some studies suggest that nicotine binds to nAChR and activates intracellular transduction, altering calcium signalling which may help reduce neuronal injuries [46]. In addition, nicotine has been found to reduce L-dopa-induced dyskinesia [LID] in a number of animal studies but not in humans [47].

2.3. Curcumin

Curcumin is an active ingredient of turmeric. It is non-toxic and able to cross the BBB. Curcumin is capable of binding to mutant α -synuclein, thereby preventing β -aggregation [48, 49]. In doing so curcumin can help reduce LBs formation and reduce risk of developing PD. Its precise mechanism of action is not known but some suggest that curcumin acts as an amyloid breaker with anti-inflammatory effects in which it is able to reduce generation of ROS [50].

2.4. Caffeine

Caffeine 3,7-trimethylpurine-2,6-dione is a psychoactive methylxanthine extracted mainly from plant sources. It is found in a number of beverages such as coffee and tea [51]. 87% of caffeine is consumed in coffee, soft drinks and tea with an average consumption of 193mg of caffeine per day 1.2mg caffeine per kg of body per day [52]. Men and women aged 35-64 were reported to be the highest caffeine consumers [52]. Caffeine is a potent neurostimulant and also has neuroprotective effects. Its effects on the CNS arise through the competitive non-selective inhibition of adenosine A₁ and A_{2A} receptors. Epidemiological studies suggest that consumption of 3-4 cups of coffee daily (~200mg) can reduce the risk of PD [53, 54].

This is further supported by randomized trials and *in-vitro* studies that highlight caffeine's ability to antagonise adenosine receptors, in particular A_{2A} receptors found in striatopallidal neurons. Blockade of A_{2A} receptors improves mobility in PD patients [55, 56].

Blockade of adenosine receptors improves motor functions of PD marginally in both animals and humans. Moreover, long-term exposure to caffeine in hemiparkinsonian mice may alter L-dopa responses in PD *via* unknown mechanisms [57]. The benefits of caffeine and its use in adjuvant therapy are outlined in the rest of this literature review.

3. DIFFERENT EFFECTS OF CAFFEINE IN PD

3.1. Adenosine A_{2A} Receptors

Adenosine A_{2A} receptors are distributed in the BG, which is heavily innervated by dopaminergic neurons. Adenosine is a ligand for all adenosine receptors and at low concentrations it shows a greater affinity for A₁ receptors where it decreases glutamate release from striatal neurons. At high concentrations, adenosine has a higher affinity for A_{2A} receptors and exerts excitatory effects through stimulation of glutamate and Ach release in the striatum. Activation of adenosine A_{2A} receptors also enhances GABA-A mediated inhibitory effects on the GP_e [58].

Adenosine A_{2A} receptors are primarily expressed on striatopallidal neurons and form functional heteromeric complexes with D₂ receptors and metabotropic glutamate receptors [mGLU₅]. In so doing, A_{2A} receptors can control functions of indirect efferent pathways of the BG. It is important to note that A_{2A} receptors are functionally equivalent to D₂ receptors since they form heteromers with D₂ receptors by interacting with the arginine-rich domain on D₂ receptors. A similar interaction is also seen between D₁-NMDA receptors [N-methyl-D-aspartate receptors] heteromerisation [59, 60]. Upon forming a heteromeric complex with D₂ receptors, A_{2A} and D₂ receptors become reciprocal antagonists and form the basis of motor control, learning, motivation, and reward systems. D₂ receptors normally antagonise A_{2A} receptors' activity by inhibiting adenylyl cyclase activation. However, when D₂ receptor interacts with A_{2A} receptor in heteromeric configuration, A_{2A} receptor can reduce the binding affinity of dopamine to D₂ receptor, preventing inhibition of motor activity [60]. In addition, co-stimulation of A_{2A} with mGLU₅ enhances A_{2A} receptors ability to inhibit D₂ receptors as well as to override the inhibitory tone imposed by endogenous dopamine *via* D₂ receptors [61]. The underlying antagonistic properties of A_{2A} receptor on D₂ receptors with deficits in dopamine levels translates to akinesia, which is commonly seen in PD [61]. Moreover, loss of striatal dopamine in PD results in unopposed adenosine stimulation resulting in hyperactivity of indirect pathway and motor dysfunction, which are also features observed in PD.

3.1.1. Caffeine as an Adenosine A_{2A} Receptor Antagonist

Caffeine is a potent neurostimulant and might protect against a number of neurodegenerative disease and may affect cell cycle function in certain types of cancer [62, 63]. Nevertheless, the role of caffeine in cardiovascular diseases remains inconclusive according to a prospective cohort study [64]. Emerging evidence from *in vivo* and *in vitro* studies suggests that caffeine may protect against development of PD by keeping the BBB intact, which is essential for neuronal survival and CNS functioning [65, 66]. In addition, caffeine may improve motor functions in PD patients by modulating adenosine A_{2A} receptors at different dosages [10]. Caffeine is structurally similar to adenosine and is a non-selective antagonist at A₁, A_{2A}, A_{2B} and A₃ adenosine receptor subtypes [67]. A_{2A} receptors are predominately found on striatopallidal neurons and are antagonised by caffeine, which results in decreased intracellular cAMP levels and a diminished release of GABA in the GP [67]. A_{2A} antagonists like caffeine thus may decrease the release of inhibitory neurotransmitters [GABA] and increases excitatory activity [serotonin, noradrenaline] in the striatopallidal neurons by stimulation of dopamine D₁ and D₂ receptors, promoting motor activities in rats [68]. Nevertheless, rapid tolerance to the effects of caffeine develops in long-term use [69]. Animal studies by Popoli *et al.* suggests that chronic caffeine administration combined with a selective A_{2A} receptor antagonist such as SCH 58261 results in less tolerance towards adenosine A_{2A} receptor antagonists compared to when adenosine A_{2A} receptor antagonists are administered on their own [70].

Several studies have reported a reduction in MPTP-induced destruction of dopaminergic neurons in mice administered with oral A_{2A} receptor antagonists. This protected against the loss of nigral dopaminergic neuronal cells induced by 6-hydroxydopamine in mice. A_{2A} antagonists also prevented the functional loss of dopaminergic nerve

terminals in the striatum and the ensuing gliosis caused by MPTP in mice [71]. The mechanism underlying the neuroprotective effects of A_{2A} antagonists such as caffeine is still not fully understood.

On the contrary, some studies have reported that the activation of A_{2A} receptors may protect tissues from reperfusion injury since A_{2A} receptors can inhibit release of pro-inflammatory cytokines [TNF- α , IL-6, IL-8 and IL-12] and increase the release of anti-inflammatory cytokines such as IL-10 [72 - 74]. Moreover, A_{2A} receptor activation can reduce the release of a number of ROS from neutrophils and also reduces neutrophilic infiltration into endothelial cells.

In light of the contradictory findings reported in the literature, further evaluation of the role that A_{2A} activation and antagonism plays in neuroprotection is needed.

3.2. Astrocytes and Their Role in PD

Aside from being an A_{2A} antagonist, caffeine may help stabilise the blood-brain barrier by regulating astrocytes and inflammatory processes in animal models [66]. Astrocytes are glial cells found in the mammalian brain that play a pivotal role in maintaining various neuronal functions by transporting nutrients and metabolites to neurons through a number of transporters such as the malate-aspartate shuttle. There are two types of astrocytes found in substantia nigra: [75].

1. Protoplasmic astrocytes which encapsulate neuronal bodies and synapses
2. Fibrous astrocytes which interact with oligodendroglia and node of Ranvier

Research on astrocytes in mice has suggested that α -synuclein affects only protoplasmic astrocytes but not fibrous astrocytes [76].

Astrocytes are significant players in the central nervous system as they develop and maintain the BBB and promote neurovascular coupling. One extreme of astrocytes faces endothelial cells to maintain BBB and control cerebral blood flow [CBF] while the other extreme is in communication with numerous neurons [77]. Moreover, astrocytes can release chemokines, gliotransmitters and help in the uptake of glutamate and GABA from the synaptic cleft. Other functions of astrocytes include pH regulation, K⁺ buffering, metabolising dopamine, and producing antioxidants such as glutathione [GSH] and superoxide dismutases [SOD 1, and 2] [78].

In neurodegenerative diseases such as PD, astrocyte functions are compromised. Neurons are more susceptible to injury due to their limited capacity to counteract free radical-induced damage. Furthermore, neurons require a number of nerve growth factors [NGF] which are derived from astrocytes [79]. In the absence of astrocytes, neuronal functions are compromised and cell death slowly ensues during brain degeneration in PD [79]. In addition, during the process of reactive astrogliosis in the substantia nigra [an abnormal increase in astrocyte numbers secondary to neuronal death], astrocytes release a number of proinflammatory cytokines that act on neighbouring neurons. The increased production of ROS and activation of apoptotic mechanisms is often observed in PD, and may lead to death of dopaminergic neurons. Reactive astrogliosis is one of the main features seen in astrocytes following brain injuries, and in neurodegenerative diseases caused by trauma, α -synuclein aggregation, and ischemia [79]. A number of researchers have suggested that neurotoxins such as MPTP are capable of inducing astrogliosis and microgliosis which are often accompanied by mitochondrial dysfunction, nuclear karyolysis, and eventually neuronal death [80].

It is widely reported in a number of animal studies that astrocytes actively participate in the death of dopaminergic neurons by activating apoptosis through cytokines [TNF α , IL-1B and IL-6] and nitric oxide [NO] [28]. NO is believed to promote the release of cytochrome C from dysfunctional mitochondria while cytokines bind to specific receptors in dopaminergic neurons to activate the pro-apoptotic caspase pathway [81, 82]. Oxidative damage is a common feature observed in PD brain samples [83]. The increased uptake of α -synuclein by astrocytes leads to up-regulation of inflammatory molecules causing further production of ROS which may then lead to DNA damage [84]. Further depletion of GSH and SOD 1 and 2 may facilitate production of ROS and reactive nitrogen species [RNS] in astrocytes. This then leads to structural alteration and aggregation of normal α -synuclein proteins [85, 86]. Recent findings in mice models of PD have found that the nitration (nitrogen group is introduced to tyrosine residue on α -synuclein) of α -synuclein induced by oxidative injury (activation of induced nitric oxide synthase [iNOS] and focal adhesion kinase [FAK]) can enhance α -synuclein fibril formation causing cell death [87, 88]. Therefore one may conclude that inhibiting glial inflammatory processes could reduce neuronal damage during PD [89].

3.2.1. Caffeine Inhibits Astrocyte-induced Inflammation in PD

Even though caffeine can antagonise all forms of adenosine receptors it remains ineffective in antagonising purinergic-7 receptors [P2X7R] which are ligand-gated cation channels that open in response to ATP binding and lead to cell depolarization. P2X7 receptors are present on neurons and astrocytes throughout the CNS and PNS [90]. Recent studies suggest that P2X7R mediates platelet aggregation as well as microglial activation during inflammation. Moreover, P2X7R are also responsible for mediating apoptosis [91]. The use of a P2X7R antagonist, in combination with an adenosine receptor antagonist such as caffeine may reduce the inflammatory processes that are triggered by both receptors and their subtypes [92].

In addition, a number of studies in newborn rat brain cultures suggest that high doses of caffeine can reduce glial cell proliferation. This may be useful in reducing the inflammatory process in PD brains [93]. By actively preventing glial cell proliferation, specifically astroglyosis, the progression of neurodegeneration may be stalled due to mechanisms that prevent the breakdown of the BBB and the excessive release of gliotransmitters such as glutamate that lead to neurotoxicity.

Caffeine is also reported to suppress the production of TNF- α through protein kinase A dependent mechanisms in human blood according to Horrigan *et al.* [94]. Furthermore, caffeine is reported to mediate glutathione synthesis, reducing the ROS and RNS induced inflammatory response in the hippocampus of rodents leading to neuroprotection [95].

Astrocytes also express adenosine A_{2A} receptors that may be antagonised by adenosine A_{2A} antagonists, favouring the increased uptake of glutamate into astrocytes and thus reducing glutamate levels at the synaptic cleft. This can reduce the incidence of neurotoxicity induced by the excessive stimulatory effects of neurotransmitters [95, 96]. In addition to increased gliotransmitter uptake, chronic caffeine intake in non-habitual caffeine consumers may reduce CBF acutely through an unknown mechanism, which may involve vasoconstriction of cerebral vessels [97, 98].

3.3. Overexpression of Ca_v 1.3 Ca²⁺ Channels and Inhibition by Caffeine in PD

Voltage-gated calcium channels 1.3 [Ca_v 1.3 Ca²⁺] are L-type calcium channels that are selectively permeable to calcium ions. Ca_v 1.3 Ca²⁺ channels have an α 1 subunit which is a prime target for calcium-channel blockers such as the dihydropyridines [99]. In addition to astrocyte dysregulation, the activity of Ca_v 1.3 Ca²⁺ channels has been implicated in the neurodegeneration of dopamine-producing neurons in PD. Dopaminergic neurons found in SN_c continuously generate low frequency basal activity using Ca_v 1.3 Ca²⁺ channels. These L-type calcium channels have low expression levels [100]. Other L-type calcium channels found in SN_c include Ca_v 1.2 Ca²⁺ channels. These are exclusively expressed in dopaminergic neurons located in the SN_c and are only activated at higher membrane potentials. Conversely, Ca_v 1.3 Ca²⁺ channels are activated at sub-threshold membrane potentials, making them ideal pacemaker cells in neurons. However, they require significantly more ATP to sustain their transient activity. Moreover, Ca_v 1.3 Ca²⁺ channels generate oxidative stress mediated by mitochondria during periods of pacemaker activity *via* unknown mechanisms [101].

Recent studies suggest that there is an increased density of Ca_v 1.3 Ca²⁺ channels in dopaminergic neurons in mice models of PD [102]. The increased expression of dopaminergic neurons in Ca_v 1.3 Ca²⁺ channels generates more ROS that leads to neurodegeneration of dopaminergic neurons as seen in PD. These changes in the expression of voltage gated calcium channel subtypes in PD are of particular therapeutic interest [102]. A number of studies have reported the potential benefits of dihydropyridine [DHP] calcium channel blockers in reducing pathogenesis of PD as well as mortality in elderly PD patients [103]. Other studies have reported that caffeine can inhibit ryanodine receptors [RyRs] in certain parts of the body [104]. RyRs play a central role in microglial-mediated inflammation, and are believed to contribute to pathogenesis of neurodegenerative disorders such as PD. Inhibition of RyRs shows therapeutic promise in the management of PD [105]. Studies have also shown that protein kinase C [PKC] activation may lead to phosphorylation of serine residue 81 on the amino-terminus of Ca_v 1.3 Ca²⁺ channels, leading to their inactivation [106]. However, no studies have reported possible effects of caffeine on protein kinase C pathways in the CNS or on inhibition of Ca_v 1.3 Ca²⁺ channels.

3.4. Efficacy of Caffeine in Different People and Genders

Caffeine may show promise in the pharamcotherapy of PD since it antagonises adenosine receptors, inhibits the

inflammatory response, and possibly inactivates calcium channels. The effect of caffeine was reported to be dose-dependent in a randomised controlled trial showing that a daily dose of 100-200mg of caffeine can reduce the risk of PD motor symptoms, while a daily dose of above 200mg caffeine was ineffective [107]. The reason behind this discrepancy includes genetic variation and gender differences. Genetic variation in cytochrome *CYP1A2* gives rise to fast and slow metabolisers of caffeine. *CYP1A2* is a primary enzyme responsible for metabolism of 95% of caffeine into three active metabolites: paraxanthine, theophylline, and theobromine. Each of these metabolites exert different effects. Certain variations in *CYP1A2* gene result in a reduced ability to metabolise caffeine slow metabolisers [108]. These individuals are prone to hypertension as high levels of caffeine in serum can activate systemic epinephrine release. Moreover, sustained hypertension leads to vascular disease and possibly early myocardial infarction and cerebrovascular accidents, that may then induce secondary neurodegenerative disorders such as vascular parkinsonism [109, 110].

Conversely, fast metabolisers overexpress *CYP1A2*. This results in rapid and efficient metabolism of caffeine [108]. These individuals may require higher doses of caffeine in order to achieve the same therapeutic effects of caffeine in reducing the prevalence of PD. However, conflicting studies have suggested that slow metabolisers are not necessarily at higher risk of developing PD when compared to fast metabolisers and this warrants further long-term clinical trials [108].

Apart from variation in caffeine metabolism, gender may also account for differences in the incidence of PD and response to caffeine. PD is more prevalent in males than in females due to differences in physiological adaptation as well as differences in toxicant exposure rate [111]. Nevertheless, in a number of epidemiological studies, it is reported that male caffeine users were more likely to have lower incidence of PD compared to female caffeine users. This led to a number of large prospective studies which highlighted the possible link between exogenous oestrogen in Hormone Replacement Therapy [HRT] and reduced efficacy in postmenopausal women with higher PD incidence [112, 113]. It seems that exogenous oestrogen present in HRT reduces half-life of caffeine similar to those seen in fast metabolisers of caffeine, suggesting that exogenous oestrogen may put females on HRT at a greater risk of developing neurodegenerative diseases such as PD. *In vitro* animal studies have compared the effects of a 20mg caffeine dose administration in normal vs ovariectomised mice exposed to MPTP. Ovariectomised mice had reduced parkinsonian symptoms compared to normal mice. In addition, when male mice were injected with exogenous oestrogen and MPTP, they showed the same parkinsonian symptoms as female mice with ovaries. This suggests that administration of exogenous oestrogen may somehow inhibit the neuroprotective effects of caffeine [114, 115].

3.5. Caffeine as an Adjuvant Therapy in PD

The evidence outlined above highlights a number of beneficial effects of caffeine that could possibly justify its use as an adjuvant drug in the management of PD. There are reported experimental uses of caffeine in treating PD. For instance, trihexyphenidyl [THP] is an anti-parkinsonian agent that has been clinically used to treat PD [116]. It is a potent M_1 receptor antagonist that inhibits M_1 receptors in different parts of the cerebral cortex. Its use is accompanied by side effects that include memory impairment [116]. When THP is co-administered with low doses of caffeine in mice, there is increased THP potency without memory impairment. Observational studies on reserpine-induced hypokinesia in rats reported no reversal of symptoms when caffeine or THP were used on their own [117, 118]. Conversely, when caffeine and THP were co-administered, they restored locomotion in reserpine-treated rats.

The authors concluded that low doses of caffeine co-administered with THP might help reduce a number of motor deficiencies in animals, which are seen in PD. Nevertheless, long-term randomized clinical trials are needed to assess such claims and therapeutic benefits of such co-administration [116].

Other studies have investigated the effects of caffeine co-administration with bromocriptine. This improves a number of motor functions in rats under an observational study. Bromocriptine is a dopamine agonist commonly used to treat PD. When Bromocriptine is administered individually in denervated rats, subthreshold motor activities were recorded [119, 120]. When Bromocriptine was co-administered with caffeine, levels of motor activity were slightly higher than with the isolated use of Bromocriptine. Nonetheless, tolerance to caffeine was reported in rats administered only caffeine for a significant period of time but no tolerance to Bromocriptine was reported. In addition, no tolerance to chronic use of caffeine was reported when it was co-administered with Bromocriptine, and sustained levels of motor activity was observed in these rats [120]. Therefore, it is important to explore the relationship between caffeine and other parkinsonian drugs to determine the underlying mechanisms that allow caffeine to enhance efficacy and reduce drug tolerance [120].

A number of novel approaches have been adopted by the scientific community in an attempt to find better and more efficacious treatment for PD. Currently, Parkinsonian symptoms are treated through medications such as L-dopa and dopamine agonists like pramipexole and rotigotine, which themselves are accompanied by significant side effects that range from hypotension to gastrointestinal disturbances [121, 122]. Moreover, most of these drugs undergo first pass metabolism through the liver and are metabolised by circulating enzymes such as Monoamine oxidases [MAO] which reduces their bioavailability in the body. Hence they do not offer a long-term solution in treating PD.

In summary, caffeine is a cheap and widely-available neuromodulator which can be used in the treatment of PD in conjunction with current therapy. Nevertheless the precise dosage and safety of caffeine is of particular concern especially when it is administered at high doses for a significant period of time; a number of adverse effects such as resting tremors have been reported in chronic caffeine users. Therefore, more long-term double-blinded randomized controlled trials in humans are necessary in order to evaluate the efficacy and possible uses of caffeine in treating conditions like PD.

CONCLUSION

It was not until recently that the scientific community began to notice the neuroprotective effects of caffeine following the observation of lower incidences of PD reported in chronic caffeine users. Recent animal and human studies have suggested that caffeine may help to treat PD since it can antagonise the adenosine A_{2A} receptors and inflammatory changes in the BG. Moreover, caffeine can prevent apoptotic cell death by lowering caspase-3 activity that is induced by neurotoxins such as MPP⁺, as well as decrease the number of apoptotic fragmented nuclei. In doing so it may halt dopaminergic neuronal destruction and act as cytoprotective agent [123].

As a result, caffeine might be able to prevent neurotoxicity induced by LBs and LNs. The beneficial functions of caffeine may not be limited solely to neuronal cells as it may be able to decrease the activity and proliferation of non-neuronal cells like astrocytes, which are primary regulators of the BBB and produce a number of inflammatory molecules in early and late stages of PD. Further randomised trials are needed to support this hypothesis.

The physiological role of caffeine may also help reduce the incidence of drug tolerance that often develops in advanced stages of PD where patients are given higher dosage to achieve the same therapeutic effects. Nevertheless, the beneficial effects of caffeine are not observed in all individuals due to genetic and gender differences. Individuals with variants in *CYP1A2* are known to metabolise caffeine at a much slower rate than normal, this potentially predisposes them to cardiovascular diseases.

Aside for genetic variation, gender difference may also limit the neuroprotective effects of caffeine since females on HRT are not protected from neurodegenerative diseases even when caffeine is consumed chronically [124]. Physiological studies have suggested that exogenous oestrogen in HRT may inactivate neuroprotective effects of caffeine which may predispose females on HRT at higher risk of developing neurodegenerative disorders like PD [115]. In addition, most studies are currently conducted on animals, and may not have similar effects in humans.

Furthermore there is no single universally agreed dosage of caffeine which has been agreed by the scientific community. Doses used in most studies varied from 20mg to over 200mg [107, 125]. Moreover, long-term double-blinded randomised trials are needed to show the correlation between caffeine usage and low incidence of PD. Nonetheless, caffeine remains an interesting area of active research and may one day prove to be a cheap alternative to current PD treatment. In addition, a number of caffeine analogues are under investigation in clinical trials and have shown some promising results. Therefore, it is not long before caffeine will be used as an adjuvant therapy in treating neurodegenerative diseases like PD.

LIST OF ABBREVIATIONS

α-synuclein	=	Alpha-synuclein
A_{2A}R	=	Adenosine A _{2A} Receptor
Ach	=	Acetylcholine
AMP	=	Adenosine Monophosphate
APOE gene	=	Apolipoprotein E gene
ATP	=	Adenosine Triphosphate
BBB	=	Blood-Brain Barrier

BG	=	Basal ganglia
cAMP	=	Cyclic Adenosine Monophosphate
Ca_v 1.3 Ca²⁺	=	Voltage gated Calcium channels 1.3
CBF	=	Cerebral blood flow
CNS	=	Central nervous system
CYP1A2	=	Cytochrome-P450 1A2
D1/D2R	=	Dopamine-1/Dopamine-2 receptors
DHP	=	Dihydropyridine
ETC	=	Electron transport chain
GABA	=	Gamma-aminobutyric acid
GP_i	=	Globus pallidus internal
GP_e	=	Globus pallidus external
G_{sa}	=	G-protein α - stimulatory subunit
G_{ia}	=	G-protein α -inhibitory subunit
HRT	=	Hormone replacement therapy
IL-6/IL-8	=	Interleukins-6/Interleukins-8
IL10/IL-12	=	Interleukin-10/Interleukins-12
LBS/ LNs	=	Lewy bodies/ Lewy neurites
L-dopa	=	Levodopa
LID	=	L-Dopa induced dyskinesia
M₁R	=	Muscarinic-1 Receptor
MAO	=	Monoamine oxidases
mGLU₅	=	Metabotropic glutamate-5 receptors
MRI	=	Magnetic Resonance Imaging
MPP⁺	=	1-methyl-4-propionoxypiperidine
MPTP	=	1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine
nAChR	=	Nicotinic Acetylcholine Receptors
NGF	=	Nerve growth factors
NMDA	=	N-methyl-D-aspartate receptor
NO	=	Nitric Oxide
PD	=	Parkinson's disease
PGE-2	=	Prostaglandin-E2
PINK-1	=	PTEN induced kinase-1
PKC	=	Protein kinase C
PNS	=	Peripheral nervous system
P2X7R	=	Purinergic-7-receptors
RNS	=	Reactive nitrogen species
ROS	=	Reactive Oxygen Species
SN_c	=	Substantia nigra pars compacta
STN	=	Subthalamic nucleus
SNr	=	Substantia Nigra pars reticulata
TNF-α	=	Tumour necrosis factor-alpha
THP	=	Trihexyphenidyl
UVB	=	Ultraviolet-B
VA	=	Ventral anterior
VL	=	Ventral lateral
X-chromosomes	=	Sex chromosomes

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Kempster PA, Hurwitz B, Lees AJ. A new look at James Parkinson's essay on the shaking palsy. *Neurology* 2007; 69(5): 482-5. [<http://dx.doi.org/10.1212/01.wnl.0000266639.50620.d1>] [PMID: 17664408]
- [2] Pankratz N, Foroud T. Genetics of Parkinson disease. *Genet Med* 2007; 9(12): 801-11. [<http://dx.doi.org/10.1097/GIM.0b013e31815bf97c>] [PMID: 18091429]
- [3] Van Den Eeden SK, Tanner CM, Bernstein AL, *et al.* Incidence of Parkinson's disease: variation by age, gender, and race/ethnicity. *Am J Epidemiol* 2003; 157(11): 1015-22. [<http://dx.doi.org/10.1093/aje/kwg068>] [PMID: 12777365]
- [4] Xia R, Mao Z-H. Progression of motor symptoms in Parkinson's disease. *Neurosci Bull* 2012; 28(1): 39-48. [<http://dx.doi.org/10.1007/s12264-012-1050-z>] [PMID: 22233888]
- [5] Lesage S, Brice A. Parkinson's disease: from monogenic forms to genetic susceptibility factors. *Hum Mol Genet* 2009; 18(R1): R48-59. [<http://dx.doi.org/10.1093/hmg/ddp012>] [PMID: 19297401]
- [6] Baba M, Nakajo S, Tu PH, *et al.* Aggregation of alpha-synuclein in Lewy bodies of sporadic Parkinson's disease and dementia with Lewy bodies. *Am J Pathol* 1998; 152(4): 879-84. [PMID: 9546347]
- [7] Costa J, Lunet N, Santos C, Santos J, Vaz-Carneiro A. Caffeine exposure and the risk of Parkinson's disease: a systematic review and meta-analysis of observational studies. *J Alzheimers Dis* 2010; 20(Suppl. 1): S221-38. [PMID: 20182023]
- [8] Ross GW, Abbott RD, Petrovitch H, *et al.* Association of coffee and caffeine intake with the risk of Parkinson disease. *JAMA* 2000; 283(20): 2674-9. [<http://dx.doi.org/10.1001/jama.283.20.2674>] [PMID: 10819950]
- [9] Seidl SE, Potashkin JA. The promise of neuroprotective agents in Parkinson's disease. *Front Neurol* 2011; 2: 68. [<http://dx.doi.org/10.3389/fneur.2011.00068>] [PMID: 22125548]
- [10] Hsu CW, Wang CS, Chiu TH. Caffeine and a selective adenosine A2A receptor antagonist induce sensitization and cross-sensitization behavior associated with increased striatal dopamine in mice. *J Biomed Sci* 2010; 17: 4. [<http://dx.doi.org/10.1186/1423-0127-17-4>] [PMID: 20074377]
- [11] Alves J, Petrosyan A, Magalhães R. Olfactory dysfunction in dementia. *World J Clin Cases* 2014; 2(11): 661-7. [<http://dx.doi.org/10.12998/wjcc.v2.i11.661>] [PMID: 25405189]
- [12] Baba T, Kikuchi A, Hirayama K, *et al.* Severe olfactory dysfunction is a prodromal symptom of dementia associated with Parkinson's disease: a 3 year longitudinal study. *Brain* 2012; 135(Pt 1): 161-9. [<http://dx.doi.org/10.1093/brain/awr321>] [PMID: 22287381]
- [13] Wood H. Parkinson disease: Severe olfactory dysfunction may herald cognitive decline in Parkinson disease. *Nat Rev Neurol* 2012; 8(3): 122-2. [PMID: 22371285]
- [14] Doty RL. Olfactory dysfunction in Parkinson disease. *Nat Rev Neurol* 2012; 8(6): 329-39. [<http://dx.doi.org/10.1038/nrneurol.2012.80>] [PMID: 22584158]
- [15] Gillies GE, Pienaar IS, Vohra S, Qamhawi Z. Sex differences in Parkinson's disease. *Front Neuroendocrinol* 2014; 35(3): 370-84. [<http://dx.doi.org/10.1016/j.yfrne.2014.02.002>] [PMID: 24607323]
- [16] Lu H, Liu X, Deng Y, Qing H. DNA methylation, a hand behind neurodegenerative diseases. *Front Aging Neurosci* 2013; 5: 85. [<http://dx.doi.org/10.3389/fnagi.2013.00085>] [PMID: 24367332]
- [17] Haaxma CA, Bloem BR, Borm GF, *et al.* Gender differences in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2007; 78(8): 819-24. [<http://dx.doi.org/10.1136/jnnp.2006.103788>] [PMID: 17098842]
- [18] Sawada H, Ibi M, Kihara T, *et al.* Estradiol protects dopaminergic neurons in a MPP+Parkinson's disease model. *Neuropharmacology* 2002; 42(8): 1056-64. [[http://dx.doi.org/10.1016/S0028-3908\(02\)00049-7](http://dx.doi.org/10.1016/S0028-3908(02)00049-7)] [PMID: 12128007]
- [19] Puschmann A. Monogenic Parkinson's disease and parkinsonism: clinical phenotypes and frequencies of known mutations. *Parkinsonism Relat Disord* 2013; 19(4): 407-15. [<http://dx.doi.org/10.1016/j.parkreldis.2013.01.020>] [PMID: 23462481]

- [20] Kumar S, Sarkar A, Sundar D. Controlling aggregation propensity in A53T mutant of alpha-synuclein causing Parkinson's disease. *Biochem Biophys Res Commun* 2009; 387(2): 305-9. [<http://dx.doi.org/10.1016/j.bbrc.2009.07.008>] [PMID: 19580781]
- [21] Kotzbauer PT, Giasson BI, Kravitz AV, *et al.* Fibrillization of alpha-synuclein and tau in familial Parkinson's disease caused by the A53T alpha-synuclein mutation. *Exp Neurol* 2004; 187(2): 279-88. [<http://dx.doi.org/10.1016/j.expneurol.2004.01.007>] [PMID: 15144854]
- [22] Gorell JM, Johnson CC, Rybicki BA, Peterson EL, Richardson RJ. The risk of Parkinson's disease with exposure to pesticides, farming, well water, and rural living. *Neurology* 1998; 50(5): 1346-50. [<http://dx.doi.org/10.1212/WNL.50.5.1346>] [PMID: 9595985]
- [23] Semchuk KM, Love EJ, Lee RG. Parkinson's disease and exposure to agricultural work and pesticide chemicals. *Neurology* 1992; 42(7): 1328-35. [<http://dx.doi.org/10.1212/WNL.42.7.1328>] [PMID: 1620342]
- [24] Fitzmaurice AG, Rhodes SL, Cockburn M, Ritz B, Bronstein JM. Aldehyde dehydrogenase variation enhances effect of pesticides associated with Parkinson disease. *Neurology* 2014; 82(5): 419-26. [<http://dx.doi.org/10.1212/WNL.0000000000000083>] [PMID: 24491970]
- [25] Goldman SM, Tanner CM, Oakes D, Bhudhikanok GS, Gupta A, Langston JW. Head injury and Parkinson's disease risk in twins. *Ann Neurol* 2006; 60(1): 65-72. [<http://dx.doi.org/10.1002/ana.20882>] [PMID: 16718702]
- [26] Jafari S, Etminan M, Aminzadeh F, Samii A. Head injury and risk of Parkinson disease: a systematic review and meta-analysis. *Mov Disord* 2013; 28(9): 1222-9. [<http://dx.doi.org/10.1002/mds.25458>] [PMID: 23609436]
- [27] Maia PD, Kutz JN. Compromised axonal functionality after neurodegeneration, concussion and/or traumatic brain injury. *J Comput Neurosci* 2014; 37(2): 317-32. [<http://dx.doi.org/10.1007/s10827-014-0504-x>] [PMID: 24916135]
- [28] Rappold PM, Tieu K. Astrocytes and therapeutics for Parkinson's disease. *Neurotherapeutics* 2010; 7(4): 413-23. [<http://dx.doi.org/10.1016/j.nurt.2010.07.001>] [PMID: 20880505]
- [29] Miller JW. Homocysteine, folate deficiency, and Parkinson's disease. *Nutr Rev* 2002; 60(12): 410-3. [<http://dx.doi.org/10.1301/002966402320964089>] [PMID: 12521146]
- [30] Chen H, Zhang SM, Schwarzschild MA, *et al.* Folate intake and risk of Parkinson's disease. *Am J Epidemiol* 2004; 160(4): 368-75. [<http://dx.doi.org/10.1093/aje/kwh213>] [PMID: 15286022]
- [31] Wichmann T, DeLong MR. Functional neuroanatomy of the basal ganglia in Parkinson's disease. *Adv Neurol* 2003; 91: 9-18. [PMID: 12442660]
- [32] Chen JF, Moratalla R, Impagnatiello F, *et al.* The role of the D(2) dopamine receptor (D(2)R) in A(2A) adenosine receptor (A(2A)R)-mediated behavioral and cellular responses as revealed by A(2A) and D(2) receptor knockout mice. *Proc Natl Acad Sci USA* 2001; 98(4): 1970-5. [<http://dx.doi.org/10.1073/pnas.98.4.1970>] [PMID: 11172060]
- [33] Wichmann T, DeLong MR. Pathophysiology of Parkinson's disease: the MPTP primate model of the human disorder. *Ann N Y Acad Sci* 2003; 991: 199-213. [<http://dx.doi.org/10.1111/j.1749-6632.2003.tb07477.x>] [PMID: 12846988]
- [34] Schiffmann SN, Fisone G, Moresco R, Cunha RA, Ferré S. Adenosine A2A receptors and basal ganglia physiology. *Prog Neurobiol* 2007; 83(5): 277-92. [<http://dx.doi.org/10.1016/j.pneurobio.2007.05.001>] [PMID: 17646043]
- [35] Chung KK, Zhang Y, Lim KL, *et al.* Parkin ubiquitinates the alpha-synuclein-interacting protein, synphilin-1: implications for Lewy-body formation in Parkinson disease. *Nat Med* 2001; 7(10): 1144-50. [<http://dx.doi.org/10.1038/nm1001-1144>] [PMID: 11590439]
- [36] Kawamata H, McLean PJ, Sharma N, Hyman BT. Interaction of alpha-synuclein and synphilin-1: effect of Parkinson's disease-associated mutations. *J Neurochem* 2001; 77(3): 929-34. [<http://dx.doi.org/10.1046/j.1471-4159.2001.00301.x>] [PMID: 11331421]
- [37] Perez RG, Waymire JC, Lin E, Liu JJ, Guo F, Zigmund MJ. A role for alpha-synuclein in the regulation of dopamine biosynthesis. *J Neurosci* 2002; 22(8): 3090-9. [PMID: 11943812]
- [38] Lemkau LR, Comellas G, Lee SW, *et al.* Site-specific perturbations of alpha-synuclein fibril structure by the Parkinson's disease associated mutations A53T and E46K. *PLoS One* 2013; 8(3): e49750. [<http://dx.doi.org/10.1371/journal.pone.0049750>] [PMID: 23505409]
- [39] Li J, Uversky VN, Fink AL. Effect of familial Parkinson's disease point mutations A30P and A53T on the structural properties, aggregation, and fibrillation of human alpha-synuclein. *Biochemistry* 2001; 40(38): 11604-13. [<http://dx.doi.org/10.1021/bi010616g>] [PMID: 11560511]

- [40] Olanow CW, Brundin P. Parkinson's disease and alpha synuclein: is Parkinson's disease a prion-like disorder? *Mov Disord* 2013; 28(1): 31-40.
[<http://dx.doi.org/10.1002/mds.25373>] [PMID: 23390095]
- [41] Leverenz JB, Quinn JF, Zabetian C, Zhang J, Montine KS, Montine TJ. Cognitive impairment and dementia in patients with Parkinson disease. *Curr Top Med Chem* 2009; 9(10): 903-12.
[PMID: 19754405]
- [42] Dawson TM. Parkin and defective ubiquitination in Parkinson's disease. *J Neural Transm Suppl* 2006; (70): 209-13.
[PMID: 17017531]
- [43] Dias V, Junn E, Mouradian MM. The role of oxidative stress in Parkinson's disease. *J Parkinsons Dis* 2013; 3(4): 461-91.
[PMID: 24252804]
- [44] Bachurin SO, Shevtsova EP, Kireeva EG, Oxenkrug GF, Sablin SO. Mitochondria as a target for neurotoxins and neuroprotective agents. *Ann N Y Acad Sci* 2003; 993: 334-44.
[<http://dx.doi.org/10.1111/j.1749-6632.2003.tb07541.x>] [PMID: 12853325]
- [45] Caruana M, Högen T, Levin J, Hillmer A, Giese A, Vassallo N. Inhibition and disaggregation of α -synuclein oligomers by natural polyphenolic compounds. *FEBS Lett* 2011; 585(8): 1113-20.
[<http://dx.doi.org/10.1016/j.febslet.2011.03.046>] [PMID: 21443877]
- [46] Quik M, Perez XA, Bordia T. Nicotine as a potential neuroprotective agent for Parkinson's disease. *Mov Disord* 2012; 27(8): 947-57.
[<http://dx.doi.org/10.1002/mds.25028>] [PMID: 22693036]
- [47] Huang LZ, Grady SR, Quik M. Nicotine reduces L-DOPA-induced dyskinesias by acting at beta2* nicotinic receptors. *J Pharmacol Exp Ther* 2011; 338(3): 932-41.
[<http://dx.doi.org/10.1124/jpet.111.182949>] [PMID: 21665941]
- [48] Ahsan N, Mishra S, Jain MK, Surolia A, Gupta S. Curcumin Pyrazole and its derivative (N-(3-Nitrophenyl)pyrazole) Curcumin inhibit aggregation, disrupt fibrils and modulate toxicity of Wild type and Mutant α -Synuclein. *Sci Rep* 2015; 5: 9862.
[<http://dx.doi.org/10.1038/srep09862>] [PMID: 25985292]
- [49] Ahmad B, Lapidus LJ. Curcumin prevents aggregation in α -synuclein by increasing reconfiguration rate. *J Biol Chem* 2012; 287(12): 9193-9.
[<http://dx.doi.org/10.1074/jbc.M111.325548>] [PMID: 22267729]
- [50] Pandey N, Strider J, Nolan WC, Yan SX, Galvin JE. Curcumin inhibits aggregation of alpha-synuclein. *Acta Neuropathol* 2008; 115(4): 479-89.
[<http://dx.doi.org/10.1007/s00401-007-0332-4>] [PMID: 18189141]
- [51] Nehlig A, Daval JL, Debry G. Caffeine and the central nervous system: mechanisms of action, biochemical, metabolic and psychostimulant effects. *Brain Res Brain Res Rev* 1992; 17(2): 139-70.
[[http://dx.doi.org/10.1016/0165-0173\(92\)90012-B](http://dx.doi.org/10.1016/0165-0173(92)90012-B)] [PMID: 1356551]
- [52] Frary CD, Johnson RK, Wang MQ. Food sources and intakes of caffeine in the diets of persons in the United States. *J Am Diet Assoc* 2005; 105(1): 110-3.
[<http://dx.doi.org/10.1016/j.jada.2004.10.027>] [PMID: 15635355]
- [53] Qi H, Li S. Dose-response meta-analysis on coffee, tea and caffeine consumption with risk of Parkinson's disease. *Geriatr Gerontol Int* 2014; 14(2): 430-9.
[<http://dx.doi.org/10.1111/ggi.12123>] [PMID: 23879665]
- [54] Hu G, Bidel S, Jousilahti P, Antikainen R, Tuomilehto J. Coffee and tea consumption and the risk of Parkinson's disease. *Mov Disord* 2007; 22(15): 2242-8.
[<http://dx.doi.org/10.1002/mds.21706>] [PMID: 17712848]
- [55] Zahniser NR, Simosky JK, Mayfield RD, et al. Functional uncoupling of adenosine A(2A) receptors and reduced response to caffeine in mice lacking dopamine D2 receptors. *J Neurosci* 2000; 20(16): 5949-57.
[PMID: 10934242]
- [56] Cieślak M, Komoszyński M, Wojtczak A. Adenosine A(2A) receptors in Parkinson's disease treatment. *Purinergic Signal* 2008; 4(4): 305-12.
[<http://dx.doi.org/10.1007/s11302-008-9100-8>] [PMID: 18438720]
- [57] Yu L, Schwarzschild MA, Chen J-F. Cross-sensitization between caffeine and L-dopa-induced behaviors in hemiparkinsonian mice. *Neurosci Lett* 2006; 393(1): 31-5.
[<http://dx.doi.org/10.1016/j.neulet.2005.09.036>] [PMID: 16236444]
- [58] Shen H-Y, Chen J-F. Adenosine A(2A) receptors in psychopharmacology: modulators of behavior, mood and cognition. *Curr Neuropharmacol* 2009; 7(3): 195-206.
[<http://dx.doi.org/10.2174/157015909789152191>] [PMID: 20190961]
- [59] Schwarzschild MA, Agnati L, Fuxe K, Chen J-F, Morelli M. Targeting adenosine A2A receptors in Parkinson's disease. *Trends Neurosci* 2006; 29(11): 647-54.
[<http://dx.doi.org/10.1016/j.tins.2006.09.004>] [PMID: 17030429]
- [60] Ferre S, von Euler G, Johansson B, Fredholm BB, Fuxe K. Stimulation of high-affinity adenosine A2 receptors decreases the affinity of dopamine D2 receptors in rat striatal membranes. *Proc Natl Acad Sci USA* 1991; 88(16): 7238-41.

- [http://dx.doi.org/10.1073/pnas.88.16.7238] [PMID: 1678519]
- [61] Ferré S, Quiroz C, Woods AS, *et al.* An update on adenosine A2A-dopamine D2 receptor interactions: implications for the function of G protein-coupled receptors. *Curr Pharm Des* 2008; 14(15): 1468-74. [http://dx.doi.org/10.2174/138161208784480108] [PMID: 18537670]
- [62] Ascherio A, Zhang SM, Hernán MA, *et al.* Prospective study of caffeine consumption and risk of Parkinson's disease in men and women. *Ann Neurol* 2001; 50(1): 56-63. [http://dx.doi.org/10.1002/ana.1052] [PMID: 11456310]
- [63] Bode AM, Dong Z. The enigmatic effects of caffeine in cell cycle and cancer. *Cancer Lett* 2007; 247(1): 26-39. [http://dx.doi.org/10.1016/j.canlet.2006.03.032] [PMID: 16709440]
- [64] Lopez-Garcia E, van Dam RM, Willett WC, *et al.* Coffee consumption and coronary heart disease in men and women: a prospective cohort study. *Circulation* 2006; 113(17): 2045-53. [http://dx.doi.org/10.1161/CIRCULATIONAHA.105.598664] [PMID: 16636169]
- [65] Chen X, Ghribi O, Geiger JD. Caffeine protects against disruptions of the blood-brain barrier in animal models of Alzheimer's and Parkinson's diseases. *J Alzheimers Dis* 2010; 20(1): S127-41. [PMID: 20164568]
- [66] Chen X, Lan X, Roche I, Liu R, Geiger JD. Caffeine protects against MPTP-induced blood-brain barrier dysfunction in mouse striatum. *J Neurochem* 2008; 107(4): 1147-57. [PMID: 18808450]
- [67] Fredholm BB, Bättig K, Holmén J, Nehlig A, Zvartau EE. Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Pharmacol Rev* 1999; 51(1): 83-133. [PMID: 10049999]
- [68] Garrett BE, Holtzman SG. D1 and D2 dopamine receptor antagonists block caffeine-induced stimulation of locomotor activity in rats. *Pharmacol Biochem Behav* 1994; 47(1): 89-94. [http://dx.doi.org/10.1016/0091-3057(94)90115-5] [PMID: 7906891]
- [69] Garrett BE, Holtzman SG. Caffeine cross-tolerance to selective dopamine D1 and D2 receptor agonists but not to their synergistic interaction. *Eur J Pharmacol* 1994; 262(1-2): 65-75. [http://dx.doi.org/10.1016/0014-2999(94)90029-9] [PMID: 7813580]
- [70] Popoli P, Reggio R, Pèzzola A. Effects of SCH 58261, an adenosine A(2A) receptor antagonist, on quinpirole-induced turning in 6-hydroxydopamine-lesioned rats. Lack of tolerance after chronic caffeine intake. *Neuropsychopharmacology* 2000; 22(5): 522-9. [http://dx.doi.org/10.1016/S0893-133X(99)00144-X] [PMID: 10731627]
- [71] Ikeda K, Kurokawa M, Aoyama S, Kuwana Y. Neuroprotection by adenosine A2A receptor blockade in experimental models of Parkinson's disease. *J Neurochem* 2002; 80(2): 262-70. [http://dx.doi.org/10.1046/j.0022-3042.2001.00694.x] [PMID: 11902116]
- [72] Cronstein BN, Levin RI, Philips M, Hirschhorn R, Abramson SB, Weissmann G. Neutrophil adherence to endothelium is enhanced *via* adenosine A1 receptors and inhibited *via* adenosine A2 receptors. *J Immunol Baltim Md* 1950; 148(7): 2201-6.
- [73] Harada N, Okajima K, Murakami K, *et al.* Adenosine and selective A(2A) receptor agonists reduce ischemia/reperfusion injury of rat liver mainly by inhibiting leukocyte activation. *J Pharmacol Exp Ther* 2000; 294(3): 1034-42. [PMID: 10945856]
- [74] Haskó G, Kuhel DG, Chen JF, *et al.* Adenosine inhibits IL-12 and TNF- α production *via* adenosine A2a receptor-dependent and independent mechanisms. *FASEB J* 2000; 14(13): 2065-74. [http://dx.doi.org/10.1096/fj.99-0508com] [PMID: 11023991]
- [75] Ricardo C, Marco FA, Daniel T. Astrocytes Role in Parkinson: A Double-Edged Sword. In: Kishore U, Ed. *Neurodegenerative Diseases*. USA: InTech 2013. [http://dx.doi.org/10.5772/54305]
- [76] Lin C-C, Deneen B. Astrocytes The missing link in neurological disease? *Semin Pediatr Neurol* 2013; 20(4): 236-41.
- [77] Abbott NJ, Rönnbäck L, Hansson E. Astrocyte-endothelial interactions at the blood-brain barrier. *Nat Rev Neurosci* 2006; 7(1): 41-53. [http://dx.doi.org/10.1038/nrn1824] [PMID: 16371949]
- [78] Barreto GE, Gonzalez J, Torres Y, Morales L. Astrocytic-neuronal crosstalk: implications for neuroprotection from brain injury. *Neurosci Res* 2011; 71(2): 107-13. [http://dx.doi.org/10.1016/j.neures.2011.06.004] [PMID: 21693140]
- [79] Maragakis NJ, Rothstein JD. Mechanisms of Disease: astrocytes in neurodegenerative disease. *Nat Clin Pract Neurol* 2006; 2(12): 679-89. [http://dx.doi.org/10.1038/ncpneu0355] [PMID: 17117171]
- [80] Sian J, Youdim MB, Riederer P, Gerlach M. MPTP-Induced Parkinsonian Syndrome. In: George JS, Bernard WA, Wayne AR, Stephen KF, Michael DU, Eds. *Basic Neurochemistry*. 6th ed. Philadelphia: Lippincott-Raven 1999.
- [81] Boyd CS, Cadenas E. Nitric oxide and cell signaling pathways in mitochondrial-dependent apoptosis. *Biol Chem* 2002; 383(3-4): 411-23. [http://dx.doi.org/10.1515/BC.2002.045] [PMID: 12033432]

- [82] Horn TF, Wolf G, Duffy S, Weiss S, Keilhoff G, MacVicar BA. Nitric oxide promotes intracellular calcium release from mitochondria in striatal neurons. *FASEB J* 2002; 16(12): 1611-22. [<http://dx.doi.org/10.1096/fj.02-0126com>] [PMID: 12374784]
- [83] Varçin M, Bentea E, Michotte Y, Sarre S. Oxidative stress in genetic mouse models of Parkinson's disease. *Oxid Med Cell Longev* 2012; 2012; 624925.
- [84] Braidy N, Gai W-P, Xu YH, *et al.* Uptake and mitochondrial dysfunction of alpha-synuclein in human astrocytes, cortical neurons and fibroblasts. *Transl Neurodegener* 2013; 2(1): 20. [<http://dx.doi.org/10.1186/2047-9158-2-20>] [PMID: 24093918]
- [85] Martin HL, Teismann P. Glutathione-a review on its role and significance in Parkinson's disease. *FASEB J* 2009; 23(10): 3263-72. [<http://dx.doi.org/10.1096/fj.08-125443>] [PMID: 19542204]
- [86] Mytilineou C, Kramer BC, Yabut JA. Glutathione depletion and oxidative stress. *Parkinsonism Relat Disord* 2002; 8(6): 385-7. [[http://dx.doi.org/10.1016/S1353-8020\(02\)00018-4](http://dx.doi.org/10.1016/S1353-8020(02)00018-4)] [PMID: 12217624]
- [87] Liu Y, Qiang M, Wei Y, He R. A novel molecular mechanism for nitrated alpha-synuclein-induced cell death. *J Mol Cell Biol* 2011; 3(4): 239-49. [<http://dx.doi.org/10.1093/jmcb/mjr011>] [PMID: 21733982]
- [88] Przedborski S, Chen Q, Vila M, *et al.* Oxidative post-translational modifications of alpha-synuclein in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) mouse model of Parkinson's disease. *J Neurochem* 2001; 76(2): 637-40. [<http://dx.doi.org/10.1046/j.1471-4159.2001.00174.x>] [PMID: 11208927]
- [89] Lull ME, Block ML. Microglial activation and chronic neurodegeneration. *Neurotherapeutics* 2010; 7(4): 354-65. [<http://dx.doi.org/10.1016/j.nurt.2010.05.014>] [PMID: 20880500]
- [90] Chen JF, Xu K, Petzer JP, *et al.* Neuroprotection by caffeine and A(2A) adenosine receptor inactivation in a model of Parkinson's disease. *J Neurosci* 2001; 21(10): RC143. [PMID: 11319241]
- [91] Monif M, Burnstock G, Williams DA. Microglia: proliferation and activation driven by the P2X7 receptor. *Int J Biochem Cell Biol* 2010; 42(11): 1753-6. [<http://dx.doi.org/10.1016/j.biocel.2010.06.021>] [PMID: 20599520]
- [92] Chrovian CC, Rech JC, Bhattacharya A, Letavic MA. P2X7 antagonists as potential therapeutic agents for the treatment of CNS disorders. *Prog Med Chem* 2014; 53: 65-100. [<http://dx.doi.org/10.1016/B978-0-444-63380-4.00002-0>] [PMID: 24418608]
- [93] Marret S, Delpech B, Girard N, *et al.* Caffeine decreases glial cell number and increases hyaluronan secretion in newborn rat brain cultures. *Pediatr Res* 1993; 34(6): 716-9. [<http://dx.doi.org/10.1203/00006450-199312000-00004>] [PMID: 8108181]
- [94] Horrigan LA, Kelly JP, Connor TJ. Caffeine suppresses TNF-alpha production *via* activation of the cyclic AMP/protein kinase A pathway. *Int Immunopharmacol* 2004; 4(10-11): 1409-17. [<http://dx.doi.org/10.1016/j.intimp.2004.06.005>] [PMID: 15313438]
- [95] Matos M, Augusto E, Santos-Rodrigues AD, *et al.* Adenosine A2A receptors modulate glutamate uptake in cultured astrocytes and gliosomes. *Glia* 2012; 60(5): 702-16. [<http://dx.doi.org/10.1002/glia.22290>] [PMID: 22298379]
- [96] Orr AG, Orr AL, Li X-J, Gross RE, Traynelis SF. Adenosine A(2A) receptor mediates microglial process retraction. *Nat Neurosci* 2009; 12(7): 872-8. [<http://dx.doi.org/10.1038/nn.2341>] [PMID: 19525944]
- [97] Addicott MA, Yang LL, Peiffer AM, *et al.* The effect of daily caffeine use on cerebral blood flow: How much caffeine can we tolerate? *Hum Brain Mapp* 2009; 30(10): 3102-14. [<http://dx.doi.org/10.1002/hbm.20732>] [PMID: 19219847]
- [98] Field AS, Laurienti PJ, Yen Y-F, Burdette JH, Moody DM. Dietary caffeine consumption and withdrawal: confounding variables in quantitative cerebral perfusion studies? *Radiology* 2003; 227(1): 129-35. [<http://dx.doi.org/10.1148/radiol.2271012173>] [PMID: 12616005]
- [99] Berger SM, Bartsch D. The role of L-type voltage-gated calcium channels Cav1.2 and Cav1.3 in normal and pathological brain function. *Cell Tissue Res* 2014; 357(2): 463-76. [<http://dx.doi.org/10.1007/s00441-014-1936-3>] [PMID: 24996399]
- [100] Kang S, Cooper G, Dunne SF, *et al.* CaV1.3-selective L-type calcium channel antagonists as potential new therapeutics for Parkinson's disease. *Nat Commun* 2012; 3: 1146. [<http://dx.doi.org/10.1038/ncomms2149>] [PMID: 23093183]
- [101] Putzier I, Kullmann PH, Horn JP, Levitan ES. Cav1.3 channel voltage dependence, not Ca2+ selectivity, drives pacemaker activity and amplifies bursts in nigral dopamine neurons. *J Neurosci* 2009; 29(49): 15414-9. [<http://dx.doi.org/10.1523/JNEUROSCI.4742-09.2009>] [PMID: 20007466]

- [102] Schapira AH. Calcium dysregulation in Parkinson's disease. *Brain* 2013; 136(Pt 7): 2015-6. [http://dx.doi.org/10.1093/brain/awt180] [PMID: 23788521]
- [103] Pasternak B, Svanström H, Nielsen NM, Fugger L, Melbye M, Hviid A. Use of calcium channel blockers and Parkinson's disease. *Am J Epidemiol* 2012; 175(7): 627-35. [http://dx.doi.org/10.1093/aje/kwr362] [PMID: 22387374]
- [104] Sei Y, Gallagher KL, Daly JW. Multiple effects of caffeine on Ca²⁺ release and influx in human B lymphocytes. *Cell Calcium* 2001; 29(3): 149-60. [http://dx.doi.org/10.1054/ceca.2000.0175] [PMID: 11162852]
- [105] Klegeris A, Choi HB, McLarnon JG, McGeer PL. Functional ryanodine receptors are expressed by human microglia and THP-1 cells: Their possible involvement in modulation of neurotoxicity. *J Neurosci Res* 2007; 85(10): 2207-15. [http://dx.doi.org/10.1002/jnr.21361] [PMID: 17526017]
- [106] Baroudi G, Qu Y, Ramadan O, Chahine M, Boutjdir M. Protein kinase C activation inhibits Cav1.3 calcium channel at NH₂-terminal serine 81 phosphorylation site. *Am J Physiol Heart Circ Physiol* 2006; 291(4): H1614-22. [http://dx.doi.org/10.1152/ajpheart.00095.2006] [PMID: 16973824]
- [107] Postuma RB, Lang AE, Munhoz RP, *et al.* Caffeine for treatment of Parkinson disease: a randomized controlled trial. *Neurology* 2012; 79(7): 651-8. [http://dx.doi.org/10.1212/WNL.0b013e318263570d] [PMID: 22855866]
- [108] Tan E-K, Chua E, Fook-Chong SM, *et al.* Association between caffeine intake and risk of Parkinson's disease among fast and slow metabolizers. *Pharmacogenet Genomics* 2007; 17(11): 1001-5. [http://dx.doi.org/10.1097/FPC.0b013e3282f09265] [PMID: 18075470]
- [109] Qiu C, Hu G, Kivipelto M, *et al.* Association of blood pressure and hypertension with the risk of Parkinson disease: the National FINRISK Study. *Hypertension* 2011; 57(6): 1094-100. [http://dx.doi.org/10.1161/HYPERTENSIONAHA.111.171249] [PMID: 21536985]
- [110] Sibon I, Tison F. Vascular parkinsonism. *Curr Opin Neurol* 2004; 17(1): 49-54. [http://dx.doi.org/10.1097/00019052-200402000-00009] [PMID: 15090877]
- [111] Wooten GF, Currie LJ, Bovbjerg VE, Lee JK, Patrie J. Are men at greater risk for Parkinson's disease than women? *J Neurol Neurosurg Psychiatry* 2004; 75(4): 637-9. [http://dx.doi.org/10.1136/jnnp.2003.020982] [PMID: 15026515]
- [112] Palacios N, Gao X, McCullough ML, *et al.* Caffeine and risk of Parkinson's disease in a large cohort of men and women. *Mov Disord* 2012; 27(10): 1276-82. [http://dx.doi.org/10.1002/mds.25076] [PMID: 22927157]
- [113] Palacios N, Fitzgerald KC, Hart JE, *et al.* Particulate matter and risk of Parkinson disease in a large prospective study of women. *Environ Health* 2014; 13: 80. [http://dx.doi.org/10.1186/1476-069X-13-80] [PMID: 25294559]
- [114] Ascherio A, Weisskopf MG, O'Reilly EJ, *et al.* Coffee consumption, gender, and Parkinson's disease mortality in the cancer prevention study II cohort: the modifying effects of estrogen. *Am J Epidemiol* 2004; 160(10): 977-84. [http://dx.doi.org/10.1093/aje/kwh312] [PMID: 15522854]
- [115] Xu K, Xu Y, Brown-Jermyn D, *et al.* Estrogen prevents neuroprotection by caffeine in the mouse 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine model of Parkinson's disease. *J Neurosci* 2006; 26(2): 535-41. [http://dx.doi.org/10.1523/JNEUROSCI.3008-05.2006] [PMID: 16407551]
- [116] Moo-Puc RE, Villanueva-Toledo J, Arankowsky-Sandoval G, Alvarez-Cervera F, Góngora-Alfaro JL. Treatment with subthreshold doses of caffeine plus trihexyphenidyl fully restores locomotion and exploratory activity in reserpinized rats. *Neurosci Lett* 2004; 367(3): 327-31. [http://dx.doi.org/10.1016/j.neulet.2004.06.022] [PMID: 15337259]
- [117] Gevaerd MS, Miyoshi E, Silveira R, Canteras NS, Takahashi RN, Da Cunha C. L-Dopa restores striatal dopamine level but fails to reverse MPTP-induced memory deficits in rats. *Int J Neuropsychopharmacol* 2001; 4(4): 361-70. [http://dx.doi.org/10.1017/S1461145701002619] [PMID: 11806861]
- [118] Fredduzzi S, Moratalla R, Monopoli A, *et al.* Persistent behavioral sensitization to chronic L-DOPA requires A2A adenosine receptors. *J Neurosci* 2002; 22(3): 1054-62. [PMID: 11826134]
- [119] Bata-García JL, Villanueva-Toledo J, Gutiérrez-Ospina G, Alvarez-Cervera FJ, Heredia-López FJ, Góngora-Alfaro JL. Sustained improvement of motor function in hemiparkinsonian rats chronically treated with low doses of caffeine or trihexyphenidyl. *Pharmacol Biochem Behav* 2007; 86(1): 68-78. [http://dx.doi.org/10.1016/j.pbb.2006.12.009] [PMID: 17250882]
- [120] Casas M, Prat G, Robledo P, Barbanj M, Kulisevsky J, Jané F. Repeated co-administration of caffeine and bromocriptine prevents tolerance to the effects of caffeine in the turning behavior animal model. *Eur Neuropsychopharmacol* 1999; 9(6): 515-21. [http://dx.doi.org/10.1016/S0924-977X(99)00037-1] [PMID: 10625120]
- [121] Holloway RG, Shoulson I, Fahn S, *et al.* Pramipexole vs. levodopa as initial treatment for Parkinson disease: a 4-year randomized controlled

trial. *Arch Neurol* 2004; 61(7): 1044-53.
[PMID: 15262734]

- [122] Poewe WH, Rascol O, Quinn N, *et al.* Efficacy of pramipexole and transdermal rotigotine in advanced Parkinson's disease: a double-blind, double-dummy, randomised controlled trial. *Lancet Neurol* 2007; 6(6): 513-20.
[[http://dx.doi.org/10.1016/S1474-4422\(07\)70108-4](http://dx.doi.org/10.1016/S1474-4422(07)70108-4)] [PMID: 17509486]
- [123] Nakaso K, Ito S, Nakashima K. Caffeine activates the PI3K/Akt pathway and prevents apoptotic cell death in a Parkinson's disease model of SH-SY5Y cells. *Neurosci Lett* 2008; 432(2): 146-50.
[<http://dx.doi.org/10.1016/j.neulet.2007.12.034>] [PMID: 18201823]
- [124] Ascherio A, Chen H, Schwarzschild MA, Zhang SM, Colditz GA, Speizer FE. Caffeine, postmenopausal estrogen, and risk of Parkinson's disease. *Neurology* 2003; 60(5): 790-5.
[<http://dx.doi.org/10.1212/01.WNL.0000046523.05125.87>] [PMID: 12629235]
- [125] Kaasinen V, Aalto S, Nägren K, Rinne JO. Dopaminergic effects of caffeine in the human striatum and thalamus. *Neuroreport* 2004; 15(2): 281-5.
[<http://dx.doi.org/10.1097/00001756-200402090-00014>] [PMID: 15076753]

© Roshan *et al.*; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution-Non-Commercial 4.0 International Public License (CC BY-NC 4.0) (<https://creativecommons.org/licenses/by-nc/4.0/legalcode>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.