

ARTICLE

# Power of QTL detection using association tests with family controls

Jules Hernández-Sánchez<sup>\*1</sup>, Chris S Haley<sup>1</sup> and Peter M Visscher<sup>2</sup>

<sup>1</sup>Roslin Institute (Edinburgh), Roslin, Midlothian EH25 9PS, Scotland, UK; <sup>2</sup>Institute of Cell, Animal and Population Biology, University of Edinburgh, West Mains Road, Edinburgh EH9 3JT, Scotland, UK

The power of testing for a population-wide association between a biallelic quantitative trait locus and a linked biallelic marker locus is predicted both empirically and deterministically for several tests. The tests were based on the analysis of variance (ANOVA) and on a number of transmission disequilibrium tests (TDT). Deterministic power predictions made use of family information, and were functions of population parameters including linkage disequilibrium, allele frequencies, and recombination rate. Deterministic power predictions were very close to the empirical power from simulations in all scenarios considered in this study. The different TDTs had very similar power, intermediate between one-way and nested ANOVAs. One-way ANOVA was the only test that was not robust against spurious disequilibrium. Our general framework for predicting power deterministically can be used to predict power in other association tests. Deterministic power calculations are a powerful tool for researchers to plan and evaluate experiments and obviate the need for elaborate simulation studies.

*European Journal of Human Genetics* (2003) 11, 819–827. doi:10.1038/sj.ejhg.5201042

**Keywords:** TDT; association; power; robustness; linkage disequilibrium

## Introduction

Geneticists have been successful in mapping genes underlying rare, monogenic disorders with clear patterns of Mendelian inheritance.<sup>1–3</sup> However, mapping genes underlying complex traits, such as common multifactorial diseases, has been more difficult.<sup>4,5</sup> Genes can be mapped via linkage or association tests. Although both strategies exploit the cosegregation of markers with phenotypes, there are some striking differences between them. For example, in humans, genome-wide searches may require testing 30 000–500 000 single-nucleotide polymorphisms (SNPs) to detect significant associations, compared to 200–400 microsatellites in a linkage analysis.<sup>6–8</sup> Moreover, theoretical work suggests that finding associations between markers and complex diseases is more powerful than

searching for linkage, even if many SNPs have to be tested and significance thresholds are raised to compensate for multiple testing.<sup>9</sup> However, not all association tests are robust to spurious associations.<sup>10</sup> This explains why association tests using family controls, such as the transmission/disequilibrium test (TDT), have been favoured over association tests using random controls, for example, case-control, because the former are robust to spurious associations caused by population stratification, or recent admixture.<sup>11–13</sup>

Power studies of association tests will help researchers to design appropriate experiments, and to choose the most powerful test for the analysis of data. In this study, we investigate the power of association tests for quantitative traits, with and without family controls. Allison<sup>14</sup> proposed five TDTs (TDT<sub>Q1–Q5</sub>) for analysing quantitative traits under different ascertainment conditions, and we have included the most powerful, TDT<sub>Q5</sub>, in this study. Long and Langley<sup>15</sup> compared the power of five random controls association tests and the TDT<sub>Q5</sub>, and found that TDT<sub>Q5</sub> was always the least powerful test. Nevertheless,

\*Correspondence: Dr J Hernández-Sánchez, Roslin Institute (Edinburgh), Roslin, Midlothian EH25 9PS, Scotland, UK. Tel: 44131 527 4462; Fax: 44 131 440 0434; E-mail: jules.Hernandez@bbsrc.ac.uk  
Received 22 October 2002; revised 17 March 2003; accepted 16 April 2003

these authors acknowledged that under population stratification, the type-I error rate of association tests using random controls can rise above the nominal level set for the experiment. Xiong *et al*<sup>16</sup> proposed the TDT<sub>G</sub>, an extension of Allison's TDT<sub>Q1</sub> that accounts for any number of sibs per family, families with one or two heterozygous parents, and any number of alleles at the marker locus. They found that TDT<sub>G</sub> is more powerful than TDT<sub>Q1</sub>, the Haseman–Elston linkage test,<sup>17</sup> and an extreme discordant sib pair test.<sup>16</sup> Lastly, Rabinowitz<sup>18</sup> developed a TDT (referred to here as TDT<sub>R</sub>) to model explicitly the correlation between a quantitative trait and marker segregation.

In this study, several tests of association have been compared in terms of power, both empirically and deterministically. Our deterministic approximation for predicting power was based on the calculation of noncentrality parameters (NCPs).<sup>19</sup> The accuracy of this methodology, which can be used to predict power of other association tests, was validated via simulation.

**Materials and methods**

**Definition of the evaluated association tests**

The power of five tests to detect both linkage and/or association between a marker locus and a Quantitative Trait Locus (QTL) was studied empirically (simulations) and deterministically (calculation of NCP). Table 1 shows all the tests used in this study: the one-way analysis of variance (one-way ANOVA), the nested analysis of variance (nested ANOVA), the TDT<sub>Q5</sub>, the TDT<sub>R</sub>, and the TDT<sub>G</sub>.<sup>14,16,18,20</sup> A general deterministic method for predicting power at a linked marker is proposed in this study, and implementation examples are given for one-way ANOVA, TDT<sub>R</sub>, and TDT<sub>Q5</sub>. The one-way ANOVA test uses a sample of unrelated individuals who have been both genotyped and phenotyped, whereas the other four tests use the same but with the inclusion of the genotypes of the parents. Recombination rate and linkage disequilibrium were denoted *c* and *D*, respectively.

**One-way ANOVA** The one-way ANOVA contrasts marker genotype means among the progeny. This is the simplest and the most powerful test of association, although it is prone to high type-I error rates in the presence of spurious

association, viz. disequilibrium without linkage.<sup>15</sup> This is so because the null hypothesis (H<sub>0</sub>) being tested by one-way ANOVA is *no association*, regardless of linkage. Therefore, H<sub>0</sub> could be rejected when testing unlinked marker loci ( $c = \frac{1}{2}$ ) if there was a sufficiently strong population-wide association ( $D \neq 0$ ). This lack of robustness is common to tests that do not use family controls, for example, case–control studies.<sup>12,13</sup> The test statistic follows an F<sub>2,n'-3</sub> distribution under H<sub>0</sub>, given a total sample size *n'* and three different genotype groups.<sup>20</sup> The sum of squares between genotype groups, after subtracting the overall mean effect, reflects differences between marker genotypes. Hence, a significant statistic suggests greater differences between genotypes than would be expected under the assumption of linkage equilibrium between the QTL and the marker. Under the alternative hypothesis (H<sub>1</sub>) of association, the distribution of the test statistic is a noncentral F with NCP equal to  $\lambda_0$ , or a noncentral  $\chi^2_{2,\lambda_0}/2$  for large *n*.

**Nested ANOVA** A way of overcoming the lack of robustness of one-way ANOVA is to contrast marker genotype means of progeny within parental types, using a nested ANOVA design.<sup>20</sup> Parental type represents a particular combination of parental marker genotypes, and family type a particular combination of marker genotypes across all family members (Table 2). Thus, the H<sub>0</sub> being tested by

**Table 2** Probability of each family type given a biallelic marker and assuming Hardy–Weinberg equilibrium

FT	PT	Child	Probability
1	MM × MM	MM	P <sub>M</sub> <sup>4</sup>
2	MM × Mm	MM	2P <sub>M</sub> <sup>3</sup> P <sub>m</sub>
3		Mm	2P <sub>M</sub> <sup>3</sup> P <sub>m</sub>
4	MM × mm	Mm	2P <sub>M</sub> <sup>2</sup> P <sub>m</sub> <sup>2</sup>
5	Mm × Mm	MM	P <sub>M</sub> <sup>2</sup> P <sub>m</sub> <sup>2</sup>
6		Mm	2P <sub>M</sub> <sup>2</sup> P <sub>m</sub> <sup>2</sup>
7		mm	P <sub>M</sub> <sup>2</sup> P <sub>m</sub> <sup>2</sup>
8	Mm × mm	Mm	2P <sub>M</sub> P <sub>m</sub> <sup>3</sup>
9		mm	2P <sub>M</sub> P <sub>m</sub> <sup>3</sup>
10	mm × mm	mm	P <sub>M</sub> <sup>4</sup>

FT: family Type, PT: parental Type; P<sub>M</sub> (P<sub>m</sub>): frequency of allele M (m).

**Table 1** Main features of the tests compared in this study assuming family trios

Abbreviation	Testing	H <sub>0</sub>	Reference
One-way ANOVA	Genotype effects	Association	20
TDT <sub>Q5</sub>	Genotype effects after correcting for parental type effects	Linkage and association	14
TDT <sub>R</sub>	Allele–phenotype correlation	Linkage and association	18
TDT <sub>G</sub>	Allele means difference	Linkage and association	16
Nested ANOVA	Genotype effects within parental type	Linkage and association within parental type	20

nested ANOVA is *no association within parental types*. There must be at least two progeny with different marker genotypes within each parental type for there to be a contrast; therefore, only those families with at least one heterozygous parent, that is, informative families, are used. This type of family ascertainment increases the degrees of freedom (df) between groups and reduces the df within groups, resulting in a loss of power compared to one-way ANOVA. The appropriate F-test in nested ANOVA is a ratio of between to within genotype mean squares within parental types. The test follows an  $F_{\alpha,\beta}$  distribution under  $H_0$ , where  $\alpha = \sum_{i=1}^{\gamma} ng_i - 1$ , and  $ng_i$  is the observed number of genotypes within parental type  $i$ ,  $\gamma$  the observed number of parental types, and  $\beta = n - (\alpha + \gamma)$ , where  $n$  is the number of informative families.

**TDT<sub>Q5</sub>** The original statistic for TDT<sub>Q5</sub> is  $[(SS_F - SS_R)/2] / [(SS_T - SS_F)/(n - 5)]$ , where  $SS_T$  is the total sum of squares,  $SS_R$  is the sum of squares explained by a reduced model that fits an overall mean and two (out of three) informative parental types as fixed factors, and  $SS_F$  is the sum of squares explained by a full model that fits, in addition to the reduced model, two more fixed factors to estimate additive and dominant effects.<sup>14</sup> The total number of informative families is  $n$ . TDT<sub>Q5</sub> is testing whether a significant amount of phenotypic variation can be explained by marker genotypes in the progeny, over and above the variation already explained by parental type. TDT<sub>Q5</sub> follows an  $F_{2,n-5}$  distribution under  $H_0$  if residuals are normally distributed, or  $\chi^2_2/2$  for large  $n$ . Under  $H_1$ , TDT<sub>Q5</sub> follows a noncentral  $F_{2,n-5,\lambda_{Q5}}$ , or a noncentral  $\chi^2_{2,\lambda_{Q5}}/2$  for large  $n$ , with NCP  $\lambda_{Q5}$ . The TDT<sub>Q5</sub> is equivalent to a two-way ANOVA with a cross-classified design where the factors are parental type and progeny marker genotype (Appendix B).

**TDT<sub>R</sub>**

Although Rabinowitz<sup>18</sup> derived a NCP, he used parameters not included in his simulations, leading to some confusion in interpreting and calculating  $\lambda_R$ . Therefore, we developed a neater NCP for TDT<sub>R</sub>. The TDT<sub>R</sub> is calculated as  $T/\sigma_T$  for a biallelic marker.  $T$  measures the strength of the covariance between the transmission of a marker allele, from heterozygous parents to progeny, and the phenotype of progeny, and  $\sigma_T$  is the standard deviation of  $T$ . We will next describe TDT<sub>R</sub> in detail, as this information will be needed for further statistical developments. The numerator is  $T = \sum_i^n (y_i - \bar{y})w_i$ , where  $y_i$  is the phenotype of the  $i$ th child,  $\bar{y}$  is the overall mean (or the mean among informative families), and  $w_i$  are weights given to each family type (Table 3). The sum is over  $n$  informative and unrelated family trios randomly drawn from a population. The variance of  $T$  is  $\sigma_T^2 = \frac{1}{4} \sum_i^n (y_i - \bar{y})^2 h_i$  where  $h_i$  is the number of heterozygous parents in the family (Table 3). Under  $H_0$ , TDT<sub>R</sub> follows a  $t_{n-1}$  distribution, so  $(TDT_R)^2$  follows an  $F_{1,n-1}$  distribution. Under the alternative

**Table 3** Variables in TDT<sub>R</sub>

FT	t	w	h	Effect
1	*	0	0	b <sub>1</sub>
2	1	1/2	1	b <sub>2</sub>
3	0	-1/2	1	b <sub>3</sub>
4	*	0	0	b <sub>4</sub>
5	1	1	2	b <sub>5</sub>
6	*	0	2	b <sub>6</sub>
7	0	-1	2	b <sub>7</sub>
8	1	1/2	1	b <sub>8</sub>
9	0	-1/2	1	b <sub>9</sub>
10	*	0	0	b <sub>10</sub>

FT: family type;  $t$  is 1(0) if a parent Mm transmits M(m), or \* if the family is uninformative;  $w = \sum \text{het}(t-1/2)$ , sum over both parents;  $\text{het}$  is 1(0) if a parent is Mm(not Mm); Effect: Expected marker effect within families;  $h$ : number of heterozygous parents.

hypothesis,  $(TDT_R)^2$  follows a noncentral F with NCP  $\lambda_R$ , or a noncentral  $\chi^2_{1,\lambda_R}$  for large  $n$ .

**TDT<sub>G</sub>** The last test being considered is TDT<sub>G</sub>.<sup>16</sup> For a biallelic marker  $TDT_G = (\bar{y}_M - \bar{y}_m)^2 / (\frac{1}{n_M} + \frac{1}{n_m})s^2$  where  $\bar{y}_M$  ( $\bar{y}_m$ ) is the mean among progeny having inherited allele M (m) from heterozygous parents, and  $n_M$  ( $n_m$ ) is the number of times allele M (m) is transmitted. The variance of  $(\bar{y}_M - \bar{y}_m)$  is

$$\left(\frac{1}{n_M} + \frac{1}{n_m}\right)s^2$$

where

$$s^2 = \frac{\sum_{k=1}^{2n} [(y_{Mk} - \bar{y}_M)^2 + (y_{mk} - \bar{y}_m)^2]}{n_M + n_m - 2}$$

and  $y_{Mk}$  is the phenotype of the child of the  $k$ th parent. The latter sum is over the  $2n$  parents in a sample of  $n$  family trios. If all family members are Mm heterozygous, then the same information is included in both allele categories. For a normally distributed trait and large  $n$ , TDT<sub>G</sub> follows a  $\chi^2_1$  distribution under  $H_0$ . The asymptotic distribution under  $H_1$  is a noncentral  $\chi^2_{1,\lambda_G}$  with NCP  $\lambda_G$ .

**Empirical power**

Power was calculated empirically as the proportion of significant results out of 1000 analyses of independent data sets, simulated under specific combinations of parameter values. Each sample consisted of  $n = 200$  unrelated family trios (father, mother, and a single child). The frequencies of the positive allele (Q) from a biallelic QTL were  $p_Q = [0.5, 0.3, 0.1]$ , and the same frequencies were assigned to allele M from a biallelic marker linked to the QTL. The recombination rates between the marker and the QTL were  $c = [0, 0.1, 0.3, 0.4, 0.5]$ . QTL and marker genotypes were generated for all individuals. Phenotypes were generated

**Table 4** Conditional QTL genotype probabilities in a child, given the family type (FT), and population parameters  $D$ ,  $c$ ,  $P_M$ ,  $P_m$ ,  $P_Q$ , and  $P_q$ 

FT	QQ	Qq	qq
1	$h_1^2 / p_M^2$	$2 h_1 h_3 / p_M^2$	$h_3^2 / p_M^2$
2	$h_1(h_1 p_m - cD) / p_M^2 p_m$	$[2h_1 h_3 p_m + (h_1 - h_3)cD] / p_M^2 p_m$	$h_3(h_3 p_m + cD) / p_M^2 p_m$
3	$h_1(h_2 p_M + cD) / p_M^2 p_m$	$[p_M (h_1 h_4 + h_2 h_3) + cD(h_3 - h_1)] / p_M^2 p_m$	$h_3(h_4 p_M - cD) / p_M^2 p_m$
4	$h_1 h_2 / p_M p_m$	$(h_1 h_4 + h_2 h_3) / p_M p_m$	$h_3 h_4 / p_M p_m$
5	$[(h_1 p_m - cD) / (p_M p_m)]^2$	$2[h_1 h_3 p_m + cD(h_1 - h_3)p_m - c^2 D^2] / (p_M p_m)^2$	$[(h_3 p_m + cD) / (p_M p_m)]^2$
6	$[h_1 h_2 p_M p_m + c(1 - c)D^2] / (p_M p_m)^2$	$[(h_1 h_4 + h_2 h_3)p_M p_m - 2c(1 - c)D^2] / (p_M p_m)^2$	$[h_3 h_4 p_M p_m + c(1 - c)D^2] / (p_M p_m)^2$
7	$[(h_2 p_M + cD) / (p_M p_m)]^2$	$2[h_2 h_4 p_M + cD(h_4 - h_2)p_M - c^2 D^2] / (p_M p_m)^2$	$(h_4 p_M - cD)^2 / (p_M p_m)^2$
8	$h_2(h_1 p_m - cD) / (p_M p_m^2)$	$[(h_1 h_4 + h_2 h_3)p_m + cD(h_2 - h_4)] / (p_M p_m^2)$	$h_4(h_3 p_m + cD) / (p_M p_m^2)$
9	$h_2(h_2 p_M + cD) / (p_M p_m^2)$	$[2p_M h_2 h_4 - (h_2 - h_4)cD] / (p_M p_m^2)$	$h_4(h_4 p_M - cD) / (p_M p_m^2)$
10	$h_2^2 / p_m^2$	$2h_2 h_4 / p_m^2$	$h_4^2 / p_m^2$

$D$ : linkage disequilibrium;  $c$ : recombination rate;  $p_M(p_m)$ : frequency of marker allele M (m);  $p_Q(p_q)$ : frequency of QTL allele Q (q);  $h_1, h_2, h_3, h_4$ : frequencies of haplotypes QM, Qm, qM, and qm, respectively, where,  $h_1 = p_Q p_M + D$ ;  $h_2 = p_Q p_m - D$ ;  $h_3 = p_q p_M - D$ ; and  $h_4 = p_q p_m + D$ . Note that, for example, to calculate the second row of probabilities, we divide by  $p_M^2 p_m$  instead of by  $2p_M p_m$  (see Table 2) as we consider one FT, MM  $\times$  Mm, but not the reciprocal Mm  $\times$  MM.

only for the progeny by adding a normally distributed error with variance  $\sigma_e^2 = 1$ , plus  $\frac{-1}{2}$ , 0, or  $\frac{1}{2}$  for QTL genotypes qq, qQ, or QQ, respectively. Neither dominance nor polygenic effects were simulated. The level of association between allele Q at the QTL and allele M at the marker was given by the standardised linkage disequilibrium parameter  $D' = [0, \frac{1}{2}, 1]$ .<sup>21</sup>

### Deterministic power

We have developed a compound method with two parts for predicting power of association tests deterministically. The first part consisted in calculating the expected effect of marker genotypes as functions of underlying QTL genotypes, conditional on population parameters and family type. This part can be used to predict power in other association tests, in addition to the ones in this study. The second part consisted in calculating the NCP as a function of marker contrasts specific to each test.

**Expected marker effects** Consider the 10 different family types at a biallelic marker (Table 2), and let  $X_j$  be a vector with the marker genotypes of child, father, and mother in a family of type  $j$ , for example,  $X_1 = [MM, MM, MM]$ . Let  $G_i$  denote the  $i$ th QTL genotype of the child, that is,  $G_1 = QQ$ ,  $G_2 = Qq$ , and  $G_3 = qq$ . The expected phenotype ( $y$ ) of a child given the  $i$ th family type, assuming no dominance, is

$$E[y|X_i] = a[P(G_1|X_i) - P(G_3|X_i)] \quad (1)$$

where  $a$  is the effect of substituting allele q for Q, assumed to be  $\frac{1}{2}$ . The conditional probabilities  $P(G_1|X_i)$  and  $P(G_3|X_i)$  can be calculated using Tables W1–W4, available on the web ([www.nature.com/ejhg/5201042](http://www.nature.com/ejhg/5201042)).<sup>22,23</sup> For example,

the probability of QTL genotype QQ given  $X_1$  is

$$P(G_1|X_1) = \frac{P(G_1 \cap X_1)}{P(X_1)} = \frac{h_1^2 p_M^2}{p_M^4} = \frac{h_1^2}{p_M^2}$$

where  $P(G_1 \cap X_1)$  is the joint probability of QTL genotype QQ in the child and marker genotype MM in all members of the family,  $P(X_1)$  is the probability of family type 1 (Table 2), and  $h_1$  is the probability of drawing haplotype QM from the population which, assuming random mating and no segregation distortion, is  $h_1 = h_{QM} = p_Q p_M + D_{QM}$ . (Note:  $D_{QM} = D' D_{\max}$ , and if  $D' > 0$  then  $D_{\max} = \min\{p_q p_M, p_Q p_m\}$ .)<sup>24</sup> The joint probability  $P(G_1 \cap X_1)$  can be obtained from Table W4 by multiplying the third and the sixth columns and adding up all. The conditional probabilities  $P[G_i|X_j]$ , for  $i = 1, 2, 3$  and  $j = 1 \dots 10$ , are all summarised in Table 4.

**Noncentrality parameters (NCP)** The NCP for the one-way ANOVA ( $\lambda_O$ ) can be obtained applying the formula<sup>25</sup>

$$\lambda_O \sim \frac{B'X'XB}{\sigma_e^2} = \frac{\sum \mu_i^2 n_i}{\sigma_e^2} \quad (2)$$

The sum in Equation (2) is over all three marker genotype classes, the vector  $B'$  contains the three marker genotype means  $[\mu_{MM}, \mu_{Mm}, \mu_{mm}]$ , and  $X'X$  is a matrix with diagonal elements  $[n_{MM}, n_{Mm}, n_{mm}]$  and zeroes elsewhere, where  $n_i$  is the sample size corresponding to marker genotype  $i$ . Equation (2) represents the sum of squares due to both the marker locus and the sample mean ( $\mu$ ). The appropriate  $\lambda_O$  can be obtained after subtracting from Eq. (2) the sum of squares due to the sample mean, that is,  $n'\mu^2$ , where  $n' = n_{MM} + n_{Mm} + n_{mm}$ . When testing the QTL (ie conditioning on  $c = 0$ ,  $D' = 1$ , and  $p_Q = p_M$ ), and assuming no

dominance, Eq. (2) simplifies to

$$\lambda_O = n' \frac{\sigma_{QTL}^2}{\sigma_e^2} \quad (3)$$

where  $\sigma_{QTL}^2 = 2p_Q p_q a^2$ .<sup>26</sup>

In Appendix B, we have shown that TDT<sub>Q5</sub> is equivalent to a two-way ANOVA analysis, where data are modelled fitting parental type and progeny genotype as fixed factors, in addition to  $\mu$ . Taking this equivalence into account, the NCP  $\lambda_{Q5}$ , derived in Appendix A, is

$$\lambda_{Q5} = \frac{\sum_{i=1}^{10} b_i^2 n_i I_i - \sum_{j=1}^6 F_j^2 f_j I_j}{\sigma_e^2} \quad (4)$$

where  $b_i$  is the expected marker genotype effect in progeny of family type  $i$  (Table 3),  $n_i$  is the number of type  $i$  families,  $I_{i(j)}$  is an indicator variable that takes the value 1 when the family is informative (viz. at least one heterozygous parent), and 0 otherwise,  $F_j$  is the mean value of the  $j$ th parental type, and  $f_j$  the number of  $j$  parental types. Eq. (4) measures, in  $\sigma_e^2$  units, the amount of total sum of squares explained by the marker, after subtracting the parental type effect. When testing the QTL, Eq. (4) reduces to

$$\lambda_{Q5} = n \frac{\sigma_{QTL}^2}{2\sigma_e^2} = \frac{\lambda_O}{2} \quad (5)$$

The NCP for TDT<sub>R</sub> ( $\lambda_R$ ) is approximately

$$\lambda_R \approx [p_M^2(b_2 - b_3) + p_M p_m(b_5 - b_7) + p_m^2(b_8 - b_9)] \times \sqrt{\frac{np_M p_m}{\sigma_e^2 + \sigma_{QTL}^2}} \quad (6)$$

(Appendix C). When testing the QTL, Eq. (6) simplifies to

$$\lambda_R \approx a \sqrt{\frac{np_Q p_q}{\sigma_e^2 + \sigma_{QTL}^2}} = \sqrt{\frac{n}{2 + 2\sigma_e^2/\sigma_{QTL}^2}} \quad (7)$$

Finally, the NCP for TDT<sub>G</sub> ( $\lambda_G$ ) is<sup>16</sup>

$$\lambda_G \approx n \frac{[(1 - 2c)Da]^2}{p_M p_m (\sigma_e^2 + \sigma_{QTL}^2)} \quad (8)$$

where  $n$  is the number of informative families. When testing the QTL in family trios, the appropriate NCP is<sup>16</sup>


$$\lambda_G = \frac{n}{2} \frac{\sigma_{QTL}^2}{\sigma_e^2 + \sigma_{QTL}^2/2} = \frac{n}{1 + (2\sigma_e^2/\sigma_{QTL}^2)} \quad (9)$$

The differences between the four NCPs  $\lambda_O$ ,  $\lambda_{Q5}$ ,  $\lambda_G$ , and  $\lambda_R$  are easily appreciated in Table 5, for both large and small sample sizes. In all cases, the QTL allele frequency and effect size only affect  $\lambda$  through the QTL variance.

**Table 5** Noncentrality parameters ( $\lambda$ ) given  $c=0$  and  $D'=1$ , and distribution under  $H_0$  for small and large sample sizes

Noncentrality parameter	Small sample	Large sample
$\lambda_O = n' \sigma_{QTL}^2 / \sigma_e^2$	$F_{2, n' - 3}$	$\chi_2^2 / 2$
$\lambda_{Q5} = n \sigma_{QTL}^2 / 2 \sigma_e^2$	$F_{2, n - 5}$	$\chi_2^2 / 2$
$\lambda_R = n \sigma_{QTL}^2 / 2(\sigma_e^2 + \sigma_{QTL}^2)$	$t_{n-1}^2$	$\chi_1^2$
$\lambda_G = n \sigma_{QTL}^2 / (2\sigma_e^2 + \sigma_{QTL}^2)$	$t_{n-1}^2$	$\chi_1^2$

<sup>a</sup> $\lambda_O$ : one-way ANOVA,  $\lambda_{Q5}$ : TDT<sub>Q5</sub>,  $\lambda_R$ : (TDT<sub>R</sub>)<sup>2</sup>,  $\lambda_G$ : TDT<sub>G</sub>;  $n'(n)$ : total number of (informative) family trios;  $\sigma_{QTL}^2$ : QTL variance;  $\sigma_e^2$ : error variance.



**Figure 1** Empirical (lines) versus deterministic (points) power for one-way ANOVA, TDT<sub>Q5</sub>, and TDT<sub>R</sub>, across  $c$  and three allele frequencies: 0.5 (circles), 0.3 (triangles), and 0.1 (squares).  $D'$  was averaged out.

## Results

### Empirical versus deterministic power

We have developed formulae to calculate NCPs for one-way ANOVA ( $\lambda_O$ ), TDT<sub>Q5</sub> ( $\lambda_{Q5}$ ), and TDT<sub>R</sub> ( $\lambda_R$ ), assuming that

**Table 6** Empirical power (%) of tests per single parameter

	One-way ANOVA	TDT <sub>Q5</sub>	TDT <sub>G</sub>	TDT <sub>R</sub>	Nested ANOVA
<i>(a) Averaging across D' and p</i>					
<i>c</i>					
0	46.5	39.2	38.4	39.7	28.5
0.1	42.8	31.4	29.3	28.4	20.7
0.2	37.6	22.4	19.2	19.6	13.5
0.3	33.3	13.5	11.8	10.9	8.4
0.4	26.7	7.8	6.8	6.4	6.1
0.5	20.5	5.5	5.4	4.8	4.9
<i>(b) Averaging across c and p</i>					
<i>D'</i>					
1	71.9	38.6	35.5	34.1	25
0.5	27.2	15	14	13.1	9.2
0	5	5.1	5.2	4.7	5.2
<i>(c) Averaging across D' and c</i>					
<i>p</i>					
0.5	41.4	23.6	21.9	21.4	15.8
0.3	38.6	21.7	19.9	19.5	14.1
0.1	24.1	13.3	12.9	11.1	9.6

*c*: recombination rate; *D'*: standardised linkage disequilibrium; *p*: frequency of alleles Q and M (assumed equal). The sample consisted of 200 unrelated family trios. The QTL effects were 0,  $\frac{1}{2}$ , or 1 for qq, qQ, or QQ genotypes, respectively.

the sample consists of family trios. Once these  $\lambda$ 's are obtained, power can be calculated from the appropriate noncentral distributions. Xiong *et al*<sup>16</sup> derived the equation for the NCP of TDT<sub>G</sub> ( $\lambda_G$ ). Figure 1 shows that predictions of power using our deterministic method (lines) match very well the simulation results (points). Power is shown as a function of *c* for three different allele frequencies denoted with circles (*p*=0.5), triangles (*p*=0.3), and squares (*p*=0.1), while averaging out *D'*. The NCP of nested ANOVA can also be calculated following this method; however, simulation results showed that nested ANOVA is the least powerful method by far, and therefore we concentrated on deriving the other NCPs. In addition to the close match between deterministic and empirical power, two other features in Figure 1 are worth mentioning. First, power decayed more when *p* dropped from 0.3 to 0.1, than when it dropped from 0.5 to 0.3. This is because the loss of information is relatively more important in the former than in the latter drop. Second, TDT<sub>Q5</sub> was less powerful than TDT<sub>R</sub>, whereas the contrary was true in Table 6. This can be explained by the fact that, in Figure 1, TDT<sub>Q5</sub> was implemented as described by Allison,<sup>14</sup> that is, using only informative families, and estimating both additive and dominant effects. The NCP  $\lambda_{Q5}$  was obtained assuming this model. However, the power of TDT<sub>Q5</sub> increases when the dominant parameter need not be estimated.

### Power ranking with more powerful models via simulations

The power of TDT<sub>Q5</sub> increases after removing the dominance parameter from the model when it is redundant, that is, the QTL has additive effects only. A further improvement in power, albeit slight, can be achieved by using all six parental types, whether informative or not. In doing so, TDT<sub>Q5</sub> follows an  $F_{1,n'-4}$  distribution under  $H_0$ , as opposed to an  $F_{2,n-5}$ , where *n'* (*n*) is the total number of (informative) families. Likewise, one-way ANOVA can become more powerful, fitting a simple regression line across genotypes to estimate additive QTL effects. Thus, one-way ANOVA will be distributed as  $F_{1,n'-2}$  under  $H_0$ , as opposed to  $F_{2,n'-3}$ . All other tests remained unchanged, and power was estimated for all via simulations.

Table 6 shows empirical power across tests, focusing on each parameter at a time (*c*, *p* or *D'*), averaging across the other two parameters. The ranking of the tests in terms of power was the same across scenarios: first the one-way ANOVA, followed by TDT<sub>Q5</sub>, TDT<sub>G</sub>, and TDT<sub>R</sub> (the last two with similar power), and lastly nested ANOVA. Table 6(a) shows power of the tests for a given *c*, averaging across values of *D'* and *p*. The last row in Table 6(a) corresponds to the empirical type-I error for each test, ie,  $c=\frac{1}{2}$ . The one-way ANOVA was the only test for which the empirical error exceeded the nominal 5%. This is caused by the fact that one-way ANOVA is testing whether *D'* is significantly different from zero, regardless of *c*.<sup>15</sup> Power declined steadily as *c* increased, because the amount of  $\sigma_{QTL}^2$  explained by the marker decreased as interloci distance increased.

Table 6(b) shows power for a given *D'*, averaging across values of *p* and *c*. The power of one-way ANOVA reached ~72% when *D'* = 1, being approximately twice as powerful as the TDTs. Undoubtedly, if spurious association is not an issue, significant extra power can be obtained by testing genotype differences directly, as opposed to using robust tests. All tests showed ~5% type-I error when *D'* = 0, even for *c* = 0.

Finally, Table 6(c) shows power for a given *p*, averaging across values of *c* and *D'*. Power decays as allele frequency becomes more extreme because (1) there are less informative families, and (2) the proportion of informative families with two heterozygous parents decreases. The first point directly causes a reduction in sample size. The second point means that less  $\sigma_{QTL}^2$  is available to TDTs. TDTs owe their robustness to the fact that they use only within-family genetic variation, which is greater in families with two heterozygous parents. These results contrast with those of Allison,<sup>14</sup> who concluded that power increases as *p* decreases. However, Allison<sup>14</sup> kept  $\sigma_{QTL}^2$  constant, so as *p* became more extreme, the QTL effect, and the mean difference between marker genotypes, increased, resulting in more powerful contrasts.

## Discussion

A comprehensive review of methodology developed in the 1990s provided more than 60 references of association tests for monogenic diseases with Mendelian inheritance, and only about a dozen references of association tests for complex diseases.<sup>27</sup> Nevertheless, complex diseases are by far the commonest human ailments; for example, infectious and parasitic diseases, psychiatric disorders, and cardiovascular diseases affect ~44% of the world population, compared to just 0.05% of Caucasians being affected by cystic fibrosis, the commonest of the monogenic diseases.<sup>28,29</sup>

TDTs are increasingly used to identify QTLs underlying complex diseases because they can be more powerful than other tests, for example, linkage analysis, when markers are tightly linked to responsible QTLs, and because they are robust to spurious associations generated by common demographic events such as population stratification and/or admixture.<sup>8,10</sup>

We have developed and verified deterministic power calculations for a range of association tests for quantitative traits, that is, three TDTs and two ANOVAs, and shown how the power depends on the effect of a QTL, the recombination rate between a QTL and a marker, and the amount of linkage disequilibrium between marker and QTL. In this study, we have assumed that both loci were biallelic, and shared the same allele frequencies. Moreover, we considered a continuously distributed trait genetically determined by a single additive QTL, without polygenic component or dominance. This simplistic scenario was chosen to facilitate the derivation of NCPs for predicting power. Nonetheless, we recognise that a more comprehensive picture of the properties of these tests requires analyses of more realistic situations, for example, including dominance and polygenic effects, which is possible within the framework presented here.

The deterministic method proposed in this study consists in deriving NCPs ( $\lambda$ 's) as functions of marker genotype contrasts specific to each test. These  $\lambda$ 's can subsequently be used to obtain power. A common feature across all  $\lambda$ 's was the use of expected marker genotype means, conditional on family information, under the assumptions of random mating and no segregation distortion. The marker effects were functions of the standardised linkage disequilibrium ( $D'$ ), the recombination rate ( $c$ ), the allele frequencies ( $p_Q$ ,  $p_M$ ), and the size of the QTL ( $a$ ). Allison<sup>14</sup> derived  $\lambda_{Q5}$  for TDT<sub>Q5</sub> when the marker is the trait locus, and we have obtained an alternative prediction of  $\lambda_{Q5}$  for any recombination rate, and linkage disequilibrium in the parent population.

Power was also predicted empirically via stochastic simulations, and results confirmed the accuracy of our deterministic predictions. The advantages of deterministic over stochastic methods are (1) ease of implementation, (2) instant predictions, and (3) direct appreciation of the relationship between population parameters and power. However, deriving NCPs

becomes cumbersome in complex scenarios. Thus, in these cases, empirical simulations are invaluable.

The tests ranked as follows in terms of power. The one-way ANOVA was the most powerful test of association across all scenarios, but also the only test not robust to spurious disequilibrium. The TDTs had similar, and intermediate, power. However, we showed how to increase the power of TDT<sub>Q5</sub> compared to the original version, if there is no dominance. Lastly, the nested ANOVA was the least powerful test of association.

The power of TDT<sub>Q5</sub> may have been previously over-emphasised because complete linkage and linkage disequilibrium between marker and QTL were assumed, and family trios were sampled from a population of informative families.<sup>14</sup> This sampling scheme means that the variance explained by the QTL is larger in the sample of informative trios than in the population at large, which would include both informative and noninformative families, and led to the counter-intuitive conclusion that the more extreme the allele frequency, the higher the power of TDT<sub>Q5</sub> to detect associations. In addition, Allison's<sup>14</sup> comparison between TDT<sub>Q5</sub> and the Haseman–Elston linkage test<sup>17</sup> favours TDT<sub>Q5</sub> because this is a test for association, and a perfect association was assumed, whereas the Haseman–Elston test is for linkage.

In summary, a new and accurate deterministic method has been developed to predict the power of QTL detection for TDTs and ANOVAs, as a function of population parameters. We have obtained specific formulae for the NCPs of the tests, when the marker is the QTL, as functions of sample size and QTL heritability. The method contains a general part (Table 4) that can be used to calculate NCPs for other association tests. Moreover, our method can also model dominant QTL effects, and a polygenic component. Extensions to cope with multiallelic markers are theoretically possible, although future association studies in human populations are more likely to employ vast arrays of SNPs than multiallelic markers.<sup>30,31</sup> Therefore, further developments of these approaches ought to be directed to coping with the problem of simultaneous testing of several loci, and the study of haplotypes.

## Acknowledgements

We are grateful to Ian White and Dr O Southwood for helpful comments on earlier versions of this manuscript. This work has been supported by Sygen International, and by the Biotechnology and Biological Sciences Research Council of UK.

## References

- 1 Kerem B, Rommens JM, Buchanan JA *et al*: Identification of the cystic fibrosis gene: genetic analysis. *Science* 1989; **245**: 1073–1079.
- 2 Hastbäckä J, de la Chapelle A, Kaitila I, Sistonen P, Weaver A, Lander E: Linkage disequilibrium mapping in isolated founder

- populations: diastrophic dysplasia in Finland. *Nat Genet* 1992; 2: 204–211.
- 3 Hastbäck J, de la Chapelle A, Mahtani MM *et al*: The diastrophic dysplasia gene encodes a novel sulfate transporter: positional cloning by fine-structure linkage disequilibrium mapping. *Cell* 1994; 78: 1073–1087.
  - 4 Terwilliger JD, Weiss KM: Linkage disequilibrium mapping of complex disease: fantasy or reality? *Curr Opin Biotech* 1998; 9: 578–594.
  - 5 Schork NJ, Cardon LR, Xu X: The future of genetic epidemiology. *Trends Genet* 1998; 14: 266–272.
  - 6 Kruglyak L: Prospects for whole-genome linkage disequilibrium mapping of common disease genes. *Nat Genet* 1999; 22: 139–144.
  - 7 Ott J: Predicting the range of linkage disequilibrium. *Proc Natl Acad Sci USA* 2000; 97: 2–3.
  - 8 Neale MC, Cherny SS, Sham PC *et al*: Distinguishing population stratification from genuine allelic effects with MX: association of ADH2 with alcohol consumption. *Behav Genet* 1999; 29: 233–243.
  - 9 Risch N, Merikangas K: The future of genetic studies of complex human diseases. *Science* 1996; 273: 1516–1517.
  - 10 Wright AF, Carothers AD, Pirastu M: Population choice in mapping genes for complex diseases. *Nat Genet* 1999; 23: 397–404.
  - 11 Spielman RS, McGinnis RE, Ewens WJ: Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am J Hum Genet* 1993; 52: 506–516.
  - 12 Clayton D: Population association; in Balding DJ, Bishop M, Cannings C (eds) *Handbook of statistical genetics*. New York: John Wiley & Sons Ltd., 2001, pp 519–540.
  - 13 Schork NJ, Fallin D, Thiel B *et al*: The future of genetic case-control studies; in Rao DC, Province MA (eds): *Genetic dissection of complex traits (Advances in genetics, Vol 42)*. US: Academic Press, 2000, pp 191–212.
  - 14 Allison DB: Transmission-disequilibrium tests for quantitative traits. *Am J Hum Genet* 1997; 60: 676–690.
  - 15 Long AD, Langley CH: The power of association studies to detect the contribution of candidate genetic loci to variation in complex traits. *Genome Res* 1999; 9: 720–731.
  - 16 Xiong MM, Krushkal J, Boerwinkle E: TDT statistics for mapping quantitative trait loci. *Ann Hum Genet* 1998; 62: 431–452.
  - 17 Haseman JK, Elston RC: The investigation of linkage between a quantitative trait and a marker locus. *Behav Genet* 1972; 2: 3–19.
  - 18 Rabinowitz D: A transmission disequilibrium test for quantitative trait loci. *Hum Hered* 1997; 47: 342–350.
  - 19 Sham PC, Cherny SS, Purcell S, Hewitt JK: Power of linkage *versus* association analysis of quantitative traits, by use of variance-components models, for sibship data. *Am J Hum Genet* 2000; 6: 1616–1630.
  - 20 Sokal RR, Rohlf FJ: *Biometry*. New York US: WH Freeman and Company, 1995.
  - 21 Lewontin RC: On measures of gametic disequilibrium. *Genetics* 1988; 120: 849–852.
  - 22 Jayakar SD: On the detection and estimation of linkage between a locus influencing a quantitative character and a marker locus. *Biometrics* 1970; 26: 451–464.
  - 23 Hill AP: Quantitative linkage: a statistical procedure for its detection and estimation. *Ann Hum Genet* 1975; 38: 439–449.
  - 24 Weir BS: *Genetic data analysis II*. Sunderland, US: Sinauer Associates, Inc. 1996.
  - 25 Searle SR: *Linear models*. New York: John Wiley & Sons, 1971.
  - 26 Falconer DS, Mackay TFC: *Introduction to quantitative genetics*. England: Longman Group Ltd, 1996.
  - 27 Zhao H: Family-based association studies. *Stat Methods Med Res* 2000; 9: 563–587.
  - 28 The World Health Report. *Part three: statistical annex*. WHO, 1999, www.who.int/whr/1999/en/report.htm.
  - 29 Underwood JCE: Genetic and environmental causes of disease; in Underwood JCE (ed): *General and systematic pathology*. London Churchill Livingstone, 1996, pp 31–60.
  - 30 Weiss KM, Terwilliger JD: How many diseases does it take to map a gene with SNPs? *Nat Genet* 2000; 26: 151–157.

- 31 Miller RD, Kwok PY: The birth and death of human single-nucleotide polymorphisms: new experimental evidence and implications for human history and medicine. *Hum Mol Genet* 2001; 20: 2195–2198.
- 32 Lynch M, Walsh B: *Genetics and analysis of quantitative traits*. Sunderland, US: Sinauer Associates, Inc., 1998.

## Appendix A

The NCP ( $\lambda$ ) of two-way ANOVA can be expressed as<sup>25</sup>

$$\lambda = \frac{(K'B)'[K'(X'X)^{-1}K]^{-1}(K'B)}{\sigma_e^2} \quad (A1)$$

Let  $\sigma_e^2$  be unity. Let  $B'$  be the vector  $[\mu, f_1, f_2, f_3, g_1, g_2, g_3]$  of parameters in the model, where  $\mu$  is the sample mean,  $f_i$  is the mean of the  $i$ th parental type, and  $g_j$  the mean of the  $j$ th marker genotype across all parental types. Let  $K$  be a matrix of parameter contrasts reflecting the  $H_0$  being tested; for example, if  $H_0: g_1 = g_2$  and  $g_2 = g_3$ , then

$$K' = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

The matrix  $X'X$  is

$$\begin{bmatrix} n_{..} & n_{1.} & n_{2.} & n_{3.} & n_{.1} & n_{.2} & n_{.3} \\ n_{1.} & n_{11} & 0 & 0 & n_{11} & n_{12} & n_{13} \\ n_{2.} & 0 & n_{2.} & 0 & n_{21} & n_{22} & n_{23} \\ n_{3.} & 0 & 0 & n_{3.} & n_{31} & n_{32} & n_{33} \\ n_{.1} & n_{11} & n_{21} & n_{31} & n_{.1} & 0 & 0 \\ n_{.2} & n_{12} & n_{22} & n_{32} & 0 & n_{.2} & 0 \\ n_{.3} & n_{13} & n_{23} & n_{33} & 0 & 0 & n_{.3} \end{bmatrix}$$

where  $n_{ij}$  is the number of records in the  $i$ th family and  $j$ th marker genotype class.  $X'X$  is a matrix of order 7 and rank 5; hence, there are seven unknowns and only five df. An appropriate generalisation of  $X'X$  is obtained deleting the first row and column, hence setting  $\mu = 0$ , and the last row and column, hence setting  $g_3 = 0$ .<sup>25</sup> Let  $G$  be the reduced  $X'X$  matrix. This  $G$  matrix can be partitioned as follows:

$$G = \begin{bmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{bmatrix} = \begin{bmatrix} n_{1.} & 0 & 0 & n_{11} & n_{12} \\ 0 & n_{2.} & 0 & n_{21} & n_{22} \\ 0 & 0 & n_{3.} & n_{31} & n_{32} \\ n_{11} & n_{21} & n_{31} & n_{.1} & 0 \\ n_{12} & n_{22} & n_{32} & 0 & n_{.2} \end{bmatrix}$$

Then, if  $C = G^{-1}$ ,  $K^*$  is the matrix  $K'$  with the first and last columns deleted, and  $B^*$  is the vector  $B$  with the first and last elements deleted, then

$$\lambda = (K^*B^*)'C_{22}^{-1}K^*B^* \quad (A2)$$

where  $C_{22} = K^*CK^* = (G_{22} - G_{21}G_{11}^{-1}G_{12})^{-1}$  and  $(K^*B^*)' = [g_1 - g_2, g_2]$  When testing the QTL, Eq. (A2) gives

$$\lambda = \sum_j^3 n_j g_j^2 - \sum_i^3 \frac{\left(\sum_j^3 n_{ij} g_j\right)^2}{n_i} \quad (A3)$$

where the first part of (A3) corresponds to the sum of squares due to genotype, and the second part



**Table A1** Family mean ( $F_i$ ) and number ( $f_i$ )

$i$	$F_i$	$f_i$
1	$b_1$	$n_1$
2	$(b_2+b_3)/2$	$n_2+n_3$
3	$b_4$	$n_4$
4	$b_6/2+(b_5+b_7)/4$	$n_5+n_6+n_7$
5	$(b_8+b_9)/2$	$n_8+n_9$
6	$b_{10}$	$n_{10}$

of (A3) corresponds to the sum of squares due to parental type.

However, it is a linked marker, rather than the QTL, what usually is being tested. Thus, Eq. (A3) needs to accommodate this fact. Using Tables 2 and 3 in the Materials and Methods section, the new  $\lambda$  can be written as

$$\lambda = \sum_{i=1}^{10} b_i^2 n_i I_i - \sum_{j=1}^6 F_j^2 f_j I_j \quad (A4)$$

where  $b_i$  is the expected marker genotype effect among progeny in the  $i$ th trio class,  $n_i$  the number of trios in class  $i$ ,  $I_{i(j)}$  an indicator variable equal to 1 if the trio is informative and 0 otherwise. Table A1 shows  $F_j$ , the mean value of the  $j$ th parental type, and  $f_j$ , the number of these parental types.

It is also possible to use all trios, thus setting  $I_{i(j)} = 1$  for all  $i$  ( $j$ ), without increasing the type-I error rate. By doing so, power increases slightly, through augmenting the residual df, and ascertainment of informative families becomes unnecessary.

This method of obtaining  $\lambda$  can be applied to derive the NCP for nested ANOVA; however, the algebra becomes more tedious. Finally, the NCP  $\lambda_O$  can be derived through Eq. (3), although a simpler method was described in the Materials and Methods section.

### Appendix B

Let us consider two fixed effects,  $\alpha$  and  $\beta$ , where  $\alpha$  could represent the factor parental type, and  $\beta$  could represent the genotype of the progeny. Thus, the model can be written as  $y_{ij} = \mu + \alpha_i + \beta_j + e_{ij}$ , which corresponds to a two-way ANOVA model without interaction. We will now show that the original statistic  $F_{2,n-5}$  for  $TDT_{Q_s}$ <sup>14</sup> is equivalent to the  $F$ -ratio for testing the effects of  $\beta$  after having corrected for the effects due to  $\mu$  and  $\alpha$ , using the previous model.

For a constant  $k = 2/n - 5$ , we can see that

$$kF_{2,n-5} = \frac{SS_F - SS_R}{SS_T - SS_F} = \frac{(SS_\mu + SS_\alpha + SS_\beta - SS_\mu - SS_\alpha)/SS_T}{1 - (SS_\mu + SS_\alpha + SS_\beta)/SS_T} = \frac{R_{\beta|\mu,\alpha}^2}{R_e^2} \quad (B1)$$

where  $SS_{\mu\alpha}$  and  $SS_{\mu\alpha\beta}$  are the sum of squares explained by a model that fits  $\mu$  and  $\alpha$ , and by a model that fits  $\mu$ ,  $\alpha$ , and  $\beta$ , respectively;  $SS_T$  is the total sum of squares; and  $R_{\beta|\mu,\alpha}^2$  and  $R_e^2$  are the proportions of the total variance explained by  $\beta$ , after taking into account the effects of  $\mu$  and  $\alpha$ , and the proportion of unexplained variance, respectively. The null hypothesis of interest is whether factor  $\beta$  explains a significant amount of phenotypic variance over and above the amount explained by  $\mu$  and  $\alpha$  jointly. The  $F$ -ratio that appropriately reflects this null hypothesis is given in Eq. (B1).

### Appendix C

Let assume  $T$  is a random variable following a t-distribution, and let  $\sigma_T$  be the standard deviation of  $T$ . A first-order Taylor's approximation for  $\lambda$  is  $\lambda = E(T/\sigma_T) \approx E[T]/E[\sigma_T]$ .<sup>32</sup> In order to derive  $E[T]$  and  $E[\sigma_T]$ , we used the probabilities of the 10 different types of trios and the expected effects of marker genotypes in the progeny contained in Tables 2 and 3. Hence, conditional on  $p_M, p_Q, c$ , and  $D'$ ,  $E[T] = E \times [\sum_i^n (y_i - \bar{y})w_i]$ , and because all family trios are independent (ie unrelated)  $E[T] = NE[(y - \bar{y})w]$ , where  $y$ , the phenotype, and  $w$ , a weighting factor, are expectations for a single trio (Table 3). Thus, the expected value of the numerator of  $TDT_R$  is approximately  $E[T] = Np_M p_m [p_M^2 (b_2 - b_3) + p_M p_m (b_5 - b_7) + p_m^2 (b_8 - b_9)]$ . When analysing the QTL, and assuming no dominance, the previous equation simplifies to  $E[T] = Np_Q p_q a$ .

The expected variance of  $T$ ,  $E[\sigma_T^2]$ , is the same regardless of whether the locus being tested is the QTL or a marker. Equation (A1.23a) in Reference<sup>32</sup> is  $E[\sqrt{v}] \approx \sqrt{\mu_v} - \sigma_v^2 \mu_v^{-3/2} / 8$ , which reduces to  $E[\sqrt{\sigma_T^2}] \approx \sqrt{E[\sigma_T^2]}$  if the second term can be ignored. Hence,  $E[\sigma_T^2] = E[1/4 \sum_i^n (y_i - \bar{y})^2 H_i] = 1/4 E[(y - \bar{y})^2 H]$  and, as the expectation of a random variable  $X$  given another random variable  $Y$  is  $E[X] = E[E[X|Y]]$ ,  $E[(y - \bar{y})^2 H] = \sum_{H=0}^2 HP_H E[(y - \bar{y})^2] = p_M p_m (\sigma_e^2 + \sigma_{QTL}^2)$ . Finally, dividing  $E[T]$  by  $\sqrt{E[\sigma_T^2]}$  we obtain  $E[TDT_R] = \lambda_R \approx [p_M^2 (b_2 - b_3) + p_M p_m (b_5 - b_7) + p_m^2 (b_8 - b_9)] \sqrt{Np_M p_m / \sigma_e^2 + \sigma_{QTL}^2}$ .